



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





N 1027

N. N. 1027


C Syx



UNIVERSITEITSBIBLIOTHEEK GENT



90



HN 1027

C. 347 104





135 348 9844



HN 1027

*C. Linnaeus*  
*& Carolus Linnaeus*

CAROLI LINNÆI

EQUITIS URNE STILLE LARI,

ARCHIATRI REGII, MED. & BOTAN. PROFESS. UPSAL. ACAD.

PARIS. UPSAL. HOLMENS. PETROPOL. BEROL. IMPERIAL.

BERNENS. LONDIN. ANGLIC. EDENBURG. MONSPEL.

TOLOS. FLORENT. SOC.

# SPECIES PLANTARUM,

EXHIBENTES

PLANTAS RITE COGNITAS

AD

GENERA RELATAS,

CUM

DIFFERENTIIS SPECIFICIS, NOMINIBUS

TRIVIALIBUS, SYNONYMIS SELECTIS,

LOCIS NATALIBUS,

SECUNDUM

SYSTEMA SEXUALE

DIGESTAS

TOMUS I.

---

Editio Tertia.

---

VINDOBONAE.

Typis JOANNIS THOMÆ DE TRATTNER,

CÆS. REG. AULÆ. TYPOGRAPHI ET BIBLIOP.

M DCC LXIV.



Ignoto Genere proprio

nulla descriptio, *quamvis accurate tradita,*

*certam demonstrat,*

*sed plerumque fallit.*

CÆSALP. *præfat:*

## LECTORI ÆQUO.

**H**omo sui conscius observat **MUNDUM** Omnipotentis theatrum, undique adornatum summis omniscia Sapienciae miraculis, se vero in hunc tamquam hospitem introductum, ut hisce deliciis sese delectando magnificentiam Domini agnoscat. Indignus profecto is censendus hospes, qui, pecoris instar, tantum gulæ inservit, nec magnalia Possessoris intueri atque æstimare novit.

**HOSPITES** ut digni evadamus orbis, opera hæc Creatoris scrutari nobis necesse est, quæ ita cum nostris commodis combinavit summum Ens, ut eis, unde omnia nostra bona desumimus, carere nulla ratione queamus; & quo magis hæc intelligimus, eo etiam plura in usum generis nostri humani cedunt.

**COGNITIONEM** horum ut rite acquiramus, singula distincta *Idea* & distincto *Nomine* complecti oportet, quibus sepositis copia rerum nos obruat necesse est, & commercium omne, deficiente communi lingua, cessabit.

**SCIENTIA NATURÆ** inde inter Homines nata est, quæ comprehendit sub *Physica* &

## LECTORI.

*Chemia* Elementa; sub *Zoologia*, *Botanica*,  
*Lithologia* tria Naturæ regna.

REM HERBARIAM heic libare lubet, quæ  
quondam paucarum plantarum erat Scientia;  
hodie vero copia detectorum evasit omnium  
amplissima.

INVENTORUM recentioris ævi indefessis la-  
boribus notitiam debemus plurium Planta-  
rum, quam omnibus priscorum temporum  
eruditis, inter quos imprimis nominandi *Clu-  
sius*, *Columna*, *Baubini*, *Hermannus*, *Rbee-  
de*, *Rumphius*, *Sloane*, *Sherardus*, *Rajus*,  
*Plukenetius*, *Tournefortius*, *Flumierus*, *Vail-  
lantius*, *Dillenius*, *Burmamnus*, *Gronovius*,  
*Rayenus*, *Gmelinus*, *Millerus*, *Brownius*,  
*Jacquinius*.

NOTITIA plantarum constabat quondam No-  
minibus arbitrariis, *memoriæ* tabulæ infixis,  
quam vacillantem *Icones* sublevabant.

SYSTEMATICORUM sapientia indagavit  
ordinem, qui memoriæ succurreret, & scien-  
tiæ solidis fundamentis inædificaret, quem  
debemus Principibus Viris *Gesnero*, *Cæsalpi-  
no*,

## LECTORI

no, *Baubino*, *Morifono*, *Hermanno*, *Tournefortio*, *Vaillantio*, *Dillenio*, aliisque.

**ARIADNEUM** hoc filum Systematicorum terminatur *Generibus*, quod tentavi ad *Species*. usque extendere, confectis legitimis harum differentiis, ut etjam in hisce sua sibi constaret certitudo, quum *omnis vera cognitio cognitione specifica innitatur*; hac enim deficiente vacillat & historia, ut in plerisque *Peregrinatorum* diariis.

**SPECIES** plantarum ut mihi familiares redderem, peragravi *Alpes Lapponicas*; totam *Sveciam*; partem *Norvegiæ*, *Daniæ*, *Germaniæ*, *Belgii*, *Angliæ*, *Galliæ*; perquisivi deinde **HORTOS** Botanicos: *Parisiinum*, *Oxoniensem*, *Chelsejanum*, *Hartecampensem*, *Leidensem*, *Ultrajectinum*; *Amstelædamensem*, *Upsaliensem* aliosque: evolvi etjam **HERBARIA** *Burseri*, *Hermanni*, *Cliffortii*, *Burmanni*, *Oldenlandi*, *Gronovii*, *Royeni*, *Sloanei*, *Sberardi*, *Bobarti*, *Milleri*, *Tournefortii*, *Vaillantii*, *Jussiei*, *Suriani*, *Backii*, *Brownii*, &c. Meo suasu **DISCIPULI** quon-



## L E C T O R I.

dam dilectissimi: *P. Kalmius* Canadam, *F. Hasselquist* Ægyptum, *P. Osbeck* Chinam, *Toren* Suratte, *Solander* Angliam, *Alstræmer* Europam australem; *Martin* Spitsbergam, *Pontin* Malabariam, *Kæbler* Italiam, *Forskæhl* Orientem, *P. Læfving* Hispaniam, *Montin* Lapponiam adierunt, plantasque lectas mecum communicarunt. E variis terris demum misere Amici BOTANICI femina plantasque siccas non paucas, uti *B. Jussieu*, *Royenus*, *J. Gesnerus*, *Wachendorffus*, *Sibtorpius*, *Montius*, *Gleditschius*, *Krafccheninikoffius*, *Minuartius*, *Velezius*, *Kleinboff*, *Ortega*, *Ellis*, *Seguier*, *Allionius*, *Zinnius*, *Gouanus*, *Gerardus*, nec non *O. Munchhausen* L. B; *S. C. Bielke*, L. B; *J. Rathgeb* Equ. *Demidoff* Nob; *Collinson*, *Braad*, alique. *Clifford* omnes, quas in duplo habuit; *Lagerström* indiæ orientalis plurimas; *Gronovius* plerasque Virginicas; *Gmelinus* Sibiricas facile omnes; *Burmannus* præter plurimas proprias & intuendas misit capenses Oldenlandi; at *Sauvagesius* integrum suum thesaurum  
plan-

## LECTORI

plantarum dedit, exemplo raro & inaudito, unde factum, ut copiam plantarum obtinuerim non mediocrem.

**DIFFERENTIALS** Specificas antehac Plantis non paucis imposui in Flora *Lapponica*, *Suecica*, *Zeylanica*; in Horto *Cliffortiano*, *Upsaliensi*. Iisdem principiis institere præstantissimi Botanici *Gronovius*, *Royenus*, *Wachendorff*, *Gorter*, *Jaquin*, *Colden*, *Hill*, *Hudson*, *Gouan*, *Dalibard*, *Kramer*, *Leyser*, *Meesse*, *Le Monier*, *Gættard*, *Sauvages*, inque non paucis *Hallerus*, *Gmelinus*, *Allionius*, *Seignier*, *Scopoli*, *Gerard*, *Burmantus*, alique, quorum opera plurimæ Species evasere ratæ & evidentes.

**NOMINA** hæc, passim dispersa, in gratiam studiosorum consultum duxi colligere, acquiritas dein plantas addere, inque unum Systema reducere; at vero pluribus *Speciebus* observatis, præstantioribus notis detectis, aptioribus *Terminis* cufis, debui interdum differentias emendare, antea quamvis optimas.

## LECTOR

Characteres essentialis pro NOMINE SPECIFICCO constituere non levidense opus est; requirit enim *Specierum* plurium accuratam collationem, harum attentissimam *partium* indagationem, *differentium* selectionem, *Terminorum* denique artis propriam applicationem, ut evadant compendiosissimi tutissimique.

NON VISAS plantas heic omisi, toties elusus ab auctoribus, ne dubia certissimis miscerem; si vero aliquando contigerit non sufficienter inspexisse plantam, vel specimen imperfectum obtinuisse, signo † hoc notavi, ut alii eandem accuratius examinent.

NUMERUM plantarum totius orbis longe pareciorem esse, quam vulgo creditur, satis certo calculo intellexi, utpote qui vix 10000 attingat.

TRIVIALIA nomina in margine apposui, ut missis ambagibus, uno quamlibet Herbarum nomine complecti queamus; hæc vero absque selectu posui, quippe quem alius dies poscit. Caveant autem quam sanctissime omnes sani

Bo-

## L E C T O R I.

Botanici, unquam proponere nomen triviale sine sufficiente differentia specifica, ne ruat in pristinam barbariem scientia. Mutatio horum, licet in melius, plus nocet quam proderit.

**SYNONYMA** paucissima in **EUROPÆIS** plantis adhibui, contentus *C. Baubino & Iconographo* præstantiore; in **EXOTICIS** vero plura, quum difficiliora minusque trita sint.

**DESCRIPTIONES** \* tantum in obscuris adhibere necessum fuit, easque sine ambagibus, ut obtinerem compendium tironibus gratum.

**LOCA** natalia more meo adjeci, inque notissimis *Fruticantes* †, *Perennes* ‡, *Biennes* §, *Annua* ○ notavi.

**GENERA** nonnulla *nova*, nonnulla *immutata* adhibui; quæ in nova editione Generum plantarum propediem sistere animus est.

**ADVERSARIORUM** tela nunquam retorſi, acerrima convitia, infimulationes, cavillationes, buccinationes (præstantiorum longe Virorum, omni ævo, laboris præmia), tranquillo animo sustinui, nec suis auctoribus hæc

## LECTORI.

invideo, si inde ipsis apud vulgus gloria major. Fero hæc, quæ in me ne capillum quidem mutant, & cur non ferrem iniquos, qui a veris Botanicis & quidem consummatissimis, quibus assurgere debent illi, summis cum latus sim encomiis? Parem deinde adversariis referre gratiam, nec anni ingravescentes, nec munus, quo fungor, nec mei mores patiuntur. Quod mihi superest brevioris ævi, tranquillius utilioribus observationibus impendam. Res naturales enim proprio privilegio munitæ persistunt, ut quemadmodum errores in his commissi a nullo defendi, ita nec veritates observationibus innixæ a toto eruditorum orbe conculcari possint; iudices itaque provoco feros Nepotes.

*Pascitur in vivis livor, post fata quiescit,  
Tum suus ex merito quemque tuetur bonos.*

Dabam 1762. d. 1. Septembris Upsaliæ.

A U.



# A U C T O R E S.

## \* R E F O R M A T O R E S.

PROPRIA	Philosophia	Botanica.	Holmiae.	1751.	off.
	Critica	Botanica.	Lugdb.	1737.	off.
	Classes	Plantarum.	Lugdb.	1738.	off.
	Syſtema	Naturæ. II.	Holmiae.	1759.	off.
	Genera	Plantarum.	Holmiae.	1754.	off.
	Species	Plantarum.	Holmiae.	1753.	off.
	Hortus	Cliffortianus.	Amſtel.	1737.	fol.
	Hortus	Upſalienſis.	Holmiae.	1748.	off.
	Flora	Lapponica.	Amſtel.	1737.	off.
	Flora	Svecia.	Holmiae.	1755.	off.
	Flora	Zeylanica.	Holmiae.	1747.	off.
	Materia	Medica.	Holmiae.	1749.	off.
	Amœnitates	Academic. I.	Holmiae.	1749.	off.
	Amœnitates	Academ. II.	Holmiae.	1751.	off.
	Amœnitates	Academ. III.	Holmiae.	1756.	off.
	Amœnitates	Academ. IV.	Holmiae.	1759.	off.
	Amœnitates	Academ. V.	Holmiae.	1760.	off.
	Amœnitates	Academ. VI.	Holmiae.	inedit.	off.
	Iter	Oelandicum.	Holmiae.	1745.	off.
	Iter	Gotlandic.	Holmiae.	cum prioræ.	
	Iter	Westrogoth.	Holmiae.	1747.	off.
	Iter	Scanicum.	Holmiae.	1751.	off.
* * *					
KALM. P.	Iter	American.	Holmiae.	1753.	off. vol. 3.
OSBECK. P.	Iter	Ind. orient.	Holmiae.	1757.	off.
TORÉN. O.	Iter	Surattænſe.	Holmiae.	cum prioræ.	
HASSELQUIST. F.	Iter	Palæſtinum.	Holmiae.	1757.	off.
LOEFLING. P.	Iter	Hiſpanicum.	Holmiae.	1758.	off.
* * *					
JACQUIN. N.	Enum.	plantar.	Lugdbat.	1760.	off.
BROWN.	Hiſt. nat.	Jamaicæ.	London.	1756.	fol.
COLDEN. C.	Plant.	Coldingh.	Amſt. Upſ.	1743.	quart.
GRÖNOVIUS J. F.	Flora	Virginica.	Leydæ.	1739.	off. vol. 2.
	Flora	Orientalis.	Leydæ.	1755.	off.
GORTER. D.	Flora	Geldrica.	Harder.	1757.	off. vol. 2.
	Flora	Ingrica.	Petrop.	1761.	off.
MEESE. D.	Flora	Frificæ.	Francæq.	1760.	off.
HILL J.	Flora	Britannica.	London.	1760.	off.
HUDSON.	Flora	Anglica.	London.	1762.	off.
DALIBARD.	Flora	Parifiſis.	Parifiis.	1749.	quod.
GÖLLAN.	Flora	Monſpel.	Monſp.	1762.	off.
KRAMER.	Flora	Auſtriaca.	Wien.	1756.	off.
JACQUIN. N.	Flora	Vindelic.	Wien.	1762.	off. LEY-

LEYSER. F. W.	Flora Halensis.	Hala.	1761. off.
LEE. J.	Introduct. in Botan.	London.	1760. off.
MILLER. PH.	Dictionar. Hortulan.	London.	1760. fol.
ROYEN. A.	Flora Leydensis.	Lugdb.	1740. off.
WACHENDORFF.	Hort. Ultraject.	Tayest.	1747. off.
SAUVAGES. F.	Flora Monspel.	Haga.	1751. off.
GERARD. L. C.	Flora Galloprov.	Parisiis.	off.
GUETTARD.	Observat. plantar.	Parisiis.	1747. off. vol. 2.
MONIER. L. G.	Cat. plant. Alvern.	Parisiis.	1745. quart.
HALLER A.	Stirpes Helveticæ.	Gatting.	1742. fol.
	Hortus Goetting.	Gatting.	1749. off.
	Opusc. Botanica.	Gatting.	1753. off.
ZINN. J. G.	Hortus Goetting.	Gatting.	1757. off.
GMELIN. J. G.	Flora Sibirica.	Petropol.	1750. quart. vol. 2.
SEGUIER. J. F.	Plantæ Veronenf.	Verone.	1745. off. vol. 3.
ALLIONIUS. C.	Stirpes Pedem.	Taurin.	1755. quart.
	Stirpes Nicæenses.	Paris.	1757. off.
	Hortus Taurinens.	-----	off.
SCOPOLI. J.	Flora Carniolica.	Wien.	1760. off.
BÜTTNER. D. G.	Planta Cunonis.	Amstel.	1750. off.
MARTYN J.	Decades rarior.	London.	1628. fol.
EHRET. D.	Icones pl. select.	London.	fol.
TREW.	Tabul. Ehretii.	Noriberg.	1750. fol.
ORDER.	Flora Danica.	Hassnia.	1761. fol.
MILLER. Ph.	Icones ad Dict.	London.	fol.
PLUMIER. C.	Icones per Burman.	Amstelod.	1755. fol.

## \* \* RESTAURATOIRES.

MORISON. R.	Histor. plantarum.	Oxon.	1680. fol. vol. 2.
	Hortus Blesensis.	London.	1669. off.
RAJUS. J.	Histor. plantar.	London.	1686. fol. vol. 2.
	Synopf. Anglica.	London.	1724. off.
HERMANNUS. P.	Hortus Lugdbat.	Lugdb.	1687. off.
	Paradis. Batavus.	Lugdb.	1688. quart.
TOURNEFORT. J.	Institut. Herbariæ.	Parisiis.	1719. quart.
	Corollarium	Parisiis.	conjunctim.
PLUMIER. C.	Genera pl. Amer.	Parisiis.	1703. quart.
	Species pl. Amer.	Parisiis.	cum priore.
	Plantæ Americ.	Parisiis.	1693. folio.
	Filices Americ.	Parisiis.	1705. folio.
VAILLANT.	Flores compos.	Act. Par.	1718. 8 <sup>vo</sup> .
	Botan. Pariliens.	Paris.	1747. fol.
FEWILL. L.	Journ. de Peru.	Paris.	1714. quart. v. 2.

BOER.

BOERHAAVE	Hortus Lugd. bat.	Lugdb.	1720. quart. vol. 2.
BURMANNUS J.	Theſaurus Zeylan.	Amſtel.	1737. quart.
	Decades African.	Amſtel.	1738. quart.
N. L.	Gerania diff. acad.	Lugdb.	1759. quart.
DILLENIUS	Hortus Eltham.	London.	1732. fol.
	Hiſtor. Muſcor.	Oxonii	1741. quart.
AMMANUS J.	Stirpes rariores.	Petrop.	1739. quart.
VOLCKAMER J. G.	Flora Norib.	Norib.	1760. quart.
SCHREUCHZER. J.	Hiſtor. Agroſt.	Tiguri.	1719. quart.
	Itinera alpina.	Tiguri.	1702. quart.
MONTI J.	Prodr. Gramin.	Bonon.	1719. quart.
MICHEL J.	Gener. nova.	Florent.	1729. quart.
TILLI	Hortus. Piſanus.	Florent.	1723. folio.
BUXBAUM	Centur. rarior.	Petropol.	1728. oct.


### \* \* \* FUNDATORES.

FLUKENET	Almag. Bôtanic.	London.	1696. quart.
	Mant. Almag.	London.	1700. quart.
	Amalt. Botanic.	London.	1705. quart.
	Phytographia.	London.	1690. quart.
PETIVER	Gazoph. naturæ.	London.	1702. fol.
SLOANE	Cat. pl. Jamaic.	London.	1696. oct.
	Hiſt. pl. Jamaic.	London.	1707. fol. vol. 2.
CATESBY	Hiſt. nat. Carol.	London.	1731. fol. vol. 2.
RHEEDE	Hortus Malabar.	Amſtel.	1678. fol. vol. 12.
RUMPHIUS.	Herbar. Amboin.	Amſtel.	1740. fol. vol. 6.
COMNELINI	Hortus Amſtel.	Amſtel.	1697. fol. vol. 2.
	Plantæ rariores.	Lugdb.	1706. quart.
	Prælect. Botanic.	Lugdb.	1703. quart.
BREYNIUS J.	Centur. plantar.	Gedan.	1678. fol.
	Prodr. per filium.	Gedan.	1739. quart.
KEMPFERUS	Amœn. exoticæ.	Lemgo.	1712. quart.
DODART	Memor. de plant.	Parif.	1676. fol.
MAGNOL P.	Botanic. Monſp.	Monſp.	1686. oct.
BARRELIER J.	Icons.	Parifis.	1714. fol.
BOCCO P.	Muſeum	Venet.	1697. quart.
	Plantæ Siculæ.	Oxon.	1674. quart.
ALPINUS	Exoticæ. plantæ.	Venet.	1627. quart.
LCESLIUS	Flora Pruſſic.	Regio.	1703. quart.
REYNALME	Specim. plantar.	Parif.	1611. quart.
CORRETTI J.	Canad. plantæ.	Parif.	1635. quart.
MARCGRAVE J.	Hiſt. nat. Braſil.	Lugab.	1648. fol.
PISO G.	Hiſt. nat. Braſil.	Amſtel.	1658. fol.

BAY.



<b>BAUHINUS C.</b>	Pinax Theatr.	<i>Bafil.</i>	1671. quart.
	Prodr. Theatr.	<i>Bafil.</i>	1671. quart.
	Theatr. Botanic.	<i>Bafil.</i>	1658. fol.
<b>BAUHINUS J.</b>	Histor. Plantar.	<i>Elrod.</i>	1650. fol. vol. 3.
<b>COLUMNA.</b>	Ecphrasis	<i>Roma.</i>	1616. quart. vol. 3.
	Phytobasanos.	<i>Flor.</i>	1744. quart.
<b>CLUSIUS C.</b>	Histor. rarior.	<i>Antw.</i>	1601. fol.
	Exotica.	<i>Antw.</i>	1605. fol.
<b>DODONÆUS</b>	Pempt. stirpium.	<i>Antw.</i>	1616. fol.
<b>LOBELIUS</b>	Jcones.	<i>Antw.</i>	1591. quart.
<b>FUCHSIUS</b>	Historia stirpium.	<i>Basel.</i>	1542. fol.
<b>GESNERUS</b>	Historia plantar.	<i>Norib.</i>	1759. fol.
<b>CAMERARIUS J.</b>	Epitome Matth.	<i>Francf.</i>	1586. quart.
	Hortus. Medic.	<i>Francf.</i>	1588. quart.
<b>TABERNÆM.</b>	Histor. plantar.	<i>Bafil.</i>	1664. folio.
<b>DALECHAMP.</b>	Histor. plantar.	<i>Lugd.</i>	1587. folio.



## Classis I.

## MONANDRIA

## MONOGYNIA.

## CANNA.

1. CANNA foliis ovatis utrinque acuminatis nervosis. *Indica.*  
*Roy. lugdb. 11. Fl. zeyl. 1. Hort. upf. 1.*  
*Canna spatulis bifloris. Hort. cliff. 1.*  
*Cannacorus. Rumph. amb. 5. p. 177. t. 71. f. 2.*  
*Arundo indica latifolia. Bauh. pin. 19.*  
*Katu Bala. Rheed. mal. 11. p. 85. t. 43.*  
*Habitat inter tropicos Asiæ, Africæ, Americæ. 24*
2. CANNA foliis lanceolatis petiolatis nervosis. *angustifolia.*  
*Canna foliis lanceolatis petiolatis. Hort. cliff. 1.*  
*Arundo indica florida angustifolia. Moris. hist. 3. p. 250.*  
*f. 8. t. 14. f. 6.*  
*Habitat inter tropicos Americæ: umbrosis spongiosis. 24*  
*Species hæc vix distinguenda est a præcedente, quamvis humilior & angustior evadat, qui itaque pro varietatibus utramque assumunt, vix errant.*
3. CANNA foliis petiolatis lanceolatis enervibus. *Roy. glauca.*  
*lugdb. 11.*  
*Canna foliis enervibus. Virid. cliff. 104.*  
*Cannoides. Hort. cliff. 488.*  
*Cannacorus glaucophyllus, ampliore flore, iridis palustris facie. Dill. elih. 69. t. 59. f. 69.*  
*Habitat in Carolina? 24*

## AMOMUM.

1. AMOMUM scapo nudo, spica ovata. *Hort. cliff. 3. Zingiber.*  
*Hort. upf. 1. Flor. zeyl. 3. Mat. med. 1. Roy. lugdb. 12.*  
*Zingiber majus. Rumph. amb. 5. p. 156. t. 66. f. 1.*  
*Zingiber. Bauh. pin. 35. Brown. jam. 119.*  
*Intsch. Rheed. mal. 11. p. 21. t. 12.*  
*Habitat in Indiis inter tropicos. 24*
2. AMOMUM scapo nudo, spica oblonga obtusa. *Hort. Zerumbet.*  
*cliff. 3. Hort. upf. 1. Fl. zeyl. 2. Roy. lugdb. 12. Ehrh.*  
*pic. t. 14. f. 1. Mill. ic. 17. t. 25. Brown. jam. 113.*  
*Linæi Spec. plant. pars. I. A Zin-*



## MONANDRIA MONOGYNIA.

*Zingiber latifolium* sylvestre. *Herm. lugdb.* 636. t. 637.

*Lampujum*. *Rumph. amb.* 5. p. 148. t. 64. f. 1.

*Katou-Infschi-kua*. *Rheed. mal.* 11. p. 27. t. 13.

*Habitat in India.* 24

Cardamom.

3. *AMOMUM* scapo simplicissimo brevissimo bracteis alternis laxis. *Fl. zeyl.* 4. *Mat. med.* 2.

*Cardamomum minus*. *Rumph. amb.* 5. p. 152. t. 65. f. 1.

*Elettari* 2. *Rheed. mal.* 11. p. 9. t. 6.

*Habitat in Indiæ umbrosissimis, cænosis montium radicibus.* 24

Gran. parad.

4. *AMOMUM* scapo ramoso brevissimo.

*Amomum caule ramoso.* *Mat. med.* 3.

*Grana paradisi officinarum.* *Bauh. pin.* 413.

*Elettari.* *Rheed. mal.* 11. t. 6.

*Habitat in Madagascar, Guinea.* 24

## COSTUS.

arabicus.

1. *COSTUS.* *Hort. cliff.* 2. \* *Hort. upf.* 2. *Fl. zeyl.* 5.

*Mat. med.* 4. *Roy. lugdb.* 11. *Hill. plant.* 149. t. 3.

*Costus arabicus.* *Bauh. pin.* 36. *Mer. surin.* 36. t. 36.

*Alpinia floribus spicatis: bracteis ovalibus.* *Jacq. amer.* 11.

*Zingiber sylvestre majus, fructu in pediculo singulari.*

*Sloan. jam.* 61. *hist.* 1. p. 165. t. 105. f. 2.

*Amomum minus, scapo vestito, floribus spicatis.* *Brown. jam.* 113.

*Paco Caatinga.* *Marcgr. bras.* 48.

*Tiana kua.* *Rheed. mal.* 11. p. 15. t. 8?

*Habitat in India uraque.* 24

## ALPINIA.

racemosa.

1. *ALPINIA.* *Roy. lugdb.* 12.

*Alpina racemosa alba, cannatori foliis.* *Plum. ic.* 10. t. 20.

*Zingiber sylvestre minus, fructu e caullum summitate.*

*Sloan. jam.* 61. *hist.* 1. p. 165. t. 105. f. 1.

*Paco seroca.* *Marcgr. bras.* 48?

*Habitat in America calidiore.* 24

## MARANTA.

arundinacea

1. *MARANTA* culmo ramoso.

*Maranta.* *Hort. cliff.* 2. \* *Roy. lugdb.* 11.

Ma-

## MONANDRIA MONOGYNIA.

3

*Maranta arundinacea*, cannaçori folio. *Plum.* gen. 16.

*Mart.* cent. 39. t. 39.

*Habitat in America calidiore.* 24.

1. *MARANTA* culmo simplici.

Galanga.

Galanga. *Rumph.* amb. 5. p. 143. t. 63.

*Habitat in India aquosa.* 24.

Galanga hæc officinarum est.

## CURCUMA.

1. *CURCUMA* foliis lanceolato-ovatis: nervis laterali. rotunda.  
bus rarissimis.

*Curcuma* foliis ovatis utrinque acuminatis: nervis lateralibus paucissimis. *Roy.* lugdb. 12. *Fl.* zeyl. 6.

*Curcuma.* *Rumph.* amb. 5. p. 162. t. 67.

*Manja-kua.* *Rheed.* mal. 11. p. 19. t. 10.\*

*Habitat in India.* 24.

2. *CURCUMA* foliis lanceolatis: nervis lateralibus numerosissimis.

*Curcuma* foliis lanceolatis utrinque acuminatis: nervis lateralibus numerosissimis. *Roy.* lugdb. 12. *Fl.* zeyl. 7.

*Mat.* med. 5.

*Curcuma* radice longa. *Herm.* lugdb. 208. t. 209.

*Manjella-kua.* *Rheed.* mal. 2. p. 21. t. 11.

*Habitat in India.* 24.

## KAEMPFERIA.

1. *KAEMPFERIA* foliis ovatis sessilibus. *Fl.* zeyl. 8. Galanga.

*Kaempferia.* *Hort.* cliff. 2. t. 3. *Roy.* lugdb. 12.

*Sonchorus.* *Rumph.* amb. 5. p. 173. t. 69. f. 2.

*Wanbom.* *Kaempf.* amoen. 901. t. 902.

*Katsjula-kelengu.* *Rheed.* mal. 11. p. 81. t. 41.

*Habitat in India.* 24.

2. *KAEMPFERIA* foliis lanceolatis petiolatis. *Fl.* zeyl. 9. rotunda

*Zedoaria* rotunda. *Bauh.* pin. 36.

*Malan-kua.* *Rheed.* mal. 11. p. 17. t. 9.

*Habitat in India.* 24.

## THALIA.

1. *THALIA.* *Roy.* lugdb. 11.

*Curtia* arundinacea, amplis cannaçori foliis. *Plum.*

geniculata.

pa. 26. ic. 198. f. 1.

A 2

Ma-

- Maranta petiollis ganglinosis. Brown. jam. 112.*  
*Caná indica, radice alba alexipharmaca. Sloan. jam. 22.*  
*hist. 1. p. 253. t. 149. f. 2. Raj. suppl. 573.*  
*Canna indica angustifolia, pediculis longis ad imum*  
*folium nodo singulari geniculata. Pluk. alm. 79.*  
*Radix quædam in malacca, quæ adversus vulnera sa-*  
*gittis toxico illitis præsentaneum remedium. Bauh.*  
*pin. 301.*  
*Aguti guepo-obi. Marcgr. bras. 53. Raj. hist. 1202.*  
*Habitat in America meridionali. 24*  
*Folia alterna, ovato-oblonga: nervis transversis parallelis.*  
*Petioli teretes, geniculo instructi, inferne membranacei*  
*vaginales.*

## BOERHAVIA.

- erecta,* 1. BOERHAVIA caule erecto glabro.  
*Boerhavia, solani folio, erecta glabra. Amm. herb.*  
*241.*  
*Habitat in Vera Cruce. 24*
- diffusa,* 2. BOERHAVIA caule diffuso.  
*Boerhavia diffusa. Fl. zeyl. 10. Hort. upf. 2.*  
*Boerhavia foliis ovatis. Hort. cliff. 17. Roy. lugdb. 234.*  
*Valerianella curassavica, semine aspero viscoso. Herm.*  
*par. 237. t. 237.*  
*Talu-dama. Rheed. mal. 7. p. 105. t. 56.*  
*Habitat in India. 24*
- scandens.* 3. BOERHAVIA caule scandente, floribus diandris.  
*Boerhavia sarmentosa. Brown. jam. 123.*  
*Solanum bacciferum americanum, fructu corymbofo.*  
*Pluk. alm. 349. t. 226 f. 7.*  
*Valerianella, alines folio, scandens. Sloan. jam. 91.*  
*hist. 1. p. 210. Raj. suppl. 244.*  
*Valerianella curassavica, semine aspero viscoso. Pluk.*  
*alm. 381. t. 113. f. 7.*  
*Antanizophyllum scandens, alines majoris folio. Vaill.*  
*act. 1722. p. 258.*  
*Habitat in Jamaica ad urbem jago de la vega. 24*
- diandra,* 4. BOERHAVIA glabra diffusa, floribus diandris.  
*Habitat in India. Ott. Munchausen L. B. 24*  
*Statura B. diffusæ, sed Caulis & fructus glabri. Pe-*  
*dunculi folio breviores, sæpius triflori. Flores mi-*  
*nores*

# MONANDRIA MONOGYNIA.

5

nores, albi, pedicellati. Fructus sulcati, pentagoni, truncati. Simillimam accepi ex Asia caule pubescente.

## 5. BOERHAVIA caule repente.

repens.

Boerhavia nubica minor. Vaill. sex. 53.

## β Boerhavia nubica minima. Vaill. sex. 55.

Habitat in Nubia inter Mocho & Tangos. 24

# SALICORNIA.

## 1. SALICORNIA patula, articulis apice compressis emarginato-bifidis.

herbacea.

Salicornia herbacea. Fl. suec. 1. Mat. med. 8.

Salicornia annua. Sauv. monsp. 7.

Salicornia. Hort. cliff. 490. Roy. lugdb. 205. Dod. pempt. 82.

## β Salicornia caulium ramorumque articulis emarginatis bifidis. Gron. virg. 129.

virginica.

Habitat in Europæ littoribus maritimis. ☉

Virginica β. ad Salinas Saxonie frequentissima, vix ac ne vix distincta est species; articuli in salsis enim magis emarginati evadunt.

## 2. SALICORNIA caule erecto fruticoso.

fruticosa.

Salicornia sempervirens. Sauv. monsp. 7.

Kali geniculatum majus. Bauh. pin. 289.

Habitat in Europæ maritimis. ☿

Sempervirens & erecta longeque major a priori videtur diversa, præterquam quod etiam spicæ differant; hæc in Suecia non visa, ubi prior notissima; differentia levis erit, si ulla.

## 3. SALICORNIA articulis obtusis basi incrassatis, spicis ovatis.

arabica.

Kali geniculatum minus. Bauh. pin. 289. Moris. hist. 2. p. 610. f. 5. t. 33. f. 7. Raj. hist. 211.

Habitat in Arabia.

## 4. SALICORNIA articulis cylindricis, spicis filiformibus.

caspica.

Kali arborescens, tamarisci facie. Buxb. cent. 1. t. 11. f. 1.

Habitat in squalidis maris Caspii & Mediæ. ☿

## HIP PURIS.

vulgaris,

1. HIP PURIS. *Fl. lapp.* 1. *Fl. svec.* 2. 2. *Hort. cliff.* 3.  
*Roy. lugdh.* 205. *Dalib. parif.* 1.  
 Equisetum palustre, brevioribus foliis, polyspermum.  
*Bauh. pin.* 15.  
 Polygonum femina. *Cam. epit.* 689.  
*Habitat in Europæ fontibus.* 2.

## DIGYNIA.

## CORISPERMUM.

hyssopifo-  
lium,

1. CORISPERMUM floribus lateralibus. *Hort. upf.* 2.  
 Corispermum floribus alternis. *Hort. cliff.* 3. *It. goth.*  
 316. *Roy. lugdh.* 205. *Sauv. monsp.* 52.  
 Corispermum hyssopifolium. *Juss. act.* 1712. p. 244.  
*Habitat ad Wolgam Tatarizæ, Gillau Borussia, Mon-*  
*speli locis arenosis.* ☉

squamrosum,

2. CORISPERMUM spicis squarrosis. *Hort. upf.* 3. \*  
 Rhagrostis foliis arundinaceis. *Buxb. cent.* 3. p. 30. t. 55.  
*Habitat ad Wolgam Tatarizæ, inque desertis Cossac-*  
*orum.* ☉

## CALLITRICHE.

verna,

1. CALLITRICHE foliis superioribus ovalibus, floribus androgynis. *Fl. svec.* 3.  
 Corispermum foliis oppositis. *Fl. lapp.* 2. *Gron. virg.* 3.  
 Stellaria aquatica. *Bauh. pin.* 141.  
 2. Lenticula palustris bifolia, fructu tetragono. *Bauh. pin.*  
 362.

*Callitriche plinii. Column. ecclæ,* 1. p. 315. t. 316.  
*Habitat in Europæ fossis aqueis, Vere florens.*

autumnalis,

3. CALLITRICHE foliis omnibus linearibus apice bifidis, floribus hermaphroditis. *Fl. svec.* 4.  
 Lenticula palustris angustifolia, folio apice dissecto. *Læs. pruss.* 140. t. 38.  
*Habitat in Europæ fossis aqueis, Autumno florens.*

## BLITUM.

capitatum.

1. BLITUM capitellis spicatis terminalibus. *Hort. upf.* 3.  
 Blitum foliis triangularibus dentatis, calycibus baccatis, capitulis summis nudis, *Virid. cliff.* 1. *Roy. lugdh.* 206.  
 Bl.

## MONANDRIA DIGYNIA.

7

*Blitum* foliis triangularibus dentatis, *Hort. cliff.* 28.

*Atriplex* baccifera. *Raj. hist.* 197.

*Atriplex* sylvestris lappulas habens. *Bauh. pin.* 119.

*Habitat in* Europa: *præsertim in comit. Tyrolensi.* ☉

2. *BLITUM* capitellis sparsis lateralibus. *Hort. ups.* 3.

virgatum.

*Blitum* foliis triangularibus dentatis, calycibus baccatis, capitulis summis foliosis. *Hort. cliff.* 495. *Roy. lugdb.* 206.

*Atriplex* sylvestris, mori fructu. *Bauh. pin.* 519.

*Habitat in* Tataria, Hispania, G. Narbonensi. ☉

## CINNA.

### I. CINNA.

*Habitat in* Canada, unde semina per *D. Kalm.* 2.

arundinacea

*Gramen* magnitudine *Avenæ.* Culmi plurimi, glabri:

Folia latiuscula, glabra, - margine scabra. Panicula

oblonga, attenuata, compressiuscula, secunda. Calycis

Gluma bivalvis, uniflora, carinata carina scabra: val-

vula majore mucrone aristiformi. Corollæ Gluma bi-

valvis: valvula exteriori intra minorem calycis, calyce

toto paullo longior, infra apicem mucronata. Anthera

purpurea. Styli hirsuti.

## Classis II.

DIANDRIA  
MONOGYNIA.

## NYCTANTHES.

- arbor tristis, 1. NYCTANTHES caule tetragono, foliis ovatis acuminatis, pericarpis membranaceis compressis. *Fl. zeyl.* 11.  
Arbor tristis myrto similis. *Bauh. pin.* 469.  
Manja-pumeran. *Rheed. mal.* 1. p. 35. t. 21.  
*Habitat in India.* §
- Sambac, 2. NYCTANTHES foliis inferioribus cordatis obtusis, superioribus ovatis acutis. *Hort. upf.* 4.\*  
Nyctanthes caule volubili, foliis subovatis acutis. *Hort. cliff.* 5. *Flor. zeyl.* 12. *Roy. lugdb.* 398.  
Syringa arabica, foliis mali aurantii. *Bauh. pin.* 398.  
Jasminum limonii folio conjugato. *Burm. zeyl.* 128. t. 58. f. 2. *Angl. hortul.* t. 27.  
Jasminum f. Sambac arabum, folio acuminato. *Till. pif.* 87. t. 31.  
Jasminum arabicum. *Clus. cur.* 3. t. 3.  
Flos manoræ. *Rumph. amb.* 5. p. 52. t. 30.  
*Habitat in India.* §
- undulata, 3. NYCTANTHES foliis ovatis acuminatis undulatis, ramis teretibus.  
Jasminum indicum, flore polypetalo exalbido, fructu minori. *Raj. hist.* 1601.  
Tsieregam mulla. *Rheed. mal.* 6. p. 97. t. 55.  
*Habitat in Malabar.* §
- hirsuta, 4. NYCTANTHES petiolis pedunculisque villosis.  
Rava-pon. *Rheed. mal.* 4. p. 99. t. 48.  
*Habitat in India.* §
- angustifolia, 5. NYCTANTHES foliis obtusis lanceolatis ovatisque.  
Jasminum indicum angustifolium, fructu gemino. *Tournef. inst.* 598.

Katu-pittiegam mulla. *Rheed. mal.* 6. p. 93. t. 53. *Raj. hist.* 1602.

*Habitat in Malabariz arenosis. ‡*

## J A S M I N U M.

1. JASMINUM foliis oppositis pinnatis. *Hort. cliff.* 5. officinale.  
*Hort. ups.* 5. *Mat. med.* 9. *Roy. lugdb.* 397.

*Jasminum vulgatus*, flore albo. *Bauh. pin.* 397.

*Habitat in India. ‡*

2. JASMINUM foliis oppositis pinnatis: foliolis ext- *grandiflo-*  
mis confluentibus. *rum.*

*Jasminum humillus*, magno flore. *Bauh. pin.* 397. *Raj.*

*hist.* 1600.

*Gelseminum catalanicum. Cam. epit.* 37.

*Pittiegam-mulla Rheed. mal.* 6. p. 91.

*Habitat in Malabariz. ‡*

*Simillimum J. officinali*, sed *Caulis rectus, nec scandens.*

*Foliola cauli propiora magis ovata, nec minora; extrema tria confluentia ex petiolo dilatato, ut distincta cadere nequeant. Corolla majores, subtus rubra.*

3. JASMINUM foliis oppositis ternatis. *Hort. cliff.* 5. azoricum.

*Fl. zeyl.* 13. *Roy. lugdb.* 397.

*Jasminum azoricum trifoliatum*, flore albo, odoratissimo. *Comm. hort.* 1. p. 159. t. 82.

*Jasminum sylvestre triphyllum*, floribus rubellis umbellatis. *Burm. zeyl.* 127. t. 58. f. 1.

*Jasminum album trifoliatum*, flore magno. *Pluk. alm.* 195. t. 303. f. 1.

*Habitat in India. ‡*

4. JASMINUM foliis alternis ternatis simplicibusque, fruticans, ramis angulatis. *Hort. cliff.* 5. *Hort. ups.* 5. *Roy.*

*lugdb.* 397. *Sauv. monsp.* 174.

*Jasminum luteum*, vulgo dictum bacciferum. *Bauh. pin.* 398.

*Trifolium fruticans. Ded. pempt.* 571.

*Habitat in Europa australi, & toto Oriente. ‡*

5. JASMINUM foliis alternis ternatis pinnatisque, *humile.*  
mis angulatis. *Hort. ups.* 5.

*Jasminum foliis alternis ternatis acuminatis. Hort. cliff.*

6. *Roy. lugdb.* 398.



*Jasminum humile luteum. Bauh. pin. 397.*

*Jasminum luteum. Bess. cyst. est. 140. f. 2.*

*Habitat - - - - - b*

odoratissi-  
mum.

6. **JASMINUM** foliis alternis ternatis pinnatisque, ramis teretibus. *Hort. upf. 5.*

*Jasminum foliis alternis ternatis obtusis. Hort. cliff. 5. Roy. lugdb. 397.*

*Jasminum indicum flavum odoratissimum. Ferr. cult. 93.*

*Jasminum flavum odoratum. Barr. ic. 62.*

*Habitat in India? b*

## LIGUSTRUM

vulgare.

1. **LIGUSTRUM.** *Hort. cliff. 6. Fl. svec. 4. 5. Roy. lugdb. 398. Dalib. parif. 2.*

*Ligustrum germanicum. Bauh. pin. 475.*

*Phillyrea. Dod. pempt. 775.*

*Habitat in Europæ collibus glareosis. b*

## PHILLYREA.

media.

1. **PHILLYREA** foliis ovato-lanceolatis subintegerrimis. *Hort. upf. 5.*

*Phillyrea foliis ovato-lanceolatis vix crenatis. Hort. cliff. 4. Roy. lugdb. 398.*

*Phillyrea ligustri folio. Bauh. pin. 476.*

*Phillyrea 3. Clus. hist. 1. p. 52.*

*Habitat in Europæ australioris collibus. b*

angustifolia

2. **PHILLYREA** foliis lineari-lanceolatis integerrimis. *Hort. cliff. 4. Roy. lugdb. 398.*

*Phillyrea angustifolia prima. Bauh. pin. 476.*

*Phillyrea angustifolia secunda. Bauh. pin. 476.*

*Phillyrea 4. 5. Clus. hist. 1. p. 52.*

*Habitat in Italia, Hispania. b*

latifolia.

3. **PHILLYREA** foliis cordato-ovatis serratis. *Hort. cliff. 4. Hort. upf. 6. Roy. lugdb. 398.*

*Phillyrea latifolia spinosa. Bauh. pin. 476.*

*Phillyrea folio leviter serrato. Bauh. pin. 476.*

*Phillyrea 1, 2. Clus. hist. 1. p. 51. 52.*

*Habitat in Europa australi. b*

## OLEA.

## OLEA.

## 1. OLEA foliis lanceolatis.

*Olea foliis lanceolatis, ramis teretiufculis. Hort. cliff.*

europæa.

4. *Mat. med.* 10. *Roy. lugdb.* 398.

*Olea sativa. Bauh. pin.* 472.

*Olea. Dod. pempt.* 821.

β. *Olea foliis lanceolatis, ramis tetragonis. Hort. cliff.* 4.  
*Roy. lugd.* 399.

*Olea sylvestris, folio duro subtus incano. Bauh. pin.* 472.

*Habitat in Europa australi.* ♀

capensis.

α. OLEA foliis ovatis. *Hort. cliff.* 4. *Roy. lugdb.* 399.

*Ligustrum capense sempervirens, folio crasso subrotundo. Dill. elih.* 193. t. 170. f. 194.

*Habitat ad Cap. b. Spei.* ♀

## CHIONANTHUS.

## 1. CHIONANTHUS pedunculis trifidis trifloris.

virginica.

*Chionanthus. Hort. cliff.* 17. *Gron. virg.* 10. *Roy. lugdb.* 17.

*Amelanchir virginiana, laurocerasi folio. Pet. sic.* 241. *Catesb. carol.* 1. p. 68. t. 68.

*Habitat in America septentrionali.* ♀

zeylonica.

2. CHIONANTHUS pedunculis paniculatis multifloris.

*Fl zeyl.* 14.

*Arbor zeylanica, cotini foliis subtus lanugine villosis.*

*Burm. zeyl.* 31.

*Arbuscula zeylanica, cotini foliis subtus lanugine villosis, floribus albis cuculli modo laciniatis. Pluk.*

*alm.* 44. t. 241. f. 4.

*Habitat in Zeylona.* ♀

## SYRINGA.

1. SYRINGA foliis ovato cordatis. *Hort. cliff.* 6. *Hort.*

vulgaris.

*ups.* 6. *Roy. lugdb.* 397. *Dalib. parif.* 2.

*Syringa cærulea. Bauh. pin.* 398.

*Habitat versus Persiam.* ♀

2. SYRINGA foliis lanceolatis.

persica.

α. *Syringa foliis lanceolatis integris. Hort. cliff.* 6. *Hort.*

*ups.* 6. *Roy. lugdb.* 397. *Mill. dict.* 1. 164. f. 1.

*Syringa babylonica, indivisis densioribus foliis. Pluk.*

*alm.* 359. t. 227. f. 8.

Agem

- laciniata.* Agem Lilac perfarum. *Corn. canad.* 190.  
 β. *Syringa* foliis lanceolatis integris dissectisque. *Herr. cliff.* 6. *Hort. upf.* 6. *Roy. lugdb.* 397.  
*Ligustrum* foliis laciniatis. *Bauh. pin.* 476. *Mill. dict.* 1. 164. f. 2.  
*Habitat in Persia ?* b

## ERANTHEMUM.

- capense.* 1. ERANTHEMUM. *Fl. zeyl.* 15. \* *Amæn. acad.* 385.\*  
*Ephemerum*, *lychnidis flore*, africanum. *Herm. parad.* 153. *Amm. herb.* 232.  
*Centaurium minus*, foliis oblongis acutis, flore patulo puniceo majori. *Raj. suppl.* 527.  
*Habitat in Æthiopia.*  
*Habitus Chironiæ*, unicum specimen minus perfectum vidi, ideoque accuratius examinanda.

## CIRCÆA.

- lutetiana.* 1. CIRCÆA caule erecto, racemis pluribus. *Fl. svec.* 6.  
*Circæa.* *Hort. cliff.* 7. *Roy. lugdb.* 303. *Dalib. parif.* 3.  
*It. scan.* 213. \*  
*Solanifolia* *Circæa dicta major.* *Bauh. pin.* 168. *It. wgoth.* 213.  
*Circæa lutetiana.* *Lob. ic.* 266.  
*canadensis.* β. *Circæa canadensis latifolia*, flore albo. *Tournef. inst.* 301.  
*Habitat in Europæ & Americæ borealis nemoribus.* 2.  
*Hæc sesquipedalis Caule erecto, Racemis terminali & lateralibus; Perianthium consistentia calycina, s. corolla longe crassius rudiisque, nec corollæ concolor.*  
*alpina.* 2. CIRCÆA caule adscendente, racemo unico. *Fl. svec.* 7.  
*Circæa calyce colorato.* *Fl. lapp.* 3.  
*Solanifolia* *Circæa alpina.* *Bauh. pin.* 168.  
*Circæa minima.* *Column. ecphr.* 2. p. 79. t. 80.  
*Habitat ad radices montium in frigidis Europæ.* 2.  
*Planta vix digiti longitudine, magis procumbens, calyce colorato, uti corolla; differt quidem a priori, attenu nimis affinis.*

## VERONICA.

\* *Spicata.*

- sibirica.* 1. VERONICA spicis terminalibus, foliis septenis verticillatis, caule subhirsuto.

Ve-

*Veronica spicata altissima, foliis verticillatim dispositis.*

*Amm. ruth. 20. t. 4.*

*Habitat in Dauria. 24*

*Caulis quadripedalis, hirtus. Folia 7. f. 6. verticillata, V. virginicæ duplo latiora. Pedunculi terminales solitarii; laterales foliolis duobus oppositis ovalibus. Corollæ cæuleæ tubo oblongo; limbo acuto parvo. Stamina 8 Pistillum corollæ duplo longiora.*

2. *VERONICA* spicis terminalibus, foliis quaternis quinisve. virginica.

*Veronica foliis quaternis quinisve. Hort. cliff. 7. \* Gron. virg. 4. Roy. lugdb. 301. Cold. novch. 3.*

*Veronica virginiana procerior, foliis ternis, quaternis & etiam quinis caulem amplexantibus, spicis florum candidissimis. Pluk. alm. 383. t. 70. f. 2.*

*Habitat in Virginia. 24*

3. *VERONICA* spicis terminalibus, foliis ternis æqualiter ferratis. spuria.

*Veronica spicata angustifolia. Bauh. pin. 246.*

*Veronica mas surrecta elatior. Barr. rar. 155. t. 89r.*

*Habitat in Europa australiore, Sibiria, Thuringia. 24*

4. *VERONICA* spicis terminalibus, foliis ternis inæqualiter ferratis. maritima.

*Veronica floribus spicatis, foliis ternis Fl. suec. 1. p. 6.*

*Veronica foliis sæpius ternis. Virid. cliff. 2. Hort. cliff. 7. Roy. lugdb. 301.*

*Veronica caule erecto, spicis pluribus, foliis lanceolatis, ferratis. Fl. lapp. 4.*

*Lyfimachia spicata cærulea. Bauh. pin. 246.*

*Pseudo-Lyfimachium cæruleum. Dod. pempt. 86.*

*Habitat in maritimis Europæ macris apricis. 24*

5. *VERONICA* spicis terminalibus, foliis oppositis lanceolatis ferratis acuminatis. longifolia.

*2. n. 9. \**

*Veronica spicata latifolia. Bauh. pin. 246.*

*Veronica 1. erectior latifolia. Clus. hist. 1. p. 346.*

6. *Veronica spicata longifolia altera. Amm. ruth. 26.*

7. *Veronica spicata, urticæ folio. Amm. ruth. 26.*

*Habitat in Tataria, Austria, Suecia. 24*

- incana.* 6. **VERONICA** spicis terminalibus, foliis oppositis crenatis obtusis, caule erecto tomentoso. *Hort. upf.* 7. Veronica caule fruticoso, foliis oblongis: inferioribus crenatis petiolatis; superioribus sessilibus integerrimis. *Roy. lugdb.* 530.  
Veronica spicata lanuginosa & incana, floribus cæruleis. *Ann. rut.* 30.  
*Habitat in Ucraina, Samara.* 24.
- spicata.* 7. **VERONICA** spica terminali, foliis oppositis crenatis obtusis, caule ascendente simplicissimo. *Fl. suec.* 2. n. 10.  
Veronica floribus spicatis, foliis oppositis, caule erecto. *Fl. suec.* 1. n. 7. *Dalib. parif.* 3.  
Veronica foliis oppositis, caule spica terminato. *Hort. cliff.* 8. *Roy. lugdb.* 302.  
Veronica spicata minor. *Bauh. pin.* 247. *Vaill. parif.* t. 33. f. 4.  
*Habitat in Europæ borealis campis.* 24.
- hybrida.* 8. **VERONICA** spicis terminalibus, foliis oppositis obtuse serratis scabris, caule erecto. *Fl. suec.* 2. n. 11.\*  
Veronica spicata cambrobritannica, bugulæ subhirsuto folio. *Raj. angl.* 3. p. 278. t. 11. f. 1.  
*Habitat in Europa rarius.* 24.  
*Folia veronicæ sequentis, Caulis præcedentis, sed spicæ majores pallidiores.* 24.
- officinalis.* 9. **VERONICA** spicis lateralibus pedunculatis, foliis oppositis, caule procumbente. *Mat. med.* 11. *Fl. suec.* 2. n. 12.  
Veronica floribus spicatis, foliis oppositis, caule procumbente. *Fl. suec.* 1. n. 8. *Dalib. parif.* 3.  
Veronica caule repente, scapis spicatis, foliis oppositis ovatis strigosis. *Fl. lapp.* 5. *Hort. cliff.* 8. *Roy. lugdb.* 302.  
Veronica mas supina & vulgatissima. *Bauh. pin.* 246.  
Veronica. *Cam. epit.* 461.  
β. Veronica mas repens pyrenaica, folio longiori glabro. *Pluk. phyt.* 223. f. 1.  
*Habitat in Europæ sylvestribus sterilibus.* 24.

\* \* *Corymbo-racemosa.*

- aphylla.* 10. **VERONICA** corymbo terminali, scapo nudo.

Vero-

*Veronica foliis ovatis radicalibus, caule nudo. Roy. lugdb. 302.*

*Veronica alpina pumila, caule aphylo. Bocq. mus. 2. p. 17. t. 1. & 9.*

*Veronica parva saxatilis, cauliculis nudis. Pluk. alm. 384. t. 114. f. 3. Segu. ver. 241. t. 3. f. 2.*

*Chamædrys alpina minima hirsuta. Bauh. pin. 243.*

*Habitat in alpidis Europæ australioris. 21*

11. **VERONICA** corymbo terminali, caule adscenden- bellidoides.  
te diphylo.

*Veronica caule non ramoso, floribus congestis terminato, foliis ovatis subasperis. Hall. helv. 531. t. 12. f. 1.*

*Veronica alpina, bellidis folio. Bauh. pin. 247. prod. 116. Burf. XIV. 27.*

*Habitat in Helvetia, Pyrenæis. 21*

12. **VERONICA** corymbo terminali, foliis lanceolatis fruticulosa.  
obtusiusculis, caulibus fruticulosiss.

*Veronica foliis ovatis crenatis, fructu ovali, floribus in summo caule purpurascens. Hall. helv. 532. t. 9. f. 1.*

*Veronica alpina frutescens. Bauh. pin. 247.*

*Habitat in Alpibus Austriæ, Helvetiæ, Pyrenæis. 21*

*Planta spithamea: furculis perennantibus, caulibus erectis, simplicibus. Folia superiora majora, lanceolata, obtusa, glabra. Differt a V. alpina caule erectiusculo, magis lignoso, foliis angustioribus, Calycibus glabris.*

13. **VERONICA** corymbo terminali, foliis oppositis, alpine.  
calycibus hispidis. Fl. suec. 2. n. 15. \* Oed. dan. t. 16.

*Veronica floribus corymbosis terminalibus, calycibus hispidis. Fl. suec. 1. n. 13.*

*Veronica caule floribus terminato, foliis ovatis crenatis. Fl. lapp. 7. t. 9. f. 4. \**

*Veronica caule inferne procumbente fruticoso, foliis ovatis oppositis, racemo terminali. Roy. lugdb. 302.*

*Veronica saxatilis Bauh. hist. 3. p. 284. Raj. extr. 261. \**

*Habitat in alpidis Europæ. 21*

14. **VERONICA** racemo terminali subspicato, foliis serpyllifolia  
ovatis glabris crenatis. Fl. suec. 2. n. 16.

Vero.

- Veronica floribus solitariis subcorymbosis, foliis ovatis glabris crenatis. Fl. fvec. 1. n. 14. Dalib. parif. 4.*  
*Veronica foliis inferioribus oppositis ovatis, superioribus alternis lanceolatis, floribus solitariis. Hort. cliff. 9. Gron. virg. 4. Roy. lugdb. 302. Cold. noveb. 1.*  
*Veronica floribus sparsis, foliis ovatis crenatis glabris. Fl. lapp. 6.*  
*Veronica pratensis serpyllifolia. Bauh. pin. 247.*  
*Veronica femina. Bauh. hist. 3. p. 285.*  
*Veronica; nummulariæ folio, pyrenaica. Tournes. inf. 145. Pluk. alm. 384. t. 233. f. 4.*  
*Habitat in Europa & America septentrionali ad vias, agros. 24.*  
 Beccabunga. 15. *VERONICA* racemis lateralibus, foliis ovatis planis, caule repente. *Fl. fvec. 11. 14. Mat. med. 12. Dalib. parif. 7.*  
*Veronica foliis oppositis lævibus crenatis, floribus racemosis lateralibus. Roy. lugdb. 302.*  
*Veronica foliis oppositis lævibus crenatis, floribus laxè spicatis ex alis. Hort. cliff. 8. Gron. virg. 4.*  
*Anagallis aquatica major (minorque), folio subrotundo. Bauh. pin. 252.*  
*Anagallis aquatica. Dod. pempt. 823.*  
*Habitat in Europa ad scaturigines vix congelandas.*  
 Anagall. ▽ 16. *VERONICA* racemis lateralibus, foliis lanceolatis ferratis, caule erecto. *Fl. fvec. 10. 13. Dalib. parif. 7.*  
*Veronica foliis oppositis lævibus crenatis, floribus laxè spicatis ex alis. Gron. virg. 4.*  
*Anagallis aquatica major, folio oblongo. Bauh. pin. 252.*  
*Berula major. Tabern. hist. 1094.*  
*Habitat in Europa ad fossas & in Oriente. ☉*  
 scutellata. 17. *VERONICA* racemis lateralibus pedicellis pendulis, foliis linearibus integerrimis. *Fl. fvec. 9. 17. Dalib. parif. 7.*  
*Veronica foliis lineari-lanceolatis integris, racemis laxè floriferis. Fl. lapp. 9. Hort. cliff. 10. Roy. lugdb. 302.*  
*Anagallis aquatica angustifolia scutellata. Bauh. pin. 252.*  
*Anagallis aquatica angustifolia. Bauh. hist. 3. p. 780.*  
*Habitat in Europæ inundatis. 24.*  
 Teucrium. 18. *VERONICA* racemis lateralibus longissimis, foliis ovatis rugosis dentatis obtusiusculis, caule erecto.

*Veronica foliis imis cordatis superioribus angustioribus.*

*Hall. herb. 529.*

*Chamædrys spuria major altera f. frutescens. Bauh. pin. 248.*

*Chamædrys spuria major angustifolia. Bauh. pin. 248.*

*Teucris IV. tertia species. Clus. hist. 1. p. 349.*

*Habitat in Germania. 24.*

*Racemi densi, stricti, subspicati. Folia obtusius serrata.*

19. *VERONICA* racemis lateralibus, foliis lineari-lanceolatis pinnato-dentatis. austriaca.

*Chamædrys austriaca, foliis tenuissime laciniatis. Bauh. pin. 248. prodr. 117.*

*Chamædrys spuria tenuissime laciniata. Bauh. hist. 3. p. 287.*

β. *Veronica cappadocica, foliis laciniatis. Tournef. cor. Habitat in Austria. 24.*

*Folia sere Coronopi glabra. Præcedenti nimis affinis.*

20. *VERONICA* racemis lateralibus, foliis multipartitis laciniatis. multifida.

*Veronica montana, folio vario. Buxb. cent. 1. p. 24. t. 38.*

*Habitat in Armeniæ, Iberiæ graminosis. 24.*

21. *VERONICA* racemis lateralibus paucifloris, calycibus hirsutis, foliis ovatis rugosis crenatis petiolatis, caule debili. *Amæn. acad. 4. p. 263.* montana.

*Chamædrys spuria affinis rotundifolia scutellata. Bauh. pin. 249.*

*Allison dioscoridis montanum. Column. ecphr. 1. p. 286. t. 288.*

*Habitat in Italiæ, Germaniæ umbrosis. 24.*

22. *VERONICA* racemis lateralibus, foliis ovato-oblongis serratis, caulibus prostratis. prostrata.

*Chamædrys incana spuria minor angustifolia. Bauh. pin. 249. prodr. 117. Bauh. hist. 1. p. 287.*

*Habitat in Germaniæ, Italiæ collibus. 24.*

*Caules duriusculi, canescentes, subtomentosi. Folia obtusa, subpetiolata, exceptis superioribus, e quorum alis Racemi longi. Flores late violacei, conferti, calycibus inæqualibus.*

23. *VERONICA* racemis lateralibus, foliis ovatis sessilibus rugosis dentatis, caule debili. *Pl. succ. 13, 18.* Chamædrys.

*Dalib. paris. 4.*

*Lineæ. Spec. plant. pars I.*

B

Ve-

U. S. GENT



*Veronica foliis oppositis plicatis dentatis, scapis ex alis inferioribus laxè spicatis. Hort. cliff. 8. Roy. lugdb. 303.*

*Veronica foliis cordatis sessilibus oppositis, racemis laxè floriferis. Fl. lapp. 8.*

*Chamædrys spuria minor rotundifolia. Rarb. pin. 249.*

*Hierobotane mas. Dalech. hist. 1337.*

*Habitat in Europæ pratis. 21*

latifolia.

24. *VERONICA* racemis lateralibus, foliis cordatis rugosis dentatis, caule stricto.

*Veronica foliis oppositis plicatis dentatis, scapis ex alis superioribus laxè spicatis. Hort. cliff. 9. Roy. lugdb. 303.*

*Veronica pratensis omnium maxima. Buxb. cent. 1. p. 23. t. 34.*

*Chamædrys spuria major latifolia. Bauh. pin. 248.*

*Habitat in Helvetia, Bithynia, Austria. 21*

paniculata,

25. *VERONICA* racemis lateralibus longissimis, foliis lanceolatis ternis ferratis, caule ascendente.

*Veronica angustifolia, floribus paniculatis. Amm. ruth. 32.*

*Habitat in Tartaria. 21*

*Statura V. latifoliae, sed Folia angustissime lanceolata, plerumque ternæ. Racemi longissimi, saturate cerulei; ad tria superiora genicula terni.*

\*\*\* *Pedunculis unifloris.*

agrestis.

26. *VERONICA* floribus solitariis, foliis cordatis incisiss pedunculo brevioribus. *Fl. suec. 17. 20. Dalib. parif. 6.*

*Veronica foliis cordatis crenatis, floribus solitariis pedunculatis. Roy. lugdb. 303.*

*Veronica foliis alternis cordatis crenatis, floribus solitariis. Hort. cliff. 9.*

*Alfine chamædryfolia, flosculis pediculis oblongis insidentibus. Bauh. pin. 250.*

*Alfine foliis triflaginis. Tabern. hist. 1089.*

*Habitat in Europæ cultis arvis. ☉*

arvensis.

27. *VERONICA* floribus solitariis, foliis cordatis incisiss pedunculo longioribus. *Fl. suec. 16, 19. Dalib. parif. 5.*

*Veronica foliis oppositis cordatis crenatis, floribus solitariis sessilibus. Hort. cliff. 9. Gron. virg. 4. Roy. lugdb. 303.*

Al-

*Aline veronicæ foliis, flosculis cauliculis adhærentibus. Bauh. pin. 250.*

*Elatine polychides. Dalech. hist. 1239.*

*Habitat in Europæ arvis cultis. ☉*

28. VERONICA floribus solitariis, foliis cordatis planis quinquelobis. *Fl. suec. 18, 21. Dalib. parif. 6.* hederifolia

*Veronica foliis alternis cordatis quinquelobis, floribus solitariis. Hort. cliff. 9. Roy. lugdb. 303.*

*Aline hederulæ folio. Bauh. pin. 250. Tabern. hist. 1088.*

- β. Veronica chia, cymbalaræ folio, verna, flore albo: umbilico virescente. *Tournef. cor. 7. Buxb. cent. 1. p. 25. t. 39. f. 1.*

*Habitat in Europæ ruderatis. ☉*

*Semina umbilicata sunt huic propria.*

29. VERONICA floribus solitariis, foliis digitato-partitis pedunculo brevioribus. *Fl. suec. 19. 22. Dalib. parif. 6.* triphyllos.

*Veronica foliis alternis inferioribus quinquepartitis; superioribus tripartitis, floribus solitariis. Hort. cliff. 9. Roy. lugdb. 302.*

*Aline triphyllos cærulea. Bauh. pin. 250*

*Aline parva erecta, folio alines hederaceo. Lob. ic.*

464

*Habitat in Europæ agris. ☉*

30. VERONICA floribus solitariis, foliis digitato-partitis pedunculo longioribus. *Fl. suec. 20. 23.* verna

*Veronica humilis erecta montana, flore parvo cæruleo. Dill. app. 38.*

*Habitat in Sueciæ, Germaniæ, Hispaniæ aridis apricis. ☉*

31. VERONICA floribus solitariis subseffilibus, foliis oblongis subdentatis, caule erecto. romana.

*Habitat in Europæ australioris agris. ☉*

32. VERONICA floribus solitariis pedunculatis, foliis ovatis glabris crenatis, caule erecto subpiloso. acinifolia.

*Veronica minima, clinopodii minoris folio glabro, romana. Bocc. mus. 2. p. 29. t. 102. Raj. suppl. 423.*

*Vaill. parif. 201. t. 23. f. 3.*

*Veronica erecta, acini folio glabro, floribus cæruleis. Dill. app. 39.*

*Habitat in Europa australi.*

- peregrina. 33. *VERONICA* floribus solitariis sessilibus, foliis lanceolato-linearibus glabris obtusis integerrimis, caule erecto. *Fl. suec.* 15, 24. \*  
*Veronica terrestris annua, folio polygoni, flore albo, Moris. hist. 2. p. 322. f. 3. t. 34. f. 19.*  
*Habitat in Europæ hortis, arvisque. ☉*
- parilandica. 34. *VERONICA* floribus solitariis sessilibus, foliis linearibus, caulibus diffusis.  
*Veronica caulibus procumbentibus, foliis linearibus, floribus sessilibus lateralibus. Gron. virg. 4.*  
*Habitat in Virginia.*

## PÆDEROTA.

- bonæ spei. 1. *PÆDEROTA* foliis pinnatifidis. *Aman. acad. 6.*  
*af. 1. \**  
*Veronica africana, floribus ad genicula pedicellis biuncialibus. Herm. afr. 783. Pluk. phyt. 320. f. 5.*  
*Anagallis foliis sinuatis. Roy. lugdb. 416.*  
*Habuat ad Cap. bon. spei.*
- Bonarota. 2. *PÆDEROTA* foliis serratis.  
*Veronica Bonarota corymbo terminali, foliis oppositis serrato-dentatis, calycibus linearibus. Sp. pl. 1. p. 11.*  
*Veronica petræa sempervirens. Pon. bald. 336.*  
*Chamædrys alpina saxatilis. Bauh. pin. 248.*  
*Bonarota. Mich. gen. 19. t. 15. f. 12.*  
*Habitat in Alpibus Italicis, Austriacis. 24*  
*Corollæ tubus longior, quam in priore, ut in Veronicis spicatis, sed Calyx 5partitus.*

## JUSTICIA.

- Adhatoda. 1. *JUSTICIA* arborea, foliis lanceolato-ovatis, bracteis ovatis persistentibus, corollarum galea concava. *Fl. zeyl. 16. \* Hort. upf. 7.*  
*Justicia foliis ovato-lanceolatis, spicis foliosis, florum galea concava. Hort. cliff. 9. Roy. lugdb. 291.*  
*Adhatoda zeylanensium. Herm. lugdb. 642. t. 643.*  
*Pluk. alm. 9. t. 173. f. 3.*  
*Ecbolium Riv. mon. 129.*  
*Habitat in Zeylona. 5*
- Ecbolium. 2. *JUSTICIA* arborea, foliis lanceolato-ovatis, bracteis ovatis deciduis mucronatis, corollarum galea reflexa. *Fl. zeyl. 17. \**

Ad-

*Adhatoda* spica longissima, flore reflexo. *Burm. Zeyl.*

7. t. 4. f. 1.

*Carim-curini. Rheed. mal. 2. p. 31. t. 20. Pluk. alm.*

126. t. 171. f. 4.

*Habitat in Malabaria, Zeylona. b*

*Simillima J. infundibuliformi, sed huic Spica terminalis sessilis.*

3. *JUSTICIA* fruticosa, foliis lanceolato-ovatis quaternis, bracteis lanceolatis ciliatis. *infundibuliformis.*

*Manja-Kurini. Rheed. mal. 9. p. 121. t. 62.*

*Habitat in India. b*

*Frutex ramis teretibus. Folia quaterna, lanceolato-ovata, integerrima, glabra, longius petiolata. Pedunculi ad singula caulis genicula solitarii, longitudine folii, leves: Spica oblonga, dislice imbricata: bracteis lanceolatis ciliatis: inter quas Bracteolæ 4, lineares & Calyx apice pilosus. Corolla speciosa, alba; tubo filiformi, stamina includente; Limbo magnitudine Narcissi poetici, lobis 5, expansis: infimo majore.*

4. *JUSTICIA* fruticosa, foliis lanceolato-ovatis hirsutis sessilibus, spicis recurvatis. *scorpioides.*

*Justicia foliis oblongo-ovatis hirsutis sessilibus, floribus spicatis alaribus, caule fruticoso. Mill. dict. n. 1.*

*Habitat in Vera Cruce. b*

5. *JUSTICIA* fruticosa, foliis lanceolato-ovatis pictis, corollis fauce inflatis. *picta.*

*Folium bracteatum. Rumph. amb. 4. p. 73. t. 30.*

*Tinde-maram. Rheed. mal. 6. t. 60.*

*Habitat in Asia. b*

*Calyx minimus, quinquedentatus. Corolla ringens faucibus inflata: Lab. superius incurvum, emarginatum; L. inferius patens, trifidum, æquale. Stamina 2 sub labio superiore.*

6. *JUSTICIA* fruticosa, foliis lanceolato-ovatis, bracteis ovatis acuminatis venoso-reticulatis coloratis. *Betonica.*

*Fl. Zeyl. 18. \**

*Betonica frutescens. Bont. jav. 146.*

*Bem-curini. Rheed. mal. 2. p. 126. t. 21.*

*Habitat in India. b*

7. *JUSTICIA* fruticosa, foliis lanceolatis integerrimis, hyssopifolia.

pedunculis trifloris ancipitibus, bracteis calyce brevioribus *Mill. dict. t. 13.*

*Justicia* foliis lineari-lanceolatis, floribus sæpius solitariis. *Hort. cliff. 10. Roy. lugdb. 291.*

*Ecbolii indicii* & *Adhatodæ* cucullatis floribus æmula, hyssopifolia planta ex insulis fortunatis. *Pluk. alm. 132. t. 280. f. 1.*

*Habitat in insulis Fortunatis. ♀*

procumbens 8. *JUSTICIA* procumbens, foliis lanceolatis integerrimis, spicis terminalibus lateralibusque alternis, bracteis setaceis. *Fl. zeyl. 19. \**

*Euphrasia*, albescentes angustiore folio, rubiæ modo spicata, golcondeæ. *Pluk. alm. 142. t. 56. f. 3. ♂ t. 393. f. 4.*

*Habitat in Zeylona. 24.*

pectinata, 9. *JUSTICIA* diffusa, spicis axillaribus sessilibus tomentosis secundis dorso imbricatis: bracteis semilanceolatis. *Amæn. acad. 4. p. 299. \**

*Habitat in Asia.*

repens, 10. *JUSTICIA* repens, foliis ovatis subcrenatis, spicis terminalibus, bracteis lanceolatis. *Fl. zeyl. 20. \**

*Adhatoda* spicata, flosculos ex foliolis membranaceis. *Burm. zeyl. 7. t. 3. f. 2.*

*Habitat in Zeylona. 24.*

*Bracteæ* margine utroque late membranaceæ.

chinensis, 11. *JUSTICIA* herbacea, foliis ovatis, floribus lateralibus, pedunculis trifloris, bracteis ovalibus.

*Habitat in China.*

*Caules angulati, pedales, inferne ramosi, folia opposita, petiolata, ovata, acuminata, obtusiuscula, vix crenata. Ex alis pedunculi duo s. tres, singuli breves, apice triflori, involucri diphylo, setaceo, nudo, bracteis ovatis, obtusis, longitudine floris.*

echioides, 12. *JUSTICIA* foliis lanceolato-linearibus obtusis sessilibus, racemis ascendenti-secundis, bracteis setaceis. *Fl. zeyl. 21. \**

*Euphrasiæ* affinis indica echioides. *Herm. lugdb. 668. t. 669.*

*Peetumba*, *Rheed. mal. 9. p. 87. t. 46.*

*Habitat in India.*

13. **JUSTICIA** foliis ovatis integerrimis, bracteis cu- sexangularis  
neiformibus, ramis sexangularibus

*Justicia* annua, hexangulari caule, foliis circæ conjugatis, flore miniato. *Amm. herb.* 274. *Hort. cliff.* 10.

*Euphrasia*, alfinæ majori folio, flore galeato pallide luteo, jamaicensis. *Pluk. alm.* 142. t. 279. f. 6.

*Habitat in Vera cruce, Jamaica.* ☉

14. **JUSTICIA** foliis ovatis integerrimis, bracteis su- assurgens.  
bulatis, ramis hexagonis. *Am. acad.* 5. p. 391. \*

*Justicia* herbacea assurgens, ad alas alternas nodos & summitates florida. *Brown. jam.* 118. t. 2. f. 1. \*

*Habitat in Jamaica.*

15. **JUSTICIA** foliis lanceolato-ovatis integerrimis, pe- natata.  
dunculis dichotomis.

*Puicollis.* *Rheed. mal.* 9. p. 135. t. 69.

*Habitat in India. Torén.*

*Rami obsole angulati. Folia lato-lanceolata f. lanceolata - ovata, obtuse acuminata, integerrima, minus glabra. Pedunculi dichotomi, longitudine foliorum. Flores sessiles, bracteis minimis, lineatis. Calyces minimi. Corollæ tubus filiformis, longus. Limbi labium superius lineare, angustum, brevius, reflexum; L. inferius æquale, trifidum. Stamina extra fauces prominentia. Stylus capillaris, persistens.*

16. **JUSTICIA** foliis ovatis utrinque mucronatis inte- purpurea  
gerrimis glabris, caule geniculato, spicis secundis.  
*Orb. it.* 230.

*Folium tinctorum.* *Rumph. amb.* 6. p. 51. t. 22. f. 1.

*Habitat in China. Osbeck.*

*Caulis herbaceus, brachiatus. Folia lato-ovata, apice ☿ basi mucronata, glabra, petiolata, integerrima. Articulati caulis ☿ rami supra-basin Geniculo tumido, quod supra angustius evadit, notantur. Spicæ ad latera ☿ apices ramorum, secundæ. Bractææ angustæ, lanceolatae, longitudine capsularum. Corollæ purpureæ: labiis erectis; altero lineari angustissimo; altero lato trilobo. Stamina duo, purpurea, longitudine corollæ.*

17. **JUSTICIA** foliis lanceolato-ovatis, pedunculis sex- bivalvis,  
floris, pedicellis lateralibus bifloris, bracteis ovatis parallelis.

*Bungum.* *Rumph. amb.* 6. p. 55. t. 22. f. 1.

*Adel-odagam.* *Rheed. mal.* 9. p. 21. t. 43.

*Habitat in Asia indica. J. Burmannus.*

*Caulis brachiatus. Folia opposita, integerrima. Pedunculi breves, terminati involucrio diphylo, setaceo Pedunculisque 4 parvis, parallelis: mediis duobus unifloris; lateralibus vero iterum similiter diphyllis bifloris. Calyces omnes inter Bracteas duas ovatas se ipsis majores. Corollæ Labium superius lanceolatum; inferius ovatum. Stamina 2, longitudine corollæ. Rheedii planta foliis serratis ♂ flore albo differt, eandem tamen separare nequeo.*

- gangetica. 18. **JUSTICIA** foliis ovatis, racemis simplicibus longis, floribus alternis secundis, bracteis obsoletis. *Aman. acad. 4. p. 299. \**  
*Habitat in Asia indica.*

### DIANTHERA.

- americana. 1. **DIANTHERA** spicis solitariis alternis.  
*Dianthera. Gron. virg. 6. \**  
*Gratiolæ affinis floridana, foliis floribus & capsulis in spica brevi, pediculis e foliorum alis innixis. Pluk. amalth. 114. t. 423. f. 5.*  
*Habitat in Virginia, Florida. 24*  
*Caulis simplicissimus. Folia linearia. Pedunculi alterni, longitudine foliorum. Spicæ ovatæ.*
- comata. 2. **DIANTHERA** spicis filiformibus verticillatis: inferioribus umbellatis.  
*Dianthera foliis lanceolato-ovatis, racemo spatioso assurgente; spicillis verticillatis. Brown. jam. 118. Antirrhinum minus angustifolium, flore dilute cæruleo. Sloan. jam. 49. hist. 1. p. 160. t. 103. f. 2.*  
*Habitat in Jamaica.*

### GRATIOLA.

- officinalis. 1. **GRATIOLA** floribus pedunculatis, foliis lanceolatis serratis. *Mat. med. 18.*  
*Gratiola. Riv. mon. 157. Hort. cliff. 9. Roy. lugdb. 292. Dalib. parif. 8. Sauv. monsp. 137.*  
*Gratiola centauroides. Bauh. pin. 279.*  
*Habitat in Lusatia, Gallia & australioris Europæ humidiusculis. 24*
- Monieria. 2. **GRATIOLA** foliis ovali-oblongis, pedunculis unifloris, caule repente. *Aman. acad. 4. p. 306. \**

Mo-

*Monieria minima repens*, foliis subrotundis, floribus singularibus alaribus. *Brown. jam.* 269.

*Monieria ramosa repens*, foliis linearibus oppositis. *Ehret. pict.* t. 14. f. 2.

*Anagallis aquatica*, portulacæ aquaticæ caule & foliis. *Sloan. jam.* 87. *hist.* 1. p. 203. t. 129. f. 1.

*Habitat in Jamaica.*

3. *GRATIOLA* foliis lanceolatis obtusis subdentatis. *virginiana.* *Gron. virg.* 6. \*

*Theria Maya Nari.* *Rheed. mal.* 9. p. 165. t. 85.

*Habitat in Virginia, Malabar.*

4. *GRATIOLA* floribus subsessilibus.

*Gratiola latiore folio*, flore albo. *Fewill. peruv.* 3. t. 47.

*peruviana.*

*Habitat in Peru.*

## PINGUICULA.

1. *PINGUICULA* nectarii apice incrassato.

*lusitanica.*

*Viola palustris*, *Pinguicula dicta*, *lusitanica.* *Grif. lusit.*

*Habitat in Lusitania.* 2.

2. *PINGUICULA* nectario cylindræo longitudine vulgaris.

petali. *Fl. lapp.* 11. *Fl. suec.* 21, 25. *Roy. lugdb.* 304.

*Sanicula montana*, flore calcari donato. *Bauh. pin.* 243.

*Pinguicula.* *Clus. hist.* 1. p. 311.

*Habitat in Europæ uliginosis.* 2.

3. *PINGUICULA* nectario conico petalo brevior. *alpina.*

*Fl. lapp.* 12. t. 12. f. 3. \*. *Fl. suec.* 22, 26.

*Pinguicula* flore albo minore, calcari brevissimo. *Raj.*

*hist.* 752.

*Habitat in Alpibus Lapponicis.* 2.

4. *PINGUICULA* scapo villoso. *Fl. lapp.* 13. t. 12. f. villosa.

2. *Fl. suec.* 23, 27.

*Pinguicula cornubiensis*, flore minore carneo. *Raj.*

*hist.* 752.

*Habitat in Lapponia, Sibiria.* 2.

## UTRICULARIA.

1. *UTRICULARIA* nectario subulato, foliis integer- *alpina.*

*rimis.* *Jacq. amer.* 11.

*Habitat in Alpibus Martinicæ insule.*



foliosa.

2. **UTRICULARIA** nectario conico, fructibus cernuis, radicibus utriculo destitutis. *Læfl. it.* 281. \*  
*Linaria palustris*, fœniculi folio. *Plum. spec.* 6. icon. 165. f. 2.

*Habitat in America australi.*

*Simillima U. vulgari*, sed *Radix repens*; Flores in scapo plures 6-10, ♂ nulli utriculi ad radículas.

vulgaris.

3. **UTRICULARIA** nectario conico, scapo paucifloro. *Fl. lapp.* 14. *Fl. fœc.* 44. 28. *Fl. Zeyl.* 22. *Roy. lugdb.* 304.

*Lenticularia. Riv. mon.* 78. \*

*Millefolium aquaticum lenticulatum. Bauh. pin.* 141.

*Habitat in Europæ fossis, paludibus profundioribus.* 24

*Nectarium subulatum, longitudine labii inferioris, cui appropinquatum; faux clausa palato gibbo. Planta nobis duplex: altera MAJOR a Rivino delineata, calyce postice rotundato. Altera MINOR calyce postice transverso, ♂ quasi truncato; crescunt conjunctim.*

minor.

4. **UTRICULARIA** nectario carinato. *Fl. fœc.* 25. 29. \*

*Millefolium palustre galericulatum minus. Pluk. alm.* 251. t. 99. f. 6.

*Aparine aquis innatans trevisana, foliis perchepter capreolis donata. Bocc. mus.* 1, p. 23. t. 4.

*Habitat in Europæ fossis variis.*

*Nectarium obsoletum, deorsum spectans; faux, absque gibbo palato, pervia ♂ hians.*

subulata.

5. **UTRICULARIA** nectario subulato. *Gron. virg.* 6. *Habitat in Virginia.*

gibba.

6. **UTRICULARIA** nectario gibbofo. *Utricularia florum nectario gibbofo, scapo nunc unifloro nunc bifloro. Gron. virg.* 129. *Habitat in Virginia.*

bifida.

7. **UTRICULARIA** scapo nudo bifido. *Osbeck. it.* 243. t. 3. f. 2.

*Habitat in China. Osbeck.*

*Caulis filiformis, digiti longitudine, bifidus s. duplici racemo, nudus. Flores alterni, lutei: labio superiore lateribus acuto.*

caerulea.

8. **UTRICULARIA** scapo nudo, squamis alternis vagis subulatis. *Fl. Zeyl.* 23. \*

Nell.

Nelipu. *Rheed. mal.* 9. p. 137. t. 70.  
*Habitat in Zeylona.*

## VERBENA.

\* *Diandra disperma.*

1. VERBENA diandra, spicis longissimis foliosis. orubica.  
*Verbena foliis ovatis acuminatis, spica foliola. Hort. cliff.* 10. \*  
*Verbena orubica, teucril folio, primulae veris flore, filiquis & feminibus longissimis. Pluk. alm.* 383. t. 228. f. 4. *Et* t. 327. f. 7.  
*Sherardia urticae folio subtus incano, floribus violaceis. Ehret. pict.* t. 5. f. 1.  
*Habitat in Oruba insula americana septentrionalis.*
2. VERBENA diandra, spicis longissimis carnosis nudis, foliis lanceolato-ovatis oblique dentatis, caule laevi. indica.  
*Habitat in Zeylona.* ☉. *Dav. Rozen.*  
*Smillima V. jamaicensis, sed Caulis levis absque pilis. Folia glabra nec retrorsum scabra, magis lineata, non serrata, sed dentibus obtusis obliquis dentata; basi sensim angustata, nec petiolis late marginatis. Flores cerulei antheris tantum duabus, reliquis Filamentis 2 castratis.*
3. VERBENA diandra, spicis longissimis carnosis nudis, foliis spatulato-ovatis serratis, caule hirtio. jamaicensis.  
*Verbena foliis obtuse ovalibus, spica carnosula nuda. Hort. cliff.* 10. *Roy. lugdb.* 327.  
*Verbena folio subrotundo serrato, flore caeruleo. Sloan. hist.* 171. t. 107. f. 1.  
*Habitat in Jamaica Et Caribæis.* 2. ☉
4. VERBENA diandra, spicis ovatis, foliis lanceolatis serrato-plicatis, caule fruticoso. stoechadifolia. *Roy. lugdb.* 327.  
*Sherardia nodiflora, stoechadis serratifolii folio. Vaill. sex.* 49.  
*Lavandula, foliis crenatis latioribus, americana frutescens. Plum. spec.* 6. ic. 162. f. 1.  
*Habitat in America Gallia aequinoctialis.*
5. VERBENA diandra, spicis laxis, calycibus alternis prismaticis truncatis aristatis, foliis ovatis obtusis. prismatica.  
*Verbena minima, chamædryos folio. Sloan, jam.* 64. *hist.* 1. p. 172. t. 107. f. 2.

Vet-

*Verbena*, *scutellariæ* f. *caſſidæ* folio, diſpermos a-  
mericana. *Pluk. alm.* 382. t. 70. f. 1.

*Habitat in Jamaica.*

mexicana.

6. *VERBENA* diandra, ſpicis laxis, calycibus fructus  
reflexis rotundato didymis hiſpidis.

*Verbena mexicana*, trachelii folio, fructu aparines.

*Dill. elh.* 407. t. 302. f. 389.

*Habitat in Mexico.* 2.

*Caulis tetragonus*, marginibus ſcaber : *Rami oppoſiti  
ſuperne dichotomi*, *Racemi dichotomiæ longi*. *Folia  
cordata*, oblonga, ſcabra, petiolis breviffimis.

lappulacea.

7. *VERBENA* diandra ? calycibus ſubrotundis erecti-  
uſculis, ſeminibus echinatis.

*Scorodonia* floribus ſpicatis purpureſcentibus pentape-  
taloidibus. *Sloan. jam.* 66. hiſt. 1. p. 174. t. 110. f. 1

*Blairia*. *Houſt. Ann. herb.* 277.

*Habitat in Jamaica.*

curaffavica.

8. *VERBENA* diandra, ſpicis longis, calycibus ari-  
ſtatis, foliis ovatis argute ferratis.

*Veronica* ſimilis fruticola curaffavica. *Herm. parad.* 240.

*Kämpferi*. *Houſt. mſſ.*

*Habitat in Curaſſao Americæ.*

\* \* *Tetrandra.*

nodiflora.

9. *VERBENA* tetrandra, ſpicis capitato-conicis, foli-  
is ferratis, caule repente. *Fl. Zeyl.* 399.

*Verbena* foliis verticaliter ovatis, ſpicis globosis. *Hort.  
cliff.* 11.

*Verbena* foliis verticaliter ovatis, ſpicis ſolitariis ova-  
tis. *Roy. lugdb.* 327.

*Verbena* caule repente, foliis oblongis ſuperne crena-  
tis, pedunculis ſolitariis capitatis. *Gron. virg.* 7.

*Verbena* nodiflora. *Bauh. pin.* 269. *prodr.* 125.

*Habitat in Virginia.* ○

bonarienſis.

10. *VERBENA* tetrandra, ſpicis ſcaſculatis, foliis lan-  
ceolatis amplexicaulibus. *Hort. apſ.* 8.

*Verbena* foliis lanceolatis, floribus congeſtis ſaſtigia-  
tis. *Hort. cliff.* 11. *Roy. lugdb.* 326.

*Verbena* bonarienſis altiffima, *lavendulæ canarienſis* fo-  
lio, ſpica *lavendulæ*. *Dill. elh.* 406. t. 300. f. 387.

*Habitat in agro Bonarienſi.* 2.

11. VER-

11. VERBENA tetrandra, spicis longis acuminatis, foliis hastatis. *Hort. upf.* 8. hastata.  
 Verbena foliis lanceolatis ferratis, spicis filiformibus paniculatis. *Roy. lugdb.* 337.  
 Verbena americana altissima, spica multiplici, urticæ foliis angustis, floribus cæruleis. *Herm. parad.* 242. t. 242.  
*Habitat in Canadæ humidis.* 24.
12. VERBENA tetrandra, spicis filiformibus, foliis indivisis lanceolatis ferratis obtusiusculis subsessilibus. Carolina.  
 Verbena carolinensis, melissæ folio aspero. *Dill. elth.* 407. t. 301. f. 388.  
 Verbena caroliniana, folio integro ferrato scabro. *Raj. suppl. app.* 249. n. 10.  
*Habitat in America septentrionali.* 24.
13. VERBENA tetrandra, spicis filiformibus paniculatis, foliis indivisis ovatis ferratis acutis petiolatis. *Hort.* urticifolia.  
*upf.* 9.  
 Verbena foliis ovatis, caule erecto, spicis filiformibus paniculatis. *Hort. cliff.* 11. *Gron. virg.* 7. *Roy. lugdb.* 327.  
 Verbena recta canadensis, urticæ foliis. *Morif. hist.* 3. p. 418. f. 11. t. 25. f. 3.  
*Habitat in Virginie, Canadæ aridis.* 24.
14. VERBENA tetrandra, spicis filiformibus, foliis multifido-laciniatis, caulibus numerosis. *Hort. upf.* 8. spuria.  
 Veronica humilior, foliis incis. *Clayt. virg.* 8.  
*Habitat in Canada, Virginia.* 24.
15. VERBENA tetrandra, spicis filiformibus paniculatis, foliis multifido-laciniatis, caule solitario. *Fl. fvec.* 2. n. 30. officinalis.  
 Verbena foliis multifido-laciniatis, spicis filiformibus. *Hort. cliff.* 11. *Fl. fvec.* 1. n. 26. *Roy. lugdb.* 327. *Dalib. parif.* 9. *Sauv. monsp.* 279.  
 Verbena communis, flore cæruleo. *Bauh. pin.* 269.  
 Verbena recta *Dod. pempt.* 150.  
*Habitat in Europæ mediterraneæ ruderatis.* ☉
16. VERBENA tetrandra, spicis filiformibus solitariis, foliis bipinnatifidis. supina.  
 Verbena tenuifolia. *Bauh. pin.* 269.  
 Verbena supina. *Clus. hist.* 2. p. 46.  
*Habitat in Hispania.*

## LYCOPUS.

- europæus.** 1. **LYCOPUS** foliis sinuato-ferratis. *Fl. fvec. 2. n. 31.*  
*Lycopus foliis indivisis. Fl. fvec. 1. n. 27. Dalib. pæ-*  
*ris. 9.*  
*Lycopus. Hort. cliff. 11. Roy. lugd. 326.*  
*Marrubium palustre glabrum. Bauh. pin. 230.*  
*Sideritis matthioli. Dalech. hist. 1117.*  
*Habitat in Europæ ripis humentibus. 24*
- virginicus.** 2. **LYCOPUS** foliis æqualiter serrulatis.  
*Lycopus foliis lanceolatis tenuissime ferratis. Gron.*  
*virg. 8.*  
*Habitat in Virginia. 24*

## AMETHYSTEA.

- cærulea.** 1. **AMETHYSTEA.** *Hort. upf. 9. Amoen. acad. 386. \**  
*Amethystina montana erecta, foliis exiguis digitatis*  
*trifidis serratis, flosculis cum coma e cæruleo-jan-*  
*thinis. Amm. ruth. 4. Hall. act. upf. 1742. p. 51. f. 1.*  
*Habitat in Sibiria montosis. ☉*

## CUNILA.

- mariana.** 1. **CUNILA** foliis ovatis ferratis, corymbis terminali-  
 bus dichotomis.  
*Satureja origanoides. Sp. pl. 1. p. 568.*  
*Hedyosmos. Mitch. E. N. C. 1748. app. 211.*  
*Thymus foliis ovatis acuminatis ferratis, corymbis la-*  
*teralibus terminalibus pedunculatis. Gron. virg. 64.*  
*Calamintha mariana, mucronatis rigidioribus & cre-*  
*natis foliis, flosculorum calyculis villis argenteis.*  
*Pluk. mant. 34. t. 344. f. 1.*  
*Calamintha erecta virginiana, mucronato folio glabro.*  
*Morif. hist. 3. p. 413. f. 11. t. 19. f. 7.*  
*Habitat in Virginia. 24*
- pulegioides.** 2. **CUNILA** foliis oblongis unidentatis, floribus verti-  
 cillatis.  
*Melissa pulegioides. Sp. plant. 1. p. 593.*  
*Melissa floribus verticillatis glomeratis secundum lon-*  
*gitudinem caulis, foliis tomentosis. Gron. virg. 167.*  
*Kalm. it. 2. p. 314.*  
*Habitat in Virginia, Canadæ siccis. ☉*  
*Planta spithamea, brachiata. Folia lanceolato-ova-*  
*ta,*

ta, scabriuscula, uno alterove dente notata: superiora angustiora. Verticilli secundum totam longitudinem plantæ. Calyx decemstriatis scaber: Labio superiore trifido acuminato; inferiore setaceo. Corolla alba fauce violacea: Labio superiore vix emarginato. Stamina duo corolla breviora, fertilia & Filamenta alia duo minoræ castrata.

3. CUNILA foliis ovalibus integerrimis, floribus verticillatis, caule tetragono. thymoides.

Thymus pulegioides. Sp. pl. 1. p. 592.

Acinos thymi folio & facie, floribus inexpandis. Morif. hist. 3. p. 404. f. 11. t. 19. f. 6.

Habitat Monipellii. ☉. D. Sauvages.

Habitus præcedentis. Caulis erectus, spithameus. Ramis paucis simplicibus brevibus. Folia ovalia, obtusa, glabra, subtus striata. Verticilli secundum totam caulis longitudinem. Stamina inquirant in viva ejus contraneis.

## ZIZIPHORA.

1. ZIZIPHORA capitulis terminalibus, foliis ovatis. capitata.

Ziziphora foliis lanceolatis, floribus terminatricibus.

Hort. cliff. 305. Roy. lugdb. 313.

Clinopodium fistulosum pumilum Indiæ occidentalis, summo caule floridum. Plu. alm. 111. t. 164. f. 4.

Thymus humilis latifolius. Buxb. cent. 3. p. 28. t. 51. f. 1.

Habitat in Syria, Armeria. ☉

2. ZIZIPHORA foliis ovatis, floribus racemoso-spicatis, bracteis obovatis nervosis acutis. hispanica.

4. p. 263. \*

Habitat in Hispania. ☉

3. ZIZIPHORA floribus lateralibus, foliis lanceolatis. tenuior.

Ziziphora foliis lanceolatis, floribus ex alis. Hort. cliff. 305.

Ziziphora foliis lanceolatis, floribus lateralibus. Roy. lugdb. 313. Hort. upf. 9.

Acinos syriaca, folio mucronato, capsulis hirsutis.

Morif. hist. 3. p. 404. f. 11. t. 19. f. 3. 4.

Habitat in Syria? ☉

4. ZIZIPHORA floribus lateralibus, foliis ovatis. acinoides.

*Clinopodium supinum incanum. Amm. ruth. 66.*

*Habitat in Sibiria. ☉*

*Habitus Thymi Acinos, sed majora quintuplo omnia. Folia Z. capitata; Flores laterales ut in Z. tenuiore, sed copiosiores. Calyces similiter hispidi; stamina extra corollæ tubum protrusa.*

### MONARDA.

*fistulosa.*

1. MONARDA capitulis terminalibus, caule obtusangulo. *Hort. upf. 12. Mill. dict. t. 122. f. 2.*

*Monarda floribus capitatis, caule obtuso. Vir. cliff. 3. Roy. ludgb. 313.*

*Monarda. Hort. cliff. 11.*

*Origanum fistulosum canadense. Corn. canad. 13. t. 14.*

- β. *Monarda mollis. Amoen. acad. 3. p. 399.*

*Clinopodium majus virginienſe, foliis minus hirtis acutioribus, floribus fistulosis. Morif. hist. 3. p. 374.*

*Habitat in Canada. 24*

*didyma.*

2. MONARDA floribus capitatis subdidynamis, caule acutangulo. *Mill. ic. 122. f. 1.*

*Monarda caule acute angulato, capitulis terminalibus. Hort. cliff. 495. Cold. noveb. 7.*

*Monarda floribus capitatis verticillatisque, caule acute angulato, foliis lanceolato-serratis glabris. Bött. cun. 226.*

*Habitat in Pennsylvania, Noveboraco.*

*clinopodia.*

3. MONARDA floribus capitatis, foliis lævissimis serratis.

*Monarda foliis ovato-lanceolatis, verticillis lateralibus dichotomis corymbosis, foliis inæqualiter serratis Gron. virg. 9.*

*Habitat in Virginia. 24*

*Radix repens. Facies plantæ præcedentis quoad staturam, at folia gerit Clinopodii incani omnino; neque huic spica rubra, sed purpurea, nec flores tetrandri; foliisque glaberrimis distincta a prima; Folia ovato-oblonga, acuminata, profunde serrata, petiolata, utrinque lævia. Caulis quadrangularis, magis quam in prima, minus argute quam in secunda. Flores in capitulum bracteis radiatum, pallidi.*

*punctata.*

4. MONARDA floribus verticillatis, corollis punctatis. *Hort. upf. 12.*

*Monarda floribus verticillatis. Hort. cliff. 495. Gron. virg. 9. Roy. lugdb. 313.*

*Clinopodium virginianum angustifolium, quovis verticillo duodecim foliolis rubentibus cincto. Pluk. alm. 111. t. 24. f. 1.*

*Habitat in Virginia. ☉*

5. **MONARDA floribus verticillatis, corollis involu-** *cillata,*  
*cro longioribus.*

*Monarda spica interrupta, involucris longitudine verticillorum lanceolatis. Gron. virg. 9. \**

*Clinopodium angustifolium non ramosum, flore cæruleo: labio trifido atro-purpureis maculis ornato.*

*Pluk. alm. 10. t. 164. f. 3. Moris. hist. 3. p. 374. f. 11. t. 8. f. 6.*

*Habitat in Virginia.*

## ROSMARINUS.

1. **ROSMARINUS.** *Hort. cliff. 14. Hort. upf. 11. officinalis. Mat. med. 16. Roy. lugdb. 310.*

*Rosmarinus spontaneus, latiore folio. Bauh. pin. 217.*

2. *Rosmarinus hortensis, angustiore folio. Bauh. pin. 217.*

*Habitat in Hispaniæ, G. narbonensis, Italiæ, orientus, Galilææ collibus. †*

## SALVIA.

1. **SALVIA foliis lineari-lanceolatis denticulatis, flori-** *ægyptiaca,*  
*bis pedunculatis. Hort. cliff. 13. \* Roy. lugdb. 309.*

*Horminum ægyptium minimum ramosissimum. Lippi.*

*Bœrh. lugdb. 1. p. 66.*

*Habitat in Ægypto.*

2. **SALVIA foliis lanceolatis, calycibus diphyllis:** *cretica.*

*Habitat in Creta.*

3. **SALVIA foliis radicalibus lyratis dentatis, corol-** *lyrata,*  
*larum galea brevissima.*

*Salvia corollarum labio superiore brevior; fauce patente. Gron. virg. 8.*

*Salvia foliis sinuato-ferratis glabris, calyce piloso vincto, corolla vix calycem superante. Zinn. get. 298?*

*Horminum virginianum, caule aphylo, foliis quercinis, tubuloso longo flore. Moris. hist. 3. p. 395.*

*f. 11. t. 13. f. 27.*

*Habitat in Virginia.*

*Linæi Spec. plant. pars. I.*

C

4. SAL-



- officinalis.** 4. *SALVIA* foliis lanceolato-ovatis integris crenulatis, floribus spicatis, calycibus acutis. *Hort. cliff.* 12. *Hort. upf.* 10. *Mat. med.* 13. *Roy. lugdb.* 307. *Sauv. monsp.* 143.  
*Salvia major.* *Bauh. pin.* 237.  
 5. *Salvia minor aurita & non aurita.* *Bauh. pin.* 237.  
*Habitat in Europa australi.* ☿
- pomifera,** 5. *SALVIA* foliis lanceolato-ovatis integris crenulatis, floribus spicatis, calycibus obtusis. *Hort. cliff.* 12. *Roy. lugdb.* 307.  
*Salvia cretica frutescens pomifera, foliis longioribus incantans & crispis.* *Tournef. cor.* 10. *itin.* 1. p. 92. t. 92.  
*Salvia baccifera.* *Bauh. pin.* 237.  
*Habitat in Creta.*
- urticifolia.** 6. *SALVIA* foliis ovato-oblongis duplicato-ferratis, calycibus tridentatis: lacinia summa tridentata. *Gron. virg.* 8.  
*Horminum minus, betonicae folio, flore purpureo.* *Pluk. amalrh.* 119. t. 420. f. 1.  
*Horminum virginianum erectum, urticae foliis, flore minore.* *Morif. hist.* 3. p. 395. f. 11. t. 13. f. 31.  
*Habitat in Virginia, Florida.*
- viridia.** 7. *SALVIA* foliis oblongis crenatis, corollarum galea semi-orbiculata, calycibus fructiferis reflexis.  
*Salvia foliis ovato-oblongis obtusis æqualiter crenatis, corollarum galea semi-orbiculata.* *Hort. upf.* 11. \*  
*Horminum coma viridi.* *Tournef. inst.* 178.  
*Habitat - - - - -* ☉
- Horminum.** 8. *SALVIA* foliis obtusis crenatis, bracteis summis sterilibus majoribus coloratis. *Vir. cliff.* 4. *Hort. cliff.* 12. *Mat. med.* 14. *Roy. lugdb.* 310.  
*Horminum sativum.* *Bauh. pin.* 238.  
*Orminum verum matthioli.* *Gesn. fasc.* 17. t. 11. f. 21.  
*Habitat in Græcia, Apulia.* ☉
- sylvestris.** 9. *SALVIA* foliis cordato-lanceolatis undatis biserratis maculatis acutis, bracteis coloratis flore brevioribus.  
*Horminum sylvestre salvifolium majus maculatum.* *Bauh. pin.* 239.  
*Horminum sylvestre v. altera species.* *Clus. hist.* 2. p. 31.  
*Habitat in Austriæ inferioris, Bohemæ agrorum marginibus, vineis.* 24

10. **SALVIA** foliis cordato-lanceolatis ferratis planis, nemorosa, bracteis coloratis, corollæ labio infimo reflexo.  
*Salvia* foliis cordato-lanceolatis simpliciter ferratis, staminibus galeam æquantibus. *Zinn. gat.* 297.  
*Horminum sylvestre salvifolium minus.* *Bauh. pin.* 239.  
*Sciarea*, folio *salviæ*, minor f. glabra. *Tournef. inst.* 180.  
*Habitat in Austria, Tartaria.* ♂  
*Folia Radicalia cordata; Caulina ovato-lanceolata, æqualiter ferrata, supra nuda, subtus subtomentosa. Spicæ ante florescentiam exacte tetragonæ bracteis coloratis flore brevioribus. Corollæ punctis impalpabilibus adpersæ; Labii inferioris medius lobus arcuæ sub calyce reflexus.*
11. **SALVIA** foliis cordato-ovatis rugosis tomentosis, hæmatodes, calycibus hispidis, radice tuberosa. †  
*Horminum sanguineum, asphodeli radice.* *Triumph. obs.* 69. t. 69.  
*Horminum sylvestre maculatum tuberosum.* *Honophr.*  
*Horminum sylvestre majus hæmatodes glabrum, flore cæruleo, italicum.* *Barr. rar.* 234. t. 185.  
*Habitat in Italia.* 24  
*Habitus Phlomidis, sed teneritudo Salviæ; Calyces pilis rectis exstantibus undique hispidi sunt.*
12. **SALVIA** foliis cordato-oblongis: summis amplexicaulibus, verticillis subnudis, corollis galea glutinosis. *Fl. fvec.* 2. n. 32.  
*Salvia* foliis ovatis inciso-crenatis, verticillis subnudis. *Hort. cliff.* 12. *Fl. fvec.* 1. n. 28. *Hort. upf.* 10. *Roy. lugdb.* 310. *Dalib. parif.* 10.  
*Horminum pratense, foliis ferratis.* *Bauh. pin.* 238.  
*Horminum sylvestre.* *Camer. epit.* 629.
13. **Salvia agrestis** foliis cordatis: summis amplexicaulibus, corollarum galea labium æquante. *Aman. acad.* 3. p. 399.  
*Horminum pratense niveum, foliis incanis.* *Bauh. pin.* 238.  
*Habitat in Europæ pratis.* 24
14. **SALVIA** foliis cordatis obtusis crenatis subtomentosis, corollis calyce angustioribus.  
*Habitat in Domingo.*
15. **SALVIA** foliis ferratis sinuatis læviusculis, corollis verbenacæ caly-

- calyce angustioribus. *Virid. cliff.* 17. *Gron. virg.* 2.  
*Roy. lugdb.* 309. *Dalib. parif.* 9. *Sauv. monsp.* 278.  
*Salvia* foliis pinnatim incisis glabris. *Hort. cliff.* 12.  
*Horminum* sylvestre, lavendulæ flore. *Bauh. pin.* 239.  
*Raj. hist.* 545.  
*Horminum* sylvestre minus, inciso folio, flore azureo. *Barr. ic.* 208.  
*Horminum*, verbenacæ lacinialis, angustifolium. *Triumph. obs.* 66. t. 66.  
*Habitat in Europæ & Orientis pascuis.* 2.
- clandestina.** 15. *SALVIA* foliis ferratis pinnatifidis rugosissimis, spica obtusa, corollis calyce angustioribus.  
*Horminum* sylvestre, inciso folio, cæsis flore, italicum. *Barr. rar.* 24. t. 220.  
*Habitat in Italia.* ♂  
*Caules* tetragoni, obtusi, villosi, ascendentes, spithamæi: *Ramis* superne sæpius duobus. *Folia* oblonga, pinnato-sinuata, rugosissima, crenata, obtusa, petiolata, in caule a. s. 3. paribus. *Spica* truncata verticillis remotis. *Bractææ* cordatæ, trifloræ. *Calyces* pilis glutinosi, labio superiore plicato. *Corolla* violacea, vix calyce duplo longior; labii inferioris medio lobo albo.
- pyrenaica.** 16. *SALVIA* foliis obtusis erosis, staminibus corolla duplo longioribus. *Roy. lugdb.* 309.  
*Horminum* pyrenaicum glutinosum anguriæ folio. *Herm. parad.* 187. t. 187.  
*Habitat in Pyrenæis.*
- diserma.** 17. *SALVIA* foliis cordato-oblongis erosis, staminibus corollam æquantibus.  
*Horminum* sylvestre majus, flore albo, integris foliis. *Barr. ic.* 187. bona.  
*Habitat in Syria.* 2.  
*Caules* pedales, villosi. *Folia* cordata, oblonga, tomentosa, rugosa, margine erosa, grato gravi odore. *Bractææ* cordatæ, villosæ, piloso-viscidæ. *Corollæ* albæ. *Stamina* longitudine corollæ, *Antheris* luteis. *Calyces* bilabiati.
- syriaca.** 18. *SALVIA* foliis cordatis dentatis: inferioribus repandis, bracteis glabris acutis cordatis, calycibus tomentosis.  
*Horminum* syriacum. *Bauh. pin.* 238. *prodr.* 114. t. 114.  
*Ha-*

*Habitat in Oriente, Palestina. Hasselquist.*

*Satura S. pratensis.* Caules obtuse angulati. Folia cordata, rugosissima, subtomentosa, repanda, dentata, obtuse crenata. Panícula brachiata, virgata. Bractæ calyce breviores, cordatæ. Calyces quinquesidi, æquales, acuminati, tomentosi. Corollæ albæ. Stylus galea duplo longior.

19. *SALVIA* foliis ovatis utrinque acuminatis serra- mexicana.  
tis. *Hort. cliff.* 13. *Roy. lugdb.* 308.

*Sclarea mexicana altissima, facie heliotropii. Willd.*  
*elsh.* 339. t. 254. f. 330.

*Habitat in Mexici humentibus. L.*

20. *SALVIA* foliis ovatis, petiolis utrinque mucronatis, hispanica.  
spicis imbricatis, calycibus trifidis. *Kniph. orig. t. 2.*

*Homninum sylvestre, lavendulæ flore. Bauh. pin.* 239?  
*Sclarea hispanica. Tabern. hist.* 764. ic. 374.

*Habitat in Italia. D. Rathgeb; in Hispania. Læsting. O*

*Camis sesquipedalis, tetragonus, obtusus, quadrifurcatus,*  
*superne pilis reflexis albis vestitus. Folia ovata, a-*  
*cuta, rugosa, serrata, basi acuta; petiolis basi utrin-*  
*que obtuso mucrone prominentibus. Spica tetragona,*  
*imbricata Bracteis ovatis, longitudine calycum, cilia-*  
*tis. Calyces sub singula squama sepe 5, ore tridentati,*  
*acuti, dente superiore solitario; inferioribus duobus.*  
*Corolla cerulea: limbo longitudine calycis: galea bre-*  
*viore, subvillosa; labio inferiore trifido, obtuso, inte-*  
*gerrimo, basi maculis 2, pallidis.*

21. *SALVIA* foliis cordatis acute crenatis: summis sessi- indica.  
libus, verticillis subnudis remotissimis. *Roy. lugdb.* 15.

*Sclarea indica, floribus variegatis. Tournesf. inst.* 179.

*Habitat in India.*

22. *SALVIA* foliis cordatis crenato-dentatis, verticil- verticillata.  
lis subnudis, stylo corollarum labio inferiore in-  
cumbentibus. *Hort. upf.* 11. \*

*Salvia foliis cordato-sagittatis dentatis. Hort. cliff.* 495.

*Roy. lugdb.* 309.

*Homninum sylvestre latifolium verticillatum. Bauh.*  
*pin.* 238.

*Homninum sylvestre tertium. Clus. hist.* 2. p. 29.

*Habitat in Austria, Misnia cultis & ruderalis. O*

23. *SALVIA* foliis cordato-sagittatis serratis acutis. *Hort. glauca,*  
*cliff.* 13. *Hort. upf.* 11. *Roy. lugdb.* 308. *Sauv. monsp.* 146.

*Horminum luteum glutinosum. Bauh. pin. 238.*

*Horminum sylvestre secundum. Clus. hist. 2. p. 29.*

*Habitat in Europæ lutosi. 2.*

- canariensis. 24. *SALVIA* foliis hastato-triangularibus oblongis crenatis obtusis. *Hort. cliff. 13. Hort. upf. 10. Roy. lugdb. 308.*

*Horminum canariense tomentosum, hastato folio.*

*Morif. hist. 3. p. 394 f. 11. t. 13. f. 17.*

*Habitat in Canariis. 5*

- africana. 25. *SALVIA* foliis subrotundis serratis: basi truncatis dentatis. *Hort. cliff. 13. Roy. lugdb. 308.*

*Salvia africana frutescens, folio scorodoniæ, flore violaceo. Comm. hort. 2. p. 181. t. 91.*

*Horminum hastatis amplioribus foliis, caulibus & pediculis araneosa lanugine. Pluk. alm. 185. t. 301. f. 2.*

*Habitat in Cap. b. Spei, locis argillofis. 5*

- aurea. 26. *SALVIA* foliis subrotundis integerrimis: basi truncatis dentatis. *Hort. cliff. 13. Roy. lugdb. 308.*

*Salvia africana frutescens, folio subrotundo glauco flore magnò aureo. Comm. hort. 2. p. 183. t. 92.*

*Habitat ad Cap. b. Spei juxta rivulos. 5*

- Sclarea. 27. *SALVIA* foliis rugosis cordatis oblongis villosis, serratis, bracteis floralibus calyce longioribus concavis acuminatis. *Hort. cliff. 12. Hort. upf. 10.*

*Mat. med. 15. Roy. lugdb. 309. Dalib. parif. 9.*

*Salvia foliis cordato-ovatis obscure undulatis, foliolis floralibus lanceolato-concavis. Guett. stamp. 263.*

*Horminum Sclarea dictum. Bauh. pin. 228.*

*Orvala. Dod. pempt. 292.*

*Habitat in Syria, Italia. 3*

- argentea. 28. *SALVIA* foliis oblongis dentato-angulatis lanatis, verticillis summis sterilibus, bracteis concavis.

*Sclarea ficula, folio argenteo subrotundo. Boerh. lugdb. 1. p. 163.*

*Æthiopis tota argentea cretica lanuginosa. Cup. cath.*

*Habitat in Creta. 5*

*Media inter S. Sclaream & Æthiopem. Folia erosa in sterili herba maxime tomentosa. Panicula minus brachiata, pubescens, viscosa, ramorum summitatibus sterilibus, quasi spinosis. Bracteæ concavæ, muticæ. Corolla alba galea purpurascenti-pubescente; labio flavicane. Stylus corolla longior. 29.*

29. *SALVIA* foliis oblongis erosis lanatis, verticillis *Æthiopis*.  
lanatis, bracteis recurvatis subspinosis.

*Salvia* foliis lanceolatis sinuato-dentatis: floralibus verticillos comprimentibus. *Hort. cliff.* 13. *Roy. lugdb.* 308.

*Æthiopis* foliis sinuosis. *Bauh. pin.* 241.

- β. *Æthiopis* laciniatis foliis. *Barr. rar.* 24. t. 188.

- γ. *Marum ægypticum*. *Vesl. alp.* 212. t. 212.

*Habitat in Illyria, Græcia, Africa.* ♂

*Labium inferius antice coheret & saccum format.*

30. *SALVIA* foliis rugosis pinnatifidis lanatis: verticillis summis sterilibus. *ceratophylla.*

*Salvia* foliis simplicibus dentato-pinnatis asperis. *Hort. cliff.* 13. *Roy. lugdb.* 308.

*Salvia* foliis simplicibus dentato-pinnatis rugosis verrucosis. *Hort. upf.* 10.

*Horminum* rugoso verrucosoque folio cornu cervi exprimente. *Pluk. alm.* 186. t. 194. f. 5. *Herm. parad.*

186. t. 186. *Morif. hist.* 3. p. 393. f. 11. t. 13. f. 6.

*Habitat in Persia.* ♂

31. *SALVIA* foliis compositis pinnatis. *Hort. cliff.* 13. pinnata. *Roy. lugdb.* 309.

*Salvia* orientalis latifolia hirsutissima viscosa pinnata, flore & calyce purpureis, inodora. *Boerh. lugdb.*

1. p. 167. t. 167.

*Horminum* arabicum, alatis foliis, flore rubello. *Pluk. alm.* 186. t. 194. f. 6.

*Habitat in Oriente & Arabia.*

32. *SALVIA* foliis cordatis inæqualiter basi excisis, *nutans,*  
caule subgundo, spicis ante florescentiam cernuis.

*Salvia* foliis cordatis obtuse crenatis, spicis florum nutantibus. *Afl. petrop.* 1747. p. 378. t. 14.

*Habitat in Imperio Rutheno.* 24

## COLLINSONIA.

1. *COLLINSONIA*. *Hort. cliff.* 14. t. 5. *Cold. novch. canadensis.*  
2. *Kalm. it.* 2. p. 317.

*Habitat in Virginia, Canada sylvis.* 24

## MORINA.

1. *MORINA*. *Hort. cliff.* 14. *Roy. lugdb.* 236. *persica.*

*Morina* orientalis, carlinæ folio. *Tournef. cor.* 48.

*ita.* 3. p. 132. t. 132.

*Habitat in Persia ad Hispahan.* 24

# DIGYNIA

## ANTHOXANTHUM.

- odoratum.** 1. **ANTHOXANTHUM** spica ovato-oblonga, flosculis subpedunculatis arista longioribus. Anthoxanthum flosculis diandris. *Hort. cliff.* 25. *Roy. lugdb.* 58. Anthoxanthum. *Fl. svec.* 29. 33. *Dalib. parif.* 10. Gramen pratense, spica flavescente. *Bauh. pin.* 3. *Scheuchz. gram.* 88. *Habitat in Europæ pratis.* 24
- indicum.** 2. **ANTHOXANTHUM** spica lineari, flosculis sessilibus arista brevioribus. *Fl. zeyl.* 25. \* *Habitat in India.*
- paniculatum.** 3. **ANTHOXANTHUM** floribus paniculatis. *Amen. acad.* 145. Gramen sparteum, panicula flavescente. *Rudb. clyf.* 1. f. 14. *Habitat in Europa australiore: ut in Horto Dei Monspelienfi.* Gramen in vivo ulterius examinandum iis, quibus adest. *Facies est Anth. odorati. Calyces quadriflori, Flores mutici.*

# TRIGYNIA.

## PIPER.

- nigrum.** 1. **PIPER** foliis ovatis subseptemnerviis glabris, petiolis simplicissimis. *Fl. zeyl.* 26. \* *Mat. med.* 26. Piper rotundum nigrum. *Bauh. pin.* 411. *Pluk. alm.* 297. t. 437. f. 1. *Morif. hist.* 3. p. 602. f. 15. t. 1. f. 1. Malago codi. *Rheed. mal.* 7. p. 23. t. 12. *Habitat in India.* 5
- Beetle.** 2. **PIPER** foliis ovatis oblongiusculis acuminatis septinerviis, petiolis bidentatis. *Fl. zeyl.* 27. \* Piper qui Saururus foliis septinerviis oblongo-acuminatis. *Burm. zeyl.* 193. t. 82. f. 2. Beetla-codi. *Rheed. mal.* 7. p. 29. t. 15. *Habitat in India.* 5

3. PIPER foliis ovatis acutiusculis subtus scabris: ner- Malamiri,  
vis quinque subtus elevatis. *Fl. zeyl.* 26. \*  
Sirium. *Rumph. amb.* 5. p. 336. t. 116. f. 2.  
Amalago. *Rheed. mal.* 7. p. - - - t. 16.  
*Habitat in India utraque.*
4. PIPER foliis cordatis subseptinerviis venosis. *Fl. zeyl.* Siriboa.  
29. \*  
Siriboa. *Rumph. amb.* 5. p. 340. t. 117. f. 2.  
Betela quem Sirii-boa vocant. *Bont. jav.* 91. t. 91.  
*Habitat in India.*
5. PIPER foliis cordatis petiolatis sessilibusque. *Fl. zeyl.* longum.  
30. \* *Mat. med.* 20.  
Piper longum orientale. *Bauh. pin.* 412.  
Piper longum. *Rumph. amb.* 5. p. 333. t. 116. f. 2.  
Piper longum, pistolochiæ foliis absque pediculis,  
maderaspatanum. *Pluk. alm.* 297. t. 104. f. 4.  
Cattu-tirpali. *Rheed. mal.* 7. p. 27. t. 14.  
*Habitat in India.*
6. PIPER foliis lanceolato-ovatis quinquenerviis rugosis. Amalago.  
Piper longum arboreum altius, folio nervoso minore,  
spica gracili & brevior. *Sloan. hist.* 1. p. 134. t. 87. f. 1.  
Piper frutex, spica longa gracili. *Pluk. alm.* 297. t.  
215. f. 2.  
Piper longum. *Rumph. amb.* 5. p. 333. t. 116. f. 1.  
Saururus foliis lanceolato-ovatis quinquenerviis ru-  
gosis. *Hort. cliff.* 140.  
*Habitat in Jamaica.*
7. PIPER foliis cordatis novem-nerviis reticulatis. decumanum  
Saururus frutescens foliis plantagineis, fructu brevior.  
*Plum. amer.* 59. t. 76.  
Sirium decumanum. *Rumph. amb.* 5. p. 45. t. 27.  
*Habitat in Indiis.* §  
*Affinis nimis P. reticulato, ut fere varietas.*
8. PIPER foliis cordatis septemnerviis reticulatis. reticulatum.  
Saururus botryoides major arborescens, foliis planta-  
gineis. *Plum. amer.* 57. t. 75.  
Jaborandi. *Marsgr. bras.* 37. *Pis. bras.* 97.  
*Habitat in Martinica, Brasilia.*
9. PIPER foliis ovato-lanceolatis: nervis alternis, spi- aduncum.  
cis uncinatis.



- calyce angustioribus. *Virid. cliff.* 17. *Gron. virg.* 8.  
*Roy. lugdb.* 309. *Dalib. parif.* 9. *Sauv. monsp.* 278.  
*Salvia* foliis pinnatim incilis glabris. *Hort. cliff.* 12.  
*Horminum* sylvestre, lavendulæ flore. *Bauh. pin.* 239.  
*Raj. hist.* 545.  
*Horminum* sylvestre minus, inciso follo, flore azureo. *Barr. ic.* 208.  
*Horminum*, verbenacæ laciniis, angustifolium. *Triumph. obs.* 66. t. 66.  
*Habitat in Europæ & Orientis pascuis.* 2.
- clandestina.* 15. *SALVIA* foliis serratis pinnatifidis rugosissimis, spica obtusa, corollis calyce angustioribus.  
*Horminum* sylvestre, inciso folio, cæcis flore, italicum. *Barr. rar.* 24. t. 220.  
*Habitat in Italia.* ♂  
*Caules* tetragoni, obtusi, villosi, adscendentes, spithamæi: *Ramis* superne sapius duobus. *Folia* oblonga, pinnato-sinuata, rugosissima, crenata, obtusa, petiolata, in caule 2. s. 3. paribus. *Spica* truncata verticillis remotis. *Bractææ* cordatæ, trifloræ. *Calyces* pilis glutinosi, labio superiore plicato. *Corolla* violacea, vix calyce duplo longior; labii inferioris medio lobo albo.
- pyrenaica.* 16. *SALVIA* foliis obtusis erosis, staminibus corolla duplo longioribus. *Roy. lugdb.* 309.  
*Horminum* pyrenaicum glutinosum anguriæ folio. *Herm. parad.* 187. t. 187.  
*Habitat in Pyrenæis.*
- disformis.* 17. *SALVIA* foliis cordato-oblongis erosis, staminibus corollam æquantibus.  
*Horminum* sylvestre majus, flore albo, integris foliis. *Barr. ic.* 187. bona.  
*Habitat in Syria.* 2.  
*Caules* pedales, villosi. *Folia* cordata, oblonga, tomentosa, rugosa, margine erosa, grato gravi odore. *Bractææ* cordatæ, villosæ, piloso-viscidæ. *Corollæ* albæ. *Stamina* longitudinæ corollæ, *Antheris* luteis. *Calyces* bilabiati.
- syriaca.* 18. *SALVIA* foliis cordatis dentatis: inferioribus repandis, bracteis glabris acutis cordatis, calycibus tomentosis.  
*Horminum* syriacum. *Bauh. pin.* 238. *prodr.* 114. t. 114. *Ha.*

*Habitat in Oriente, Palæstina. Hasselquist.*

*Satura S. prætensis. Caules obtuse angulati. Folia cordata, rugosissima, subtomentosa, repanda, dentata, obtuse crenata. Panicula brachiata, virgata. Bractææ calyce breviores, cordatæ. Calyces quinquesidi, æquales, acuminati, tomentosi. Corollæ albæ. Stylus galea duplo longior.*

19. *SALVIA* foliis ovatis utrinque acuminatis serra- mexicana.  
tis. *Hort. cliff. 13. Roy. lugdb. 308.*

*Sclarea mexicana altissima, facie heliotropii. Dill.  
elh. 339. t. 254. f. 330.*

*Habitat in Mexici humentibus. §*

20. *SALVIA* foliis ovatis, petiolis utrinque mucronatis, hispanica.  
spicis imbricatis, calycibus trifidis. *Kniph. orig. t. 2.*

*Horminum sylvestre, lavendulæ flore. Bauh. pin. 239?*

*Sclarea hispanica. Tabern. hist. 764. ic. 374.*

*Habitat in Italia. D. Rathgeb; in Hispania. Læfing. ☉*

*Caulis sesquipedalis, tetragonus, obtusus, quadrifidus, superne pilis reflexis albis vestitus. Folia ovata, acuta, rugosa, serrata, basi acuta; petiolis basi utrinque obtuso mucrone prominentibus. Spica tetragona, imbricata Bracteis ovatis, longitudine calycum, ciliatis. Calyces sub singula squama sæpe 5, ore tridentati, acuti, dente superiore solitario; inferioribus duobus. Corolla cærulea: limbo longitudine calycis: galea brevior, subvillosa; labio inferiore trifido, obtuso, integerrimo, basi maculis 2, pallidis.*

21. *SALVIA* foliis cordatis acute crenatis: summis sessi- indica.  
libus, verticillis subnudis remotissimis. *Roy. lugdb. 15.*

*Sclarea indica, floribus variegatis. Tournesf. inst. 179.*

*Habitat in India.*

22. *SALVIA* foliis cordatis crenato-dentatis, verticil- verticillata.  
lis subnudis, stylo corollarum labio inferiore incumbentibus. *Hort. upf. 11. \**

*Salvia foliis cordato-sagittatis dentatis. Hort. cliff. 495.  
Roy. lugdb. 309.*

*Horminum sylvestre latifolium verticillatum. Bauh.  
pin. 238.*

*Horminum sylvestre tertium. Clus. hist. 2. p. 29.*

*Habitat in Austris, Misnia cultis & ruderalis. ☉*

23. *SALVIA* foliis cordato-sagittatis serratis acutis. *Hort. glutinosa,  
cliff. 13. Hort. upf. 11. Roy. lugdb. 308. Sauv. monsp. 146.*

Horminum luteum glutinosum. *Bauh. pin.* 238.

Horminum sylvestre secundum. *Clus. hist.* 2. p. 29.

*Habitat in Europæ luteis.* 24

- canariensis. 24. *SALVIA* foliis hastato-triangularibus oblongis crenatis obtusis. *Hort. cliff.* 13. *Hort. upf.* 10. *Roy. lugdb.* 308.

Horminum canariense tomentosum, hastato folio.

*Morif. hist.* 3. p. 394 f. 11. t. 13. f. 17.

*Habitat in Canariis.* 5

- africana. 25. *SALVIA* foliis subrotundis ferratis: basi truncatis dentatis. *Hort. cliff.* 13. *Roy. lugdb.* 308.

*Salvia africana* frutescens, folio scorodoniæ, flore violaceo. *Comm. hort.* 2. p. 181. t. 91.

Horminum hastatis amplioribus foliis, caulibus & pediculis araneosa lanugine. *Pluk. alm.* 185. t. 301. f. 2.

*Habitat in Cap. b. Spei, locis argillofis.* 5

- aurea. 26. *SALVIA* foliis subrotundis integerrimis: basi truncatis dentatis. *Hort. cliff.* 13. *Roy. lugdb.* 308.

*Salvia africana* frutescens, folio subrotundo glauco flore magnò aureo. *Comm. hort.* 2. p. 183. t. 92.

*Habitat ad Cap. b. Spei juxta rivulos.* 5

- Sclarea. 27. *SALVIA* foliis rugosis cordatis oblongis villosis, ferratis, bracteis floralibus calyce longioribus concavis acuminatis. *Hort. cliff.* 12. *Hort. upf.* 10.

*Mat. med.* 15. *Roy. lugdb.* 309. *Dalib. parif.* 9.

*Salvia* foliis cordato-ovatis obscure undulatis, foliis floralibus lanceolato-concavis. *Guett. stamp.* 263.

Horminum *Sclarea* dictum. *Bauh. pin.* 228.

Orvala. *Dod. pempt.* 292.

*Habitat in Syria, Italia.* 5

- argentea. 28. *SALVIA* foliis oblongis dentato-angulatis lanatis, verticillis summis sterilibus, bracteis concavis.

*Sclarea ficula*, folio argenteo subrotundo. *Boerh. lugdb.* 1. p. 163.

*Æthiopis* tota argentea cretica lanuginosa. *Cup. cath.*

*Habitat in Creta.* 5

*Media inter S. Sclaream & Æthiopem.* Folia erosa in sterili herba maxime tomentosa. Panicula minus brachiata, pubescens, viscosa, ramorum summitatibus sterilibus, quasi spinosis. Bracteæ concavæ, muticæ. Corolla alba galea purpurascenti-pubescente; labio flavicaneus. Stylus corolla longior. 29.

29. **SALVIA** foliis oblongis erosis lanatis, verticillis *Æthiopis*.  
 lanatis, bracteis recurvatis subspinosis.  
*Salvia* foliis lanceolatis sinuato-dentatis: floralibus verticillos comprimentibus, *Hort. cliff.* 13. *Roy. lugdb.* 308.  
*Æthiopis* foliis sinuosis. *Bauh. pin.* 241.  
 30. *Æthiopis* laciniatis foliis. *Barr. rar.* 24. t. 188.  
 31. *Marum ægypticum*. *Vesl. alp.* 212. t. 212.  
*Habitat in Illyria, Græcia, Africa.* ♂  
*Labium inferius antice cohaeret & saccum format.*
30. **SALVIA** foliis rugosis pinnatifidis lanatis: verticillis summis sterilibus, *ceratophylla*.  
*Salvia* foliis simplicibus dentato-pinnatis asperis. *Hort. cliff.* 13. *Roy. lugdb.* 308.  
*Salvia* foliis simplicibus dentato-pinnatis rugosis verrucosis. *Hort. upf.* 10.  
*Horminum* rugoso verrucosoque folio cornu cervi exprimente. *Pluk. alm.* 186. t. 194. f. 5. *Herm. parad.* 186. t. 186. *Morif. hist.* 3. p. 393. f. 11. t. 13. f. 6.  
*Habitat in Persia.* ♂
31. **SALVIA** foliis compositis pinnatis. *Hort. cliff.* 13. *pinnata*.  
*Roy. lugdb.* 309.  
*Salvia* orientalis latifolia hirsutissima viscosa pinnata, flore & calyce purpureis, inodora. *Boerh. lugdb.* 1. p. 167. t. 167.  
*Horminum arabicum*, alatis foliis, flore rubello. *Pluk. alm.* 186. t. 194. f. 6.  
*Habitat in Oriente & Arabia.*
32. **SALVIA** foliis cordatis inæqualiter basi excisis, *nutans*.  
 canle subnudo, spicis ante florescentiam cernuis.  
*Salvia* foliis cordatis obtuse crenatis, spicis florum nutantibus. *Aff. petrop.* 1747. p. 378. t. 14.  
*Habitat in Imperio Rutheno.* 21

## COLLINSONIA.

1. **COLLINSONIA**. *Hort. cliff.* 14. t. 3. *Cold. novch. canadensis*.  
 8. *Kalm. it.* 2. p. 317.  
*Habitat in Virginia, Canadae sylvis.* 21

## MORINA.

1. **MORINA**. *Hort. cliff.* 14. *Roy. lugdb.* 236. *persica*.  
*Morina* orientalis, carlinæ folio. *Tournef. cor.* 48.  
*ita.* 3. p. 132. t. 132.  
*Habitat in Persia ad Hispaham.* 21

# DIGYNIA

## ANTHOXANTHUM.

- odoratum.** 1. **ANTHOXANTHUM** spica ovato-oblonga, flosculis subpedunculatis arista longioribus. Anthoxanthum flosculis diandris. *Hort. cliff.* 25. *Roy. lugdb.* 58. Anthoxanthum. *Fl. svec.* 29. 33. *Dalib. parif.* 10. Gramen pratense, spica flavescente. *Bauh. pin.* 3. *Scheuchz. gram.* 88. *Habitat in Europæ pratis.* 2.
- indicum.** 2. **ANTHOXANTHUM** spica lineari, flosculis sessilibus arista brevioribus. *Fl. zeyl.* 25. \* *Habitat in India.*
- paniculatum.** 3. **ANTHOXANTHUM** floribus paniculatis. *Aman.* 1. *acad.* 145. Gramen sparteum, panicula flavescente. *Rudb. elys.* 1. f. 14. *Habitat in Europa australiore: ut in Horto Dei Monspelienfi.* Gramen in vivo ulterius examinandum iis, quibus adest. *Facies est Anth. odorati. Calyces quadriflori, Flores mutici.*

# TRIGYNIA

## PIPER.

- nigrum.** 1. **PIPER** foliis ovatis subseptemnerviis glabris, petiolis simplicissimis. *Fl. zeyl.* 26. \* *Mat. med.* 26. Piper rotundum nigrum. *Bauh. pin.* 411. *Pluk. alm.* 297. t. 437. f. 1. *Morif. hist.* 3. p. 602. f. 15. t. 1. f. 1. Malago codi. *Rheed. mal.* 7. p. 23. t. 12. *Habitat in India.* 5
- Beetle.** 2. **PIPER** foliis ovatis oblongiusculis acuminatis septinerviis, petiolis bidentatis. *Fl. zeyl.* 27. \* Piper qui Saururus foliis septinerviis oblongo-acuminatis. *Burm. zeyl.* 193. t. 82. f. 2. Beetla-codi. *Rheed. mal.* 7. p. 29. t. 15. *Habitat in India.* 5

3. **PIPER** foliis ovatis acutiusculis subtus scabris: ner- Malamiri,  
vis quinque subtus elevatis. *Fl. zeyl.* 26. \*  
Sirium. *Rumph. amb.* 5. p. 336. t. 116. f. 2.  
Amalago. *Rheed. mal.* 7. p. - - - t. 16.  
*Habitat in India utraque.*
4. **PIPER** foliis cordatis subseptinerviis venosis. *Fl. zeyl.* Siriboa:  
29. \*  
Siriboa. *Rumph. amb.* 5. p. 340. t. 117. f. 2.  
Betela quem Sirii-ba vocant. *Bont. jav.* 91. t. 91.  
*Habitat in India.*
5. **PIPER** foliis cordatis petiolatis sessilibusque. *Fl. zeyl.* longum.  
30. \* *Mat. med.* 20.  
Piper longum orientale. *Bauh. pin.* 412.  
Piper longum. *Rumph. amb.* 5. p. 333. t. 116. f. 2.  
Piper longum, pistolochiæ foliis absque pediculis,  
maderaspatanum. *Pluk. alm.* 297. t. 104. f. 4.  
Cattu-tirpali. *Rheed. mal.* 7. p. 27. t. 14.  
*Habitat in India.*
6. **PIPER** foliis lanceolato-ovatis quinquenerviis rugosis. Amalago.  
Piper longum arboreum altius, folio nervoso minore,  
spica gracili & brevior. *Sloan. hist.* 1. p. 134. t. 87. f. 1.  
Piper frutex, spica longa gracili. *Pluk. alm.* 297. t.  
215. f. 2.  
Piper longum. *Rumph. amb.* 5. p. 333. t. 116. f. 1.  
Saururus foliis lanceolato-ovatis quinquenerviis ru-  
gosis. *Hort. cliff.* 140.  
*Habitat in Jamaica.*
7. **PIPER** foliis cordatis novem-nerviis reticulatis. decumanum  
Saururus frutescens foliis plantagineis, fructu brevio-  
ri. *Plum. amer.* 59. t. 76.  
Sirum decumanum. *Rumph. amb.* 5. p. 45. t. 27.  
*Habitat in Indiis.* §  
*Affinis nimis P. reticulato, ut fere varietas.*
8. **PIPER** foliis cordatis septemnerviis reticulatis. reticulatum.  
Saururus botryoides major arborecens, foliis planta-  
gineis. *Plum. amer.* 57. t. 75.  
Jaborandi. *Marsgr. bras.* 37. *Pisf. bras.* 97.  
*Habitat in Martinica, Brasilia.*
9. **PIPER** foliis ovato-lanceolatis: nervis alternis, spi- aduacum.  
cis uncinatis.

## DIANDRIA TRIGYNIA.

*Piper longum*, folio nervoso pallide viridi, humiliss.

*Sloan. hist.* 1. p. 135. t. 87. f. 2.

*Saururus foliis ovato-lanceolatis* : nervis alternis.

*Hort. cliff.* 140. *Roy. lugdb.* 8.

*Saururus arborescens*, fructu adunco. *Plum. amer.*

58. t. 77.

*Habitat in Jamaica.*

pellucidum. 10. *PIPER* foliis cordatis petiolatis, caule herbaceo.

*Piper foliis cordatis*, caule procumbente. *Hort. cliff.*

6. t. 4. \* *Roy. lugdb.* 8.

*Saururus minor* procumbens botryitis, folio crasso cordato. *Plum. amer.* 54. t. 72.

*Habitat in America calidiore.* ☉

acuminatum. 11. *PIPER* foliis lanceolato-ovatis nervosis carnosiss.

*Saururus alius humilis*, folio carnosso & acuminato.

*Plum. amer.* 54. t. 71.

*Habitat in America calidiore.*

obtusifolium. 12. *PIPER* foliis obovatis enerviis.

*Saururus humilis*, folio carnosso subrotundo. *Plum. amer.* 53. t. 70.

*Habitat in America calidiore.*

rotundifolium. 13. *PIPER* foliis orbiculatis solitariis carnosiss.

*Saururus repens*, folio orbiculari nummulariæ facie.

*Plum. amer.* 52. t. 69.

*Piper longum* minimum herbaceum scandens rotundifolium. *Sloan. jam.* 45. *hist.* 1. p. 137.

*Habitat in America calidiore.*

maculosum. 14. *PIPER* foliis peltatis ovatis.

*Saururus hederaceus*, caulibus maculosis, major.

*Plum. amer.* 60. t. 66.

*Habitat in Domingo.*

peltatum. 15. *PIPER* foliis peltatis orbiculato-cordatis obtusis repandis, spicis umbellatis.

*Saururus arborescens*, foliis amplis rotundis & umbellatis. *Plum. amer.* 56. t. 74.

Lomba. *Rumph. amb.* 6. p. 133. t. 59. f. 1.

*Habitat in Indiis.*

distachyon. 16. *PIPER* foliis ovatis, spicis conjugatis.

*Saururus hederaceus*, caulibus maculosis, minor.

*Plum. amer.* 51. t. 67.

*Ha-*

*Habitat in Americæ Gallia æquinoctiali.*

17. PIPER foliis cordatis subrotundis acutis venosis, umbellatum  
spicis umbellatis.

*Saururus arborefcens*, foliis amplis cordatis non umbilicatis. *Plum. amer. 53. t. 73.*

*Habitat in Domingo.*

18. PIPER foliis ternis subrotundis.

trifolium.

*Saururus hederaceus triphyllus*, folio rotundo. *Plum. amer. 52. t. 68.*

*Habitat in Americæ Gallia æquinoctiali.*

19. PIPER foliis quaternis cuneiformibus sessilibus.

quadri-

*Saururus repens tetraphyllus*, folio crassiore. *Plum. ic. 218. t. 242. f. 2.*

lum.


*Habitat in America meridionali.*

20. PIPER foliis verticillatis ovatis trinerviis. *Amæn. acad. 5. p. 391. \**

verticilla-  
tum.

*Saururus erectus minor*, foliis orbiculatis verticillatis tumentibus, spicis terminalibus. *Brown. jam. 204.*

*Habitat in Jamaica. ☉*





## Classis III.

TRIANDRIA  
MONOGYNIA

## VALERIANA.

- rubra.* 1. VALERIANA floribus monandris caudatis, foliis lanceolatis integerrimis. *Hort. cliff.* 15. *Hort. upf.* 14. *Roy. lugdb.* 235. *Dalib. parif.* 12.  
Valeriana rubra. *Bauh. pin.* 165.  
Valeriana marina latifolia major rubra. *Morif. hist.* 3. p. 102. f. 7. t. 14. f. 15.  
2. Valeriana rubra angustifolia. *Bauh. pin.* 165. *prodr.* 88. *Bauh. hist.* 3. p. 211.  
*Habitat in Gallia, Helvetia, Italia, Orientis rud-ratis.* 24
- calcitrapa.* 2. VALERIANA floribus monandris, foliis pinnatifidis. *Hort. upf.* 14.  
Valeriana foliis pinnato-laciniatis, floribus monandris. *Virid. cliff.* 4. *Hort. cliff.* 16. *Roy. lugdb.* 235. *Sauv. monsp.* 275.  
Valeriana foliis calcitrapæ. *Bauh. pin.* 164. *Morif. hist.* 3. p. 101. f. 7. t. 14. f. 7.  
Valeriana sylvestris, foliis tenuissime divisis. *Bauh. pin.* 165.  
Valeriana annua f. æstiva. *Clus. hist.* 2. p. 54.  
*Habitat in Lusitania, Oriente.* ☉
- cornucopia.* 3. VALERIANA floribus diandris ringentibus, foliis ovatis sessilibus. *Hort. cliff.* 15. *Hort. upf.* 13. *Roy. lugdb.* 235.  
Valeriana peregrina purpurea albave. *Bauh. pin.* 164. *prodr.* 87.  
Valeriana indica. *Clus. hist.* 2. p. 54.  
Pseudo-Valeriana cornucoplioides annua purpurea. *Morif. hist.* 3. p. 104. f. 7. t. 16. f. 27.  
*Habitat in America, Mauritania, Sicilia, Hispania arvis.* ☉
- dioica.* 4. VALERIANA floribus triandris dioicis, foliis pinnatis integerrimis.

Vale-

- Valeriana dioica*. *It. æl.* 46. *Fl. fœc.* 31. 35. *Dalib. parif.* 11.  
*Valeriana*, foliis caulinis pinnatis, sexu distincta. *Hort. cliff.* 16.  
*Valeriana*, foliis caulinis pinnatis, polygama. *Vir. cliff.* 5. *Roy. lugdb.* 235.  
*Valeriana palustris minor*. *Bauh. pin.* 164.  
*Valeriana palustris inodora parum laciniata*. *Bauh. pin.* 86.  
*Valeriana alpina minor*. *Bauh. pin.* 165. *prodr.* 87.  
*Valeriana pratensis minor*. *Morif. umb. t.* 10. d. e.  
*Habitat in Europæ & Orientis campis uliginosis.* 24
5. *VALERIANA* floribus triandris, foliis omnibus pinnatis. *Mar. med.* 21. *Hort. cliff.* 15. *Fl. fœc.* 30; 34. *Roy. lugdb.* 234. *Dalib. parif.* 11.  
*Valeriana palustris major*. *Bauh. pin.* 164. *prodr.* 86.  
*Valeriana sylvestris major*. *Bauh. pin.* 164. *Fl. lapp.* 13.  
*Valeriana sylvestris*. *Dod. pempt.* 349.  
*Habitat in Europæ nemoribus paludosis.* 24
6. *VALERIANA* floribus triandris, foliis caulinis pinnatis; radicalibus indivisis. *Hort. upf.* 13. *Mar. med.* 22. *Phu.*  
*Valeriana* foliis infimis integris; proximis laciniatis; caulinis pinnatis. *Hort. cliff.* 15. *Roy. lugdb.* 234.  
*Valeriana hortensis*. *Bauh. pin.* 164.  
*Habitat in Alsatia.* 24  
*Caule non sulcato & foliolis exterioribus majoribus a V. officinali etiam differt.*
7. *VALERIANA* floribus triandris, foliis dentatis: radicalibus cordatis; caulibus ternatis ovato-oblongis. *tripteris.*  
*Valeriana* floribus triandris, foliis radicalibus cordatis, aliis laciniatis. *Sauv. monsp.* 276.  
*Valeriana alpina prima*. *Bauh. pin.* 165. *prodr.* 86. t. 86.  
*Valeriana alpina altera*. *Bauh. pin.* 165.  
*Valeriana alpina minor, planta palmaris*. *Pluk. alm.* 380. t. 231. f. 7.  
*Valeriana minima, planta uncialis*. *Pluk. alm.* 380. t. 231. f. 8.  
*Habitat in alpinis Helvetiæ.*  
*Involutra propria sunt diphylla, setacea.* 24
8. *VALERIANA* floribus triandris, foliis ovato-oblongis subdentatis, caule simplici. *montana.*

- Valeriana foliis omnibus integris ex ovato acuminatis leviter dentatis. *Hall. helv.* 664. *it.* 8.
- Valeriana montana, subrotundo folio. *Bauh. pin.* 165.
- Valeriana alpina, scrophulariæ folio. *Bauh. pin.* 164. *prodr.* 87. \*
- Nardo celticæ similis inodora. *Bauh. pin.* 165.
- Habitat in alpihus Helveticis, Rhæticiis, Pyrenæis.* 24.
- celtica.** 9. VALERIANA floribus triandris, foliis ovato-oblongis obtusis integerrimis. *Mat. med.* 23.
- Valeriana foliis ovatis obtusis minime dentatis. *Hall. helv.* 664.
- Nardus ex apulia. *Bauh. pin.* 165.
- Nardus celtia dioscoridis. *Bauh. pin.* 165.
- Nardus celtica altera. *Bauh. pin.* 166.
- Nardus celtica. *Bauh. hist.* 3. p. 205.
- Spica celtica fastigio flosculorum ordine differens. *Cam. epit.* 14.
- Habitat in alpihus Helvetiæ, Valesiæ.* 24.
- tuberosa.** 10. VALERIANA floribus triandris, foliis lanceolatis integerrimis: caulinis basi pinnatis.
- Nardus montana, radice olivari. *Bauh. pin.* 165.
- β. Nardus montana, radice oblonga. *Bauh. pin.* 165.
- Nardus montana, longius radicata. *Cam. epit.* 16.
- Habitat in Dalmatia, Sicilia, Galloprovincia.* 24.
- saxatilis.** 11. VALERIANA floribus triandris, foliis subdentatis: radicalibus ovatis, caulinis lineari-lanceolatis.
- Valeriana alpina nardo celticæ similis. *Bauh. pin.* 165.
- Valeriana sylvestris alpina 2. saxatilis. *Cusf. hist.* 1. p. 56. *bona.*
- Valeriana alpina nardo celticæ similis inodora. *Pluk. alm.* 380. t. 232. f. 2.
- Nardo Celticæ similis inodora. *Bauh. pin.* 165. *prodr.* 88. *Burf. VIII.* 122.
- Habitat in alpihus Siniæ, Austriæ, Baldi, Montalbani Italiæ.* 24.
- pyrenaica.** 12. VALERIANA floribus triandris, foliis caulinis cordatis serratis petiolatis: summis ternatis. *Hort. cliff.* 15. *Roy. lugdb.* 235. *Monnier. obs.* 235.
- Valeriana maxima pyrenaica, cacaliæ folio. *Tournef. inst.* 131.
- Valeriana canadensis. *Riv. mon.* 6.

*Valeriana orientalis*, alliariæ folio, flore albo. *Buxb.*

*cent. 2. p. 19. t. 11.*

*Habitat in Pyrenæis? 2.*

13. *VALERIANA* floribus triandris, foliis ternatis, scandens. caule scandente. *Læfl. it. 235. †*

*Habitat in Cumana.*

14. *VALERIANA* floribus triandris, foliis omnibus *Chinensis*. cordatis repando-lobatis.

*Habitat in China. Osbeck.*

*Caules herbacei, erecti, plus quam pedales, ramis floriferis oppositis; Folia opposita, petiolata, glabra, cordata, acuta, lobis lateralibus utrinque binis obtusis. Rami & caules terminati umbellulis involucello cinctis. Semina nuda.*

15. *VALERIANA* floribus triandris regularibus, foliis *echinatis*. dentatis, fructibus linearibus tridentatis: extimo majore recurvo.

*Valeriana foliis dentato-ferratis, pedunculis conicis, seminibus tridentatis. Sauv. monsp. 130.*

*Valerianella echinata. Bauh. pin. 165.*

*Valerianella cornucopioides echinata. Colum. ecphr.*

*1. p. t. 204. t. 206.*

*Habitat in Italiæ & Monspelli umbrosis. Sauvages. ①*

*Caulis dichotomus. Folia lanceolata, sessilia, obtusa, dentata. Flores & dichotomia, solitarii, sessiles. Fructus linearis, apice tridentatus: dente exteriori majore recurvo. Corolla infundibuliformis, regularis, quinque-fida, alba. Stigma trifidum. Summi flores in Spicas dichotomas, coadunatas, obtusas, bracteis & pedunculis incrassatas digesti; tamquam ex *V. Locusta* matre & *V. cornucopioide* patre prognata esset.*

16. *VALERIANA* floribus triandris, caule dichotomo, foliis linearibus. *Fl. svec. 32; 36. Hort. upf.*

*14. Dalsb. parif. 12.*

*Valeriana caule dichotomo, foliis lanceolatis integris. Vir. cliff. 5.*

1. *Valeriana caule dichotomo, foliis lanceolatis integris, olitoria. fructu simplici. Hort. cliff. 16. Roy. lugdb. 235.*

*Valeriana campestris inodora major. Bauh. pin. 165.*

2. *Valeriana caule dichotomo, foliis lanceolatis ferratis, vesicaria. calycibus inflatis. Hort. cliff. 16. Roy. lugdb. 235.*

*V a-*

- Valeriana cretica*, fructu vesicario. *Tournef. cor.* 6.  
*Boerh. lugdb.* 1. p. 75. t. 75.
- coronata.* γ. *Valeriana* caule dichotomo, foliis lanceolatis dentatis, fructu sexdentato. *Hort. cliff.* 16. *Roy. lugdb.* 235.  
*Valeriana* semine stellato. *Bauh. pin.* 165.  
*Valerianella* altera tenuifolia semine scabiosæ stellato.  
*Column. ecphr.* 1. p. 207. t. 209.
- dentata.* δ. *Valeriana* caule dichotomo, foliis oblongis subserratis, feminis corona tridentata. *Hall. helv.* 665. *Dalib. parif.* 12.  
*Locusta major.* *Riv. mon.* 6.
- radiata.* ε. *Valeriana* caule dichotomo, capitulis terminalibus involucri cinctis. *Gron. virg.* 10.  
*Valeriana marilandica*, foliis oblongis obtusis. *Raj. suppl.* 3. p. 244.
- discoides.* ζ. *Pseudo-Valeriana* annua, semine coronato, major lusitanica. *Morif. hist.* 3. p. 104. f. 7. t. 16. f. 29.  
*Habitat* α. in Europæ, β. Cretæ, γ. Lusitaniz, δ. Europæ australioris, ε. Marilandiz arvis. ☉
- mixta.* η. VALERIANA floribus triandris, caule quadrifido, foliis imis bipinnatifidis, feminis pappo plumoso. *Sauv. monsp.* 275. †  
*Valerianella* semine umbilicato hirsuto minore. *Morif. umb.*  
*Habitat* Monspelii.
- sibirica.* θ. VALERIANA floribus tetrandris æqualibus, foliis pinnatifidis, seminibus paleæ ovali adnatis. *Hort. upf.* 13.  
*Valeriana lutea humilis.* *Amm. ruth.* 18. n. 25. t. 3.  
β. *Anonymus cheusanensis*, sideritidis folio, fructu ulmi samaris haud absumili. *Pluk. alm.* 16. t. 443. f. 1.  
*Habitat* in Sibiriz campis. ☉

## O L A X.

- zeylanica.* ι. OLAX. *Fl. zeyl.* 34. \* *Amæn. acad.* 1. p. 387.  
*Arbor stercoraria zeylanica glandifera.* *Burm. zeyl.* 26.  
*Habitat* in Zeylona. ‡

## T A M A R I N D U S.

- indica.* κ. TAMARINDUS. *Hort. cliff.* 18. *Mat. med.* 28.  
*Hort. upf.* 15. *Fl. zeyl.* 14. *Roy. lugdb.* 465. *Kumph. amb.* 2. p. 90. t. 23. *Lxfl. it.* 210.  
*Siliqua arabica*, quæ Tamarindus. *Bauh. pin.* 403.  
*Balam-*

Balam-puli. *Rheed. mal.* 1. p. 39. t. 23.

*Habitat in India, America, Aegypto, Arabia.* ♀

RUMPHIA.

1. RUMPHIA.

*Myxa pyriformis*, officulo trispermo. *Raj. hist.* 1556.

*Thiem tani. Rheed. mal.* 4. p. 25. t. 11.

*Habitat in India.* ♀

amboinensis

CNEORUM.

1. CNEORUM. *Hort. cliff.* 18. *Roy. lugdb.* 119.

*Chamælea triccocos. Bauh. pin.* 462.

*Chamælea. Cam. epit.* 973.

*Habitat in Hispaniæ, Narbonæ glareosis.* ♀

triccocos

CAMOCLADIA.

1. CAMOCLADIA foliolis integris. *Jacq. amer.* 12. integrifolia.

*Camocladia caudice simplici, floribus confertis sessilibus, racemis alaribus. Brow. jam.* 124.

*Prunus racemosa*, caudice non ramoso, alato fraxini folio non crenato, fructu rubro subdulci. *Sloan. jam.*

184. *hist.* 2. p. 131. t. 222. f. 1. *Raj. dendr.* 44.

*Habitat in America calidiore.* ♀

2. CAMOCLADIA foliolis spinoso - dentatis. *Jacq. amer.* 12. dentata.

*Habitat in America meridionali.* ♀

MELOTHRIA.

1. MELOTHRIA. *Hort. cliff.* 490. \* *Hort. ups.* 15. pendula.

*Gron. virg.* 10. *Roy. lugdb.* 528.

*Cucumis parva repens virginiana, fructu minimo.*

*Pluk. alm.* 123. t. 85. f. 5.

*Cucumis minima, fructu ovali nigro lævi. Sloan. hist.*

1. p. 227. t. 142. f. 1.

*Bryonia, olivæ fructu, minor. Plum. spec.* 3. ic. 66.

f. 2.

*Habitat in Canada, Virginia, Jamaica.* ☉

*In meis plantis horti nunquam flores masculi.*

ORTEGIA.

1. ORTEGIA. *Læfl. it.* 112. \*

*Rubia linifolia aspera. Bauh. pin.* 333. *Raj. hist.* 1033.

*Juncaria salmantica. Clus. hist.* 1. p. 174.

*Habitat in Castilia, Bætica, Salmantica.*

*Linnei spec. plant.* P. 1.

D

1. LOEF-

hispanica.

## LOEFLINGIA.

hispanica.

1. LOEFLINGIA. *Loefl. it. 113. t. 1. f. 2. \***Habitat in Hispaniæ collibus apricis. ☉**Caules profusi Schleranthi f. Herniariæ, viscid. Folia subulata, brevissima, utrinque dente notata. Spicæ breves, squarrosæ, axillares: Bracteis imbricatis, subulatis, dentatis.*

## POLYCNEMUM.

arvense.

1. POLYCNEMUM. *Sauv. monsp. 45.**Salsola caule repente, foliis & stipulis sessilibus subulatis cartilagineis. Dalib. parif. 80.**Chenopodium foliis subulatis prismaticis, floribus solitariis sessilibus axillaribus. Gvest. stamp. 14.**Chenopodium annuum humifusum, folio breviori & capillaceo. Tournef. inst. 506.**Camphorata vaginis spinosis. Hall. helv. 183.**Camphorata congener. Bauh. pin. 486.**Herniaria foliis longis angustis acuminatis & glabris.**Magn. char. 51.**Habitat in Galliæ, Italiæ, Germaniæ arvis. ☉*

## HIPPOCRATEA.

velutilla.

1. HIPPOCRATEA. *Hort. cliff. 484. Jacq. amer. 12.**Coa scandens, fructu trigemino subrotundo. Plum. gen. 8. ic. 88.**Bejuco pendulus, floribus paniculatis. Loefl. it. 314.**Habitat in America australi. ☿*

## CROCUS.

sativus.

## 1. CROCUS spatha univalvi radicali, corollæ tubo longissimo.

*Crocus floribus fructui impositis: tubo longissimo. Roy. lugdb. 41. Hort. upf. 15. Mat. med. 27.**Crocus flore fructui imposito. Hort. cliff. 18.*

officinalis.]

a. Crocus autumnalis sativus. *Morif. hist. 2. p. 335. f. 4. t. 2. f. 1.**Crocus sativus. Bauh. pin. 65.*

vernus.

β. Crocus vernus latifolius. I-XI. & I-VI. *Bauh. pin. 65. 66.**Habitat in Alpibus Helveticis, Pyrenæis, Lusitanicis, Thracicis. 24*

IXIA.

1. IXIA scapo unifloro brevissimo, foliis linearibus. Bulbocodi-  
um.  
Crocus spatha diphylla flori approximata, tubo corollæ brevi. *Sp. pl.* 1. p. 36.  
Crocus flore fructui imposito: tubo brevi. *Roy. lugdb.* 41.  
Sisyrinchium minus angustifolium, flore majore variegato. *Bauh. pin.* 41.  
Sisyrinchium asprensium, angusto folio, alterum. *Col. ecphr.* 2. p. 5. t. 7. f. 1.  
Bulbocodium crocifolium, flore magno albo fundo luteo. *Tournef. cor.* 50.  
*Habitat in alpidibus Italicis.* 24
2. IXIA floribus capitatis, spathis laceris. africana.  
*Ixia. Hort. cliff.* 490: \* *Roy. lugdb.* 20.  
*Ixia* foliis ad radicem nervosis gramineis, floribus ac fructu convolutis. *Burm. afric.* 191. t. 70. f. 2.  
*Bermudiana capensis*, capitulis lanuginosis. *Pet. succ.* 242.  
*Gramen eriophorum africanum*, flore lanato. *Pluk. mant.* 98.  
*Habitat ad Cap. b. Spei.* 24
3. IXIA floribus corymbosis pedunculatis, caule ancipiti. corymbosa.  
*Amzn. acad.* 4. p. 300. \*  
*Habitat ad Cap. b. Spei.*  
Flores cærulei, corymbosi, calyce bivalvi instructi.  
*Planta palmaris.*
4. IXIA foliis linearibus, axillis bulbiferis, floribus alternis, staminibus lateralibus. bulbifera.  
*Amzn. acad.* 4. p. 300.  
*Habitat ad Cap. b. Spei.* 24  
Flores e flavescente albi, sæpius duo. Bulbus in alis foliorum.
5. IXIA foliis linearibus, racemo flexuoso multifloro. flexuosa.  
*Ixia* foliis linearibus, floribus spicatis sessilibus. *Mill. ic.* 156. f. 2 ?  
*Habitat ad Cap. b. Spei.* 24  
Flores albi, fundo flavo, extus subviolacei, plurimi.  
Folia levia. Scapus pedalis, teres, a flore ad florem parum flexus, sæpe etiam ramum emittens.
6. IXIA foliis linearibus, scapo spicis pluribus. polystachya.  
*Ixia* foliis lineari-lanceolatis, floribus alaribus & terminalibus. *Mill. ic.* 155. f. 2.



*Habitat ad Cap. b. spei. 24*

Flores albi filamentis ♂ pistillis violaceis. Stylum inter tripartitum stamina eriguntur, quasi Styli exteriores essent. Folia angusta, lævia. Spicæ multifloræ, præter terminalem, etiam nonnullæ inferius, alternæ.

scillaris.

7. IXIA foliis ensiformibus striatis, spica elongata.

*Habitat ad Cap. b. spei. 24*

Flores violacei ut in *Scilla amœna*, antheris flavis, alterni, remoti.

crocata.

8. IXIA foliis ensiformibus, floribus alternis, petalis basi hyalina macula.

*Ixia* foliis gladiatis glabris, floribus corymbosis terminalibus. *Mill. ic. 156. f. 1.*

*Sisyrinchium majus* afric. flore luteo macula notato. *Old. afr. 32.*

*Habitat ad Cap. b. spei. 24*

Flores crocei in spica secunda rariore. Petala obovata basi hyalina, ♂ ad ungues macula oblonga flava.

chinensis.

9. IXIA foliis ensiformibus, floribus remotis, panícula dichotoma, floribus pedunculatis. *Hort. upf. 16.*

*Bermudiana* radice carnosa, floribus maculatis, seminibus pulpa obductis. *Amm. act. petrop. XI. p. 308. t. 7.*

*Bermudiana* iridis folio majori, flore croceo eleganter punctato. *Kraus. hort. 25. t. 25.*

*Balemcanda - Schularmandi. Rheed. mal. 11. p. 73. t. 37.*

*Habitat in India. 24*

Flores flavi maculis sanguineis. Stylus inclinatus.

## GLADIOLUS.

- communis. 1. GLADIOLUS foliis ensiformibus, floribus distantibus.

*Gladiolus* foliis ensiformibus. *Hort. cliff. 20. Hort. upf. 16.*

*Gladiolus* caule simplicissimo, foliis ensiformibus. *Roy. lugdb. 19.*

*Gladiolus. Riv. mon. 163.*

*Habitat in Europa australi. 24*

- imbricatus. 2. GLADIOLUS foliis ensiformibus, floribus imbricatis.

*Habitat in Russia citeriore. 24*

Flores parvi versus unum latus imbricati.

3. GLA-

3. **GLADIOLUS** foliis linearibus, caule simplicissimo, spicatus.  
 floribus spicatis. *Roy. lugdb.* 19.  
*Habitat in Africa.* 24
4. **GLADIOLUS** foliis ensiformibus, petalis lateralibus alatus.  
 latissimis. *Amæn. acad.* 6. *afric.* 2. \*  
*Sisyrinchium viperarum. Pluk. phyt.* 224. f. 8.  
*Habitat ad Cap. b. spei.* 24
5. **GLADIOLUS** foliis ensiformibus plicatis villosis, plicatus.  
 scapo laterali, corollis regularibus.  
*Ixia plicata. Amæn. acad.* 4. p. 300. \* *absque synonymo.*  
*Mill. ic.* 155. f. 1.  
*Sisyrinchium latifolium, floribus patentibus vix diffor-*  
*mis. Breyn. prod.* 3. p. 22. t. 9. f. 2.  
*Habitat in Æthiopia.* 24  
*Bulbus tunicatus. Folia ensiformia, villosa, secundum*  
*nervos plicata. Scapus lateralis floribus 2. f. 3. viola-*  
*ceo-albis regularibus. Bractæ ternæ, angustæ: exte-*  
*riore majore.*
6. **GLADIOLUS** foliis lineari-cruciatis, corollis cam- tristis.  
 panulatis.  
*Gladiolus bifolius & biflorus, foliis quadrangulis. Trew.*  
*chret.* t. 39.  
*Habitat in Æthiopia.* 24  
*Folia singularia, subensiformia, sed utroque latere cari-*  
*nata carina elevata, obtusissima, ut fere quadrangulo-*  
*cruciata; variat a caule bifloro ad multiflorum & tum*  
*foliis rite ensiformibus. Corolla colore tristi, campanu-*  
*lata, sed æstivatione omnino Gladioli.*
7. **GLADIOLUS** foliis linearibus, floribus distantibus, angustus.  
 corollarum tubo limbis longiore. *Mill. ic.* 142. f. 2.  
*Gladiolus caule simplicissimo, foliis linearibus, flori-*  
*bus alternis. Roy. lugdb.* 19.  
*Gladiolus foliis linearibus. Hort. cliff.* 20. \* t. 6.  
*Habitat in Africa.* 24
8. **GLADIOLUS** caule ramoso, foliis linearibus. *Roy.* ramosus.  
*lugdb.* 19.  
*Habitat in Africa.* 24
9. **GLADIOLUS** caule ramoso, capitulis pedunculatis, capitatus.  
 radice tuberosa.  
*Habitat in Africa.* 24 \*  
*Planta est maxima; floribus caeruleis.*

- alopecuroides. 10. **GLADIOLUS** foliis linearibus, spica disticha imbricata. *Amén. acad.* 4. 301. \*  
*Habitat in Æthiopia.*  
*Spica ut in Plantagine oltecta floribus parvis numerosissimis, sed disticha.*

## ANTHOLYZA.

- ringens. 1. **ANTHOLYZA** corollæ labiis divaricatis faucē compressa.  
*Antholyza* stamine unico declinato. *Sp. pl.* 1. p. 37.  
*Gladiolo æthiopico similis planta angustifolia. Comm. hort.* 1. t. 41. *Rudb. elyf.* 2. p. 237. f. 15.  
*Gladiolus floridus rectum referens coccineus: suprema lacinia erecta & fistulosa. Breyn. ic.* 21. t. 8. f. 1.  
*Habitat in Æthiopia.* 24.
- cunonia. 2. **ANTHOLYZA** corollis rectis: labii quinquepartiti lobis duobus extimis latioribus adscendentibus. *Mill. ic.* 113.  
*Cunonia floribus sessilibus, spathis maximis. Buttn. cun.* 211. t. 1.  
*Habitat in Persia.* 24.  
*Corollæ Labium superius longissimum, obtusum, convolutum; inferius 5-partitum; Laciniis 3. intermediis minutissimis; lateralibus vero latis, labio superiore dimidio brevioribus. Spica longa, disticha.*
- æthiopica. 3. **ANTHOLYZA** corollis incurvatis; labii quinquepartiti lobis duobus alternis patulis majoribus lanceolatis.  
*Gladiolus æthiopicus, flore coccineo. Corn. canad.* 79. t. 79. *Morif. hist.* 2. p. 421. f. 4. t. 23. f. 1. *Rudb. elyf.* 2. p. 236. f. 14.  
*Hyacinthus africanus, foliis colchici floribus coccineis. Pluk. phyt.* 195. f. 2.  
*Habitat in Æthiopia.* 24.  
*Corolla tubulosa, coccinea; Labio superiore maximo, lanceolato; inferiore 5-partito: lobis lanceolatis brevibus; tribus alternis brevissimis. Hæc cum antecedente Gladiolis valde affinis.*
- Meriana. 4. **ANTHOLYZA** floribus infundibuliformibus subæqualibus.  
*Meriana flore rubello. Trew. chret.* t. 40.  
*Habitat ad Cap. b. spei?* 24.

Flores rubri : corolla *Polyanthis* ; *supera* , *aestivazione*  
*Gladioli*. Folia *ensiformia*.

I R I S.

\* *Barbata nectariis petalorum reflexorum*

1. IRIS corolla barbata, caulis foliis longiore unifloro. *fusiana*.

*Hort. cliff.* 18. *Roy. lugdb.* 17.

*Iris fusiana*, flore maximo ex albo nigricante. *Bauh.*  
*pin.* 31. *theatr.* 579. *Morif. hist.* 2. p. 351. f. 4. t. 6.  
 f. 6.

*Iris latifolia major.* *Clus. hist.* 1. p. 217.

*Habitat* in Oriente ; venit *Constantinopoli* in *Belgium*  
 1573. 24

*Caulis teres* ; *Germen trigono - teretiusculum* ; *Petala* 8.  
*interiora majora reflexa.*

2. IRIS corollis barbatis, caule foliis altiore subbifloro, *florentina*.  
*floribus sessilibus.* *Mill. ic.* 154.

*Iris alba florentina.* *Bauh. pin.* 31.

- 3 *Iris alba germanica.* *Bauh. pin.* 31.

*Iris flore albo.* *Raj. hist.* 1180.

*Habitat* in *Europa australi* - *Carniola.* 24

*Simillima* *I. Germanicæ* ut vix differat, sed *Corolla al-*  
*ba petalis integris : inferioribus marginum basi reflex-*  
*is ; superioribus magis erectis. Stigmata magis ere-*  
*cta & parum serrata.*

3. IRIS corollis barbatis, caule foliis longiore multiflo- *germanica.*  
 ro, floribus inferioribus pedunculatis. *Hort. cliff.* 18.

*Hort. upf.* 16. *Mat. med.* 24. *Roy. lugd.* 17.

*Iris vulgaris germanica* f. *sylvestris.* *Bauh. pin.* 30.

*Habitat* in *Germaniæ editis.* 24

*Petala inferiora plana ; interiora integerrima , nec emar-*  
*ginata.*

4. IRIS corollis barbatis, caule foliis altiore multifloro, *sambucina*.  
*petalis deflexis planis : erectis emarginatis.*

*Iris latifolia germanica*, *sambuci* odore. *Bauh. pin.* 31.

*Iris major latifolia* VIII. *Clus. hist.* 1. p. 219.

*Habitat* in *Europa australi.* 24

*Simillima* *I. Germanicæ* , sed *deflexa petala saturatius*  
*violacea , plana quidem at subemarginata. Erecta*  
*pallide , saturatius tamen , cærulea & emarginata.*  
*Stigmata serrata , acutiuscula , carina cærulescente.*

- squalens.* 5. *IRIS* corollis barbatis, caule foliis altiore multifloro, petalis deflexis replicatis; erectis emarginatis.  
*Iris* folio lato rugoso, petalis repandis ex purpureo fordido pallido & luteo variis, erectis vero squalide lutescentibus. *Boerh. lugdb.* 2. p. 125.  
*Habitat in Europa australi.* 24  
*Simillima* *I. germanicæ*, sed *Petala deflexa in medio replicata*, saturatius violacea venis albis flavescentibus  
 & in medio cærulescentibus. *Erecta petala*, uti &  
*Stigmata squalide flava*, valde emarginata.
- aphylla.* 6. *IRIS* corollis barbatis, scapo nudo longitudine foliorum multifloro. *Roy. lugdb.* 17.  
*Iris latifolia*, caule aphylo. *Bauh. pin.* 32.  
*Habitat in* - - - 24
- variegata.* 7. *IRIS* corollis barbatis, caule subfolioso longitudine foliorum multifloro. *Roy. lugdb.* 17. *Hort. upf.* 16.  
*Iris* corollis barbatis, foliis altitudine caulis multiflori.  
*Hort. cliff.* 19.  
*Iris latifolia pannonica*, colore multiplici. *Bauh. pin.* 31.  
*Iris lutea variegata.* *Lob. hist.* 34 *Ehret. pict. t.* 10. f. 3.  
*Habitat in Hungaria.* 24.
- biflora.* 8. *IRIS* corollis barbatis, caule foliis breviori trifloro.  
*Hort. upf.* 17.  
*Iris* corollis barbatis, foliis caulem multiflorum superantibus. *Hort. cliff.* 19. *Roy. lugdb.* 17.  
*Iris latifolia biflora.* *Besl. cyst. vern.* 114  
*Chamæiris major* saturate purpurea biflora. *Bauh. pin.* 33.  
*Habitat in Lusitanix rupibus.* 24
- pumila.* 9. *IRIS* corollis barbatis, caule foliis breviori unifloro.  
*Iris* corollis barbatis, foliis caulem uniflorum superantibus. *Hort. cliff.* 19. *Roy. lugdb.* 17.  
*Chamæiris minor*, flore purpureo. *Bauh. pin.* 33.  
*Chamæiris latifolia minor.* 1. 2. *Clus. hist.* 1. p. 225.  
 β. *Chamæiris minor*, flore purpureo-cæruleo. *Bauh. pin.* 33.  
 γ. *Chamæiris minor*, flore rubello. *Bauh. pin.* 33.  
 δ. *Chamæiris* flore pallido & albo. *Bauh. pin.* 33.  
 ε. *Chamæiris minor*, flore variegato. *Bauh. pin.* 34.  
*Habitat in Austriæ, Pannoniæ collibus apricis.* 24

\* \* *Imberbes* : petalis deflexis lævibus.

*Pseud Aco.* 10. *IRIS* corollis imberbibus : petalis interioribus stigmatem

- minoribus, foliis ensiformibus. *Hort. cliff.* 19. *Fl. succ.* 33. 37. *Mat. med.* 25.  
*Acorus adulterinus. Bauh. pin.* 34. *theatr.* 634.  
*Acorum falsum. Cam. epit.* 6.  
*Habitat in Europa ad ripas paludum, fossarum.* 24  
*Petala exteriora utrinque ad filamenta dentem exserunt. Germen trigonum angulis sulco bifidis.*  
11. *IRIS* corollis imberbibus, petalis interioribus patentissimis, caule uniangulato, foliis ensiformibus. *Hort. cliff.* 19. *Roy. lugdb.* 18. *Dalib. parif.* 13. *Sauv. monsp.* 41. *foetidissima.*  
*Gladiolus foetidus. Bauh. pin.* 30.  
*Spathula foetida, Xyris. Bauh. hist.* 2. p. 731. *Dod. pempt.* 247.  
*Habitat in Gallia, Anglia, Hetruria.* 24  
*Caulis teres, hinc angulatus, longitudine foliorum, obiectus Foliis foetidissimis. Germen trigonum angulis sulco bifidis. Corolla carulescenti-tristis ingratissimi coloris, nec noctu odora. Petala exteriora aenae subtus plicis rugoso; Petala interiora stigmatem majora, patentia.*  
12. *IRIS* corollis imberbibus, germinibus trigonis, caule tereti, foliis linearibus. *Hort. cliff.* 19. *Hort. upf.* 17. *Roy. lugdb.* 78. *Hall. helv.* 280. *Gmel. sibir.* 1. p. 28. *Iris pratensis angustifolia non foetida altior. Bauh. pin.* 32. *theatr.* 597. *fibrica.*  
*Iris angustifolia 2. Clus. hist.* 1. p. 227.  
*Habitat in Austriae, Helvetiae, Sibiriae pratis.* 24  
*Spathae in I. sibirica & versicolore aridae s. scariosae sunt. Huic caulis foliis altior; petala interiora erecta. Germen trigonum angulis non sulcatum.*  
13. *IRIS* corollis imberbibus, germinibus subtrigonis, caule tereti flexuoso, foliis ensiformibus. *versicolor.*  
*Iris americana versicolor, stylo crenato. Dill. elkh.* 188. t. 155. f. 188.  
*Iris americana versicolor, stylo non crenato. Dill. elkh.* 187. t. 155. f. 187.  
*Iris latifolia virginiana, florum petalis repandis purpureis. Ehret. pict.* t. 6. f. 2.  
*Habitat in Virginia, Marilandia, Pennsylvania.* 24  
*Caulis foliis non altior. Folia ad alterum latus inclinantia. Petala interiora longitudine stigmatum. Stigmata linearia.*

- virginica.* 14. IRIS corollis imberbibus, germinibus trigonis, caule ancipiti. *Gron. virg.* 11.  
*Habitat in Virginia.* 24
- martinicens.* 15. IRIS corollis imberbibus, germinibus trigonis, petalis basi foveolis glandulosis. *Jacqu. amer.* 12.  
Xiphion flore e luteo nigricante. *Plum. spec.* 8. ic. 260. t. 261. f. 1.  
*Habitat in Martinica.* 24
- spuria.* 16. IRIS corollis imberbibus, germinibus sexangularibus, caule tereti, foliis sublinearibus. *Hort. cliff.* 19.  
*Hort. upf.* 17. *Roy. lugdb.* 16.  
Iris pratenfis angustifolia, folio foetido. *Bauh. pin.* 32.  
Iris angustifolia 1. *Cus. hist.* 1. p. 228.  
*Habitat in Germaniæ pratis.* 24
- graminea.* 17. IRIS corollis imberbibus, germinibus sexangularibus, caule ancipiti, foliis linearibus. *Hort. cliff.* 19.  
*Hort. upf.* 17. *Roy. lugdb.* 18  
Iris angustifolia prunum redolens minor. *Bauh. pin.* 33.  
Iris angustifolia 3. *Cus. hist.* 1. p. 230.  
*Habitat in Austria ad radices montium.* 24
- verna.* 18. IRIS corollis imberbibus, caule unifloro foliis brevioribus, radice fibrosa. *Gron. virg.* 10.  
Iris virginiana pumila s. Chamæiris verna angustifolia, flore purpureo - cæruleo odorato. *Pluk. alm.* 198 t. 196. f. 6.  
*Habitat in Virginia.* 24
- tuberosa.* 19. IRIS corollis imberbibus, foliis tetragonis *Vir. cliff.* 6. *Hort. cliff.* 20. *Mat. med.* 25. *Roy. lugdb.* 18.  
Iris tuberosa, folio anguloso. *Bauh. pin.* 40. *Morif. hist.* 2. p. 348. f. 4. t. 5. f. 1.  
Hermodactylus folio quadrangulo. *Tournef. cor.* 50.  
*Habitat in Arabia & Oriente.* 24
- Xiphium.* 20. IRIS corollis imberbibus, floribus binis, foliis subulato-caniculatis caule brevioribus. *Hort. upf.* 17.  
Iris foliis margine conniventibus, corollis imberbibus. *Hort. cliff.* 20.  
Iris caulescens, foliis margine conniventibus, corollis imberbibus. *Roy. lugdb.* 18.  
Iris bulbosa latifolia, caule donata. *Bauh. pin.* 38.  
β. Iris bulbosa cæruleo-violacea. *Bauh. pin.* 40.  
*Habitat in Hispania.* 24

21. IRIS corolla imberbi: petalis interioribus patentissimis ferratis, foliis subulato-canalicultatis caule longioribus.

perica.

Iris acaulis, foliis margine conniventibus, corollis imberbibus. *Roy. lugdb.* 18.

Iris perica martia. *Rudb. chys.* 2. p. 10. f. 9.

Iris perica bulbosa variegata præcox. *Dodart. mem.* 83.

Xiphion perficum præcox, flore variegato. *Tournef. inst.* 363.

*Habitat in Persia.* 2.

*Bulbus exserit Gemmam e duobus paribus squamarum, intra quæ folia & flos unicus spatha simplici. Petala interiora brevissima.*

22. IRIS corollis imberbibus, foliis canaliculatis, bulbis geminis superimpositis.

sifyrinchium.

Sifyrinchium majus. *Bauh. pin.* 40. *Clus. hist.* 1. p. 216.

β. Sifyrinchium medium. *Bauh. pin.* 41.

Sifyrinchium minus. *Clus. hist.* 1. p. 216.

*Habitat in Hispania, Lusitania.* 2.

Colore est petalorum purpureo macula lutea loco barbae; bulbis (non tunicatis simplicibus ut in bulbosis Xiphiis 20. 21., sed) ex gemino bulbo altero alteri imposito, ut in Gladiolo, Croco.

## MORÆA.

1. MORÆA foliis canaliculatis. *Mill. dict.*

vegeta.

*Habitat in Africa.* 2.

2. MORÆA foliis subulatis. *Mill. dict.*

junceæ.

*Habitat in Africa.* 2.

## WACHENDORFIA.

1. WACHENDORFIA scapo simplici.

thyrsiflora.

Wachendorfia foliis lanceolatis quinquenerviis canaliculo-plicatis, floribus in thyrsum collectis. *Burm. monogr.* 2. f. 2.

*Habitat ad Cap. b. spei.* 2.

2. WACHENDORFIA scapo polystachyo.

paniculata.

Wachendorfia foliis ensiformibus trinerviis, floribus paniculatis. *Burm. monogr.* 4. f. 1.

Sifyrinchium ramosum æthiopicum. *Breyn. cent.* 85. t.

27? *Rudb. chys.* 2. p. 13. f. 10?

Aspho-



*Aphodelus latifolius*, floribus patulis. *Breyn. prodr.* 3.

*p. 22. t. 9. f. 1.*

*Habitat ad Cap. b. spei.* 4

## COMMELINA.

\* *Dipetala dista.*

- communis.** 1. **COMMELINA** corollis inæqualibus, foliis ovato-lanceolatis acutis, caule repente glabro. *Hort. upf.* 18.  
*Commelina* foliis ovato-lanceolatis, caule procumbente glabro, petalis duobus majoribus. *Virid. cliff.* 6.  
*Hort. cliff.* 21. *Roy. lugdb.* 17. *Gron. virg.* 130.  
*Commelina procumbens* annua, saponariæ folio. *Dill. elth.* 93. t. 78. f. 89.  
*Habitat in America.* ☉  
*Confer Plantam Kæmpf. jap. t. 889. cum Tinctura inde paranda. Caulis ramosus. Foliorum margines laves imberbes.*
- africana.** 2. **COMMELINA** corollis inæqualibus, foliis lanceolatis glabris, caule decumbente.  
*Commelina* radice perenni, foliis lanceolatis, caule repente glabro, petalis duobus majoribus. *Wach. ultr.* 323.  
*Commelina procumbens*, flore luteo. *Roy. lugdb.* 538  
*Habitat in Æthiopia.* 24  
*Differt a prima petalis istis majoribus reniformibus luteis, tertio ovato filamentis longioribus æquali, vaginis foliorum ciliatis, involucro non reticulato. Vaginæ foliorum subciliatæ vagina alia minima intra exteriorem.*
- benghalensis.** 3. **COMMELINA** corollis inæqualibus, foliis ovatis obtusis, caule repente.  
*Ephemerum benghalense* serpens, folio subrotundo brevi, phalangoides. *Pluk. alm.* 130. t. 27. f. 3.  
*Habitat in Benghala.*
- erecta.** 4. **COMMELINA** corollis inæqualibus, foliis ovato-lanceolatis, caule erecto scabro simplicissimo. *Hort. upf.* 18.  
*Commelina* foliis ovato-lanceolatis, caule erectiusculo scabro, petalis duobus majoribus. *Virid. cliff.* 16.  
*Hort. cliff.* 495. *Gron. virg.* 11. *Roy. lugdb.* 38.  
**Com.**

# TRIANDRIA MONOGYNIA.

63

*Commelina erecta*, ampliore subcæruleo flore. *Dill.*

*clh.* 94. p. 77. t. 88.

*Habitat in Virginia.* 2

\* \* *Tripetala* : *Zanonia Pl.*

5. *COMMELINA* corollis subæqualibus, foliis lanceolatis subpetiolatis ore barbatis, caulibus erectis.

*Ephemerum phalangoides virginianum majus erectum.*

*Pluk. alm.* 135. t. 174. f. 4.

*Habitat in Virginia.* 2

*Caules erecti, simplices, leves, bipedales. Folia subpetiolata, lanceolata, supra retrorsum scabra: vaginis nervosis, subpubescentibus, ore ciliatis f. barbatis pilis ferrugineis. Spathæ terminales, subcordatæ. Flores carulei, Petalis cordatis, integerrimis: inferiore brevius pedicellato.*

virginica.

6. *COMMELINA* corollis æqualibus, foliis sessilibus, ovato-lanceolatis subciliatis. *Hort. upf.* 18.

*Commelina foliis ovato-lanceolatis, petalis tribus majoribus æqualibus. Hort. cliff. 21. Roy. lugdb.* 38.

*Commelina radice anacampserotides. Dill. clh.* 94. t. 79. f. 90.

*Habitat in Mexico.* 2

tuberosa.

7. *COMMELINA* corollis æqualibus, pedunculis incrassatis, foliis lanceolatis: vaginis laxis margine hirsutis, bracteis geminis.

*Zanonia graminea perfoliata. Plum. gen.* 38.

*Periclymenum rectum herbaceum, gentianæ folio, folii pediculo caulem ambiente. Sloan. jam.* 115. *hist.*

*1. p.* 243. t. 147. f. 1.

*Habitat in Americæ Gallia æquinoctiali.*

*Foliorum vaginæ latissimæ & a caule distantes, margine hirsutæ Bractæe duæ lanceolatæ oppositæ.*

Zanonia.

8. *COMMELINA* corollis æqualibus, pedunculis capillaribus, foliis lineari-lanceolatis, involucro nullo. *Fl. zeyl.* 31.

*Ephemerum phalangoides maderaspatense minimum, foliis perangustis, perfoliatum. Pluk. alm.* 135. t.

*27. f.* 4.

*Habitat in India.*

nud. & ora.

9. *COMMELINA* corollis æqualibus, floribus sessilibus, foliis linearibus.

axillaris

Ephe-

*Ephemerum phalangoides maderaspatanum* minimum, secundum caulem quasi ex utriculis floridum. *Pluk. alm.* 135. t. 174. f. 3.

*Ephemerum malabaricum*, flore tripetalo in foliorum alis sessili. *Raj. suppl.* 567.

*Nir-pulli*. *Rheed. mal.* 10. p. 28. t. 13.

*Habitat in India.*

eristata. 10. **COMMELINA** corollis æqualibus, involucris spicatum imbricatis. *Fl. zeyl.* 32.

*Ephemerum zeylonicum procumbens inflatum.* *Herm. parad.* 148. *Raj. hist.* 566.

*Habitat in Zeylona.*

### CALLISIA.

repens. 1. **CALLISIA.** *Loefl. it.* 305. \*

*Hapalanthus repens.* *Jacq. amer.* 12.

*Habitat in Americæ meridionalis udis depressis umbrosis.* ☉

*Affinis Commeline sed Nectariis destituta.*

### XYRIS.

indica. 1. **XYRIS.**

*Xyris foliis gladiatis.* *Gron. virg.* 11. *Flor. zeyl.* 35.

*Gladiolo lacustri accedens malabarica, e capitulo botryoideo florifera.* *Pluk. alm.* 170. t. 416. f. 4.

*Gramen junceum brasilianum, capite ovalli squamoso florido.* *Morif. hist.* 3. p. 229. f. 8. t. 9. f. 28.

*Jupicai.* *Pis. bras.* 238.

*Katsjiletri-pullo.* *Rheed. mal.* 9. p. 139. t. 7.

*Habitat in Indiis.*

### SCHOENUS.

\* *Culmo tereti.*

mariscus. 1. **SCHOENUS** culmo tereti, foliis margine dorsoque aculeatis. *Fl. suec.* 35. 38. *It. gotl.* 170. \* *Dalib. paris.*

14.

*Pseudo-cyperus palustris, foliis & carina serratis.*

*Scheuchz. gram.* 375.

*Cyperus longus inodorus germanicus.* *Bauh. pin.* 14.

*Cyperus longus inodorus major, foliis & carina serratis.* *Morif. hist.* 3. p. 237. f. 8. t. 11. f. 24.

*Habitat in Europæ paludibus.* 24.

2. SCHOENUS culmo tereti ramoſo, capitulis terminalibus, involucri triphyſſo breviffimo rigido patente. *Loefl. it.* 114. \* aculeatus.  
Gramen album, capitulis aculeatis, italicum. *Bauh. pin.* 7. *theatr.* 108. *Scheuch. gram.* 85. *Morif. hiſt.* 3. p. 195. f. 8. t. 5. f. 3.  
Gramen ſpicatum, ſpicis in capitulum foliatum congeſtis. *Tournef. inſt.* 517.  
Gramen aculeatum. *Cam. epit.* 745.  
*Habitat in Italia, Narbona, Luſitania, Archipelagi inſulis.* 24
3. SCHOENUS culmo tereti nudo, ſpicis faſciculatis mucronatis. mucronatus.  
divaricatis, involucri triphyſſo ſubulato.  
Melanoſchoenus maritimus humilis, cyperi effigie *Mich. gen.* 46.  
Scirpus culmo tereti, foliis tribus patulis ſiſculisque aggeſtis terminato. *Sauv. monſp.* 9.  
Scirpus maritimus, capite glomerato. *Tournef. inſt.* 46. *Scheuchz. gram.* 367.  
Gramen Cyperoides maritimum. *Bauh. pin.* 6. *theatr.* 91.  
Gramen junceum maritimum, capite ſquamuloſo foliaceo. *Morif. hiſt.* 3. p. 227. f. 8. t. 9. f. 6.  
*Habitat in Tyrreni, Smyrnæ, Hiſpaniæ, G. Narbonenſis maritimis arenofis.* 24  
*Folia etiam involucri apice pungunt. Spicæ oblongæ, ſeſſiles, apicibus diſtantes; Involucrum triphyllum, interdum diphyllum.*
4. SCHOENUS culmo tereti, ſpathis alternis mucronatis paniculas involventibus, ſpicis alternis patentibus. spathaceus.  
Cyperus ferrugineus. *Sp. pl.* 44.  
Cyperus racemis ſimplicibus lateralibus ſolitariis diſtiſchis, ſpicis alternis patentibus. *Gron. virg.* 131.  
Gramen junceum elatius, caule articulado, virginianum, cyperi paniculis inter folia prope ſummitatem prodeuntibus. *Pluk. alm.* 179. t. 301. f. 1.  
Gramen fluviatile geniculatum, panicula foliacea, locuſtis tenuibus oblongis, virginianum. *Morif. hiſt.* 3. p. 183. f. 8. t. 3. f. 17.  
*Habitat in Virginia, Cap. b. ſpei.* 24  
*Magnitudo Sacchari. Spathæ magnæ, monophyllæ, alternæ, deciduæ, racemos floriferos involvunt.*

5. SCHOE-

- nigricans.** 5. **SCHOENUS** culmo tereti nudo, capitulo ovato; involucri diphylli valvula altera subulata longa. *Fl. svec.* 36. 39. *It. gotl.* 234. *Dalib. parif.* 113.  
*Scirpus flosculus spicatis. Act. stockh.* 1741. p. 182.  
*Juncus lævis minor, panicula glomerata nigricante. Morif. hist.* 3. p. 233. f. 8. t. 10. f. 28.  
*Juncus lithospermi semine. Magn. monf.* 145. t. 144.  
*Juncus affinis capitulo glomerato nigricante. Scheuch. gram.* 349.  
*Habitat in Europæ paludibus æstate exsiccat.* 24
- ferrugineus.** 6. **SCHOENUS** culmo tereti nudo, spica duplici, involucri valvula majore spicam æquante. *Fl. svec.* 37. 40.  
*Gramen Cyperoides minimum, caryophylli proliferi capitulo simplici squamato. Morif. hist.* 3. p. 245. f. 8. t. 12. f. 40.  
*Juncello accedens graminifolia plantula, capitulis armeriæ proliferæ. Raj. angl.* 4. p. 32.  
*Habitat in Gotlandiæ, Angliæ paludibus.* 24
- capensis.** 7. **SCHOENUS** culmo tereti ramossissimo vaginato, spicis ovatis nudis dichotomis pendulis. *Amæn. acad.* 4. p. 264.  
*Canna Capitis b. spei, spicis juliformibus. Scheuch. gram.* 352.  
*Juncus e Cap. b. spei, paniculis fuscis juliformibus. Breyn. cent. t. 91. Pet. gaz. t. 7. f. 5. mus.* 424.  
*Juncus afric. lignoso calamo ad nodos involucris nigris convoluto, panicula arundinacea. Pluk. mant.* 109.  
*Habitat ad Cap. b. spei. b*  
*Culmi tenaces referunt Thefii juncei caules.*
- \* \* *Culmo triquetro.*
- coloratus.** 8. **SCHOENUS** culmo triquetro, capitulo subrotundo, involucro longissimo plano variegato.  
*Schoenus florum capitulis involucro longissimo radiatis. Hort. cliff.* 495.  
*Gramen Cyperoides spica compacta alba, foliis ad spicam, partim albis partim viridibus. Sloan. hist.* 1. p. 119. t. 78. f. 1.  
*Gramen capitatum. Rumph. amb.* 6, p. 8. t. 3. f. 2.  
 β. *Scirpus glomeratus culmo triquetro folioso, capitulo glomerato involucro folioso. Spect. pl.* 1. p. 52.

*Cyperus culmo triquetro folioso, capitulo glomerato triphylo, spicis teretibus. Gron. virg. 131.*

*Pee mottenga. Rheed. mal. 2. p. 99. t. 53.*

*Habitat in Indiis.*

*Culmus triqueter, Involucrum nutans, longitudine culmi. Spiculæ plurimæ, ovato-oblongæ, sessiles, in capitulum subrotundum congestæ.*

9. **SCHOENUS** culmo triquetro folioso, panicula villosa, seminibus globosis nitidis, foliis trifariam aculeolatis.

lithospermus.

*Schoenus fecans. Amæn. acad. 5. p. 391. \* Syst. nat. 2. p. 865.*

*Scirpus lithospermus. Sp. p. 1. p. 51.*

*Cyperus maximus, americanus, lithospermi semine.*

*Morif. hist. 3. p. 237. f. 8. t. 11. f. 16.*

*Gramen Cyperoides sylvaticum maximum geniculatum asperius, semine milii folis. Sloan. jam. 36. hist. 1. p. 118. t. 77. f. 1.*

*Calamagrostis. Rumph. amb. 5. p. 16. t. 6. f. 2.*

*Kaden-pullu. Rheed. mal. 12. p. 89. t. 48.*

*Habitat in Indiis. 2.*

*Distinctissimus a reliquis seminibus globosis, margaritæ colore nitidissimis. Folia longissima, pollice latiora, margine carinaque aculeis minutissimis secantia.*

10. **SCHOENUS** culmo subtriquetro nudo, spica disticha, involucro monophyllo.

compressus.

*Cyperella montana spicata, radice repente, caulero-tundo triquetro, spica fusca compressa disticha, semine cinereo. Mich. gen. 53.*

*Gramen Cyperoides spica simplici compressa disticha.*

*Pluk. alm. 173. t. 34. f. 9. Sheuch. gram. 490. Raj. hist. 1910. Morif. hist. 3. p. 245.*

*Habitat in Anglia, Helvetia, Italia.*

11. **SCHOENUS** culmo triquetro folioso, floribus fasciculatis, foliis planis, pedunculis lateralibus geminis.

glomeratus.

*Schoenus culmo triquetro, pedunculis geminis lateralibus, floribus conglomeratis. Gron. virg. 131.*

*Habitat in Virginia.*

12. **SCHOENUS** culmo subtriquetro folioso, floribus fasciculatis, foliis setaceis. *F. succ. 34. 41. Dalib. parif. 14.*

albus.

*Schoenus flosculus fasciculatis. Hort. cliff. 22. Roy lugdb. 48.*

*Cyperus palustris hirsutus minor, paniculis albis. Morif. hist. 3. f. 8. t. 9. f. 39. Fl. lapp. 17.*

*Cyperella palustris, capitulis umbellatis primum albis, deinde fulvis. Mich. gen. 53.*

*Gramen luzulæ accedeus glabrum, in palustribus proveniens, paniculatum. Pluk. alm. 178. t. 34. f. 11.*

*Gramen Cyperoides palustre leucanthemum. Scheuchz. gram. 503.*

*Habitat in Europæ borealis paludibus turfosis. 24*

## CYPERUS.

### \* *Culmo tereti.*

articulatus. 1. **CYPERUS** culmo tereti nudo articulado.

*Cyperus culmo nudo articulofo. Roy. lugdb. 51.*

*Juncus Cyperoides creberrime geniculatus medulla farctus aquaticus, radice rubra tuberosa odorata. Sloan. jam. 36. hist. 1. p. 121. t. 81. f. 1.*

*Habitat in Jamaicæ rivulis. 24*

minimus. 2. **CYPERUS** culmo tereti nudo, spicis sub apice.

*Cyperus culmo tereti, spicas sub apice ferente. Hort. cliff. 21. Roy. lugdb. 49.*

*Gramen junceum perpusillum, capillaceis foliis, æthiopicum. Pluk. alm. 179. t. 300. f. 5.*

*Gramen Cyperoides minimum, spicis pluribus compactis ex oblongo-rotundis. Sloan. jam. 36. hist. 1. p. 120. t. 79. f. 3. Raj. hist. 3. p. 625.*

*Habitat in Jamaica, Africa.*

### \* \* *Culmo triquetro.*

squarrosus. 3. **CYPERUS** culmo triquetro folioso, umbella foliosa glomerata, spicis striatis squarrosis. *Aman. acad. 4. p. 303. \**

*Gramen Cyperoides maderaspathi minimum ischæmi paniculis cum aristis. Pluk. amalt. 114. t. 397. f. 2.*

*Habitat in Asia.*

hispidus. 4. **CYPERUS** culmo triquetro folioso, umbella supradecomposita, spiculis umbellato-sessilibus *Flor. zeyl. 37. \**

*Gramen Cyperoides maderaspatanum, panicula magis sparsa & speciosa. Pluk. alm. 179. t. 192. f. 2.*

*Habitat in India, Æthiopia. 24*

3. CYPERUS culmo triquetro folioso, umbella foliosa supradecomposita, pedunculis nudis, spicis alternis. longus.  
*Roy. lugdb. 50. Mat. med. 29. Dalib. paris. 14.*  
*Cyperus odoratus radice longa f. Cyperus officinarum Bauh. pin. 14. Scheuch. gram. 378. Moris. hist. 3. pag. 237. f. 8. t. 11. f. 13.*  
*Habitat in Italiæ, Galliæ paludibus. 21*
6. CYPERUS culmo triquetro subnudo umbella decomposita: spicis alternis linearibus. FL. zeyl. 36. rotundus.  
*Mat. med. 30.*  
*Cyperus orientalis, radice olivarii, spicis longis e spadiceo purpurascens, Scheuch. gram. 391.*  
*Habitat in India.*
7. CYPERUS culmo triquetro nudo, umbella foliosa, radicum tuberibus ovatis: zonis imbricatis. Ray. esculentus  
*lugdb. 51.*  
*Cyperus rotundus esculentus angustifolius. Bauh. pin. 14. theatr. 222. Scheuch. gram. 382. Moris. hist. 3. p. 236. f. 8. t. 11. f. 10.*  
*Cyperus rotundus. Lob. ic. 75. Dod. pempt. 238.*  
*Thrasl. Bauh. hist. 2. p. 504.*  
*Habitat Monspelli, inque Italia, Oriente. 21*
8. CYPERUS culmo triquetro subnudo, umbella diphylla simplici trifida, spicis cuspidatis intermedia difformis.  
*sessili. Aman. acad. 4. p. 302. \**  
*Gramen Cyperoides elegans vasumbo curry malabarorum. Pluk. amalt. t. 317. f. 5.*  
*Habitat in India.*
9. CYPERUS culmo triquetro seminudo, umbella foliosa decomposita, spiculis alternis, granis distinctis. Iria.  
*Gramen Cyperoides Indiæ orientalis elatius, panicula sparsa pallescente. Pluk. alm. 179. t. 191. f. 7.*  
*Iria f. Balari. Rheed. mal. 12. p. 105. t. 56.*  
*Habitat in India. Osbeck.*  
*Pedunculi in umbella plurimi, apice umbelluliferi; umbella ex tribus quatuorve radiis, qui alternatim spiculas gerunt ex granis triquetris, obtusissimis, vix manifesta palea vestitis, parum distantibus.*
10. CYPERUS culmo triquetro nudo, umbella foliosa supradecomposita, spicis digitalibus imbricatis: spiculis subulatis. Aman. acad. 4. p. 301. \* elatus.  
*Habitat in India.*



- elegans.** 11. **CYPERUS** culmo triquetro nudo, umbella foliosa, pedunculis nudis proliferis, spicis confertis mucronibus patulis. *Roy. lugdb.* 51.  
*Cyperus* panicula maxime sparsa ferruginea compressa elegantissima. *Sloan. jam.* 35. *hist.* 1. p. 117. t. 75. f. 1. *Raj. suppl.* 623.  
*Habitat in Jamaica paludosis maritimis.*
- glomeratus.** 12. **CYPERUS** culmo triquetro nudo, umbella triphylla supradecomposita, spicis glomerato-rotundatis: spiculis subulatis. *Amoen. acad.* 4 p. 301. \*  
*Cyperus* aquaticus italicus procerior, locustis tenuissimis in racemum dense congestis. *Mich. gen.* 45. *Segu. veron.* 3. p. 68. t. 2. f. 2.  
*Cyperus* capitulis glomeratim congestis, feminibus aristatis. *Mont. gram.* 14. t. 1. f. 1.  
*Habitat in Italia paludosis ex India 24 Seguier.*
- odoratus.** 13. **CYPERUS** culmo triquetro nudo, umbella decomposita simpliciter foliosa, pedicellis distiche spicatis. *Cyperus* culmo triquetro nudo, umbella duplicata foliosa, pedunculis propriis distiche spicatis. *Roy. lugdb.* 50. *Gron. virg.* 131.  
*Cyperus* longus odoratus, panicula sparsa, spicis strigosioribus viridibus. *Sloan. jam.* 35. *hist.* 1. p. 116. t. 74. f. 1. & t. 8. f. 1.  
*Habitat in America ad fluviorum ripas.*
- compressus.** 14. **CYPERUS** culmo triquetro nudo, umbella universaliter triphylla, glumis mucronatis latere membranaceis.  
*Cyperus* culmo triquetro nudo, panicula foliosa, pedunculis simplicibus, spicis alternis subulatis distichis. *Roy. lugdb.* 51. *Gron. virg.* 131.  
*Cyperus* rotundus gramineus fere inodorus, panicula sparsa compressa viridi. *Sloan. jam.* 35. *hist.* 1. p. 117. t. 76. f. 1. *Raj. suppl.* 623.  
*Habitat in America septentrionalis pratis arenosis.*  
*Dignoscitur Panicula & Spicis viridibus; Glumis mucronatis, non patulis, at marginibus lateralibus membranaceis. Umbellae partiales nude, universalis glomerata.*
- flavescens.** 15. **CYPERUS** culmo triquetro nudo, umbella triphylla, pedunculis simplicibus inæqualibus, spicis confertis lanceolatis

*Cyperus culmo triquetro nudo, panicula foliosa supradecomposita, spicis confertis distiche compressis.*  
*Dalib. parif. 14.*

*Cyperus minimus, panicula sparsa flavescente. T.*  
*Scheuch. gram. 385. Hall. helv. 246.*

*Cyperus minor pulcher, panicula lata compressa subflavescente. Morif. hist. 3. p. 239. f. 8. t. 11. f. 37.*

*Gramen Cyperoides minus, panicula sparsa subflava.*  
*Bauh. pin. 6. theatr. 88. t. 88.*

*Habitat in Germaniæ, Helvetiæ, Galliæ, Italiæ paludosis.*

- 16 **CYPERUS** culmo triquetro nudo, umbella trifida, pedunculis simplicibus inæqualibus, spicis confertis linearibus.

fuscus.

*Cyperus culmo triquetro nudo, panicula diphylla supradecomposita, spicis strigosioribus confertis distiche compressis. Dalib. parif. 15.*

*Cyperus minimus, panicula sparsa nigricante. T.*  
*Scheuch. gram. 484.*

*Cyperus minor pulcher, panicula compressa nigricante. Morif. hist. 3. p. 239. f. 8. t. 9. f. 38.*

*Gramen Cyperoides minus, panicula sparsa nigricante.*  
*Bauh. pin. 6.*

*Habitat in Galliæ, Germaniæ, Helvetiæ pratis humidis.*

*Differt a C. flavescente vix manifeste, spiculis angustioribus fuscis & foliis magis scabris.*

17. **CYPERUS** culmo triquetro nudo, umbella diphylla composita, spiculis alternis digitatis lanceolatis: glumis mucronatis. *Amæn. acad. 4. p. 302. \**

pumilus.

*Gramen Cyperoides pumilum elegans e maderaspatan.*  
*Pluk. alm. 179. t. 191. f. 8.*

*Habitat in India.*

18. **CYPERUS** culmó triquetro nudo, umbella simplici, spiculis linearibus confertissimis horizontalibus.

strigosus.

*Cyperus rotundus, panicula sparsa, spicis strigosis ferrugineis. Sloan. jam. 35. hist. 1. p. 116. t. 74. f. 2. 3.*

*Habitat in paludibus Jamaicæ, Virginæ.*

*Radix subrotunda. Umbellæ radii simplices, erectiusculi, instructi spiculis linearibus, longiusculis, ferrugineis, confertissimis, numerosissimis, horizontaliter a pedunculo communi patentibus vel etiam deorsum flexis.*

ligularis

19. **CYPERUS** culmo triquetro, umbellæ spiculis capitatis oblongis sessilibus, involucris longissimis serrato-asperis. *Amæn. acad.* 5. p. 391. \*

*Cyperus maximus*, panicula minus sparsa ferruginea, capitulis compactis crassioribus. *Sloan, jam.* 35. *hist.* 1. p. 36. t. 9.

*Habitat in Jamaica.*

Papyrus.

20. **CYPERUS** culmo triquetro nudo, umbella simplici foliosa, pedunculis simplicissimis vaginatis distiche spicatis. *Roy. lugdb.* 50.

*Cyperus omnium maximus Papyrus dictus.* (*Mont. gram.* 14.), locustis minimis. *Mich. gen.* 44. t. 19.

*Cyperus enodis nudus*, culmis e vaginis brevibus prodeuntibus, spicis tenuioribus. *Scheuch. gram.* 387. t. 8. f. 14.

*Cyperus niloticus vel syriacus maximus papyraceus.* *Morif. hist.* 3. p. 239. f. 8. t. 11. f. 41.

*Papyrus syriaca & siciliana.* *Bauh. pin.* 19. *theatr.* 332. *Habitat in Calabria, Sicilia, Syria, Ægypto.*

## SCIRPUS.

\* \* *Spica unica.*

articulatus.

1. **SCIRPUS** culmo tereti nudiusculo semigeniculato, capitulo glomerato laterali.

*Tsieli. Rheed. mal.* 12. p. 135. t. 71.

*Habitat in Malabariæ aquis arenosis.*

*Culmus cavius, geniculis cinctus, at non undique, Spicæ laterales in capitulum semiverticillatum collectæ.*

palustris.

2. **SCIRPUS** culmo tereti nudo, spica subovata terminali. *Fl. suec.* 41. 42. *Dalib. parif.* 15.

*Scirpus culmo nudo, spica terminali subovata.* *Roy. lugdb.* 48.

*Scirpus equiseti capitulo majore.* *T. Scheuchz. gram.* 360. *Fl. lapp.* 19.

*Juncus palustris, capitulo equiseti, major.* *Bauh. pin.* 12.

*Juncellus cyperoides capitulo simplici.* *Laf. pruss.* 131. t. 36.

*Habitat in Europæ fossis & inundatis.*

pitatus.

3. **SCIRPUS** culmo tereti nudo setiformi, spica subglobosa terminali.

Scir-

*Scirpus culmo setaceo nudo, spica subglobosa. Gron.*  
*virg. 12.*

*Habitat in Virginia.*

4. **SCIRPUS** culmo triquetro nudo, spica cylindrica terminali. *Amæn. acad. 5. p. 391. \** mutatus.

*Habitat in Jamaica. 24*

*Simillimus S. geniculato, sed Culmus triqueter, mollisculus, nec articulatus.*

5. **SCIRPUS** culmo tereti nudo, spica oblonga terminali. geniculatus.

*Scirpus culmo nudo, spica terminali subrotunda. Hort. cliff. 21. \* Roy. lugdb. 48.*

*Juncus aquaticus geniculatus, capitulis equiseti major & minor. Sloan. jam. 37. hist. 1. p. 122. t. 75. f. 2. 63 t. 81. f. 3. Raj. suppl. 628.*

*Habitat in Jamaica.*

6. **SCIRPUS** culmo striato nudo, spica bivalvi terminali longitudine calycis, radicibus squamula interstinctis. *Fl. svec. 42. 43. Dalib. parif. 17.* cespitosus.

*Scirpus folio culmi unico. Fl. lapp. 20. Roy. lugdb. 49.*

*Scirpus montanus, capitulo brevior, Scheuch. gram. 363. t. 7. f. 18. Pluk. phyt. 40. f. 6. Raj. angl. 4. p. 32.*

*Juncus parvus palustris, cum parvis capitulis equiseti. Raj. hist. 1306. Pluk. phyt. 40. f. 6.*

*Gramen junceum, foliis & spica junci, minus. Bauh. pin. 6.*

*Habitat in Europæ paludibus cespitosis sylvaticis. 24*

7. **SCIRPUS** culmo tereti nudo setiformi, spica ovata terminali bivalvi, seminibus nudis. *Fl. svec. 43. 44. Dalib. parif. 17.* acicularis.

*Scirpus magnitudine aciculæ. Fl. lapp. 21. Roy. lugdb. 49.*

*Juncus minimus, spica brevior squamosa spadicea. Scheuch. gram. 364. t. 7. f. 4.*

*Juncellus minimus, capitulis equiseti. Pluk. alm. 201. t. 40. f. 7. Moris. hist. 3. p. 234. f. 8. t. 10. f. 37.*

*Habitat in Europa sub aquis purioribus.*

8. **SCIRPUS** culmis teretibus nudis alternis, caule folioso flaccido. fluitans.

*Scirpus foliis linearibus planis alternatim fasciculatis, spica terminali. Guett. Jlamp. 141.*

*Scirpus caule folioso flaccido, scapis alternis capitatis,*  
*Roy. lugdb. 49 Sauv. monsp. 9,*

*Scirpus equiseti capitulo minori. T. Scheuch. gram.*  
*365. t. 7. f. 20.*

*Juncellus, capitulis equiseti, minor fluitans, Bauh. pin.*  
*prodr. 23. theatr. 187.*

*Gramen junceum clavatum minimum. Raj. hist. 1310.*  
*Morif. hist. 3. p. 230. f. 8. t. 10. f. 31. Pluk. alm.*  
*180. t. 35 f. 1.*

*Habitat in Angliæ, Galliæ udis,*

\* \* *Culmo tereti polystachio.*

lacustris.

9. SCIRPUS culmo tereti nudo, spicis ovatis pluribus  
 pedunculatis terminalibus. *Fl. suec. 40. 46. Dalib.*  
*parif. 16.*

*Scirpus spicis copiosis. Roy. lugdb. 48.*

*Juncus maximus si Scirpus major. Bauh. pin. 12. the-*  
*art. 178.*

*Habitat in Europæ aquis puris stagnantibus & fluvia-*  
*tilibus.*

Holoscœnus 10. SCIRPUS culmo tereti nudo, spicis subglobosis glo-  
 meratis pedunculatis, involucreo diphylo inæquali  
 mucronato.

*Scirpus panicula solum foliosa, spicis globosis peduncu-*  
*latis lateralibus. Sauv. monsp. 8.*

*Juncus acutus maritimus, capitulis rotundis. Bauh. pin*  
*11. theatr. 17.*

*Scirpoides maritimum, capitulis sparsis glomeratis,*  
*Scheuch. gram. 371.*

Holoscœnus. *Dalech. hist. 8. 987.*

*Habitat in Europa australi.*

romanus.

11. S. IRPUS culmo tereti nudo, capitulo laterali con-  
 globato, bractea reflexa.

*Scirpus capitulo singulari, radice tomentosa. Barr. rar.*  
*1249. t. 255. f. 3.*

*Scirpoides acutum maritimum, capitulo glomerato soli-*  
*tario. Scheuch. gram. 373. Mich. gen. 52.*

8. *Juncus minor acutus maritimus proliſer ex uno capi-*  
*tello multa. Pluk. alm. 200. t. 40. f. 5.*

*Habitat in Galloprovincia & Romæ. 24*

*Huic Calami ſili craſſitie; præcedentis vero eras-*  
*ſi ut in S. lacuſtri & apice pungentes.*

12. SCIRPUS culmo nudo setaceo, spicis lateralibus subfoliariis sessilibus. *It. scan.* 227. \* *Fl. suec.* 2. setaceus.

n. 45.

Scirpus culmo nudo setaceo, spicis pedunculatis. *Roy. lugdb.* 49.

Scirpus omnium minimus, capitulo brevior. *T. Scheuch. gram.* 358.

Juncus inutilis f. Chamæschœnus. *Bauh. pin.* 12. *prod.* 22.

Juncellus omnium minimus. *Morif. hist.* 3. p. 232. f. 9. t. 10. f. 23.

Gramen junceum minimum, capite squamoso. *Bauh. pin.* 6. *prodr.* 13.

Habitat in Europæ litoribus maritimis.

13. SCIRPUS culmo tereti nudo, spicis sessilibus in medio culmo glomeratis. *Valib. parif.* 16. † supinus.

Scirpus supinus minimus, capitulis conglobatis, foliis rotundo-teretibus. *Tournef. inst.* 528.

Habitat Parisiis.

14. SCIRPUS culmo nudo capillari, spicis peduncularis sessilibusque lateralibus. capillaris.

Scirpus culmo teretiusculo seminudo, spicis capitatis sessilibus, involucro diphylo setaceo. *Flor. zeyl.* 39.

Cyperus culmo tereti spicas infra apicem ferente. *Hort. cliff.* 21.

Gramen junceum perpusillum, capillaceis foliis, æthiopicum. *Pluk. alm.* 179. t. 300. f. 5.

Gramen pusillum, junci capitulis minimis, ad basin foliolis binis acutis. *Burm. zeyl.* 108. t. 47. f. 2.

Habitat in Virginia, Æthiopia, Zeylona.

\* \* \* Culmo triquetro, panicula nuda.

15. SCIRPUS culmo triquetro nudo acuminato laterali, spicis conglomeratis. mucronatus

Scirpo-Cyperus maritimus. *Mich. gen.* 47. *ord. c.* 2. 3.

Juncus acutus maritimus, caule triangulo. *Bauh. pin.* 11. *prodr.* 22. *Morif. hist.* 3. p. 232. f. 8. t. 10. f. 20.

Juncus acutus maritimus, caule triquetro rigido & molli. *Pluk. alm.* 200. t. 40. f. 1. 3.

Habitat in Angliæ, Italiæ, Helvetiæ, Virginæ stagnis maritimis.

- dichotomus. 16. **SCIRPUS** culmo triquetro nudo, umbella decomposita, spicis dichotomizæ sessilibus. *Flor. zeyl.* 40. \*  
Gramen Cyperoides maderaspatanum, juncelli generi capitulis sparsis. *Pluk. alm.* 179. t. 119. f. 3.  
*Habitat in India.*
- echinatus. 17. **SCIRPUS** culmo triquetro nudo, umbella simplici, spicis ovatis. *Fl. zeyl.* 38. \*  
Cyperus floribus capitatis erectis pedunculatis. *Gron. virg.* 12.  
Gramen Cyperoides americanum, spicis grandioribus oblongo-rotundis, sparganii in modum echinatis, ad summum caulem pediculis longis innitentibus. *Pluk. alm.* 179. t. 91. f. 4.  
*Habitat in India utraque.*
- retrofractus. 18. **SCIRPUS** culmo triquetro, umbella simplici: spicarum flosculis retrofractis.  
Cyperi genus indianum, panicula speciosa, spiculis propendentibus atris. *Pluk. phyt.* 415. f. 4.  
*Habitat in Virginia.* 24
- ferrugineus. 19. **SCIRPUS** culmo triquetro subnudo, involucris longitudine paniculæ ciliatis.  
Cyperus culmo triquetro nudo, panicula diphylla, pedunculis quibusdam subdivisis, spicis ovatis solitariis. *Roy. lugdb.* 50.  
Gramen Cyperoides majus, spicis ex oblongo-rotundis compactis ferrugineis. *Sloan. jam.* 36. *hist.* 1. p. 36. t. 77. f. 2.  
*Habitat in Jamaicæ paludibus maritimis.*
- radiceus. 20. **SCIRPUS** culmo triquetro nudo, umbella subnuda, spicis oblongis sessilibus terminalibusque.  
**SCIRPUS** culmo triquetro nudo, panicula laxa, spicis alternis subsessilibus, pedunculis longis terminalibus. *Gron. virg.* 132.  
Gramen Cyperoides majus aquaticum, paniculis plurimis junceis sparsis, spicis ex oblongo rotundis spadicis. *Sloan. jam.* 36. *hist.* 1. p. 118. t. 76. f. 2.  
*Habitat in Jamaicæ fluviis.*  
\* \* \* \* Culmo triquetro, Panicula foliacea.
- maritimus. 21. **SCIRPUS** culmo triquetro, panicula conglobata foliacea, spicularum squamis trifidis: intermedia subulata. *Fl. succ.* 39. 47. *Dalib. parif.* 15.

*Cyperus culmo triquetro, panicula foliacea, pedunculis simplicissimis, spicis confertis. Roy. lugdb. 50.*

*Cyperus panicula subsessili, capitulis subovatis. Guett. flamp. 2. p. 414.*

*Cyperus rotundus inodorus germanicus. Bauh. pin. 14. theatr. 215.*

β. *Cyperus rotundus inodorus anglicus. Bauh. pin. 14.*

γ. *Cyperus panicula compacta e spicis teretibus crassioribus composita Scheuch. hist. 400.*

δ. *Gramen Cyperoides panicula sparsa majus. Bauh. pin. 6.*

ε. *Scirpo-Cyperus palustris, radice repente nodosa inodora, panicula sparsa, capitulis majoribus. Mich. gen. 48.*

*Habitat in Europæ litoribus maritimis. 24*

*Anatome partium fructificationis conjunctionem. β. γ. δ. ε. iussit.*

22. **SCIRPUS** culmo triquetro nudo, umbella supradecomposita, spicis intermediis sessilibus, involucrio setaceo. miliaceus.

*Habitat in India.*

*Culmi plurimi, nudi, simplices, triquetri, læves, semipedales. Folia graminea, basin culmi obvestientia, lævia. Umbella laxa, supradecomposita, diffusa. Involucra universalia partialiaque foliaceo-setacea, umbella breviora. Spicæ pedunculatæ, at solitariæ, sessiles in ramificationibus umbellæ; omnes ovatæ, minutissimæ ut semina milii.*

23. **SCIRPUS** culmo triquetro nudo, umbella foliosa prolifera, spiculis subrotundis. lucula.

*Habitat in India.*

*Culmus plus quam pedalis, nudus, triqueter, lævis. Involucrum universale polyphyllum: foliis ensiformibus, umbella longioribus. Partialia vix ulla, nisi seta brevissima. Umbellæ pedunculi intermediarii brevissimi; omnes terminati capitello, iterum prolifero. Capitella subrotunda, glomerata ex spiculis subrotundis.*

24. **SCIRPUS** culmo triquetro folioso, umbella foliacea, pedunculis nudis supradecompositis, spicis confertis. *Fl. suec. 38. 48. Dalib. parif. 15. Raj. angl. 4. p. 33.* sylvaticus.

*Cyperus culmo triquetro, panicula foliosa, pedunculis*



- illis nudis supradecompositis, spicis confertis. *Roy. lugd.* 49.
- Gramen cyperoides miliaceum. *Bauh. pin.* 6.
- Gramen arundinaceum, foliis acutissimis, panicula multiplici, cyperi facie. *Laf. pruss.* 119. t. 33.
- Habitat in Europæ sylvis humilibus.* 4.
- corymbosus. 25. SCIRPUS culmo triquetro folioso, corymbis lateralibus simplicibus; terminali prolifero, spicis subulatis. *Amæn. acad.* 4. p. 303.
- Habitat in India.*
- michelianus 26. SCIRPUS culmo triquetro, capitulo globofo, involucri polyphylo longo.
- Cyperus italicus omnium minimus.* *Till. pis.* 51. t. 20. f. 5.
- Habitat in Italia rarius.*
- \* \* \* \* Culmo triquetro, capitulo terminali.
- cephalotes. 27. SCIRPUS culmo triquetro nudo capitulo ovato squarrofo, involucri triphylo longo.
- Bobartia spicis capitatis involucri majori folioso tectis.* *Brown. jam.* 129.?
- Habitat in India.*
- Culmus altus, triqueter, levis, striatus. Capitulum ovatum, terminale, pedicellis non conspicuis. Calyces imbricati, acuminati squarrosi f. recurvati. Involucrum universale longissimum.* 3. f. 4. phyllum, reflexum, margine scabrum.

## ERIOPHORUM.

- vaginatum. 1. ERIOPHORUM culmis vaginatis teretibus, spica scariosa. *Fl. succ.* 45. 50. \* *Dalib. paris.* 18.
- Eriophorum spica erecta, caule tereti.* *Fl. l. pp.* 23. *Roy. lugdb.* 51.
- Gramen tomentosum alpinum & minus. *Bauh. pin.* 5. *prodr.* 10. *Burf.* 1: 43.
- Juncus alpinus capitulo lanuginoso f. Schœnolagurus.* *Bauh. pin.* 12. *prodr.* 23. *theatr.* 188. *Scheuch. gram.* 302. t. 7. f. 1.
- Habitat in Europæ frigidis sterilibus.* 4.
- polystachion. 2. ERIOPHORUM culmis teretibus, foliis planis, spicis pedunculatis. *Fl. succ.* 44. 49. *Dalib. paris.* 18.
- Erio-

*Eriophorum spicis pendulis. Fl. lapp. 28, Hort. cliff. 22.  
Roy. lugdb. 51.*

*Gramen pratense tomentosum, panicula sparsa. Bauh.  
pin. 4.*

*Habitat in Europæ uliginosis turfosis. 24*

3. **ERIOPHORUM** culmis foliosis teretibus, foliis planis, spica erecta. virginicum.

*Eriophorum spica compacta erecta foliacea, caule compresso. Gron. virg. 132.*

*Gramen tomentosum virginianum, panicula magis compacta aureo colore perfusa. Pluk. alm. 179. t. 299.  
f. 4*

*Gramen tomentosum, capitulo ampliore fusco & foliaceo. Moris. hist. 3. p. 224. f. 8. t. 9. f. 2.*

*Habitat in Virginia. 24*

4. **ERIOPHORUM** culmis teretibus foliosis, panicula supradecomposita prolifera, spiculis subternis. cyperinum.

*Scirpus paniculatus, foliis floralibus paniculam superantibus. Gron. virg. 12.*

*Cyperus milliaceus ex provincia mariana, panicula villosa aurea. Pluk. mant. 62. t. 419. f. 3.*

*Cyperus milliaceus marilandicus, spicis seminiferis magis confertis rubentibus lanuginosis. Raj. suppl. 620.*

*Habitat in America septentrionali.*

*Statura omnino Cyperi, sed Spiculæ Scirpi, nisi Semina maturescantia producerent Lanam gilvam s. testaceam, vix spiculis longiorem.*

5. **ERIOPHORUM** culmis nudis triquetris, spica pappo brevior. alpinum.

*Eriophorum spica erecta, caule triquetro. Fl. lapp. 24.*

*Linagrostis juncea alpina, capitulo parvo: tomento rariore. Scheuch. gram. 305. t. 7. f. 4.*

*Juncus alpinus bombycinus. Bauh. pin. 12. prodr. 6.*

*Habitat in Europæ alpinis, locisque affinis. 24*

## N A R D U S.

1. **NARDUS** spica setacea recta. stricta.

*paris. 18.*

*Nardus spica lineari. Fl. lapp. 40. Hort. cliff. 42.*

*Gramen sparsum juncifolium. Bauh. pin. 5. Scheuch. gram. 90.*

*Gramen sparsum hollandicum, capillaceo folio, & minus*

- nus. *Bauh. pin. 5. prodr. 11. Morif. hist. 3. p. 217. f. 8. t. 7. f. 8.*  
*Habitat in Europæ asperis, sterilibus, duris. 2.*
- gangitis. 2. **NARDUS spica recurva.**  
*Nardus spuria narbonensis. Bauh. pin. 13.*  
*Nardus gangitis spuria narbonensis. Lob. ic. 84. Morif. hist. 3. p. 257. f. 8. t. 13. f. ult. Raj. hist. 1911.*  
*Habitat in G. Narbonensi.*
- aristatus. 3. **NARDUS calycibus aristatis. Gouan.**  
*Gramen exile arundinaceum minimum, acumine reflexo. Bocc. mus. 2. p. 74. t. 57. Scheuch. gram. 11.*  
*Gramen junceum nodosum minimum capillare. Barr. rar. 106. t. 117. f. 1.*  
*Habitat Romæ.*
- ciliaris. 4. **NARDUS spica incurva ciliata.**  
*Habitat in India.*  
*Folia graminea, plana. Culmus spithameus. Spica dimidiata f. altero latere tantum. Flosculis alternis, erectis, 12. ad 15. valvula exteriore ovato-oblonga, mutica, margine ciliata pilis scabris, numerosis. Calyx proprie nullus, sed ad alterum latus flosculorum seta, brevior ipsa gluma.*

## LYGEUM.

- Spartum. 1. **LYGEUM. Læfl. it. 285. t. 2. \***  
*Spartum herba, alterum. Clus. hist. 2. p. 220. f. 2.*  
*Gramen Sparteum secundum, panicula brevi folliculo inclusa. Bauh. pin. 5.*  
*Habitat in Hispaniæ campis argillofis. 2.*

## DIGYNIA.

## BOBARTIA.

- iadica. 1. **BOBARTIA.**  
*Bobartia spicis capitatis, involucri folioso. Fl. Zeyl. 41. \**  
*Aman. acad. 1. p. 388. \**  
*Scirpus maderaspatanus, capitulo squamoso subrotundo. Scheuch. gram. 369.*  
*Habitat in India.*  
*Refert ita Schœnum mucronatum, ut ejus figura pro hujus ubique venditari possit.*

COR.

CORNUCOPIÆ.

1. CORNUCOPIÆ. *Hort. cliff.* 490. *Roy. lugdb.* 52. cucullatum.  
*Hasslq. it.* 452.  
*Juncus clavatus vaginatus polycephalus. Pet. gaz. t.*  
*73. f. 5.*  
*Gramen orientale venum in udis proveniens, capitu-*  
*lo inflexo. Scheuch. gram.* 117.  
*Habitat Smyrnæ. D. Hasselqvist.*

SACCHARUM.

1. SACCHARUM floribus paniculatis. *Hort. cliff.* 26. officinarum.  
*Mat. m.d.* 33. *Roy. lugdb.* 52.  
*Arundo saccharifera. Bauh. pin.* 18. *Sloan. jam.* 31. *hist.*  
*1. p. 108. t. 66. Rumph. amb.* 5. *p. 186. t. 74. f. 1.*  
*Habitat in Indiæ utriusque locis inundatis.*
2. SACCHARUM floribus spicatis, foliis undatis. spicatum.  
*Alopecurus malabarica, foliis undulatis, spica præte-*  
*nui. Scheuch. gram.* 91.  
*Gramen paniculatum brevifolium crispum, spica pur-*  
*pureo-sericea maderaspatanum. Pluk. alm.* 177. *t. 119.*  
*f. 1. Moris. hist.* 3. *p. 191. n. 7.*  
*Tjeria kuren-pullu. Rheed. mal.* 12. *p. 117. t. 62. Mo-*  
*rif. hist.* 3. *p. 191. n. 5.*  
*Habitat in Indiæ petrosifis.*

PHALARIS.

1. PHALARIS panicula subovata spiciformi: glumis canariensis.  
*carinatis. Hort. upf.* 19.  
*Phalaris radice annua. Hort. cliff.* 23. *Roy. lugdb.* 63.  
*Dalib. parif.* 20.  
*Phalaris major, semine albo. Bauh. pin.* 28. *Moris.*  
*hist.* 3. *p. 186. f. 8. t. 3. f. 1.*
2. PHALARIS panicula cylindrica: glumis carinatis. bulbosa.  
*Amæn. acad.* 4. *p. 264. \**  
*Phalaris radice perenni. Roy. lugdb.* 63.  
*Phalaris bulbosa, semine albo. Raj. hist.* 1249. *Scheuch.*  
*gram.* 53.  
*Habitat in Oriente.*
3. PHALARIS panicula ovato-oblonga spiciformi; glu- aquatica.  
*mis carinatis lanceolatis. Amæn. acad.* 4. *p. 264. \**  
*Gramen typhinum phalaroides majus bulbosum aqua-*  
*ticum. Barr. rar.* 1200. *t. 700. f. 1.*

Gra-

- Gramen phalaroides hirsutum, spica longissima. *Euxb. cent. 4. p. 32. t. 53.*  
*Habitat ad Tyberim & in Ægypto. 2*
- utriculata. 4. PHALARIS panicula spicata, petalis arista articulata, vagina supremi folii spathiformi.  
 Gramen pratense, spica purpurea ex utriculo prodeunte, f. Gramen folio spicam amplexante. *Bauh. pin. 3. theatr. 44.*  
*Habitat in Italia.*
- phleoides. 5. PHALARIS panicula cylindrica spiciformi.  
 Phalaris spica cylindrica. *Fl. succ. 49. 54. Dalib. parif. 19.*  
 Gramen typhoides asperum primum. *Bauh. pin. 4. Scheuch. gram. 61. Hall. helv. 232.*  
*Habitat in Europæ versuris.*
- arundinacea 6. PHALARIS panicula oblonga ventricosa.  
 Phalaris panicula oblonga. *Fl. succ. 48. 53. Dalib. parif. 19.*  
 Arundo foliis planis, panicula spicata, spiculis confertis. *Hort. cliff. 26. Roy. lugdb. 66.*  
 Gramen aquaticum paniculatum, phalaridis femine. *T. Scheuch. gram. 126.*  
 Gramen arundinaceum spicatum. *Bauh. pin. 6. theatr. 94.*  
 Gramen aquaticum paniculatum latifolium. *Bauh. pin. 3.*  
 Gramen arundinaceum, acerosa gluma, jerscianum. *Raj. ang. 3. p. 400. Morif. hist. 3. p. 203. f. 8. t. 5. f. 41.*
- picca. 8. Gramen paniculatum, folio variegato. *Bauh. pin. 3.*  
*Habitat in Europæ subhumidis ad ripas lacuum*
- cruciformis 7. PHALARIS panicula lineari secunda, calycibus bifloris.  
 Dactylis spicis numerosis alternis culmo adpressis longitudine internodiorum, calycibus bifloris. *Roy. lugdb. 57. Gmel. sibir. 1. p. 130. t. 29.*  
 Gramen palustre, locustis erucaformibus. *Barr. rar. 1158. t. 2.*  
*Habitat in Sibiria, Russia, Europa australi.*  
*Paspalum refert, sed valvulae floris modo contrario dehiscunt.*

1. **PHALARIS** panicula effusa, glumarum carinis ciliatis. oryzoides.

*Oryza* glumis carina hispidis. *Gron. virg.* 153.

Gramini tremulo affine paniculatum elegans majus.

*Sloan. jam.* 34. *hist.* 1. p. 113. t. 71. f. 1.

Gramen palustre, panicula speciosa. *Bauh. pin.* 2. *prodr.* 7.

Gramen paniculatum aquaticum asperum, locustis planis & orbiculatis. *Pont. comp.* 57.

Gramen miliaceum palustre ferotinum, paniculis ex locustis compressis compositis. *Mont. prodr.* 51. *Se-gu. ver.* 1. p. 351.

Habitat in Virginiae paludibus nemorosis, hodie in Italiae agris *Oryzae*.

## PASPALUM.

1. **PASPALUM** spicis alternis: rachi membranacea. dissectum.

*Panicum* dissectum. *Sp. pl.* 1. p. 57.

Gramen paniceum spica simplici ad caulem intervallata binis granorum ordinibus uno versu constante, americanum. *Pluk. mant.* 94. t. 350. f. 2.

Habitat in America calidiore. ☉

Gramen prostratum foliorum vaginis fere spathaceis.

Spicae paucae rachi membranacea dilatata ad latitudinem spicae ipsius distichae & secundae. Flores orbiculati.

1. **PASPALUM** spiculis paniculatis alternis basi villosis. virgatum.

Gramen Daetylon majus, panicula longa, spicis plurimis nudis crassis. *Sloan. jam.* 34. *hist.* 1. p. 112. t. 69. f. 2.

Habitat in Jamaica.

Precedenti duplo vel quadruplo majus. Spiculae basi villosae ad decem usque, alternae, paniculam constituent, secundae, ordine non duplici, sed quadruplici florum.

2. **PASPALUM** spicis paniculatis verticillato-aggregatis. paniculatum.

Gramen miliaceum, panicula viridi l. purpurea. *Sloan. pin.* 34. *hist.* 1. p. 115. t. 72. f. 2.

Habitat in Jamaica.

Erectius hoc est, panicula quasi verticillata spiculis innumeris, linearibus, secundis, filiformibus, an-  
*Linzi Spec. plant. pars I.* R gu-

- guftiffimis: floribus duplici ordine digeffis, acutiusculis.*
- diffichum. 4. **PASPALUM** spicis duabus: altera subseffili, floribus acuminatis. *Amæn. acad. 5. p. 391. \**  
*Habitat in Jamaica.*  
*Gramen pedale. Foliis involuto - subulatis. Culmus terminatus Spicis duabus, erectis, approximatis, secundis: floribus alternis, acuminatis: omnibus sunt Stamina tria.*

## PANICUM.

\* *Spicata.*

- alopecu-  
deum. 1. **PANICUM** spica tereti, involucris setaceis basi ciliatis fasciculatis unifloris flosculo quadruplo longioribus.  
*Gramen geniculatum brevifolium crispum, spica purpureo-sericea. Pluk. alm. 177. t. 119. f. 1.?*  
*Gramen indicum alopecuroides holosericeum majus, spica longa pappescente. Pluk. alm. 177. t. 92. f. 5.*  
*Habitat in Jamaica.*  
*Spica longa, ferruginea; Flosculi sparsi, pedunculis brevibus, cincti setis longis, erectis. Scapus infra spicam parum villosus. Pedicelli omnes uniflori.*
- polysta-  
chyon. 2. **PANICUM** spicis teretibus, involucellis unifloris fasciculato-setosis, culmis erectis superne ramosis. *Gramen caricosum alterum. Rumph. amb. 6. p. 17. t. 7. f. 2. B.*  
*Habitat in India. ♂.*  
*Similis Spicis Panico glauco, sed dimidio minoribus, pistillis sanguineis. Culmus ramosus, pedalis ad sex-pedalem, valde ramosus. Folia pilosa, superne fere opposita.*
- verticilla-  
tum. 3. **PANICUM** spica verticillata, racemulis quaternis, involucellis unifloris bisetis, culmis diffusis. *Gramen paniceum, spica aspera. Bauh. pin. 8. theatr. 139. t. 139. Scheuch gram. 67.*  
*Habitat in Europa australi & Oriente.*  
*Simillimum P. viridi, sed Spica racemis longioribus, 3. f. 4. deorsum secundis. Flores virides, pistillis purpurascensibus. Seta 1. f. 2. scabra, ad angulum florem, eoque longior. Pedicelli a casu florum terminati cotyla truncata, concava, alba. Spica rachis*

*chis angulis 3. f. 4. bifidis. Culmi leves, diffusi, bipedales. Folia nuda.*

4. PANICUM spica tereti, involucllis bifloris fasciculato-setosis, seminibus undulato rugosis. *Fl. zeyl. 44.* glaucum.

Panicum spica simplici, aristis aggregatis flosculo subjectis. *Gron. virg. 134.*

Habitat in Indiis & Italia. ☉

Seta in spica longitudine flosculorum. Foliorum vagina oris pilosa. Dum spica recens prodit. Flosculi in series dispositi observantur. Semina striis undulatis notata.

5. PANICUM spica tereti, involucllis bifloris fasciculato-pilosis, seminibus nervosis. viride.

Græmen paniceum f. Panicum sylvestre, spica simplici. *Bauh. pin. 8. Scheuch. gram. 46.*

Habitat in Europa australi. ☉

6. PANICUM spica composita, spiculis glomeratis fetis immixtis, pedunculis hirsutis. italicum.

Panicum spica composita, aristis flosculo brevioribus. *Vir. cliff. 7. Hort. upf. 19. Roy. lugdb. 54. Gron. virg. 134.*

Panicum italicum f. panicula majore, *Bauh. pin. 27. theatr. 519.*

Panicum. *Rumph. amb. 5. p. 202. t. 75. f. 2.*

Habitat in Indiis. ☉

7. PANICUM spica composita: spiculis adpressis alternis, calycibus geminis, valvulis omnibus aristatis; hirtellam.

arista extrema longissima. *Amæn. acad. 5. p. 391. \**

Panicum sylvestre, panicula rariori, spicillis longioribus uno versu floridis. *Brown. jam. 133.*

Habitat in Indiis, etiam ibi colitur.

8. PANICUM spicis alternis conjugatisque, spiculis subdivisis, glumis aristatis hispidis, rachi quinquangulari. *Fl. junc. 2. n. 55.* crus galli.

Panicum spicis alternis remotis declinatis compositis.

*Vir. cliff. 7. Roy. lugdb. 55. Dalib. paris. 22.*

Panicum spicis alternis remotis laxis. *Hort. cliff. 27.*

Græmen paniceum, spica divisa. *Bauh. pin. 8. Scheuch. gram. 49. Morif. hist. 3. p. 189. f. 8. t. 4. f. 15.*

Dens caninus secundus. *Bauh. hist. 2. p. 434.*

5. Græmen paniceum, spica divisa, aristis longis armata.

*Bauh. pin. 8.*



*Habitat in Europæ, Virginæ cultis.* ☉

Varietas  $\beta$ . *aristis decies glumis longioribus manifeste, communi planta, cui vix sesquialongiores aristæ, differe-*

crus corvi. 9. PANICUM spicis alternis secundis spiculis subdivi-  
fis, glumis subaristatis hispidis, rachi trigona.

*Habitat in Indiis.* ☉

*Simillimum P. crus galli, sed minus, tenerius. Panicula magis nutans rachi trigona, nec pentagona. Spicæ secundæ, nec undique rachi incumbentes.*

colonum. 10. PANICUM spicis alternis secundis muticis ovatis  
scabris, rachi teretiuscula.

*Gramen paniceum minus, spica divulsa insulæ barba-  
densis. Pluk. alm. 174. t. 189. f. 5.*

*Gramen paniceum minimum humistratum, spica divisi-  
mutica, foliis variegatis. Sloan. jam. 30. hist. 1. p. 107  
t. 64. f. 3. Raj. suppl. 600. Ehret. pist. t. 3. f. 3.*

*Gramen aciculatum Rumph. amb. 6. p. 13. t. 5. f. 3.*

*Habitat in Indiæ cultis.* ☉

*Culmi spithamei, teretes, adscendentes. Folia levia,  
sepe ferrugineo-maculata, fauce purpurascens abs-  
que ligula. Spica simplex, secunda, rachi tereti,  
debili. Spiculæ alternæ, plurimæ, subsessiles, se-  
cundæ, ovatæ s. oblongiusculæ absque fasciculis pi-  
lois. Flosculi subtrianguli. Anthæræ purpurascens,  
Pistilla alba, mox purpurascens.*

compositum. 11. PANICUM spica composita, spiculis linearibus se-  
cundis, flosculis geminis remotis, calycibus aristatis  
*Fl. Zeyl. 42. \**

*Habitat in Zeylona.*

dimidiatum. 12. PANICUM spica dimidiato-secunda, rachi lineari  
membranacea extrorsum flosculis aggregatis.

*Gramen bermudense frumentaceum, tolius brevibus ob-  
tusis, spica crassiuscula compressa antrorsum gemi-  
no versus granifera. Pluk. phyt. 244. f. 6.*

*Habitat in India.*

*Folia latissima, obtusiuscula vaginis latis. Spica ex  
scapi apice membranaceo, plana, cui ad alterum  
tantum latus Flores alternatim divergentes, termi-  
nata. Structura Paspali, sed fructificatio diversa.*

sanguinale. 13. PANICUM spicis digitatis basi interiore nodosis,  
flosculis geminis muticis, vaginis foliorum punctatis

Pa-

*Panicum* spicis alternis oppositisque linearibus patentissimis muticis, flosculis alternatim binis: altero pedunculato. *Roy. lugdb.* 55. *Gron. virg.* 154. *Dalib. parif.* 22.

*Gramen daetylon majus repens, foliis hirsutissimis.*  
*Ruxb. cent.* 5. p. 24. t. 65.

*Gramen daetylon, folio latiore.* *Bauh. pin.* 8. *theatr.* 114. *Scheuch. gram.* 101.

*Gramen daetylon majus, panicula longa e spicis plurimis gracilioribus purpureis & viridibus mollibus constante.* *Sloan. jam.* 3. *hist.* 1. p. 113. t. 70. f. 2.

*Habitat in America, Europa australi, Hollandia, Media.* ☉

14. *PANICUM* spicis digitatis patentibus basi interiore villosis, floribus solitariis, sarmentis repentibus.

*Daetylon.*

*Panicum Daetylon, folio arundinaceo, majus aculeatum.* *Bauh. pin.* 7.

*Panicum Daetylon radice repente* ☿ *officinaŕum.*  
*Scheuch. gram.* 104.

*Gramen legitimum.* *Clus. hist.* 2. p. 217.

*Habitat in Europa australi, Oriente.* 24

15. *PANICUM* spicis subdigitatis approximatis erectis filiformibus; rachi flexuosa, dentibus bifloris: altero sessili.

*filiforme.*

*Habitat in America septentrionali.* *Kalm.*

*Folia lanceolata. Culmi pedales filiformes, superne nudii erecti. Spica filiformis, rara, flexuosa rachi alternatim utrinque ad dentes Flosculi 2. altero sessili, altero pedicellato; spicæ tenues, primum valde approximatae quasi in unum, quamvis alternæ.*

16. *PANICUM* spicis digitatis subquaternis linearibus, flosculis solitariis secundis muticis.

*lineare.*

*Habitat in Indiis.*

*Culmi prostrati, Leves, ramosi. Spiculæ lineares, rectæ, angustæ. Flores subius alterni. Calycis squama exterior brevior, patens, rachi adhaerens.*  
\* \* *Paniculata.*

17. *PANICUM* paniculis simplicibus, culmo ramoſo dichotomo. *Gron. virg.* 133. \*

*dichotomum.*

*Habitat in Virginia.*

*Arva refert arbusculam inferne simplicem; superne fasciculatam.*

- grosularium. 18. **PANICUM** panicula ramis simplicibus, floribus geminis: pedicello altero brevissimo, altero longitudine floris. *Amæn. acad. 5. p. 392. \**  
*Habitat in Jamaica.*
- clandestinum. 19. **PANICUM** racemis occultatis intra foliorum vaginas.  
 Gramen cyperoides polystachyon, spicis ad nodos ex foliorum alis prodeuntibus. *Sloan. jam. 36. hist. 1. p. 120. t. 80.*  
*Habitat in Jamaica, Pennsylvania. Kalm. 24*  
 Culmi dichotomi, ramosi. Folia ovato-lanceolata, vaginis punctatis, pillosis, intra quas Racemi fructificationum omnino occultantur, ut extus ne vestigium earum appareat; ceterum valde affine *P. latifolio.*
- capillare. 20. **PANICUM** panicula capillari erecta patente, caule hirsuto. *Gron. virg. 13.*  
*Panicum miliaceum viride, foliis latis brevibus, panicula capillacea. Sloan. jam. 35. hist. 1. p. 115. t. 72. f. 3.*  
*Milium indicum, panicula sparsa erecta. Tournef. injt. 515.*  
*Habitat in Virginia, Jamaica. ☉*  
*Simile Poæ capillari, sed totum hirsutum.*
- patens. 21. **PANICUM** panicula oblonga patente, calycibus bifloris, foliis lineari lanceolatis.  
*Tsiamapullu. Rheed. mal. 12. p. 75. t. 41.*  
*Habitat in India: similis e Lusitania.*
- divaricatum. 22. **PANICUM** paniculis brevibus muticis, culmo ramosissimo divaricatissimo, pedicellis bifloris: altero longiore. *Amæn. acad. 5. p. 392. \**  
*Habitat in Jamaica.*
- miliaceum. 23. **PANICUM** panicula laxaflaccida, foliorum vaginis pubescentibus. *Hort. cliff. 27. Hort. ups. 20. Nat. med. 34. Roy. lugdb. 55. Dalib. parif. 22.*  
*Milium femine luteo & albo. Bauh. pin. 26. theatr. 502.*  
*Habitat in India.*
- latifolium. 24. **PANICUM** panicula racemis lateralibus simplicibus, foliis ovato-lanceolatis collo pilosis.  
*Gramen virginianum, lato brevique folio, panicula rariore. Morif. hist. 3. p. 196. f. 8. t. 5. f. 4.*

*Gramen miliaceum sylvaticum*, maximum, semine albo. Sloan. jam. 34. hist. 1. p. 114. t. 71. f. 3.

*Habitat in America.*

*Folia latitudine Commelinæ, ad fauces amplexicaulia, extus collo circum fauces villosa, etiam basi foliorum margine pilosa. Panicula valde mediocris ex racemis lateralibus, non subdivisis, sed pedicellos proprios, nec subdivisos proferentes. Flores mutici, masculi.*

15. **PANICUM** paniculatum, foliorum vaginis longitudo brevifolium  
tudinaliter ciliatis.

*Panicum miliaceum, latiore folio, madéraspatanum.*

*Pluk. aln. 176. t. 189. f. 4.*

*Habitat in India.*

*Gramen pedule vel bipedale. Folia ovata, amplexicaulia. Panicula tenuissima, folio supremo & majore compresso. Variat foliis angustis & lanceolatis.*

- 16 **PANICUM** paniculatum ramosissimum, foliis ovato-arborescens.  
oblongis acuminatis. *Fl. Zeyl. 43.*

*Panicum panicula laxa sparsa, foliorum vaginis glabris. Hort. cliff. 27.*

*Habitat in India. b*

*Altitudine certat cum altissimis arboribus. Culmus vix penna anserina latior.*

17. **PANICUM** panicula virgata, foliis divaricatis.

*repens.*

*Habitat in Hispania? inde missum a Claud. Alstræmer. 24*

*Culmi repentes pedales adscendentes. Folia inferiora latiora, undique caulem vaginis vestientia, patentissima & fere reflexo-divaricata; mucronata vaginis subpubescentibus: Superiora folia angustiora, longiora, erecta. Panicula constans tribus quatuorve racemis erectis, vix ramosis, quibus Flores alterni, remotiusculi, ovati; valvula calycina exterior brevis, obtusa.*

18. **PANICUM** panicula virgata, glumis acuminatis virgatum.  
lævibus: extima dehiscente.

*Panicum paniculatum, glumis acutis. Gron. virg. 133.*

*Habitat in Virginia.*

## PHLEUM.

1. **PHLEUM** spica cylindrica longissima culmo recto.

*pratense.*

*F 4*

*Fl.*

- gustissimis: floribus duplici ordine digestis, acutiusculis.*
- distichum. 4. **PASPALUM** spicis duabus: altera subseffili, floribus acuminatis. *Amæn. acad. 5. p. 391. \**  
*Habitat in Jamaica.*  
*Gramen pedale. Foliis involuto - subulatis. Culmus terminatus spicis duabus, erectis, approximatis, secundis: floribus alternis, acuminatis: omnibus sunt Stamina tria.*

## PANICUM.

\* *Spicata.*

- alopecu-  
deum. 1. **PANICUM** spica tereti, involucris setaceis basi ciliatis fasciculatis unifloris flosculo quadruplo longioribus.  
*Gramen geniculatum brevifolium crispum, spica purpureo-sericea. Pluk. alm. 177. t. 119. f. 1.?*  
*Gramen indicum alopecuroides holosericeum majus, spica longa pappescente. Pluk. alm. 177. t. 92. f. 5.*  
*Habitat in Jamaica.*  
*Spica longa, ferruginea; Flosculi sparsi, pedunculis brevibus, cincti setis longis, erectis. Scapus infra spicam parum villosus. Pedicelli omnes uniflori.*
- polysta-  
chyon. 2. **PANICUM** spicis teretibus, involucellis unifloris fasciculato-setosis, culmis erectis superne ramosis.  
*Gramen caricosum alterum. Rumph. amb. 6. p. 17. t. 7. f. 2. B.*  
*Habitat in India. ♂.*  
*Similis Spicis Panico glauco, sed dimidio minoribus, pistillis sanguineis. Culmus ramosus, pedalis ad sex-pedalem, valde ramosus. Folia pilosa, superne fere opposita.*
- verticilla-  
tum. 3. **PANICUM** spica verticillata, racemulis quaternis, involucellis unifloris bifetis, culmis diffusis.  
*Gramen paniceum, spica aspera. Bauh. pin. 8. theat. 139. t. 139. Scheuch. gram. 67.*  
*Habitat in Europa australi & Oriente.*  
*Simillimum P. viridi, sed Spica racemis longioribus, 3. f. 4. deorsum secundis. Flores virides, pistillis purpurascensibus. Seta 1. f. 2. scabra, ad angulum florem, eoque longior. Pedicelli a casu florum terminati cotyla truncata, concava, alba. Spicae rachis*

*chis angulis 3. f. 4. bifidis. Culmi laeves, diffusi, bipedales. Folia nuda.*

4. **PANICUM** spica tereti, involucellis bifloris fasciculato-setosis, seminibus undulato rugosis. *Fl. zeyl. 44.* glaucum.

*Panicum spica simplici, aristis aggregatis flosculo subiectis. Gron. virg. 134.*

*Habitat in Indiis & Italia. ☉*

*Setæ in spica longitudine flosculorum. Foliorum vaginae oris pilosæ. Dum spica recens prodit. Flosculi in series dispositi observantur. Semina striis undulatis notata.*

5. **PANICUM** spica tereti, involucellis bifloris fasciculato-pilosis, seminibus nervosis. viride.

*Germen paniceum f. Panicum sylvestre, spica simplici. Bauh. pin. 8. Scheuch. gram. 46.*

*Habitat in Europa australi. ☉*

6. **PANICUM** spica composita, spiculis glomeratis setis immixtis, pedunculis hirsutis. italicum.

*Panicum spica composita, aristis flosculo brevioribus. Vir. cliff. 7. Hort. ups. 19. Roy. lugdb. 54. Gron. virg. 134.*

*Panicum italicum f. panicula majore. Bauh. pin. 27. theatr. 519.*

*Panicum. Rumph. amb. 5. p. 202. t. 75. f. 2.*

*Habitat in Indiis. ☉*

7. **PANICUM** spica composita: spiculis adpressis alternis, calycibus geminis, valvulis omnibus aristatis; hirtellum.

*arista extrema longissima. Amæn. a. ad. 5. p. 391. \**

*Panicum sylvestre, panicula rariori, spicillis longioribus uno versu floridis. Brown. jam. 133.*

*Habitat in Indiis, etiam ibi colitur.*

8. **PANICUM** spicis alternis conjugatisque, spiculis subdivisis, glumis aristatis hispidis, rachi quinque- crus galli.

*gulari. Fl. jvec. 2. n. 55.*

*Panicum spicis alternis remotis declinatis compositis.*

*Vir. cliff. 7. Roy. lugdb. 55. Dalib. paris. 22.*

*Panicum spicis alternis remotis laxis. Hort. cliff. 27.*

*Germen paniceum, spica divisa. Bauh. pin. 8. Scheuch. gram. 49. Moris. hist. 3. p. 189. f. 8. t. 4. f. 15.*

*Dens caninus secundus. Bauh. hist. 2. p. 434.*

9. **Germen paniceum, spica divisa, aristis longis armata.**

*Bauh. pin. 8.*

*Habitat in Europæ, Virginæ cultis.* ☉

Varietas  $\beta$ . *aristis decies glumis longioribus manifeste a communi planta, cui vix sesquolongiores aristæ, differt.*

crus corvi. 9. **PANICUM** spicis alternis secundis spiculis subdivisis, glumis subaristatis hispidis, rachi trigona.

*Habitat in Indiis.* ☉

*Simillimum P. crus galli, sed minus, tenerius. Panicula magis nutans rachi trigona, nec pentagona. Spicæ secundæ, nec undique rachi incumbentes.*

colonum. 10. **PANICUM** spicis alternis secundis muticis ovatis scabris, rachi teretiuscula.

*Gramen paniceum minus, spica divulsa insulæ barbadensis. Pluk. alm. 174. t. 189. f. 5.*

*Gramen paniceum minimum humistratum, spica divisa mutica, foliis variegatis. Sloan. jam. 30. hist. 1. p. 107. t. 64 f. 3. Raj. suppl. 600. Ehret. pist. t. 3. f. 3.*

*Gramen aciculatum Rumph. amb. 6. p. 13. t. 5. f. 3.*

*Habitat in Indiæ cultis.* ☉

*Culmi spithamei, teretes, ascendentes. Folia lævia, sæpe ferrugineo-maculata, fauce purpurascensia absque ligula. Spica simplex, secunda, rachi tereti, debili. Spiculæ alternæ, plurimæ, subseßiles, secundæ, ovatæ s. oblongiusculæ absque fasciculis pilosis. Flosculi substriati. Anthæræ purpurascens; Pistilla alba, mox purpurascensia.*

compositum. 11. **PANICUM** spica composita, spiculis linearibus secundis, flosculis geminis remotis, calycibus aristatis. *Fl. Zeyl. 42. \**

*Habitat in Zeylona.*

dimidiatum. 12. **PANICUM** spica dimidiato-secunda, rachi lineari membranacea extrorsum flosculis aggregatis.

*Gramen bermudense frumentaceum, foliis brevibus obtusis, spica crassiuscula compressa antrorsum gemino versus granifera. Pluk. phyt. 244. f. 6.*

*Habitat in India.*

*Folia latissima, obtusiuscula vaginis latis. Spica ex scapi apice membranaceo, plana, cui ad alterum tantum latus Flores alternatim divergentes, terni, Structura Paspali, sed fructificatio diversa.*

sanguinale. 13. **PANICUM** spicis digitatis basi interiore nodosis, flosculis geminis muticis, vaginis foliorum punctatis.

Pa-

*Panicum spicis alternis oppositisque linearibus patentissimis muticis, flosculis alternatim binis: altero pedunculato. Roy. lugdb. 55. Gron. virg. 154. Dalib. parif. 22.*

*Gramen dactylon majus repens, foliis hirsutissimis. Ruxb. cent. 5. p. 34. t. 65.*

*Gramen dactylon, folio latiore. Bauh. pin. 8. theatr. 114. Scheuch. gram. 101.*

*Gramen dactylon majus, panicula longa espicis plurimis gracilioribus purpureis & viridibus mollibus constante. Sloan. jam. 3. hist. 1. p. 113. t. 70. f. 2.*

*Habitat in America, Europa australi, Hollandia, Media. ☉*

14. *PANICUM spicis digitatis patentibus basi interiore villosis, floribus solitariis, sarmentis repentibus.*

*Dactylon.*

*Panicum Dactylon, folio arundinaceo, majus aculeatum. Bauh. pin. 7.*

*Panicum Dactylon radice repente & officinarum. Scheuch. gram. 104.*

*Gramen legitimum. Clus. hist. 2. p. 217.*

*Habitat in Europa australi, Oriente. 24*

15. *PANICUM spicis subdigitatis approximatis erectis filiformibus; rachi flexuosa, dentibus bifloris: altero sessili.*

*filiforme.*

*Habitat in America septentrionali. Kalm.*

*Folia lanceolata. Culmi pedales filiformes, superne nudi erecti. Spica filiformis, rara, flexuosa rachi alternatim utrinque ad dentes Flosculi 2. altero sessili, altero pedicellato; spicæ tenues, primum valde approximatae quasi in unum, quamvis alternæ.*

16. *PANICUM spicis digitatis subquaternis linearibus, flosculis solitariis secundis muticis.*

*lineare.*

*Habitat in Indiis.*

*Culmi prostrati, leves, ramosi. Spiculæ lineares, rectæ, angustæ. Flores subtus alterni. Calycis squama exterior brevior, patens, rachi adhaerens.*

*\* \* Paniculata.*

17. *PANICUM paniculis simplicibus, culmo ramo dichotomo. Gron. virg. 133. \**

*dichotomum.*

*Habitat in Virginia.*

*Statura refert arbusculam inferne simplicem; superne fasciculatam.*



- grosularium. 18. **PANICUM** panicula ramis simplicibus, floribus geminis: pedicello altero brevissimo, altero longitudine floris. *Amæn. acad. 5. p. 392. \**  
*Habitat in Jamaica.*
- clandestinum. 19. **PANICUM** racemis occultatis intra foliorum vaginas.  
 Gramen cyperoides polystachyon, spicis ad nodos ex foliorum alis prodeuntibus. *Sloan. jam. 36. hist. 1. p. 120. t. 80.*  
*Habitat in Jamaica, Pennsylvania. Kalm. 4*  
 Culmi dichotomi, ramosi. Folia ovato-lanceolata, vaginis punctatis, pillosis, intra quas Racemi fructificationum omnino occultantur, ut extus ne vestigium earum appareat; ceterum valde affine P. latifolio.
- capillare. 20. **PANICUM** panicula capillari erecta patente, caule hirsuto. *Gron. virg. 13.*  
 Panicum milliaceum viride, foliis latis brevibus, panicula capillacea. *Sloan. jam. 35. hist. 1. p. 115. t. 72. f. 3.*  
 Milium indicum, panicula sparsa erecta. *Tournef. injt. 515.*  
*Habitat in Virginia, Jamaica. ☉*  
*Simile Poæ capillari, sed totum hirsutum.*
- patens. 21. **PANICUM** panicula oblonga patente, calycibus bifloris, foliis lineari lanceolatis.  
 Triama-pullu. *Rheed. mal. 12. p. 75. t. 41.*  
*Habitat in India: similis e Lusitania.*
- divaricatum. 22. **PANICUM** paniculis brevibus muticis, culmo ramosissimo divaricatissimo, pedicellis bifloris: altero longiore. *Amæn. acad. 5. p. 392. \**  
*Habitat in Jamaica.*
- millaceum. 23. **PANICUM** panicula laxa flaccida, foliorum vaginis pubescentibus. *Hort. cliff. 27. Hort. ups. 20. Nat. med. 34. Roy. lugdb. 55. Dalib. parif. 22.*  
 Milium semine luteo & albo. *Bauh. pin. 26. theatr. 502.*  
*Habitat in India.*
- latifolium. 24. **PANICUM** panicula racemis lateralibus simplicibus, foliis ovato-lanceolatis collo pilosis.  
 Gramen virginianum, lato brevique folio, panicula rariore. *Morif. hist. 3. p. 196. f. 8. t. 5. f. 4.*

**Gramen miliaceum sylvaticum**, maximum, semine albo. *Sloan. jam. 34. hist. 1. p. 114. t. 71. f. 3.*

*Habitat in America.*

**Folia latitudine Commelinæ**, ad fauces amplexicaulia, extus collo circum fauces villosa, etiam basi foliorum margine pilosa. Panicula valde mediocris ex racemis lateralibus, non subdivisis, sed pedicellos proprios, nec subdivisos proferentes. Flores mutici, majusculi.

25. **PANICUM paniculatum**, foliorum vaginis longi- brevifolium  
tudinaliter ciliatis.

**Panicum miliaceum**, latiore folio, madêraspatanum.

*Pluk. aln. 176. t. 189. f. 4.*

*Habitat in India.*

**Gramen pedale vel bipedale.** Folia ovata, amplexicaulia. Panicula tenuissima, folio supremo & majore compresso. Variat foliis angustis & lanceolatis.

26. **PANICUM paniculatum ramosissimum**, foliis ovato- arborescens.  
oblongis acuminatis. *Fl. zeyl. 43.*

**Panicum panicula laxa sparsa**, foliorum vaginis glabris. *Hort. cliff. 27.*

*Habitat in India. b*

**Altitudine certat cum altissimis arboribus.** Culmus vix penna anserina latior.

27. **PANICUM panicula virgata**, foliis divaricatis.

repens.

*Habitat in Hispania? inde missum a Claud. Alstræmer. 24*

**Culmi repentes pedales adscendentes.** Folia inferiora latiora, undique caulem vaginis vestientia, patentissima & fere reflexo-divaricata; mucronata vaginis subpubescentibus: Superiora folia angustiora, longiora, erecta. Panicula constans tribus quatuorve racemis erectis, vix ramosis, quibus Flores alterni, remotiusculi, ovati; valvula calycina exteriore brevissima obtusa.

28. **PANICUM panicula virgata**, glumis acuminatis virgatum.  
lævibus: extima dehiscente.

**Panicum paniculatum**, glumis acutis. *Gron. virg. 133.*

*Habitat in Virginia.*

## PHLEUM.

1. **PHLEUM spica cylindrica longissima culmo recto.**

pratense.

F 4

Fl.

- Fl. lapp.* 26. *Fl. fvec.* 50. 56. *Hort. cliff.* 22. *Roy. lugdb.* 53. *Dalib. parif.* 20.
- Gramen thyphoides asperum primum. *Bauh. pin.* 4.  
 Gramen typhoides maximum, spica longissima. *Bauh. pin.* 4. *Scheuch. gram.* 60. *Morif. hist.* 3. p. 188. f. 8. t. 4. f. 1. *Vgill. parif.* 83.  
*Habitat in Europæ versuris* ☞ *pratis.* 24
- nodosum. 2. PHLEUM spica cylindrica basi sterili, culmo adscendente, radice bulbifera.  
 Gramen typhoides asperum alterum. *Bauh. pin.* 4. *theatr.* 52. *Scheuch. gram.* 62.  
 Gramen nodosum, spica parva. *Bauh. pin.* 2. *prodr.* 6. *theatr.* 20.  
 Gramen typhinum supinum tuberosum, spica aspera. *Bar. ic.* 53.  
*Habitat in Gallia, Helvetia, Italia.* 24  
*Simillimum P. pratensis, sed Radix bulbosa. Culmus non rectus, sed adscendens, foliorum vaginis tectus. Folia bifaria, obliqua, lævia exceptis marginibus scabris.*
- alpinum. 3. PHLEUM spica ovato-cylindracea. *Fl. lapp.* 25. *Fl. fvec.* 51. 57. *Dalib. parif.* 20.  
 Gramen typhoides alpinum, spica brevi densa & velut villosa. *Scheuch. gram.* 64. t. 3. *prodr.* 3.  
*Habitat in Alpibus Lapponicis, Helveticis aliisque.* 24
- arenarium. 4. PHLEUM spica ovata ciliata, caule ramoso. *Fl. fvec.* 2. n. 58. *Hort. cliff.* 23. *it. scan.* 109. *Roy. lugdb.* 54.  
 Gramen typhinum maritimum minus. *R. Pluk. alm.* 177. t. 33. f. 8. *Scheuch. gram.* 63.  
*Habitat in Europæ locis arenosis.* ☉  
*Spica vix absolute e vagina Folii exit.*
- schoenoides. 5. PHLEUM spicis ovatis obvolutis, foliis brevissimis mucronatis amplexicaulibus.  
 Gramen maritimum typhinum, brevi & crassiori spica ad singula genicula prodeunte. f. secundum. *Scheuch. gram.* 86.  
*Habitat in Italia, Smyrnæ, inque Hispania. Lasting.*  
*Habitus Schœni aculeati foliis floralibus amplexicaulibus, brevibus, mucronatis.*

## ALOPECURUS.

- pratensis. 1. ALOPECURUS culmo spicato erecto, glumis villosis.

lofis. *Roy. lugdb.* 54. *Fl. suec.* 52. 59. *Dalib. parif.* 21.

*Gramen phalaroides majus f. italicum. Bauh. pin.* 4. *thatt.* 55. *Morif. hist.* 3. f. 8. t. 4. f. 8.

*Gramen spicatum, spica cylindrica tenuissima longiore. Scheuch. gram.* 69.

*Gramen alopecuro simile glabrum cum pilis longiusculis in spica; Onocordon. Bauh. hist.* 2. p. 475.

*Habitat in Europæ pratis.* 2.

1. **ALOPECURUS** culmo spicato erecto, glumis nudis. agrestis.

*Gramen typhoides, spica angustiore. Bauh. pin.* 4. *thatt.* 53.

*Gramen typhinum, plantaginis spica, aristis geniculatis. Barr. ic.* 699. f. 2.

*Gramen spicatum, spica cylindracea tenuissima longiore. Scheuch. gram.* 69.

*Habitat in Europa australi.* 2.

*Simillimum A. pratensi, sed spica glabra.*

2. **ALOPECURUS** culmum spicato infracto. *Fl. suec. geniculatus.* 53. 60. *Dalib. parif.* 20.

*Alopecurus culmo infracto, aristis gluma longioribus. Roy. lugdb.* 54.

*Alopecurus aristis gluma longioribus. Fl. lapp.* 37.

*Gramen aquaticum geniculatum spicatum. Bauh. pin.* 3. *Scheuch. gram.* 72.

3. *Alopecurus aristis gluma æqualibus. Fl. lapp.* 38.

*Habitat in Europæ uliginosis.* 2.

4. **ALOPECURUS** panicula subspicata, glumis scabris, monspeliensis.  
corollis aristatis.

*Alopecurus panicula villosa oblonga folio involuta. Sauv. monsp.* 37.

*Gramen alopecurum majus, spica virescente divulsa pilis longioribus. Barr. ic.* 115. f. 2. *Scheuch. gram.* 155.

*Gramen alopecuroides anglo-britannicum maximum. Bauh. pin.* 4.

*Alopecurus altera maxima anglica paludosa. Morif. hist.* 3. p. 191. f. 8. t. 4. f. 3.

*Habitat in Angliæ, Galliæ humenibus.* ☉

*Similis A. paniceo, sed triplo major. Culmi & Folia rigida. Glumæ scabræ, ore levi, utraque valvula aristata; Corollæ valvula aristâ brevior.*

- panicetus.** 5. **ALOPECURUS** panicula subspicata, glumis villosis, corollis aristatis.  
*Cynosurus panicus. Sp. pl. 1. p. 73. Fl. suec. 54. 90.*  
*Gramen alopecurum minus, spica longiore. Bauh. pin. 4. theatr. 57. Scheuch. gram. 154.*  
*Gramen alopecurum minus, spica virescente divulsa. Barr. ic. 115. f. 1.*  
*Habitat in Europæ cultis aridis. ☉*  
*Simillimus priori, sed palmaris, mollis. Glumæ pubescentes ore villosa; Corollæ arista calycinis etiam brevior.*
- hordeiformis.** 6. **ALOPECURUS** racemo simplici, flosculis aristis circumvallatis.  
*Habitat in India.*  
*Habitus omnino Hordei murini. Culmus pedalis. Folia plano-canaliculata, nervosa, culmo sæpius longiora. Spica racemosa, longa. Flores numerosi; pedicelli florum brevissimi, simplicissimi. Calyces obvallati setis aristiformibus, numerosis, floribus triplo longioribus, gluma bivalvi, altera minore. Corolla bivalvis glumis oblongis, acuminatis. Pistillum flore longius. Stamina tria.*

## MILIUM.

- effusum.** 1. **MILIUM** floribus paniculatis dispersis muticis. *Fl. suec. 55. 61. Dalib. parisi. 23.*  
*Milium glumis diphyllis. Hort. cliff. 27.*  
*Gramen sylvaticum, panicula miliacea sparsa. Bauh. pin. 8. theatr. 141. Scheuch. gram. 133. Moris. hist. 3. f. 8. t. 5. f. 10.*  
*Habitat in Europæ nemoribus umbrosis. 24*
- confertum.** 2. **MILIUM** floribus paniculatis confertis. *Roy. lugdb. 57.*  
*Gramen paniculatum alpinum latifolium, panicula miliacea sparsa. Scheuch. gram. 134. Hall. helv. 220.*  
*Habitat in Helvetiæ sylvis.*
- paradoxum.** 3. **MILIUM** floribus paniculatis aristatis.  
*Agrostis paradoxa. Sp. pl. 1. p. 62.*  
*Gramen avenaceum paniculatum galloprovinciale, aquilegiæ semine. T. Moris. hist. 3. p. 214. Fluk. alm. 174. t. 32. f. 2.*  
*Habitat in Gallia meridionali.*  
*Gramen sesquipedale. Panicula alba. Olyx laxus. Semen-*

*men nudum, nigrum, nitidum, ovatum, apice aristata caduca.*

4. MILIUM panicula subspicata, floribus aristatis. *lendigerum.*  
 Gramen terotinum arvense, panicula pyramidalis.  
*Raj. hist. 1288. Scheuch. gram. 148.*  
 Gramen loliaceum, panicula ramosa lutea. *Magn. monsp. 297. Moris hist. 3. p. 182. f. 8. t. 3. f. 12.*  
 Gramen alopecuro accedens ex culmi geniculis spicas in petiolis longiusculis promens. *Pluk. alm. 177. t. 33. f. 6.*  
*Habitat in Monspelii, Gouan. ☉*
5. MILIUM paniculae ramis simplicissimis, floribus punctatum.  
 alternis secundis geminis. *Amæn. acad. 5. p. 392. \**  
*Habitat in Jamaica.*

AGROSTIS,

\* *Aristata.*

1. AGROSTIS petalo exteriori exferente aristam rectam strictam longissimam. *Roy. lugdb. 59. Fl. fvec. spica venti.*  
*58. 62.*  
 Gramen segetum altissimum, panicula sparsa. C. B.  
*Scheuch. gram. 144.*  
 Gramen capillatum. *Bauh. hist. 3. p. 462.*  
*Habitat in Europa inter segetes. ☉*
2. AGROSTIS petalo exteriori arista terminali recta miliacea.  
 stricta mediocri.  
*Agrostis miliacea. Roy. lugdb. 59. Sp. pl. 1. p. 61. n. 2.*  
*Habitat in Hispania. Læfing; in Sibiria. Gmelin. 24*  
*Similis A. Spica venti, sed arista flosculis longitudine tantum equalis. Culmi e radice plures. Panicula minus ampla. Cultura etiam diversissima perstitit.*
3. AGROSTIS panicula oblonga, petalo exteriori basi villosa, arista torta calyce longiore. *Fl. fvec. 59, 63. \**  
*Agrostis culmo enodi, panicula contracta. Fl. lapp. 44.*  
 Gramen miliaceum sylvestre, glumis oblongis. *Pet. gram. 121. Raj. angl. 3. p. 404.*  
 Gramen avenaceum, panicula acerosa, semine papposo. *Dill. app. 48.*

Gra-

*Gramen avenaceum montanum*, panicula angusta e dilutissimo fusco albicante & papposa. *Scheuch. gram.* 507.

*Habitat in Europæ monticulis silvosis glareosis juniperetis.* 24.

Calamagrostis.

4. *AGROSTIS* panicula incrassata, petalo exteriori toto lanato apice aristato, caule ramofo.

*Gramen arundinaceum*, panicula densa viridi-argentea splendente aristata. *Scheuch. gram.* 146. *Hall. helv.* 222. *Segu. veron.* 3. p. 163.

*Habitat in Helvetiæ, Veronæ alpinis.* 24

*Similis A. arundinaceæ sed corolla tota valde hirsuta: aristula recta non e basi, sed apice petali. Culmi ramofo. Panicula densa.*

interrupta.

5. *AGROSTIS* petalo aristato, panicula attenuata contracta interrupta. †

*Gramen arvense paniculatum*, spica interrupta. *Scheuch. gram.* 146.

*Gramen capillatum*, paniculis interruptæ angustioribus. *Vaill.* 88. t. 71. f. 4.

*Habitat in Gallia, Italia.*

rubra.

6. *AGROSTIS* paniculæ parte florente potentissima, petalo exteriori glabro terminato arista tortili recurva. *Fl. succ.* 60. 64. *Dalib. parif.* 24. non synonyma.

*Agrostis* panicula inferiore verticillatim laxa, superiore contracta. *Fl. lapp.* 46.

*Gramen alopecuro accedens.* *Pluk. alm.* 277. t. 33. f. 6.

*Habitat in Svecia vulgaris.* 24

*Paniculæ verticilli florentes horizontales, nondum florentes contracti quasi in spicam, deflorata tota rubescens.*

canina.

7. *AGROSTIS* calycibus coloratis, petalorum arista dorsalis recurva, culmis prostratis subramosis. *Fl. succ.* 1138. 65.

*Gramen supinum caninum paniculatum*, folio varians. *Bauh. pin.* 1. theatr. 12.

*Gramen paniculatum supinum ad infima culmorum genicula foliorum capillarium geniculis fasciculatis donatum.* *Scheuch. gram.* 141.

*Gramen caninum supinum paniculatum folio varians.* *Bauh. pin.* 1.

Ha-

*Habitat in Europæ pascuis humidiusculis. 24.*  
*Remota purpurascens lucens facilius observatur, quam proxima.*

\* \* *Mutica.*

8. AGROSTIS paniculæ ramulis divaricatis muticis. *stolonifera.*  
 culmo ramofo repente, calycibus æqualibus. *Fl. fvec. 2. n. 66.*

Agrostis culmo repente foliis radicalibus brevior, follii suprema vagina ventricosa, flosculis muticis.  
*Roy. lugb. 59. Fl. fvec. 1. n. 61.*

Agrostis culmo repente, vagina supremi follii ventricosa. *Roy. lugdb. 59. Dalib. parif. 23.*

Gramen caninum supinum minus. *Vaill. parif. 86. Scheuch. gram. 128.*

- β. Gramen radice repente, panicula densa spicata spadiceo-viridi, locustis exiguis muticis. *Scheuch. gram. 130.*

*Habitat in Europa. β. in arena mobilis. 24.*

9. AGROSTIS panicula capillari patente, calycibus subulatis æqualibus hispidiusculis coloratis, flosculis muticis. *capillaris.*  
*Roy. lugdb. 59. Dalib. parif. 23.*

Agrostis panicula tenuissima. *Fl. lapp. 45.*

Agrostis panicula capillari patente, spiculis coloratis muticis. *Scop. carn. 188.*

Gramen montanum, panicula spadicea delicatior.  
*Bauh. pin. 3. prodr. 12. Scheuch. gram. 129.*

*Habitat in Europæ pratis.*

Panicula vere capillaris, tenuissimisque pedicellis singularis.

10. AGROSTIS panicula laxa, calycibus muticis æqualibus. *alba.*  
*Roy. lugdb. 50.*

Gramen miliaceum majus, panicula spadicea & viridi. *Pet. gram. 118. 119. Raj. angl. 3. p. 404.*

*Habitat in Europæ nemoribus.*

11. AGROSTIS panicula mutica filiformi. *minima.*

Nardus panicula spicata, floribus solitariis alternis  
*Guett. stamp. 155. Dalib. parif. 155.*

Gramen spartum, capillaceo folio, minimum. *Dill. app. 173.*

Gramen minimum, paniculis elegantissimis. *Bauh. pin. 2. Scheuch. gram. 40. Morif. hist. 3. p. 200. f. 8. t. 2. f. 10.*

Gra-



Gramen minimum. *Bauh. hist.* 2. p. 465. *Dalech. hist.* 425.

*Habitat in Gallia, Germania.*

- virginica. 12. AGROSTIS panicula contracta mutica, foliis involuto-subulatis rigidis exstantibus. *Clayt. virg.* 507. *Habitat in Virginia.*

- indica. 13. AGROSTIS panicula contracta mutica, racemis lateralibus erectis alternis.

Gramen pratense, panicula & foliis angustissimis, spicis brevibus muticis, locustis minimis. *Sloan. jam.* 35. *hist.* 1. p. 115. t. 73. f. 1.

*Habitat in India.*

Culmus erectus. Folia plana angusta. Panicula oblonga & racemis lateralibus, alternis, rachi approximatis; glumis brevissimis, acutis.

\* \* \* *Cruciatae.*

- radiata. 14. AGROSTIS spicis subquinis cruciatis basi villosis valvulis petaloideis aristatis. *Aman. acad.* 5. p. 392. \*

Gramen cruciatum assurgens, spicis subhirsutis tenuioribus & longioribus deorsum frugiferis. *Brown. jam.* 137.

Gramen Dactylon spicis gracilioribus plerumque quatuor cruciformiter dispositis. *Sloan. jam.* 33. *hist.*

1. p. 111. t. 68. f. 3.

*Habitat in Indiis.*

- cruciata. 15. AGROSTIS spicis quaternis cruciatis basi glabris; valvulis petaloideis aristatis.

Gramen Dactylon bicornis minimum aristis longis armatum. *Sloan. jam.* 34. *hist.* 1. p. 112. t. 69. f. 1.

*Habitat in Jamaica.*

## AIRA.

\* *Mutica.*

- indica. 1. AIRIA spica subovata secunda, flosculis obtusis muticis, foliis planis.

*Habitat in India.*

Gramen procumbens, digiti longitudine. Folia plana.

Spica secunda, conglomerata, ovato-oblonga. Glumæ obtusæ, muticæ.

- crisata. 2. AIRA panicula spicata, calycibus subtrifloris peduncul-

- culo longioribus, petalis subaristatis inæqualibus.  
*Roy, lugdb. 61. Dalib. parif. 27.*  
 Gramen, spica cristata subhirsutum. *Bauh. pin. 13. prod. 8. Morif. hist. 3. p. 194. f. 8. t. 4. f. 7. Scheuch. gram. 166.*  
*Habitat in Angliæ. Galliæ; Helvetiæ fœccioribus.*
3. **AIRA** foliis planis, panicula coarctata, floribus pedunculatis muticis convoluto-subulatis. *Fl. fœc. 71. 67. \* Dalib. parif. 26.* cerulea.  
 Poa spiculis subulatis, panicula rara contracta. *Fl. lapp. 29. Roy. lugdb. 61.*  
 Gramen arundinaceum enode minus sylvaticum. *Bauh. pin. 7. theatr. 97. Scheuch. gram. 209.*  
 Gramen pratense serotinum, panicula rara purpurascens. *Morif. hist. 3. p. 201. f. 8. t. 5. f. 22.*  
*Habitat in Europæ pascuis aquosis. 24.*  
*Variat flosculis quatuor in fertiliori solo.*
4. **AIRA** panicula oblonga secunda mutica imbricata, arundinacea, foliis planis.  
 Gramen orientale paniculatum, portulacæ semine. *Tournef. cor. 39.*  
*Habitat in Oriente.*
5. **AIRA** panicula laxa subfastigiata ramosissima, flosculis muticis. *Læfl. it. 117. \** minuta.  
 Gramen paniculatum arvense minimum. *Buxb. cent. 5. p. 35. t. 67. ?*  
*Habitat in Hispania. Læfling. ☉*  
*Differt tantum a Buxbaumiano, quod longe minus, vix pollicem transversum altum.*
6. **AIRA** foliis planis, panicula patente, floribus muticis lævibus calyce longioribus. *Fl. fœc. 66. 68. Dalib. parif. 26.* aquatica.  
 Aira culmo inferiore repente, flosculis muticis calyce longioribus: altero pedunculato. *Roy. lugdb. 60.*  
 Gramen aquaticum miliaceum. *Scheuch. gram. 176. Vaill. parif. 89. t. 17. f. 7.*  
 Gramen caninum supinum paniculatum dulce. *Bauh. pin. 1.*  
*Habitat in Europæ pascuis aquosis. 24*  
*\* \* Aristata.*
7. **AIRA** foliis planis, panicula spicata, flosculis medio aristatis: aristâ reflexa laxiore. *Fl. fœc. 65. 69. \** spicata.  
 Aira

*Aira panicula spicata, pedicellis flosculo brevioribus.*  
*Fl. lapp. 47.*

*Gramen avenaceum paniculatum alpinum humile, locustis in spicam collectis varicoloribus aristatis.*

*Scheuch. gram. 221. prodr. 24. t. 6.*

*Habitat in Lapponiæ alpibus. 24.*

caespitosa. 8. *AIRA* foliis planis, panicula patente, petalis basi villosis aristatisque: arista recta brevi. *Fl. suec. 63. 70.*

*Aira panicula patentissima, flosculis subaristatis sessilibus basi villosis, foliis planis. Mor. afl. Stockh. 1742. p. 30.*

*Aira panicula longissima tenui. Fl. lap. 50.*

*Gramen fegetum, panicula arundinacea. Bauh. pin. 3. theatr. 35. Scheuch. gram. 244.*

*Gramen pratense paniculatum, locustis parvis splendentibus non aristatis. T. Vuill. parif. 26.*

*Gramen agrorum, latiore arundinacea comosa panicula. Morif. hist. 3. f. 8. t. 5. f. 17.*

*Habitat in Europæ pratis cultis & fertilibus. 24.*

flexuosa. 9. *AIRA* foliis setaceis, culmis subnudis, panicula divaricata, pedunculis flexuosis. *Fl. suec. 64. 71. Dalib. parif. 26.*

*Aira panicula rara, calycibus albis. Fl. lapp. 28.*

*Gramen alpinum nemorosum paniculatum, foliis angustissimis, locustis, splendentibus aristatis. Scheuch. gram. 218. t. 6. f. 1.*

*Gramen nemorosum, paniculis albis, capillaceo folio. Bauh. pin. 7. prodr. 14. Morif. hist. 3. p. 300. f. 8. t. 7. f. 9.*

*Habitat in Europæ petris, rupibus. 24.*

montana 10. *AIRA* foliis setaceis, panicula angustata, flosculis basi pilosis aristatis: arista tortili longiore. *Fl. suec. 68. 72.*

*Aira panicula rara, calycibus fuscis. Fl. lap. 49.*

*Gramen avenaceum paniculatum alpinum, foliis capillaceis brevibus, locustis purpureo argenteis & aristatis. Scheuch. gram. 216. t. 4. f. 4.*

*Gramen avenaceum capillaceum, minoribus glumis. Bauh. pin. 10.*

*Habitat in Europæ alpinis. 24.*

alpina. 11. *AIRA* foliis subulatis, panicula densa, flosculis basi pilosis aristatis: arista brevi. *Fl. suec. 69. 73.*

Gra-

Gramen avenaceum capillaceum, minoribus glumis argenteis vix aristatis, foliis angustioribus Scheuch. gram. 222.

Habitat in alpidibus Lapponicis, Germania.

11. AIRA foliis setaceis: summo spathaceo paniculam canescens. inferne obvolvente. Roy. lugdb. 60. Fl. succ. 73.

74. Dalib. parif. 25.

Aira foliis setaceis, arista a basi glumarum calycem æquante. Hort. cliff. 495.

Gramen foliis junceis, radice jubata. Bauh. pin. 5. Scheuch. gram. 243.

Gramen sparsum variegatum. Bauh. pin. 5.

Habitat in Scania & australioris Europæ arvis arenosis. ☉

Arista in medio cincta denticulis, infra crassior fusca, supra tenuior subclavata albidâ. Panicula ante florescentiam spicata.

12. AIRA foliis setaceis: vaginis angulatis, floribus paniculato-spicatis, flosculis basi aristatis. It. scan. 226. \* Fl. succ. 2. n. 75. præcox.

Gramen minimum, spica brevi habitiore, nostrum. Scheuch. gram. 249.

Gramen parvum præcox, panicula laxa canescente. Raj. ang. 3. p. 407. t. 22. f. 2. Pluk. alm. 177. t. 33. f. 9.

Habitat in Europæ australioris campis arenosis inundatis. ☉

14. AIRA foliis setaceis, panicula divaricata, floribus aristatis distantibus. caryophyll-  
lea.

Aira foliis subulatis, aristis a basi flosculorum tortilibus, flosculo altero pedunculato. Roy. lugdb. 60. Dalib. parif. 26.

Gramen phalaroides, sparsa panicula, minimum angustifolium. Barr. rar. 1218. t. 44. f. 1.

Gramen paniculatum, locustis purpureo argenteis, annuum. Raj. angl. 3. p. 407. Morij. hist. 3. p. 200. f. 8. t. 5. f. 11.

Gramen paniculatum minimum molle. Magn. monsp. 296. Scheuch. gram. 215.

Caryophyllus arvensis glaber minimus. Bauh. prodr. 105.

Habitat in Angliæ, Germaniæ, Galliæ glareosis. ☉

## M E L I C A.

1. MELICA flosculi inferioris petalo exteriori ciliato. ciliata.

It. gotl. 297. Fl. succ. 56. 77.

Linnei Spec. plant. pars. L. G

Me-

- Melica floribus horizontaliter patentibus. Roy. lugdb. 57.*  
*Gramen avenaceum montanum lanuginosum. Bauh. pin. 10. prodr. 20.*  
*Gramen avenaceum, spica simplici, locustis densissimis candicantibus & lanuginosis. Scheuch. gram. 174.*  
*Habitat in Europæ collibus sterilibus saxosis. 24*
- nucans. 2. *MELICA* petalis imberbibus, panicula nutante simplici.  
*Melica petalis imberbibus. Fl. suec. 57. 76. Dalib. paris. 23.*  
*Melica floribus sub culmo pendulis. Fl. lapp. 36. Roy. lugdb. 57.*  
*Gramen montanum avenaceum, locustis rubris. Bauh. pin. 10. prodr. 20.*  
*Gramen montanum avenaceum spicatum. Bauh. pin. 10.*  
*Gramen avenaceum, locustis rarioribus. Bauh. pin. 10. prodr. 20.*  
*Habitat in Europæ frigidioris rupibus. 24*
- altissima. 3. *MELICA* petalis imberbibus, panicula ramosissima.  
*Hort. ups. 20.*  
*Melica flosculis glabris: summo urceolari. Gmel. sib. 1. p. 98. t. 20.*  
*Gramen avenaceum, locustis rarioribus muticis, virginianum majus. Moris. hist. 3. p. 216. f. 8. t. 7. f. 51.*  
*Habitat in Sibiria, Canada. 24*

## P O A.

- aquatica. 1. *POA* panicula diffusa, spiculis sexfloris linearibus. *Fl. suec. 73. 78. Dalib. paris. 29.*  
*Poa panicula diffusa, spiculis sexfloris linearibus muticis compressis. Afl. Stockh. 1741. p. 185.*  
*Poa panicula contracta, spicis ovatis teretiusculis. Hort. cliff. 494. Roy. lugdb. 61.*  
*Gramen palustre paniculatum altissimum. Bauh. pin. 2. theatr. 38. Scheuch. gram. 191.*  
*Gramen aquaticum paniculatum latifolium. Bauh. pin. 3. Moris. hist. 3. p. 201. f. 8. t. 6 f. 25.*  
*Habitat in Europa ad ripas piscinarum, fluviorum. 24*
- palustris. 2. *POA* panicula diffusa, spiculis subtrifloris, foliis subtus scabris +  
*Gramen paniculatum aquaticum angustifolium, panicula speciosa. Scheuch. gram. 184.*

**Gramen palustre, panicula speciosa.** *Bauh. pin. 3. prodr.*

7 *Morif. hist. 3. p. 201. f. 8. t. 6. f. 27.*

*Habitat in Helvetiæ, Italiæ paludibus.*

*Dignoscitur, dum runcantur agri oryzaei, tenella foliis glaucis, adultior spinulis foliorum. Segu. veron.*

1. p. 351.

3. **POA panicula diffusa ramosissima, spiculis sexfloris alpina.**  
cordatis. *Fl. suec. 79. Dalib. parif. 28.*

**Gramen alpinum paniculatum majus, panicula speciosa**  
variegata. *Scheuch. gram. 186. Hall. helv. 214.*

β. **Gramen alpinum latifolium, panicula laxa foliacea, fo-** *vivipara.*  
liolis in panicula paucioribus & magis crispis. *Scheuch.*  
*gram. 212.*

*Habitat in alpihus Lapponicis, Helveticis. 24*

*An hæc α. β. sequentis tantum varietas?*

4. **POA panicula subdiffusa, spiculis trifloris basi pube-** *trivialis.*  
scentibus, culmo erecto tereti. *Fl. suec. 78. 80. Dalib.*  
*parif. 28.*

**Gramen pratense paniculatum medium.** *Bauh. pin. 2.*  
*Scheuch. gram. 180.*

*Habitat in Europæ pascuis. 24*

5. **POA panicula diffusa, spiculis quadrifloris pubescen-** *angustifolia.*  
tibus, culmo erecto tereti. *Fl. suec. 77. 81. Dalib.*  
*parif. 28.*

**Poa spiculis ovato-oblongis, foliis subulatis** *Fl. lapp. 52.*

**Gramen pratense paniculatum majus, angustiore folio.**

*Bauh. pin. 2. prodr. 5. Scheuch. gram. 178.*

*Habitat in Europa ad agrorum versuras. 24*

6. **POA panicula diffusa, spiculis quinquefloris glabris,** *pratensis.*  
culmo erecto tereti. *Fl. suec. 76. 82. Dalib. parif. 27.*

**POA spiculis ovatis compressis muticis.** *Fl. lapp. 51.*

*Hort. cliff. 27. Roy. lugdb. 61.*

**Gramen pratense paniculatum majus (latiore folio.** *Bauh.*  
*pin. 2.) Scheuch. gram. 177.*

*Habitat in Europæ pratis fertilissimis.*

*Limites pro speciebus 3. 4. 5. 6. rite discernendis vix*  
*dum perspicue dati sunt.*

7. **POA panicula diffusa angulis rectis, spiculis obtusis,** *annua.*  
culmo obliquo compresso. *Fl. suec. 75. 83. Dalib.*  
*parif. 28.*

**Gramen pratense paniculatum minus.** *Bauh. pin. 2.*  
*theatr. 31.*

- Gramen pratense minus vulgatissimum. Raj. hist. 1285. \**  
 β. *Gramen pratense paniculatum minus rubrum. Tournef. inst. 522.*  
*Habitat in Europa ad vias. ☉*
- flava.* 8. POA panicula diffusa, spiculis ovato-oblongis nitidis.  
*Gron. virg. 13.*  
*Gramen pratense majus virginianum. Pet. musf. 239.*  
*Habitat in Virginia.*
- pilosa.* 9. POA panicula patente stricta: ramificationibus primariis pilosis.  
*Gramen, paniculis elegantissimis, majus, locustis purpureo-spadiceis minoribus. Scheuch. gram. 193.*  
*Habitat in Italia.*
- amabilis.* 10. POA panicula patente, spiculis octifloris linearibus.  
*Fl. Zeyl. 46. \**  
*Habitat in India.*
- Eragrostis.* 11. POA panicula patente, pedicellis flexuosis, spiculis ferratis decemfloris.  
*Gramen, paniculis elegantissimis, minimum. Scheuch. gram. 192.*  
*Gramen Phalaroides, sparsa brizæ panicula, minus. Barr. ic. t. 44. f. 2.*  
*Habitat in Italia supra muros, D. Bæck.*  
*Affinis Brizæ Eragrosti; statura Poæ annuæ. Culmi decumbentes. Folia pilosa.*
- capillaris.* 12. POA panicula laxa patentissima capillari, foliis pilosis, culmo ramosissimo.  
*Poa culmo subcompresso, panicula tenuissima, spiculis trifloris minimis, flosculis basi pubescentibus. Kalm. it. 3 p. 291.*  
*Poa panicula laxa erecta stricta, spiculis erectis oblongis. Gron. virg. 136.*  
*Gramen paniculatum virginianum, locustis minimis. Moris. hist. 3. p. 202. f. 8. t. 6. f. 33.*  
*Habitat in Virginia, Canada. P. Kalm.*  
*Panicula capillacea, strictissima, patentissima, ramosissima. Spicula ex flosculis 4. 5. f. 6 minima. Folia ad basin pilis longis raris. Culmus ramosus.*
- malabarica.* 13. POA paniculæ ramis simplicissimis, floribus sessilibus, seminibus distantibus, culmo repente.  
*Habitat in Indiæ arenosis.*
- chunensis.* 14. POA paniculæ ramis simplicissimis, floribus sessilibus, seminibus imbricatis, culmo erecto.

*Habitat in India. Osbeck.*

*Rami paniculae simplicissimi, ut in antecedente, magis conferti, Flores minores; Flosculi 4, nec septem.*

15. POA panicula oblonga capillari subverticillata, flosculis sexfloris minutissimis. tenella.

*Gramen paniculatum ex oris malabaricis, panicula delicatior. Pluk. alm. 176. t. 300. f. 2.*

*Gramen paniculatum, locustis tenuissimis subrotundis. Burm. Zeyl. 110. t. 47. f. 3.*

*Gramen fumi. Rumph. amb. 6. t. 4. f. 3.*

*Tiama-pullu. Rheed. mal. 12. p. 75. t. 4.*

*Habitat in India.*

*Gramen bipedale, Foliis longis. Panicula angustata, ramulis confertissimis, subramosis ut fere verticillata appareat; Flores tam parvi, ut nudis oculis vix flosculi numerentur ante maturitatem.*

16. POA panicula lanceolata subramosa, floribus alternis secundis. Amæn. acad. 4. p. 265. \* Roy. lugd. 62. Guett. stamp. 62.

*Gramen exile duriusculum in muris & aridis proveniens. Raj. angl. 3. p. 410.*

*Gramen arvense, filicina duriore panicula. Barr. ic. 49.*

*Gramen Loliaceum murorum duriusculum, spica erecta rigida. Moris. hist. 3. p. 182. f. 8. t. 2. f. 9.*

*Gramen panicula multiplici. Bauh. pin. 3. prodr. 6.*

*Habitat in Galia, Angliæ siccis. ☉*

17. POA panicula secunda coarctata, culmo obliquo compresso. Fl. svec. 74. 84. It. gotl. 178. Dalib. parif. 28.

*Poa spiculis ovatis compressis muticis, panicula secunda coarctata. Aët. stockh. 1741. p. 184.*

*Gramen murorum, radice repente. Bauh. pin. 2. prodr. 2.*

*Gramen paniculatum, radice repente, culmo compresso. Vaill. parif. 91. t. 18. f. 5.*

*Gramen paniculatum, radice repente, culmo compresso, locustis distichis pulchellis. Scheuchz. gram. 198.*

*Habitat in Europæ & Americæ septentrionalis siccis, muris, tellis. 24*

*Americana flosculis sæpius ternis, Europæa vero senis, habitus tamen idem & structura.*



- nemoralls. 18. POA panicula attenuata, spiculis sub-bisfloris muoronatis scabris, culmo incurvo. *Fl. fvec. 2. n. 85.*  
 Gramen sylvaticum, panicula tenui e raris locustis composita. *Dill. spec. 57.*  
 Gramen paniculatum angustifolium alpinum, locustis rarioribus & angustioribus non aristatis. *Scheuch. gram. 164. t. 2.*  
 Gramen nemorosum, panicula laxa, radice repente. *Vaill. parif. 90.*  
 Gramen foliaceum alpinum, paniculis ex rarioribus & exiguis locustis compositum. *Mont. gram. 39.*  
 β. Gramen cirrhosa & villosa spongia ad singula genicula donatum, paniculatum minimum alpinum. *Bocc. mus. 2. p. 70. t. 59.*  
*Habitat in Europa ad radices montium umbrosas. 24.*  
 bulbosa. 19. POA panicula secunda patentiuscula, spiculis quadrifloris.  
 Poa culmorum basi tuberosa. *Guett. stamp. 1. p. 168.*  
*Dalib. parif. 29.*  
 α. Gramen xerampelinum, miliacea prætenui ramosaque sparsa panicula f. Xerampelino congener arvense æstivum, Gramen minutissimo semine. *Lob. illustr. 14. Scheuch. gram. 185.*  
 β. Gramen arvense, panicula crispa. *Bauh. pin. 3. Barr. ic. 703.*  
 Gramen arvense angustifolium panicula densa foliacea: foliolis in panicula angustissimis. *Scheuch. gram. 211.*  
 γ. Gramen vernum, radice ascalonica. *Vaill. parif. 91. t. 17. f. 8.*  
*Habitat in Gallia, Hispania, Oriente, Nycopia Sveciæ.*  
*In α. culmi semper inclinati; in γ. Glumæ inæquales: aliæ latæ planæ, aliæ contrariæ longiores.*  
 ciliaris. 20. POA panicula glomerata, glumarum valvulis interioribus piloso-ciliatis. *Amæn. acad. 5. p. 392. \**  
 Briza tenuissima, paniculis quasi lanuginosis, pediculis brevibus & tenuissimis. *Brown. jam. 135.*  
 Gramen prætenue, panicula & foliis angustissimis, spicis brevibus muticis, locustis minimis. *Sloan. jam. 35. hist. 1. p. 114. t. 73. f. 1.*  
*Habitat in Jamaica.*

## BRIZA.

- minor. 1. BRIZA spiculis triangulis, calyce flosculis (7) longiore.  
*Hort. cliff. 23. Roy. lugdb. 63.*

Gra-

*Gramen tremulum minus, panicula parva. Bauh. pin.*

2. *prodr.* 4.

*Habitat in Helvetia, Italia. ☉*

2. *BRIZA* spiculis ovatis, calyce flosculis (7) æquali. *viens.*

*Gramen paniculatum minus, locustis magnis tremulis.*

*Tournef. inst.* 523.

*Habitat in Oriente, Hispania. ☉ Alstræmer.*

*Simillima B. mediæ, sed Folia duplo latiora, planiora, decurrentia per ligulam acutam. Panicula similis, copiosior, viridis nec purpurascens, ramulis magis ramosis, divaricatis, nodis callosis eos basi deprimentibus. Flores paullo minores, dum comprimuntur facile decidui.*

2. *BRIZA* spiculis ovatis, calyce flosculis (7) brevior. *media.*

*Hort. cliff.* 23. *Fl. suec.* 85, 86. *Roy. lugdb.* 63.

*Gramen tremulum majus. Bauh. pin.* 2. *Scheuch. gram.*

204.

*Habitat in Europæ prætis fœccioribus. 24*

4. *BRIZA* spiculis cordatis, flosculis septendecim. *Hort. maxima.*

*cliff.* 23. *Hort. upf.* 20. *Roy. lugdb.* 63.

*Gramen tremulum maximum. Bauh. pin.* 2. *Scheuch. gram.* 202.

*Habitat in Italia, Lusitania,*

5. *BRIZA* spiculis lanceolatis, flosculis viginti. *Eragrostis.*

*Briza panicula spicata, spiculis lanceolatis. Guett.*

*stamp.* 1. p. 163.

*Uniola calycibus diphyllis, spiculis ovato-lanceolatis.*

*Gron. virg.* 136.

*Gramen paniculis elegantissimis. Bauh. pin.* 2. *Scheuch.*

*gram.* 194. *Morif. hist.* 3. p. 204. f. 8. t. 6. f. 52.

*Gramen Erantheum f. Eragrostis Barr. rar.* t. 43.

*Habitat in Europa australi ad agrorum versuras.*

## UNIOLA.

1. *UNIOLA* paniculata, spiculis ovatis. *paniculata.*

*Uniola calycibus polyphyllis. Gron. virg.* 136.

*Uniola. Hort. cliff.* 23. *Roy. lugdb.* 63.

*Gramen myloicophorum oxyphyllon carolinianum.*

*Pluk. alm.* 173. t. 32. f. 6. *Cartesb. car.* 1. p. 32.

t. 32.

*Habitat in Carolina.*

- bipennata. 2. **UNIOLA** panicula subspicata, racemis pinnatis subtus imbricatis.  
*Briza bipinnata. Syst. nat. 10. p. 875. Aman. acad. 4. p. 450.*  
*Habitat in Ægypto. Hasselquist.*  
*Gramen omnium speciosissimum. Culmi magnitudine Tritici simplices. Racemus simplex, pedalis et Pedunculis alternis, simplicissimis, æqualibus, utrique longitudinaliter a basi ad apicem pinnatis. Floribus sessilibus, approximatis, compressis, æqualibus, complanatis more Cyperi. Glumæ compressæ, carinatæ, acutiusculæ, ovatæ.*
- mucronata. 3. **UNIOLA** spica disticha, spiculis ovatis, calycibus subaristatis.  
*Habitat in India. Burmannus.*  
*Culmus pedalis, levis. Folia angusta, glabra vaginis striatis. Spica disticha ex floribus 11. s. 12, alternis, distichis, subsessilibus, ovatis, glabris, septemfloris. Calyx mucronatus fere in aristam.*
- spicata. 4. **UNIOLA** subspicata, foliis involutis rigidis.  
*Gramen parvum, maritimum spicatum, foliis angustis rigidis. Clayton. 507.*  
*Habitat in Americæ borealis maritimis.*  
*Culmus spithameus, Foliis alternis rigidis involuto-mucronatis. Panicula minima, coarctata, ut vix signum pedicellorum, secunda; Calyx 3 glumæ carinatæ flosculis quatuor.*

## DACTYLIS.

- cynosurol des. 1. **DACTYLIS** spicis sparsis secundis scabris numerosis.  
*Lafl. it 115. \**  
*Gramen maritimum, spica crassa dactyloide terminali, odore rancido, culmo albo. Gron. virg. 135.*  
*Dactylis spicis alternis secundis incisis erectis approximatis, calycibus unifloris subulatis. Gron. virg. 134.*  
*Spartum eslexiense, spica gemina. Raj. angl. 3. p. 393.*  
*Habitat in Virginia, Canada, nunc in Lusitania, Anglia.*  
 2. *Culmus bipedalis, arundinaceus. Folia in culmo sex, lata, culmo longiora, levissima, margine scabra incurva, pagina interiore glauciora. Spicæ sex s. plurimæ, secundæ, divergentes, paleacæ; flosculis imbricatis latere postico scabris. Calycibus unifloris carina*

*vina scabris, mucronatis, flosculo longioribus, sessilibus, secundis. Pistillis villosis longioribus.*

2. DACTYLIS panicula secunda glomerata. *Fl. succ. glomerata.*  
*1. n. 87.*  
 Cynosurus panicula secunda glomerata. *Fl. succ. 2.*  
*n. 85.*  
 Poa flosculis confertis uno versu dispositis. *Hort. cliff.*  
*28. Roy. lugdb. 62.*  
 Gramen spicatum, folio aspero. *Bauh. pin. 3. prodr.*  
*9. Scheuch. gram. 299. Moris. hist. 3. f. 8. 1. 6. f. 38.*  
*Habitat in Europæ cultis ruderatis. 24*

### CYNOSURUS.

1. CYNOSURUS bracteis pinnatifidis. *Hort. cliff. 493. cristatus.*  
*Fl. succ. 81. 88. Roy. lugdb. 64.*  
 Gramen pratense cristatum, f. spica cristata lævi. *Bauh.*  
*pin. 2. prodr. 8. Scheuch. gram. 79.*  
 Gramen cristatum. *Bauh. pin. 2. p. 468.*  
*Habitat in Europæ pratis. 24*
2. CYNOSURUS bracteis pinnato - paleaceis aristatis. *echinatus.*  
*Roy. lugdb. 64.*  
 Gramen alopecuroides, spica aspera. *Bauh. pin. 4.*  
*Scheuch. gram. 80.*  
 Gramen alopecurum, spica aspera. *Bar. rar. 1. 123.*  
*Habitat in Europa australiori, Oriente.*
3. CYNOSURUS spica secunda, calycis gluma interiore spiculis subjecta. *Læst. it. 41. Lima.*  
*Habitat in Hispania. ☉*  
*Gramen vix Spithameum. Foliorum vaginæ tumidiu-*  
*sculæ, folio ipso parvo. Spica secunda, arte imbricata,*  
*oblonga, ex floribus secundis, sessilibus, duplici serie,*  
*compresso-carinatis: calycina valvula exteriore com-*  
*pressa, majore; flores singuli constans flosculis 3.*
4. CYNOSURUS spiculis alternis secundis sessilibus durus, rigidis obtusis appressis.  
 Gramen arvense, polypodii panicula crassiore. *Bar.*  
*rar. 1213. 1. 50.*  
*Habitat in Europa australi.*  
*Spica lanceolata, obtusa: spiculis appressis, secundis,*  
*duplici serie imbricatis rachi rigida. Generis dubii*  
*gramen.*

- caeruleus.** 5. **CYNOSURUS** bracteis integris. *Hort. cliff.* 495.  
*Fl. fvec.* 82. 89. *Roy. lugdb.* 64.  
 Gramen glumis variis. *Bauh. pin.* 10. *prodr.* 31. *Scheuch.*  
*gram.* 83.  
*Selleria.* *Scop. carn.* 189.  
*Habitat in Europæ pascuis uliginosis. Halæ ad Salinas.* 24.
- virgatus.** 6. **CYNOSURUS** panicula ramis simplicibus, floribus sessilibus subsexfloris: ultimo sterili; infimis subaristatis. *Amæn. acad.* 5. p. 393. \*  
 Gramen loliaceum panicula e spicis simplicibus teretibus, spicillis minimis compressis distichis alternis. *Brown. j m.* 137.  
 Gramen Dactylon panicula longa e spicis plurimis gracilioribus mollibus constante. *Sloan. jam.* 34. *hif.* 1. p. 113. t. 70. f. 2.  
*Habitat in Jamaica.*
- egyptius.** 7. **CYNOSURUS** spicis quaternis obtusis patentibus dimidiatis, calycibus mucronatis, caule repente. *Cynosurus* spicis quaternis terminalibus horizontalibus. *Roy. lugdb.* 64.  
 Gramen ichæzum malabaricum speciosius, longioribus mucronatis foliis. *Pluk. alm.* 175. t. 300. f. 8.  
 Gramen Dactylon ægyptiacum. *Bauh. pin.* 7. *theatr.* 110. *Roy. lugdb.* 109. *Morif. hif.* 3. p. 184. f. 8. t. 3. f. 7. *Scheuch. gram.* 109.  
*Habitat in Africa, Asia, America.*
- indicus.** 8. **CYNOSURUS** spicis digitatis linearibus, culmis compressis declinatis basi nodosis, foliis alternis. *Cynosurus* spicis aggregatis terminalibus deorsum aristatis. *Roy. lugdb.* 65.  
 Gramen Dactyloides spicis deorsum aristatis. *Burm. zeyl.* 106. t. 47. f. 1.  
 Gramen vaccinum femina. *Rumph. amb.* 6. p. 10. t. 4. f. 2.  
 Kavara-pullu. *Rheed. mal.* 12. p. 131. t. 69.  
*Habitat in Indiis.* ☉  
*A casu seminum remanent Spica pectinatae glumis calycinis, quarum laterales patent carina scabra.*
- coracanus.** 9. **CYNOSURUS** spicis digitatis incurvatis, culmo erecto compresso, foliis suboppositis.

*Gramen Dactylon ægyptiacum. Bauh. pin. 7. theatr.*  
110.

*Gramen Dactylon americanum minus. Scheuch. gram.*  
107.

*Gramen Dactylon orientale majus frumentaceum, femine napi. Pluk. alm. p. 174. t. 91. f. 2. Raj. suppl. 606.*

*Panicum gramineum f. Naatsjoni. Rumph. amb. 5. p. 203. t. 76. f. 2.*

*Noem & Sabil. Vessl. ægypt. 52. t. 53.*

*Tsittipullu. Rheed. mal. 12. p. 149. t. 78.*

*Habitat in India. ☉*

*Culmi quadripedales, erecti, compressi, foliorum petiolis cæcili. Folia longiora, bifaria, margine & pagina superiore, imprimis versus fauces, pilis raris longis villosa. Vaginis oppositis, compressis, amplexicaulibus. Spicæ cruciatæ 4. f. 6. terminales, raro unica inferiore, lineares, strictæ, erectiusculæ, digiti longitudine, bifariam subtus floribus imbricatæ, supra nudæ pedunculo carinato. Maturefcentes spicæ latiores evadunt, sæpe pedunculo plano membranaceo f. explanato incurvantur. Semina subglobosa, magna, quadruplici serie tumentia, nuda.*

10. *CYNOSURUS paniculæ spiculis sterilibus pendulis ternatis, floribus aristatis. aureus.*

*Gramen barcinonense, panicula tensa aurea. Tournef. inst. 523. Shaw. afric. 232. f. 232.*

*Gramen panicula pendula aurea. Bauh. pin. 3. theatr. 33. Scheuch. gram. 149.*

*Gramen sciurum f. alopecurum minus, he eromalla panicula. Barr. rar. 1180. t. 4.*

*Habitat in Europa australi, inter saxa & rupes, & in Oriente. ☉*

*Spiculæ ternatæ, steriles; Calyce bivalvi, lineari, acuminato, involvente Glumas plurimas, solitarias, alternas ovatas, obtusas, concavas, absque valvula minore, staminibus aut pistillis; nequeunt commodè dici flores, sed potius pro involucris haberi; Flores bini ad basin spicularum, Calyce lineari, longitudine flosculorum, bifloro; Flosculo inferiori sessili: Gluma exterior ovata, infra apicem arista longa recta; interiore angustissima. Flosculus alter, pedunculo longitudine flosculi prioris, minor, simillimus; tertiis flosculi solum rudimentum apparet.*

## FESTUCA.

\* *Panicula secunda.*

ovina.

1. **FESTUCA** panicula secunda coarctata aristata, culmo tetragono nudiusculo, foliis setaceis. *Fl. suec.* 95. 91.

*Poa* foliis setaceis compressis : caulino latiore, panicula contracta secunda, corollis acuminato-aristatis. *Aët. stockh.* 1741. p. 184.

*Poa* foliis setaceis, panicula ramosa, floribus petiolatis antrorsum spectantibus, glumis subulatis. *Roy lugdb.* 62.

*Poa* spiculis ovato-angustis aristato-acuminatis. *Fl. lapp.* 55.

Gramen, foliis, junceis brevibus, majus, radice nigra. *Bauh. pin.* 5. prodr. 34. *Scheuch. gram.* 279.

Gramen cristatum, paniculis nigricantibus. *Læf. pruss.* 110. t. 24.

Gramen capillatum, locustis pennatis non aristatis. *Vaill. parif.* 62.

*Habitat* in Europæ collibus apricis aridis vulgarissimum. 21.

vivipara.

- β. **Festuca** spiculis viviparis. *Fl. suec.* 1. n. 94.

Gramen paniculatum sparteum alpinum, panicula angustis spadiceo viridi, proliferum. *Scheuch. gram.* 213. t. 1. *Raj. angl.* 3. t. 22. f. 1.

*Habitat* in Alpibus Lapponiæ, Helvetiæ, Scotiæ. 24.

reptatrix.

2. **FESTUCA** paniculæ ramis simplicibus, spiculis subsessilibus.

*Habitat* in Arabia, Palæstina. 21.

*Radix crassitie pennæ asserine, longissime reptans sub terra, vestita latis foliorum rudimentis. Culmus plus quam sesquipedalis. Folia involuto-filiformia. Panicula oblonga ramis alternis, secundis, simplicissimis, quibus insident spiculæ sessiles, alternæ, plurimæ, lanceolæ, sexfloræ, acuminatæ, muticæ.*

duriuscula.

3. **FESTUCA** panicula secunda oblonga, spiculis sexfloris oblongis lævibus, foliis setaceis.

*Festuca* panicula nutante inferne ramosa, spicis ascendentibus hispidis, foliis setaceis. *Roy. lugdb.* 68.

Gramen pratense, panicula durior laxa unam partem spectante. *Raj. hist.* 1286. *angl.* 3. p. 413. t. 19. f. 1. *Scheuch. gram.* 285.

Gra-

Gramen tenue duriusculum & pene junceum. *Bauh. hist. 2. p. 463.*

*Habitat in Europæ pratis siccis.*

4. *FESTUCA panicula spiciformi pubescente, foliis dumetorum. filiformibus.*

*Habitat in Hispania. 24.*

*Culmi pedales f. sesquipedales, filiformes, teretes gemiculis duobus tumidis. Folia Radicalia pedalia, teretia, vix ancipitia; Caulina breviora, canaliculata. Panicula parva, quasi spicata. Spiculæ 10. f. 12. oblongæ, pubescentes, canescentes: inferiores geminae, pedicellatæ; superiores sessiles, solitariae. Glumæ terminatæ arista minuta. Figura Moris. hist. 3. f. 8. 1. 2. f. ult. hanc refert. Bulbi sæpe prognascuntur intra vaginas culmi. Affinis valde F. duriusculæ.*

5. *FESTUCA panicula secunda scabra, spiculis sex-rubra. floris aristatis: flosculo ultimo mutico, culmo semitereti. Fl. succ. 93. 92. \**

*Gramen alpinum pratense, panicula duriore laxa spadicea: locustis majoribus. Scheuch. gram. 287.*

*Habitat in Europæ sterilibus siccis.*

*Magnitudine, colore maturitatis rubro, culmo tereti, sed altero latere planiusculo, distinguatur a F. ovina.*

6. *FESTUCA panicula flexuosa, spiculis secundis inclinatissimis submuticis, foliis setaceis. Roy. lugdb. 68.*

*Gramen montanum, foliis capillaribus longioribus, panicula heteromalla spadicea & velut amethystina. Scheuch. gram. 276.*

*Habitat in Italia, Gallia, Anglia.*

7. *FESTUCA panicula erecta ramosa, spiculis sessilibus carinatis muticis.*

*Habitat in Palæstina.*

*Culmus altus, ramosus. Folia e vaginis latiusculis, angusta, longitudinaliter involuta, longa, subulata. Panicula erecta, ramosa & subdivisa. Spiculæ erectæ, policares, longæ, carinatæ, muticæ, obscuræ: flosculis 16. f. 24.*

8. *FESTUCA panicula spicata nutante. calycibus minutissimis muticis, floribus scabris longius aristatis. Roy. lugdb. 68.*

Gra-



Gramen festuceum myurum, minori spica heteromalla. *Barr. 10. t. 99. f. 1. Scheuch. gram. 194.*

Gramen murorum, spica longissima. *Kaj. ang. 3 p. 415. hist. 286. Moris. hist. 3. p. 215. f. 8. t. 7. f. 43.*

*Habitat in Anglia, Italia, Barbaria.*

maritima.

9. **FESTUCA** spica lineari secunda recta, floribus adpressis subaristatis. *Læfl. it. 44.*

Gramen exile duriusculum maritimum. *Scheuch. gram. 272.*

*Habitat in Hispania. ☉*

Gr. vix spithameum. Flores omnino spicati, secundi, aristis capillaribus. Spiculæ sexfloræ. Folia subulata: vaginis friatis.

bromoides.

10. **FESTUCA** panicula secunda, spiculis erectis lævibus, calycis altera valvula integra, altera aristata.

Festuca spicis erectis ad unum latus, palea altera calycina minima, altera acuminata. *Roy. lugdb. 68.*

Gramen paniculatum bromoides minus, paniculis aristatis unam partem spectantibus. *Raj. angl. 3. p. 415. hist. 1287. Pluk. alm. 174. t. 33. f. 10. Scheuch. gram. 297.*

*Habitat in Anglia, Gallia.*

Differt a *F. myuro* panicula subspicata, glumis non ciliatis.

calycina.

11. **FESTUCA** panicula coarctata, spiculis linearibus, calyce flosculis longiore, foliis, basi barbatis. *Amæn. acad. 3. p. 400. \**

Festuca panicula contracta, spiculis linearibus muticis longitudine calycis. *Læfl. it. 116. \**

*Habitat in Hispania. ☉*

Folia Radicalia in cespitem collecta. Culmi filiformes, angusti, digitales. Panicula attenuata, contracta. Calyx spiculæ longitudine.

\* *Panicula æquali.*

decumbens.

12. **FESTUCA** panicula erecta, spiculis subovatis muticis, calyce flosculis majore, culmo decumbente. *Fl. suec. 2. 93. lt. scan. 226.*

Gramen montanum avenaceum, locustis muticis tumentibus, pilosum. *Scheuch. gram. 70.*

Gramen triticeum palustre humilius, spica mutica brevior. *Moris. hist. 3 p. 177. f. 8. t. 1. f. 6.*

Gra-

**Gramen avenaceum parvum procumbens, paniculis non aristatis.** *Raj. angl.* 3. p. 408. *hist.* 1288. *Pluk. alm.* 174 t. 34 f. 1. *Mont. prodr.* 53. t. 2. f. 1.  
*Habitat in Europæ pascuis sterilibus.*

13. **FESTUCA panicula secunda erecta, spiculis subaristatis exterioribus teretibus.** *Fl. suec.* 91. 94.

**Festuca panicula spicata, spiculis uno versu inclinatissimis muticis.** *Roy. lugdb.* 68.

**Gramen loliaceum, spica divisa, pratense majus.** *Morif. hist.* 3. p. 184. f. 8. t. 2. f. 15.

**Gramen spartum, spica brizæ paniculata & corniculata.** *Barr. rar.* 1154. t. 25.

**Gramen paniculatum elatius, spicis longis muticis squamosis.** *Vaill. parif.* 92.

**Gramen arundinaceum, spica multiplici, Calamagrostis.** 1. *Bauh. pin.* 6. ex *Vaillant.*

- β. **Gramen pratense majus, locustis tumidis.** *Buxb. cent.* 5. p. 41. t. 16.

**Gramen arundinaceum aquaticum, panicula avenacea.** *Raj. angl.* 3. p. 411.

*Habitat in Europæ pratis fertilissimis.* 24.

14. **FESTUCA panicula ramosa erecta, spiculis subaristatis teretibus muticis.** *Fl. suec.* 90. 95.

**Poa spiculis oblongis erectis.** *Hort. cliff.* 28. *Roy. lugdb.* 62.

**Gramen aquaticum fluitans, multiplici spica.** *Bauh. pin.* 2. *theatr.* 41. *Scheuch. gram.* 199.

**Gramen loliaceum fluitans, spica longissima divisa.** *Morif. hist.* 3. p. 183. f. 8. t. 3. f. 16.

*Habitat in Europæ fossis & paludibus.*

15. **FESTUCA panicula spicata lobata, spiculis ovatis cristatis, latis sexfloris hirsutis.** *Læfl.*

*Habitat in Lusitaniæ collibus sterilibus.*

**Culmi e radice plurimi, vix digiti longitudine. Panicula subspicata, fere ovata, diversa ab Aira cristata flosculis pluribus iisque hirsutis, culmo brevior, facie autem ei accedit.**

16. **FESTUCA panicula patente, spiculis quadrifloris serotina, subaristatis, culmo assurgente tecto foliis brevissimis patentibus.**

**Gramen loliaceum serotinum, panicula dispersa.** *Segu. ver.* 3. p. 146. t. 2. f. 2.

*Ha-*

*Habitat Veronæ. Seguiet.*

*Culmi decumbentes articulis numerosissimis. Folia plana. Flosculi remoti arista brevissima.*

## B R O M U S.

secalinus.

1. BROMUS panicula patente, spiculis ovatis, aristis rectis. *Fl. suec.* 84. 96.

*Festuca culmo paniculato, spicis compresso-ovatis, Fl. lapp.* 28. *Roy. lugdb.* 67.

*Festuca graminea, glumis hirsutis. Bauh. pin.* 9. *theatr.* 143. *Scheuch. gram.* 250.

*Gramen avenaceum segetale majus, gluma turgidiore. Morif. hist.* 3. f. 8. t. 7. f. 17.

- β. Bromus hordeaceus panicula erecta coarctata. *Fl. suec.* 1 n. 87. *Sp. pl.* 1. p. 77.

*Habitat in Europæ agris secalinis arenosis. ☉*

mollis.

2. BROMUS panicula erectiuscula, spicis ovatis aristis rectis, foliis mollissime villosis.

*Gramen avenaceum locustis villosis angustis candidibus & aristatis. T. Scheuch. gram.* 254.

*Gramen avenaceum pratense, panicula squamata & villosa. Morif. hist.* 3. p. 213. f. 8. t. 7. f. 18.

*Festuca avenacea hirsuta, paniculis minus sparsis. Raj. hist.* 1289.

*Habitat in Europæ australioris siccis. ♂*

*Similis B. secalino, sed albido-pubescent. Geniculi crassiores, cylindrici, subtus inprimis valde hirsuti. Folia utrinque mollissima vaginis striatis. Panicula brevius pedicellata.*

squarrosus.

3. BROMUS panicula nutante, spicis ovatis: aristis divaricatis.

*Festuca graminea, glumis vacuis. Scheuch. gram.* 251. t. 5. f. 11.

*Gramen avenaceum, locustis amplioribus candicantibus glabris aristatis. Tournes. inst.* 525.

*Gramen phalaroides acerosum, nutante spica. Barr. ic.* 24. f. 1.

- β. Gramen festuceum majus, locustis crassis lanuginosis. *Buxb. cent.* 5. p. 19. t. 38. f. 1.

*Habitat in Gallia, Helvetia, Sibiria.*

*Panicula nutans. Spicæ glabræ, ovatæ: pedicellis propriis filiformibus, versus apices incrassatis. Aristæ divaricatæ.*

4. BRO-

4. **BROMUS** panicula nutante crispa, foliis utrinque nudis: vaginis pilosis, glumis villosis. purgans.  
Gramen bromoides catharticum. *Few. peruv.* 705. t. 1?  
*Habitat in Canada. Kalm.* 24  
*Culmus sesquipedalis, firmus. Folia latitudine arundinis, carinata, viridia, supra fere nuda, subtus omnino glabra. Vaginæ foliorum retrorsum pilosæ. Panicula crispa, flexuosa, nutans. Spiculæ flosculis 8, ad 14, oblongæ. Glumæ pilis brevissimis undique extus villosæ. Arista sub apice glumæ exterioris. Differt nostra a Fevillæi radicibus fibrosis, nec squamosis; spiculis angustioribus.*
5. **BROMUS** panicula nutante, foliis utrinque vaginisque subpilosis, glumis ciliatis. ciliatus,  
*Habitat in Canada; ex semine D. Kalm.* 24  
*Culmi tenues. Folia utrinque 3 vaginæ vix manifeste pubescentes. Panicula valde nutans, non crispis pedunculis. Spiculæ oblongæ, compressæ, peralorum marginibus (non dorso) valde pilosis, qua nota facile distinguitur. Flosculi 8, sub apice aristati; Calyces nudi. Glumæ corollæ lanceolatae.*
6. **BROMUS** panicula patula, spiculis oblongis distichis, glumis subulato-aristatis. sterilis.  
*Festuca avenacea sterilis elatior. Bauh. pin.* 9. *Morif. hist.* 3. p. 212. f. 8. t. 7. f. 11.  
*Gramen avenaceum, panicula sparsa, locustis majoribus & aristatis. T. Scheuch. gram.* 258.  
*Habitat in Europæ australioris agris, sylvis.*
7. **BROMUS** panicula nutante, spiculis ovato-oblongis. *Fl. syec.* 85. 97. arvensis,  
*Bromus culmo paniculato, spicis compressis. F. lapp.* 27. *Hort. cliff.* 25.  
*Festuca graminea, juba effusa. Bauh. pin.* 9. *prodr.* 19. *Scheuch. gram.* 262.  
*Festuca avenacea sterilis elatior circa Upsaliam. Rudb. clus.* 1. p. 87. f. 7. *opt.*  
*Festuca graminea nemoralis latifolia mollis. Bauh. pin.* 9. 2  
*Gramen avenaceum dumetorum panicula sparsa. Râf. angl.* 3. p. 415. *Vaill. parif.* 93.  
*Gramen avenaceum dumetorum paniculatum majus hirsutum. Morif. hist.* 3. p. 213.  
*Binnei spec. plant. P. 1. H* *Ægi-*

- Ægilops major*, caule & foliis arundinaceis, locustis glabrioribus & angustioribus e fusco xerampelinis. *Dill. app.* 60.  
*Habitat in Europa ad versuras agrorum.* ☉
- testorum. 8. **BROMUS** panicula nutante, spiculis linearibus. *Fl. fvec.* 86, 98. \*  
 Gramen murorum, spicis pendulis angustioribus. *Tournef. parif.* 91.  
*Festuca avenacea sterilis*, spicis erectis. *Raj. angl.* 3. p. 413. *Pluk. alm.* 174. t. 299. f. 2.  
*Festuca avenacea sterilis*, pedicellis brevioribus & spicis erectis. *Morif. hist.* 3. p. 212. f. 8. t. 7. f. 13.  
*Habitat in Europæ collibus siccis & tectis terrestribus.* ♂
- madritensis. 9. **BROMUS** panicula patulo-erecta, spiculis linearibus: intermediis geminis, pedicellis incrassatis. *A. mæn. acad.* 4. p. 265. \*  
*Bromus sterilis*, erecta panicula, major. *Barr. ic.* 76. f. 1.  
 Gramen Bromoides pumilum, locustis erectis majoribus aristatis. *Scheuch. gram.* 260.  
*Habitat in Hispania, Anglia.*
- rubens. 10. **BROMUS** panicula fasciculata, spiculis subseffilibus villosis, aristis erectis. *Amæn. acad.* 4. p. 265. \*  
 Gramen panicula molli rubente. *Bauh. hist.* 2. p. 464?  
*Habitat in Hispania.*
- scoparius. 11. **BROMUS** panicula fasciculata, spiculis subseffilibus glabris, aristis patulis. *Amæn. acad.* 4. p. 266. \*  
*Habitat in Hispania.*
- giganteus. 12. **BROMUS** panicula nutante, spiculis quadrifloris: aristis brevioribus. *Fl. fvec.* 2. n. 99. *st. scan.* 394.  
 Gramen sylvaticum glabrum, panicula recurva. *Vaill. parif.* 93. t. 18. f. 3.  
 8. **Bromus** paniculæ ramis conjugato-binatis. *Fl. fvec.* 1. n. 88.  
*Festuca* foliis digiti latitudine longissimis, panicula laxa. *Roy. lugdb.* 69.  
 Gramen bromoides aquaticum latifolium, panicula sparsa tenuissima aristata. *Scheuch. gram.* 264.  
*Habitat in Europæ sylvis siccis.* 24
- racemosus. 13. **BROMUS** racemo simplicissimo, pedunculis unifloris, floribus subseffloris lævibus aristatis.

*Festuca avenacea* spicis strigofloribus e glumis glabris compactis. *Raj. angl.* 3. p. 414.

*Habitat in Anglia.*

*Panicula constat racemo simplici: pedunculis alternis, sepius solitariis, unifloris. Flores ovati, acuminati; Folia subtus villosa.*

14. **BROMUS** spiculis alternis subsessilibus teretibus, culmo indiviso. *FL. suec.* 89. 100. *it gotl.* 180. pinnatus.

*Festuca* culmo alternatim spicato, spicis teretibus. *Fl. lapp.* 29. *Roy. lugdb.* 67.

*Gramen*, spica brizæ, majus. *Bauh. pin.* 9. *prodr.* 19. *theatr.* 133.

*Gramen loliaceum corniculatum.* *Scheuch. gram.* 35.

8. *Gramen lol. cornic.* spicis teretibus angustis & glabris. *Scheuch. gram.* 36.

*Habitat in Europæ sylvis montosis asperis.* 24

*Spicularum latus planum culmo obvertitur; aristæ sunt terminales.*

15. **BROMUS** spiculis distiche imbricatis sessilibus de- cristatus.  
pressis. *Amæn. acad.* 2. p. 339. \*

*Festuca* culmo spicato, spiculis multifloris. *Gmel. fibr.*

1. p. 115. t. 23.

*Gramen triticeum*, spica latiore compacta, cristatum.

*Ruxb. cent.* 1. p. 32. t. 50. f. 3.

*Habitat in Sibiria, Tartaria.* 24

16. **BROMUS** panicula patente, floribus subtrifloris. triflorus.

*Gramen Bromoides* panicula sparsa, locustis minoribus aristatis. *Scheuch. gram.* 511.

*Habitat in Germaniæ, Daniæ nemoribus Forskâhl.*

17. **BROMUS** spicis duabus erectis alternis. *Amæn. a-* distachyos.  
*cad.* 4. p. 304. 450. \*

*Gramen* spica brizæ minus. *Bauh. pin.* 9. *prodr.* 10.

*Pluk. alm.* 173. t. 33. f. 1.

*Habitat in Europa australi, Oriente.* ☉

*Calmus ramosus geniculis tumidis pubescentibus.*

## STIPA.

1. **STIPA** aristis lanatis.

*Festuca* aristis paniculæ circiter longitudine plumosis. pennata.

*Roy. lugdb.* 68.

- Gramen spicatum, aristis pennatis. Tournef. inst. 518.  
Scheuch. gram. 153.*
- Gramen sparteum pennatum. Bauh. pin. 5. theatr. 70.  
Spartum austriacum. Clus. hist. 2. p. 221.  
Habitat in Austria, Gallia, Svecia. 2.*
- capillata. 2. STIPA aristis nudis incurvatis, calycibus semine longioribus.  
*Festuca longissimis aristis. Bauh. pin. 10. theatr. 153.  
Gramen avenaceum montanum, spica simplici, aristis recurvis. Raj. Vaill. parif. 85.  
Habitat in Germania, Gallia.  
Simillima Stipæ junceæ, sed Folia non teretia sulco longitudinali; sed rigidiora, breviora, minus scabra, latere superiore magis explanata & subpubescentia. Calyx non albidus, sed spadiceus. Aristæ breviores & adulescentes varie curvata.*
- juncea. 3. STIPA aristis nudis rectis, calycibus semine longioribus.  
*Festuca aristis circiter longitudine paniculæ nudis. Roy. lugdb. 68.  
Festuca junceo folio. Bauh. pin. 9. Theatr. 145.  
Scheuch. gram. 151.  
Habitat in Helvetia, Gallia. 3.*
- tenacissima. 4. STIPA aristis basi pilosis, panicula spicata, foliis filiformibus. *Amæn. acad. 4. p. 266. Læfl. it. 118. \**  
*Gramen Sparteum panicula comosa. Bauh. pin. 5.  
Spartum herba Plinii. Clus. hist. 2. p. 220.  
Habitat in Hispaniæ collibus arenosis. 2.*
- avenacea. 5. STIPA aristis nudis, calycibus semen æquantibus.  
*Andropogon folio superiore spathiceo, pedunculis lateralibus oppositis unifloris, aristis flexuosis. Gron. virg. 133.  
Habitat in Virginia.  
Culmi tenues, Folium summum ventricosum longitudine paniculæ. Panicula pedunculis conjugatis, unifloris. Calycis gluma uniflora, pallida, bivalvis, aequalis. Corollæ arista terminalis, contorta, basi articulata, nuda, longitudine pedunculorum. Gramen hoc Avena sativa minus est.*
- membrana-  
cea. 6. STIPA pedicellis dilatatis membranaceis.  
*Habitat in Hispania. Læfling.*

## TRIANDRIA DIGYNIA.

117

*Gramen vix pedale, facie Avenæ. Panicula simplex, vix subdivisa, laxa, debilis. Pedicelli membranacei, ancipites, obtusiusculi. Calycis arista longitudine ariste corollinæ.*

7. STIPA aristis nudis, bracteis basi barbatis, flosculis sessili-fasciculatis. arguens.

*Gramen arguens. Rumph. amb. 6. p. 15. t. 6. f. 1. Habitat in India.*

*Flores in fasciculos congesti, ut glumæ vix a bracteis distinguuntur. Bracteæ longiores, basi extus valde pilosæ. Semina basi hirsuta aristis terminalibus nudis, nigris, contortis*

## AVENA.

1. AVENA paniculata, calycibus unifloris, seminibus hirsutis, aristis calyce triplo longioribus. sibirica;

*Festuca glumis villosis, aristis calyce triplo longioribus. Gmel. sibir. 1. p. 113. t. 22.*

*Habitat in Sibiria.*

2. AVENA paniculata, calycibus bifloris, flosculo hermaphrodito mutico, masculo aristato. elatior.

*Avena calycibus bifloris, panicula laxe spicata: pedunculis prioribus fasciculatis. Guett. stamp. 1. p. 182.*

*Gramen avenaceum, panicula acerosa, semine papposo. Dill. app. 48.*

*Gramen avenaceum elatius, juba longa splendente. Raj. meth. 179. angl. 3. p. 406. Moris. hist. 3. p. 214. f. 8 t. 7. f. 37. Scheuch. gram. 239. Hall. helv. 225. Vaill. parif. 89.*

3. Avena panicula nutante, calycibus bifloris: altero flosculo aristato. Roy. lugdb. 66.

*Gramen nodosum avenacea panicula, radice tuberosa prædita. Bauh. pin. 2. prodr. 3. theatr. 18. Scheuch. gram. 237.*

*Habitat in Europæ maritimis & aprieis. 24*

*Flosculi Masculi arista major, versus basin inserta; Hermaphroditi brevissima & fere terminalis.*

4. AVENA panicula attenuata, calycibus bifloris, seminibus villosis, aristis calyce duplo longioribus. pensylvanica.

*Habitat in Pensylvania. Kalm.*

H 3

4. AVE.



- Jæfingiana. 4. *AVENA* panicula contracta, flosculis binis apice biaristatis, dorso arista reflexa. *Læfl.*  
*Habitat in Hispania.*  
*Gramen vix digiti longitudine; Culmi plures ex radice. Folia planiuscula: summum ventricosum latius. Panicula coarctata, crassa. Calyx flosculis longior. Aristæ terminales, setaceae, longitudine corollarum: dorsalis duplo longior, torta.*
- fativa. 5. *AVENA* paniculata calycibus dispermis, feminibus lævibus. *Hort. cliff. 25. Hort. upf. 20. Mat. med. 38. Roy. lugdb. 65.*  
*Avena nigra. Bauh. pin. 23.*  
*β. Avena alba. Bauh. pin. 23.*  
*Habitat in insula Ivan Fernandez versus Chilli. Anson. ☉*
- nuda. 6. *AVENA* paniculata, calycibus trifloris, receptaculo calycem excedente, petalis dorso aristatis. *Amæn. acad. 3. p. 401.*  
*Avena nuda. Bauh. pin. 23. Bauh. hist. 2. p. 339. Raj. angl. 3. p. 389.*  
*Habitat - - - - - ☉*  
*Affinis admodum A. fativæ, sed semina decorticata decidunt.*
- fatua. 7. *AVENA* paniculata; calycibus trifloris, flosculis omnibus basi pilosis aristis totis lævibus. *Fl. svec. 97, 101.*  
*Avena feminibus hirsutis. Fl. lapp. 30. Roy. lugdb. 65.*  
*Festuca utriculis lanugine flavescentibus. Bauh. pin. 10.*  
*Gramen avenaceum, locustis lanugine flavescentibus. Scheuch. gram. 239.*  
*Habitat in Europæ agris inter segetes. ☉*
- sterilis. 8. *AVENA* paniculata, calycibus quinquefloris: exterioribus flosculis aristisque basi pilosis, interioribus muticis.  
*Flabit in Hispania. ☉ Alstræmer.*  
*Simillimus A. fatuæ, ut forte sola varietas, sed triplo omnibus partibus major & cultura constans.*
- flavescens. 9. *AVENA* panicula laxa, calycibus trifloris brevibus, flosculis omnibus aristatis. *Roy. lugdb. 66. Fl. svec. 2. n. 103.*  
*Gramen avenaceum pratense elatius, panicula flavescent-*

## TRIANDRIA DIGYNIA.

119

- scente, locustis parvis. *Raj. angl. 3. p. 407. hist. 1284. Scheuch. gram. 223.*
- Gramen avenaceum, spica sparsa flavescente, locustis parvis. *Morif. hist. 3. p. 215. f. 8. t. 7. f. 42.*
- Habitat in Germania, Anglia, Gallia.*
10. AVENA spicata, flosculis subquaternis calyce longioribus. fragilis.
- Gramen loliaceum lanuginosum, spica fragili articulata, glumis pilosis aristatum. *Scheuch. gram. 32.*
- Gramen loliaceum spurium hirsutum, aristis geniculatis. *Barr. ic. 905. f. 1, 2, 3.*
- Habitat in Lusitania, Hispania, Læss.*
11. AVENA spicata calycibus quinquefloris. pratensis.
- Avena panicula spicata, spiculis cylindricis culmo appressis. *Guett. flamp. 1. p. 182.*
- Avena spicis erectis, calyce spiculis brevior. *Fl. lapp. 31. Fl. suec. 96, 104.*
- Avena calycibus trifloris, panicula nutante, foliis planis. *Gmel. sibir. 1. p. 129.*
- Gramen avenaceum montanum, spica simplici, aristis recurvis. *Raj. angl. 3. p. 252. t. 21. f. 1. Scheuch. gram. 226.*
12. Gramen avenaceum glabrum s. potius hirsutum, panicula purpureo-argentea splendens. *Raj. angl. 3. p. 406. t. 21. f. 2.*
- Festuca dumetorum. *Bauh. pin. 20. prodr. 19. ex Vaill. parif. 89.*
- Habitat in Europæ pascuis siccis apricis.*
12. AVENA spicata, calycibus spiculis sexfloris longioribus: petalo exteriori apice aristato furcatoque. spicata.
- Habitat in Pennsylvania.*
- Gramen angustum; Folia linearia; Spica composita ex spiculis 3. f. 4, remotis, erectis: Spiculæ subpedunculatæ: singulis Calyx diphyllus, subulatus, aequalis, spicula longior; Flosculi 6, sessiles, erecti: Petalo exteriori apice bifido in apices duos setaceos, e quorum divisura arista articulata, longitudine spiculæ. Facies Festucae decumbentis.

## LAGURUS.

1. LAGURUS spica ovata aristata. *Hort. cliff. 25.* ovatus.
- Gramen alopecuroides, spica rotundior. *Bauh. pin. 4. theatr. 56.*

**Gramen spicatum tomentosum longissimis aristis denatum.** *T. Scheuch. gram.* 58.

*Habitat in Italia, Gallia, Sicilia, Lusitania.* ☉

cylindricus.

2. **LAGURUS** spica cylindrica mutica.

**Gramen alopecuros, spica longa tomentosa candicante.**

*Bauh. hist.* 2. p. 474.

**Gramen tomentosum alopecuros.** *Dalech. hist.* 430.

**Gramen tomentosum spicatum.** *Bauh.* pin. 4.

**Gramen tomentosum creticum spicatum, spica purpurea.** *Scheuch. gram.* 57.

*Habitat Monipellii, in Creta, Smyrnæ, D. Gerardus.*

## ARUNDO.

Bambos,

1. **ARUNDO** calycibus multifloris, spicis ternis sessilibus.

**Arundo arbor.** *Bauh. pin.* 18, *Hort. cliff.* 25. *Fl. zeyl.* 47. *Roy. lugdb.* 67.

**Tabaxir & Mombu arbor.** *Bauh. hist.* 1, p. 222.

*Ily. Rheed. mal.* 1. p. 25, t. 16.

*Habitat in India utraque.* b

*Panicula Scapus rectus, rigidus, gerens ad alternos dentes sapius Flores tres, alternatim s. distiche positos, sessiles, r. glandulos, lineares, longos.*

Donax.

2. **ARUNDO** calycibus trifloris, panicula diffusa. *Roy. lugdb.* 66.

**Arundo sativa.** *Virid. cliff.* 7. *Hort. cliff.* 26.

**Arundo sativa, quæ Donax dioscoridis.** *Bauh. pin.* 17. *theatr.* 27. *Scheuch. gram.* 159.

β, **Arundo indica laconica versicolor.** *Morif. hist.* 3. p. 219. f. 8. t. 8. f. 9. *Hort. cliff.* 26.

*Habitat in Hispania, Galloprovincia.*

phragmitis.

3. **ARUNDO** calycibus quinquefloris, panicula laxa. *Roy. lugdb.* 66. *Gron. virg.* 137.

**Arundo panicula laxa, flosculis quintis.** *Fl. svec.* 99. 105.

**Arundo panicula, laxa, calycibus subbifloris.** *Sauv. monsp.* 38.

**Arundo vulgaris s. phragmites dioscoridis.** *Bauh. pin.* 17. *Scheuch. gram.* 161.

*Habitat in Europæ lacubus, fluviis.* 24

epigejos.

4. **ARUNDO** calycibus unifloris, panicula erecta, foliis subtus glabris.

A.

*Arundo panicula coarctata erecta integra, foliis inferne glabris. Fl. fvec. 101. 106.*

*Gramen arundinaceum paniculatum montanum, panicula spadiceo-viridi, femine papposo. Scheuch. gram. 124.*

*Habitat in Europæ collibus aridis. 24*

5. ARUNDO calycibus unifloris, culmo ramofo. *Fl. Calamogre-  
lapp. 42. Fl. fvec. 100, 107. Roy. lugdb. 66. flis.*

*Gramen arundinaceum, panicula molli spadicea, majus. Bauh. pin. 7. theatr. 64. Scheuch, gram. 122. t. 5.*

*Habitat in Europæ paludibus graminosis. 24*

6. ARUNDO calycibus unifloris, foliis involutis mucronato-pungentibus. *arenaria.*

*Arundo foliorum lateribus convolutis acumine pungente. Fl. lapp. 43. Fl. fvec. 102, 108. It. scan. 336.*

*Gramen sparteum spicatum, foliis mucronatis longioribus. Bauh. pin. 5.*

*Gramen spicatum secalinum maritimum maximum, spica longiore. T. Scheuch. gram. 138.*

*Habitat in Europa & America ad maris littora arenosa. 24*

## ARISTIDA.

1. ARISTIDA panicula ramofo, spicis sparfis. *adscensionis,*

*Gramen avenaceum, panicula minus sparfa, cujus singula grana tres aristas longiffimas habent. Sloan. jam. 35. hift. 1. p. 16. t. 2. f. 56.*

*Gramen avenaceum maderaspatanum, panicula sparfa ariftis longioribus ornata. Pluk. alm. 174. t. 191. f. 3. ?*

*Habitat in Infula Adscensionis, una ex quatuor istius pauperrimæ Floræ plantis, quarum 1 Sherardia fruticosa, 2 Euphorbia origanoides, 3 Portulaca. Osbeck. 24*

*Habitus Fefluæ ovine, fed paulo major. Radix cespitem conftituens. Culmi inferne ramofo. Folia e basi latiore fenfim anguftata, plano-canaliculata subulata. Panicula oblonga, diftincta glumis corollæ univalvibus, filiformibus, longitudinaliter convolutis, triplici arifta terminatis.*

- americana. 2. *ARISTIDA* paniculæ ramis simplicissimis, spicis alternis. *Amæn. acad.* 5. p. 393. \*  
*Habitat in Jamaica.*

## LOLIUM.

- perenne. 1. *LOLIUM* spica mutica: spiculis compressis multifloris. *Lolium* spicis muticis. *Hort. cliff.* 24. *Fl. fvec.* 104, 110. *Roy. lugdb.* 69.  
*Lolium* spicis compressis, radice perenni. *Fl. lapp.* 32.  
*Gramen loliaceum*, angustiore folio & spica. *Baub. pin.* 9. *theatr.* 127. *Scheuch. gram.* 25.  
*Habitat in Europa ad agrorum versuras solo fertili.* 24
- tenuis. 2. *LOLIUM* spica mutica tereti spiculis trifloris. *Gramen Loliaceum*; foliis & spicis tenuissimis. *Vaill. parif.* 81.  
*Graminis loliacei angustiore folio & spica varietas.*  
*Baub. pin.* 9. spica tereti angustissima. *Scheuch. gram.* 28.  
*Habitat in Gallia.*  
*Hec faciunt plures tantum varietatem prioris; diversissimum a Tritico tenello est.*
- temulentum. 3. *LOLIUM* spica aristata: spiculis compressis aristatis. *Lolium* spicis aristatis, radice annua. *Hort. cliff.* 23. *Fl. fvec.* 103, 109. *Roy. lugdb.* 69.  
*Gramen loliaceum*, spica longiore, f. *Lolium dioscoridis.* *Bauh. pin.* 9. *theatr.* 121. *Scheuch. gram.* 31.  
*Habitat in Europæ agris inter Hordeum, Linum.* ☉

## ELYMUS.

- arenarius. 1. *ELYMUS* spica erecta arcta, calycibus tomentosis flosculo longioribus.  
*Elymus* foliis mucronato pungentibus. *It. scan.* 336.  
*Secale* spiculis geminatis. *Fl. fvec.* 106, 111.  
*Triticum* foliis acuminatis pungentibus. *Roy. lugdb.* 71.  
*Gramen caninum maritimum*, spica triticea, nostras.  
*Raj. hist.* 1256. *Scheuch. gram.* 6.  
*Habitat ad Europæ littoræ marina in arena mobili.* 24  
*Spica erecta, longa, tomentosa. Spiculæ 2, rectæ, bifloræ, muticæ, calyce breviores. Folia arundinacea, glauca f. albida, involuto-mucronata.*
- philadelphicus. 2. *ELYMUS* spica pendula patula, spiculis sexfloris: inferioribus ternatis. *Amæn. acad.* 4. p. 266. \*

Ha-

*Habitat in Pennsylvania.* 24

*Smillimus E. sibirico* & *canadensi.*

3. ELYMUS spica pendula arcta, spiculis binatis ca-  
lyce longioribus. *Hort. upf.* 22. \* *Amæn. acad.* 3.  
p. 20.

Triticum radice perenni, spiculis binis longissime ar-  
statis. *Gmel. sibir.* 1. p. 123. t. 28.

*Habitat in Sibiria.* 24

4. ELYMUS spica nutante patula, spiculis inferiori-  
bus ternatis, superioribus binatis. *Amæn. acad.* 3.  
p. 20.

Gramen secalinum majus altissimum virginianum. *Mor-  
rif. hist.* 3. p. 118. f. 8. t. 10. f. 2. *Raj. suppl.* 599.

*Habitat in Canada.* *Kalm.* 24

*Convenit in plurimis cum E. Sibirico, a quo tamen  
differt 1. Spiculis villosis, nec nudis. 2. involucris  
in aristas exeuntibus longioribus ipsa spicula, dem-  
tis aristis, at in Sibirica involucri minus subula-  
tum, sed fere lanceolatum in aristas desinens, quæ,  
longe brevior sua spicula. 3. Spicæ inferne ternæ,  
quum Sibirica semper duas obtineat. 4. Spiculæ,  
dum florent, dehiscunt a scapo patulo, in Sibirica  
vero spiculæ approximatae sunt scapo. Ad singuli flo-  
sculi basin, a latere interiore spicularum, est punctum  
f. macula rufo-fusca. Folia hujus rore caerulefcente,  
praesertim subtus obducta, quod non in precedente. Ari-  
stæ corollinae cum maturascit fructus patenti reflexae  
sunt, contra ac in Sibirica.*

5. ELYMUS spica erecta, spiculis binatis involucrio virginicus.  
longioribus. *Hort. upf.* 22.

Hordeum flosculis omnibus hermaphroditis, involu-  
cris flosculos crassitie & longitudine superantibus.  
*Gron. virg.* 13.

*Habitat in Virginia.* 24

5. ELYMUS spiculis bifloris, involucris setaceis pa-  
tentissimis, caput me-  
dusæ.

Elymus involucris reflexo-patentibus. *Amæn. acad.* 3  
p. 21.

Avena lusitanica spicata, caput medusæ referens. *Mor-  
rif. hist.* 3. p. 210. *Raj. suppl.* 611.

Gramen hordaceum, spica aristas longissimis circum-  
vallata. *Scheuch. gram.* 20.?

Ha.

*Habitat in Lusitaniæ, Hispaniæ maritimis.*

*Culmus angustus, pedalis. Spica oblonga. Involucra partialia tetraphylla, patentissima s. reflexa, setacea, longitudine flosculorum cum eorum aristis. Spiculae geminae, biflorae: flosculo utroque aristato, sed interiore omnibus partibus minore.*

- Hystrix. 7. **ELYMUS** spica erecta, spiculis, involucro destitutis. Gramen avenaceum locustis aristatis, paniculis echinum referentibus. *Gron.*

*Habitat - - - -*

*Spica composita ex spiculis duabus ad singulum dentem racheos. Spiculae quadriflorae aristis longis. Involucra nulla, sed horum loco calli duo.*

- caninus. 8. **ELYMUS** spica nutante arcta, spiculis rectis involucro destitutis: infimis geminis. *Fl. svec. 2. p. 112. Triticum calycibus subulatis quadrifloris aristatis. Sp. pl. 1. p. 86.*

*Gramen caninum non repens elatius, spica aristata. Moris. hist. 3. p. 177. f. 8. t. 1. f. 2. Buxb. cent. 4 p. 29. t. 50.*

*Gramen Loliaceum, fibrosa radice, arillis donatum. Vaill. parif. 82.*

*Habitat in Europa. 21*

## SECALE.

- cereale. 1. **SECALE** glumarum ciliis scabris. *Hort. upf. 22.*

hybernum. *Secale hybernum vel majus. Bauh pin. 22. theatr. 425.*

vernum. β. *Secale vernum s. minus. Bauh. pin. 23.*

*Habitat - - - - O*

- villosum. 2. **SECALE** glumarum ciliis villosis, squamis calycinis cuneiformibus.

*Gramen spicatum secalinum, glumis villosis in aristas longissimas desinentibus. Tournesf. inst. 518.*

*Gramen secalinum maximum. Park. theatr. 1144.*

- β. *Gramen creticum spicatum secalinum, glumis ciliaribus. Tournesf. cor. 39. Buxb. cent. 5 p. 21. t. 41.*

*Habitat in Europa australi, & in Oriente.*

- orientale. 3. **SECALE** glumis hirsutis, squamis calycinis subulatis.

*Gramen orientale secalinum, spica brevi & lata. Tournesf. cor. 39.*

*Habitat ad Archipelagum*

4. SE-

4. **SECALE** glumis extrorsum ciliatis.

creticum

Gramen creticum spicatum secalinum altissimum, tuberosa radice. *Tournef. cor. 39. Scheuch. gram. 22. Habitat in Creta.*

**HORDEUM.**

1. **HORDEUM** flosculis omnibus hermaphroditis aristatis; ordinibus duobus erectioribus. *vulgare.*

Hordeum flosculis omnibus hermaphroditis, seminibus corticatis. *Hort. upf. 22. Mat. med. 35.*

Hordeum flosculis omnibus hermaphroditis. *Hort. cliff. 24. Roy. lugdb. 69.*

Hordeum polystichon vernum. *Bauh. pin. 22. theatr. celeste. 439. Morif. hist. 3. f. 8. t. 6. f. 3.*

β. **HORDEUM** flosculis omnibus hermaphroditis, seminibus decorticatis. *Hort. upf. 23.*

*Habitat* - - - - ○

2. **HORDEUM** flosculis omnibus hermaphroditis aristatis, seminibus sexfariam æqualiter positis. *hexastichon, Hort. upf. 23.*

*Habitat* - - - - ○

3. **HORDEUM** flosculis lateralibus masculis muticis, seminibus angularibus imbricatis. *distichon, Hort. upf. 23.*

Hordeum distichon. *Bauh. pin. 22. Morif. hist. 3. f. 8. t. 6. f. 1.*

β. **HORDEUM** flosculis lateralibus masculis muticis, seminibus angularibus imbricatis decorticatis. *se-mulum,*

*Habitat* - - - - ○

4. **HORDEUM** flosculis lateralibus masculis muticis, seminibus angularibus patentibus corticatis. *zeocriton, Hort. upf. 23. n. 5.*

Zeocriton f. *Oryza germanica. Bauh. pin. 22. Morif. hist. 3. f. 8. t. 6. f. 2.*

*Habitat* - - - - ○

5. **HORDEUM** flosculis omnibus fertilibus ternis aristatis, involucris fetaceis basi ciliatis. *bulbosum, Aman. acad.*

4. p. 304. \*  
Gramen secalinum bulbosa radice. *Barr. ic. 112. f. 2. Scheuch. gram. 19.*

Gramen bulbosum ex aleppo. *Bauh. pin. 2. prodr. 4. theatr. 21.*

Gra-



*Gramen secalinum chalepense, radice tuberosa. Moris. hist. 3. p. 179. f. 8. t. 6. f. 7.*

*Habitat in Italia, Oriente. 24*

- murinum.* 6. **HORDEUM** flosculis lateralibus masculis aristatis, involucris intermediis ciliatis. *Hort. cliff. 24. Fl. suec. 107, 113. Roy. lugdb. 69.*  
*Gramen hordeaceum minus & vulgare. Bauh. pin. 9. theatr. 134. Scheuch. gram. 14.*  
*β. Gramen spicatum secalinum minus. Scheuch. gram. 17. Vaill. parif. 83. t. 17. f. 6.*  
*Habitat in Europæ locis ruderalis. ☉*
- nodosum.* 7. **HORDEUM** flosculis lateralibus masculis muticis, involucellis setaceis lævibus.  
*Gramen myosuroides nodosum. Raj. angl. 3. p. 397. t. 20. f. 2.*  
*Habitat in India, Anglia.*  
*Simile H. murino, sed angustius, altius. Flosculi laterales carent & aristis, involucella minime aristata.*
- jubatum.* 8. **HORDEUM** aristis involucrisque setaceis longissimis.  
*Habitat in Canada. Kalm.*  
*Habitus Hordei murini, sed involucri & aristæ setæ quadruplo longiores, quam tota spica, unde singularis facies graminis.*

## TRITICUM.

- æstivum.* 1. **TRITICUM** calycibus quadrifloris ventricosus glabris imbricatis aristatis. *Hort. upf. 21.*  
*Triticum radice annua, spica glabra aristata. Hort. cliff. 24. Roy. lugdb. 170.*  
*Triticum æstivum. Bauh. pin. 21.*  
*Habitat apud Baschiros in campis. Heintzelman. ☉*
- hybernum.* 2. **TRITICUM** calycibus quadrifloris ventricosus lævibus imbricatis submuticis. *Hort. upf. 21. Mat. med. 36.*  
*Triticum radice annua, spica mutica. Hort. cliff. 24. Roy. lugdb. 70.*  
*Triticum hybernum, aristis carens. Bauh. pin. 21.*  
*Habitat - - - ♂*
- turgidum.* 3. **TRITICUM** calycibus quadrifloris ventricosus villosis imbricatis subaristatis. *Hort. upf. 21.*

Tri.

*Triticum radice annua, glumis villosis Hort. cliff. 24.  
Roy. lugdb. 70.*

*Triticum spica villôsa quadrata brevior & turgidior.*

*Morif. hist. 3. p. 176. f. 8. t. 1. f. 14.*

*Habitat - - - - ♂*

4. **TRITICUM** calycibus bifloris nudis, flosculis ion- polonicum.  
gissime aristatis, racheos dentibus barbatis.

*Triticum polonicum. Pluk. phyt. 231. f. 6.*

*Triticum majus, longiore grano, glumis foliaceis,  
polonicum. Morif. hist. 3. p. 3. 175. f. 8. t. 1. f. 8.*

*Habitat - - - -*

*Simile præcedentibus, sed spica major, cinerea; Glumæ  
duplo longiores, bifloræ tantum arista utriusque flosculi  
longissima.*

5. **TRITICUM** calycibus truncatis quadrifloris, flo- Spelta.  
sculis aristatis hermaphroditis: intermedio neutro.

*Hort. upf. 21.*

*Hordeum flosculis lateralibus masculis muticis invo-  
lucro destitutis. Sauv. monf. 39.*

*Zea dicoccos vel Spelta major. Bauh. pin. 22. theatr.*

*412. t. 414. Morif. hist. 3. f. 8. t. 6. f. 1.*

*Habitat - - - - ♂*

6. **TRITICUM** calycibus trifloris: flosculo altero ari- monococ-  
stato; intermedio neutro. cum.

*Triticum involucris unifloris, floribus aristatis, spica  
disticha. Sauv. monf. 39. †*

*Hordeum distichum spica nitida f. Briza nuncupatum.*

*Tournes. inst. 513.*

*Zea Briza dicta f. monococcos germanica. Bauh.  
pin. 21. Morif. hist. 3. f. 8. t. 6. f. 2.*

*Briza monococcos dodonæi. Lob. ic. 31.*

*Habitat. - - - - ⊙*

7. **TRITICUM** calycibus subquadrifloris: flosculis tenellum.  
muticis acutis, foliis setaceis.

*Gramen foliaceum minus, spica simplici. Bauh. pin.  
9. prodr. 11.*

*Gramen foliaceum, foliis & spicis tenuissimis. Morif.  
hist. 3. p. 182. f. 8. t. 2. f. 3.*

*Habitat Montpelii? Sauvages. ⊙*

*Culmus spithameus, tener, filiformis. Folia setacea,  
involuta. Spica floribus 4 f. 7. alternis.*

- juncum.* 8. **TRITICUM** calycibus truncatis quinquefloris, foliis involutis. *Amzn. acad.* 4. p. 266. \*  
Gramen tritici spica mutici simili. *Bauh. pin.* 9. *prodr.* 18. t. 17. *theatr.* 132. *Scheuch. gram.* 7. *Vaill. parif.* 81.  
*Habitat in Europæ australi, Oriente.* 24
- repens.* 9. **TRITICUM** calycibus subulatis quadrifloris acuminatis.  
*Triticum radice repente, foliis viridibus. Fl. lapp.* 33. *Fl. svec.* 105. 114. *Hort. cliff.* 24. *Mat. med.* 37. *Roy. lugdb.* 70.  
Gramen caninum arvense f. Gramen diofcoridis.  
*Bauh. pin.* 1. *Scheuch. gram.* 5.  
*Habitat in Europæ cultis.* 24
- maritimum.* 10. **TRITICUM** calycibus multifloris, flosculis mucronatis, spica ramosa.  
*Poa panicula ramosa, floribus alternis sessilibus. Roy. lugdb.* 274.  
Gramen Loliaceum, paniculâ ramosa, maritimum.  
*Bauh. theatr.* 130. *Scheuch. gram.* 274.  
Gramen maritimum, panicula loliacea. *Bauh. pin.* 8.  
*prodr.* 19. *Morif. hist.* 3. f. 8. t. 2. f. 6.  
*Habitat in Galliæ & Angliæ maritimis.*  
*Culmus spithameus, subramosus. Folia longitudine culmi vaginis purpurascens. Spica ramosa, fere paniculata ramis angulatis. Spiculæ 8. f. 10-floræ, alternæ, compressæ, basi crassiusculæ, sessiles, vel etiam solitariæ in ramificationibus.*

## TRIGYNIA.

### ERIOCAULON.

- triangulare.* 1. **ERIOCAULON** culmo triangulari, foliis ensiformibus, capitulo ovato.  
*Plantaginella aurea alopecuroides brasiliiana, foliis gramineis. Breyn. cent.* t. 50. *Morif. hist.* 3. p. 259. f. 8. t. 16. f. 17.  
*Habitat in Brasilia.*  
*Flosculos non potui perspicere in Herb. Pisonis apud Burmannum.*

a. ERI-

2. **ERIOCAULON** culmo quinquangulari, foliis ensiformibus, calyce universali pentaphyllo. *Fl. zeyl.* 48. \*  
*Scabiola graminea nudicaulis, capitulis argenteis f. Statice minima m. deraipatana. Pluk. Alm. 366. t. 221. f. 7.*  
*Habitat in India. 24.*
3. **ERIOCAULON** culmo sexangulari, foliis ensiformibus. *Fl. zeyl.* 49.  
*Gramen junceum indiae orientalis minus, capitulo rotundo ex paleaceis spiculis in cacumine caulis glomerato. Pluk. mant. 48.*  
*Habitat in India. 24.*
4. **ERIOCAULON** culmo sexangulari, foliis setaceis. *Fl. zeyl.* 50.  
*Randia malabarica, capillaceo folio. Petiv. topic. 344.*  
*Tieru-kotsijelleti pullu. Rheed. mal. 12. p. 129. t. 63.*  
*Habitat in India.*
5. **ERIOCAULON** culmo decangulari, foliis ensiformibus.  
*Eriocaulon noveboracense, capitulo albo globoso f. Globularia americana statice haud absumilis, cauliculis lana atro-nitente refertis. Pluk. amalth. t. 409. f. 5.*  
*Randia americana procerior. Pet. gaz. t. 6. f. 2.*  
*Globularia affinis aquatica, caule tenui aphylo gramineo, capitulis albicantibus parvis globosis, foliis paucis humi stratis gramineis. Gron. virg. 13.*  
*Habitat in America septentrionalis paludibus.*

# MONTIA.

1. **MONTIA.** *Fl. lapp.* 51. *Fl. fvec.* 108, 115. *Roy. fontanae lugdb.* 199.  
*Montia aquatica minor. Mich. gen. 18. t. 13. f. 2.*  
*Cameraria arvensis minor. Dill. giff. 46.*  
*Alfneformis paludosa tricarpos. Pluk. alm. 20. t. 7. f. 5.*  
*Alfnoidea annua verna. Vaill. parif. t. 3. f. 4.*  
*Alfne palustris, portulacae aquaticae similis. Raj. hist. 1035. Pet herb. t. 10. f. 12.*  
*Portulaca arvensis. Bauh. pin. 288.*  
*Habitat in Europa ad scaturigines. ☉*

# PROSERPINACA.

1. **PROSERPINACA.** *Aët. upf.* 1741. p. 81.  
*Linnaei Spec. plant. pers. I I Tri.*

galassia.

*Trixis. Mitch. E. N. C. 1748. n. 23.*

*Habitat in Virginiae paludibus.*

*Radix repens. Caules pedales, teretiusculi. Folia alterna, lanceolata, serrata, in petiolos desinentia: demersa s. ima in lacinias lineares pinnatifida. Flores axillares, solitarii. Fructus figura Polygoni.*

### TRIPLARIS.

americana. 1. **TRIPLARIS.** *Læfl. it. 256.*

*Habitat in America meridionali. b*

*Folia spithamea, ovata, integra, acuminata, petiolata. Spicæ pilosæ, bracteis ovatis, acuminatis, parvis, pilosis.*

### HOLOSTEUM.

cordatum. 1. **HOLOSTEUM** foliis subcordatis. *Amæn. ac. 3. p. 21.*  
*Alfine foliis subcordatis, stipulis utrinque quaternis.*

*Hort. upf. 24.*

*Holosteum foliis orbiculatis oppositis, racemis laxis terminalibus remotis. Brown. jam. 139.*

*Alfine americana, nummulariæ folio. Herm. parad. 11. 1. 11. Sloan. jam. 87. hist. 1. p. 203.*

*Habitat in Jamaica, Surinama.*

fuculentum. 2. **HOLOSTEUM** foliis ellipticis carnosis. *Amæn. acad. 3. p. 21.*

*Alfine foliis ellipticis succulentis. Cold. noveb. 9.*

*Habitat in Noveboraco.*

hirsutum. 3. **HOLOSTEUM** foliis orbiculatis hirsutis. *Amæn. acad. 3. p. 21.*

*Alfine foliis orbiculatis hirsutis, floribus sessilibus. Roy. miss.*

*Habitat in Malabar.*

umbellatum. 4. **HOLOSTEUM** floribus umbellatis. *Læfl. it. 120. \**

*Spergula foliis oppositis, pedunculis umbellatis. Guert. stamp. 298. Dalib. parif. 134.*

*Holosteum cariophyllæum arvense. Tabern. ic. 233.*

*Lychnis graminea hirsuta umbellifera. Morif. hist. 2. p. 346. f. 5. t. 22. f. 46.*

*Caryophyllus arvensis umbellatus, folio glabro. Bauh. pin. 210.*

*Caryophyllus arvensis. Bauh. hist. 3. p. 361.*

*Habitat in Europæ australis arvis. ☉*

POLYCARPON.

1. POLYCARPON.

Mollugo *tetraphylla* foliis quaternis obovatis, paniculis dichotomis. *Hort. cliff.* 28. *Sp. pl.* 1. p. 89. tetraphyllum.

*Anthyllis marina* alfinefolia. *Bauh. pin.* 282.

*Anthyllis alfinefolia* polygonoides major. *Barr. rar.* 103. t. 534.

*Habitat in Italix, G. Narbonensis vineis.* ☉

MOLLUGO.

1. MOLLUGO foliis oppositis lanceolatis, ramis alternis, pedunculis lateralibus confertis unifloris. *Fl. zeyl.* 52. oppositifolia

*Habitat in Zeylona.*

2. MOLLUGO foliis subquaternis lanceolatis, ramis paniculatis. stricta.

*Habitat in Asia.*

*Radix fibrosa* Caules 3. f. 4. *stricti, laeves.* Folia *verticillata, sæpe quaterna, stricta, acuminata, lævia.*

*Panicula rarior e racemis terminalibus & lateralibus.*

3. MOLLUGO foliis quinis obovatis æqualibus, floribus paniculatis. *Fl. zeyl.* 51. pentaphylla.

*Alfine ramosa procumbens quadrifolia, ad radicem polyphylla.* *Burm. zeyl.* 13. t. 8. f. 12.

*Habitat in Zeylona.*

4. MOLLUGO foliis quinis obovatis, ramis alternis, pedunculis unifloris. Spergula.

*Habitat in India.*

*Caulis debilis, filiformis, articulatus Ramis alternis.*

*Folia quina, obovata, inæqualia, obtusa.* Pedunculi 2. f. 5, *ad singula genicula, foliis longiores, unisflori.*

5. MOLLUGO foliis verticillatis cuneiformibus acutis, caule subdiviso decumbente, pedunculis unifloris. verticillata.

*Hort. upf.* 24.

*Mullugo foliis sæpius septenis lanceolatis.* *Gron. virg.* 14.

*Alfine Spergula mariana, latiori folio, floribus ad nodos pediculis curtis circa caulem insidentibus, calycibus eleganter punctatis.* *Pluk. mant.* 9. t. 332. f. 5.

*Alfine procumbens, gallii facie.* *Ehret. pict.* t. 6. f. 3.

*Habitat in Virginia.* ☉

## MINUARTIA.

- dichotoma. 1. MINUARTIA floribus confertis dichotomis.  
 Minuartia foliis setaceis, caule simplici erecto, floribus sessilibus confertim dichotomis. *Læfl. it. 121.*  
*\* t. 1. f. 5.*  
*Habitat in Hispania. ☉*  
 Plantula rigida, dura, tenax, sæpius simplex. Folia opposita, setacea. Flores cymosi in capitulum quadratum, arte dichotomum, fastigiatum, caule sæpius majus, sessiles. Bractæ setacæ, flore longiores, apice recto. Semina quinque ad octo, subrotunda, reniformia. Color plantæ subfuscus. *Læfl.*
- campestris. 2. MINUARTIA floribus terminalibus alternis bractæa longioribus. *Læfl. it. 122. \**  
*Habitat in Hispaniæ collibus planis. ☉*
- montana. 3. MINUARTIA floribus lateralibus alternis bractæa brevioribus. *Læfl. it. 122. t. 1. f. 4. \**  
*Habitat in Hispaniæ collibus altis. ☉*

## QUERIA.

- hispanica. 1. QUERIA floribus confertis.  
 Queria hispanica, bracteis patulis hamatis. *Læfl. it. 48.*  
*Habitat in Hispania. ☉*  
 Antecedente minor, fragilior. Capitulum florum obseletius tetragonum, facillime a caule discedens. Bractæ setacæ; erectæ, apice extrorsum hamatæ, vestibus adherentes. Semen 1. compressum, subrotundum. Color plantæ albidus. Habitus antecedentis & structura floris eadem; at Semina solitaria & plura constituunt genera diversissima.
- canadensis. 2. QUERIA floribus solitariis, caule dichotomo.  
 Mollugo foliis oppositis, stipulis quaternis, caule dichotomo. *Gron. virg. 14.*  
*Habitat in Canada, Virginia. 24*  
 Radix fibrosa. Caulis strictus, dichotomus, spithameus, teres, filiformis, erectus. Herba lini cathartici. Folia opposita, lanceolato-ovata, integerrima, adpersa utrinque punctis fuscis. Stipulæ geminæ utrinque, acutæ, membranacæ. Flores solitarii, subpedunculati, parvi, virides, ex singula dichotomia. Calyx quinquepartitus: foliolis obtusiusculis, subfornicatis. Petala nulla. Stamina 3, Antheris flavis. Germen subrotundum.

*tundum. Styli tres. Spontanea planta altior caule capillari; culta crassior ☿ foliis minime punctatis.*

**LECHEA.**

1. **LECHEA** foliis linearl-lanceolatis, floribus paniculatis. *Amæn. acad. 3. p. 10. t. 1. f. 4. \**

*Capraria foliis integerrimis. Gron. virg. 75.*

*Scoparia foliis tenuissimis in plurimos & tenuissimos ramulos divisa & subdivisa. Raj. suppl. 132.*

*Habitat in Canadæ sylvis glareosis. 24*

2. **LECHEA** foliis ovato-lanceolatis, floribus lateralibus majoribus. *Amæn. acad. 3. p. 11. \* f. 4. \**

*Habitat in Canadæ aridis.*





## Classis IV.

TETRANDRIA  
MONOGYNIA.

## LEUCADENDRON.

- racemosum. 1. **LEUCADENDRON** foliis setaceis, flosculis disjunctis racemosis.  
*Protea* foliis setaceis, floribus racemosis. *Hort. cliff.* 496.  
*Habitat in Aethiopia.* §
- proteoides. 2. **LEUCADENDRON** foliis subulatis, ramis determinatis, floribus terminalibus.  
*Protea* foliis linearibus simplicissimis: ramis determinatis, floribus terminalibus. *Roy. lugdb.* 186.  
*Habitat ad Caput b. spei.* §  
*Folia conferta. - Rami ad determinata spatia conferti, ceterum nulli. Calyces subrotundi, pubescentes.*
- cancellatum. 3. **LEUCADENDRON** foliis subulatis, ramis sparsis, floribus lateralibus.  
*Lepidocarpodendron* foliis angustissimis gramineis, fructu cancellato, semine coronato. *Boerh. lugdb.* 2. p. 193. t. 193.  
*Habitat ad Caput b. spei, in Ecrietis montis Tigridis.* §
- Lepidocarpodendron* 4. **LEUCADENDRON** foliis lanceolatis, floribus oblongis, calycum squamis summis hirsutis.  
*Protea* foliis lanceolatis integerrimis glabris, calycinis superne villosis. *Roy. lugdb.* 186.  
*Lepidocarpodendron* foliis angustis longioribus salignis, calycis squamis elegantissime ex flavo fusco albo nigro variegatis, florum plumis atropurpureis. *Boerh. lugdb.* 2. p. 188. t. 188.  
 β. *Protea* foliis lanceolatis integerrimis glabris, calycem succingentibus hirsutis. *Roy. prodr.* 188.  
*Lepidocarpodendron* folio saligno viridi: nervo & margine flavo, cono longo superiore parte maxime clauso. *Boerh. lugdb.* 2. p. 189. t. 189.  
*Habitat ad Cap. b. spei.* §

5. LEU-

5. **LEUCADENDRON** foliis lanceolatis , floribus repens,  
oblongis, calycum squamis glabris.  
Protea caule multifloro , calycibus oblongis , foliis  
lanceolatis integerrimis. *Roy. lugdb.* 185.  
*Lepidocarpodendron* foliis angustis brevioribus sali-  
gnis, calycis squamis elegantissime ex roseo aureo al-  
bo atrorubro variegatis, florum plumis albis. *Boerh.*  
*lugdb.* 2. p. 187.
- β. *Leucadendron* foliis longissimis obtuse trigonis, lon-  
gitudine florem superantibus. *Wach. ultraj.* 204.  
Protea caule unifloro, calyce oblongo, foliis lineari-  
bus longissimis. *Roy. lugd.* 185.  
*Lepidocarpodendron* foliis longissimis angustissimis,  
fructum elegantissime ex rubro flavo & albo varie-  
gatum succingentibus, radice repente. *Boerh. lugdb.*  
2. p. 190. t. 190.
- Habitat ad Caput b. spei, in sabuletis. h*
6. **LEUCADENDRON** foliis lanceolatis , floribus acaulon.  
subrotundis, caule suffruticoso unifloro.  
*Leucadendron acaulon.* *Wach. ultraj.* 204.  
Protea caule unifloro, foliis lanceolatis. *Roy. lugdb.*  
185.  
*Lepidocarpodendron acaulon*, foliis paucis latis, ner-  
vo & marginibus rubris ornatis, fructu parvo.  
*Boerh. lugdb.* 2. p. 191. t. 191.
- Habitat ad Caput b. spei, ad radices montium. h*
- Caulis brevis. Folia lineari-lanceolata, alterna s. spar-  
sa. Calyx patens: Limbo folioso, obtuso, concavo,  
flore longiore colorato.*
7. **LEUCADENDRON** foliis lanceolatis , floribus Scolymoce-  
phalum.  
subrotundis, caule fruticoso ramofo.  
*Lepidocarpodendron acaulon*, ramis numerosis e ter-  
ra excrescens, calyce floris immaturo extus ex ru-  
bro & flavo variegato, intus flavo. *Boerh. lugdb.* 2.  
p. 192. t. 192.
- Habitat ad Caput b. spei, in ericetis mont. Tigridis. h*
8. **LEUCADENDRON** foliis integris, floribus de- Cinaroides.  
pressis, corollarum limbis cylindricis.  
*Leucadendron* foliis subrotundis patentissimis petiolatis,  
foliolis calycinis carinatis. *Wach. ultraj.* 204.  
*Lepidocarpodendron* folio subrotundo rigido, in pe-  
dunculo longo crasso, flore maximo purpureo.  
*Boerh. lugdb.* 2. p. 184. t. 184.

*Cinaroides frutex*, folio subrotundo rigido, e monte tabulari. *Petiv. musf.* 374.

- β. *Protea* foliis lanceolatis integerrimis, flore patente glabro, styliis longissimis. *Roy. lugdb.* 186.

*Habitat ad Caput b. spei in humidis mont. Tabularis. b*

*Variat foliis cordatis, ovatis, lanceolatis.*

hirtum.

9. *LEUCADENDRON* foliis lanceolatis apice callosis, caule hirsuto, floribus sparsis axillaribus. *A-mæn. acad.* 6, *afr.* 3. \*

*Lepidocarpodendron* foliis sericeis brevibus confertissimis, cono gracili longo. *Boerh. lugdb.* 2. p. 194. t. 194.

*Habitat ad Cap. b. spei. b*

caecullatum.

10. *LEUCADENDRON* foliis calloso-tricuspidatis involventibus flores laterales.

*Leucadendron* foliis cuneiformibus apice tridentatocallofis, summis ultra florem protensis. *Wach. ultraj.* 203.

*Protea* foliis lanceolatis obtusis flores involventibus apice tridentato-callosis. *Roy. lugdb.* 184.

*Hypophyllocarpodendron* foliis inferioribus apice trifido rubro; superioribus penitus rubris glabris. *Boerh. lugdb.* 2. p. 206. t. 206.

*Habitat ad Cap. b. spei locis uliginosis. b*

Hypophyllocarpodendron.

11. *LEUCADENDRON* foliis calloso-tricuspidatis sublanceolatis, calycibus turbinatis nudis.

*Leucadendron africana* f. *Scolymocephalus* folio angustiori; apicibus tridentatis. *Pluk. alm.* 212.

*Protea* foliis lanceolato-linearibus apice tridentatocallofis, capitulis aphyllis. *Roy. lugdb.* 184.

*Protea* foliis lanceolato-linearibus apice tridentatocallofis. *Hort. cliff.* 29.

*Conocarpodendron* folio rigido angusto apice tridentato rubro, flore aurantio. *Boerh. lugdb.* 2. p. 198. t. 198.

*Habitat ad Cap. b. spei in campis sabulosis. b*

Conocarpodendron.

12. *LEUCADENDRON* foliis calloso-quinquedentatis,

*Leucadendron* foliis obverse ovatis oblongis; margine calloso-fimbriatis; ad apicem crenatis. *Wach. ultraj.* 203.

*Leucadendron africana* arbor argentea, summo folio crenato. *Pluk. alm.* 212. t. 200. f. 2.

*Protea* foliis oblongo-ovatis apice quinquedentatocallofis. *Roy. lugdb.* 184.

**Conocarpodendron** folio crasso nervo lanuginoso supra crenato ibique limbo rubro, flore aureo, cono facile deciduo. *Boerh. lugdb.* 2. p. 196. t. 196.

**Scolymocephalus africanus** latifolius lanuginosus, foliis in summitate crenatis. *Herm. afr.* 20. *Raj. dendr.* 9.

*Habitat in Æthiopia. b*

13. **LEUCADENDRON** foliis setaceis semitrifidis. cyanoides.

*Protea* foliis linearibus ramosis. *Roy. lugdb.* 186.

**Cyanus æthiopicus**, rigidis capillaceis tenuissimis foliis trifidis. *Pluk. mant.* 61. t. 345. f. 6.

*Habitat ad Cap. b. spei. b*

14. **LEUCADENDRON** foliis setaceis multifidis, capitulis glomeratis.

*Serraria* foliis tenuissime divisis, capitulis tomentosis.

*Burm. afric.* 265. t. 99. f. 2.

*Habitat ad Cap. b. spei. b*

*Simile L. Serraria, sed flores non solitarii, at glomerati & demum spicati.*

15. **LEUCADENDRON** foliis setaceis multifidis, *Serraria* capitulis solitariis.

*Abrotanoides* arboreum monomatapense, floribus in ramulorum cymis. *Pluk. mant.* 1. t. 329. f. 1.

*Serraria* foliis tenuissime divisis, floribus rubris apetalis. *Burm. afr.* 264. t. 99. f. 1.

16. **Anethifolius** frutex africanus, flore sericeo. *Seb. thes.* 2. t. 63. f. 6.

*Habitat in Æthiopia. b*

## P R O T E A.

17. **PROTEA** foliis lanceolatis obliquis acutis sparsis argentea villosa-sericeis planis: floralibus verticillatis.

*Protea* foliis lanceolatis integerrimis hirsutis nitidis. *Hort. cliff.* 29.

*Leucadendros* africana arbor tota argentea sericea, foliis integris. *Pluk. alm.* 212. t. 200. f. 1.

*Argyrodendros* africana, foliis sericeis & argenteis. *Comm. hort.* 2. p. 51. t. 26. *Raj. dendr.* 9.

**Conocarpodendron** foliis argenteis sericeis latissimis. *Boerh. lugdb.* 2. p. 195. t. 195.

**Globularia** africana frutescens thymeleæ folio lanuginoso. *Tournef. inst.* 467.

*Frutex æthiopicus conifer, foliis cneori lanuginosis salici æmulis. Breyn. prodr. 2. p. 49.*

*Arbor ferens folia argentea. Zanoni. hist. 24.*

*Habitat ad Cap. b. Spei. b*

*Frutex sylvas argenteas constituens, nitidissimus omnium, foliis floralibus verticillatis.*

conifera.

2. *PROTEA* foliis lineari-lanceolatis integerrimis acutis glabris obliquatis. *Hort. cliff. 29.*

*Frutex æthiopicus conifer, foliis cneori salici æmulis. Breyn. cent. 21. t. 9.*

β. *Protea* foliis lineari-lanceolatis glabris, capitulis squamosis, corona foliacea succinctis. *Roy. lugdb. 185.*

*Conocarpodendron folio rigido crasso angusto, cono laticis parvo. Boerh. lugdb. 2. p. 197. t. 197.*

γ. *Protea* foliis lineari-lanceolatis glabris, capitulis ad basin squamosis. *Roy. lugdb. 185.*

*Conocarpodendron folio angusto rigido brevi, cono parvo aureo: corona foliacea succincto. Boerh. lugdb. 2. p. 200.*

δ. *Protea* foliis lineari-lanceolatis glabris, caule inferne radicato. *Roy. lugdb. 185.*

*Conocarpodendron folio tenuissimo angustissimo saligno, cono calyculato. Boerh. lugdb. 2. p. 203. t. 203.*

ε. *Protea* foliis lanceolatis acuminatis flexuosis, capitulis corona foliacea succinctis. *Roy. lugdb. 185.*

*Conocarpodendron folio tenui angusto saligno, cono calyculato corona foliacea succincto. Boerh. lugdb. 2. p. 204. t. 204.*

*Habitat ad Cap. b. Spei. b*

*Planta naturalis, in patria, an ipsa? argentea; at culta & captiva extra patriam exuit decus; variat dein etiam domi mille modis vere Protea.*

## CEPHALANTHUS.

occidentalis. 1. *CEPHALANTHUS* foliis oppositis ternisque. *Gron. virg. 15.*

*Cephalanthus* foliis ternis. *Hort. cliff. 79. Roy. lugdb. 187.*

*Scabiosa dendroides americana, ternis foliis caulem ambientibus, floribus ochroleucis. Pluk. alm. 336. t. 77. f. 4.*

Ha-

*Habitat in America septentrionali. b*

*Capsule nulla. Semina solitaria, longa pyramidata: basi attenuata, angulata, lanuginosa unilocularia. Receptaculum commune globosum, villosum.*

## GLOBULARIA.

1. **GLOBULARIA** caule fruticoso, foliis lanceolatis *Alypum*, tridentatis integrisque. *Roy. lugdb.* 190.  
*Globularia fruticosa, myrti folio tridentato. T. Ga-*  
*rid. aix.* 210.  
*Alypum monspeliensium, f. Frutex terribilis. Bauh.*  
*hist.* 1. p. 598. *Niss. afl.* 1712. p. 336. t. 18.  
*Thymelæa foliis acutis, capitulo succifæ. Bauh. pin.* 463.  
*Habitat Monspeli inque Regni Valentini & Italiæ*  
*sylvis ad rupes & saxosa. b*
2. **GLOBULARIA** caule fruticoso, foliis radicalibus *bisnagarica*, cuneiformibus retusis; caulinis lanceolatis.  
*Scabiosa bisnagarica f. Globularia frutescens, rigidis*  
*foliis ad radicem rotundioribus cordatis. Pluk. alp.*  
*336. t. 58. f. 5. Moris. hist.* 3. p. 51.  
*Habitat in Bisnagarie sylvis. 24*
3. **GLOBULARIA** caule herbaceo, foliis radicalibus *vulgaris*, tridentatis; caulinis lanceolatis. *Fl. succ.* 109. 116.  
*it. æl.* 65. \* *Dalib. parif.* 43.  
*Globularia caule folioso, foliis ovatis integerrimis.*  
*Hort. cliff.* 490. *Roy. lugdb.* 190. *Hall. helv.* 667.  
*Aphyllanthes anguillare. Cam. hort.* 18. t. 7.  
*Bellis cærulea, caule folioso. Bauh. pin.* 262.
3. *Bellis cærulea apula. Tabern. hist.* 2. p. 709.
4. *Bellis cærulea monspeliaca. Tabern. hist.* 2. p. 709.  
*Habitat in Europæ apricis duris. 24*
4. **GLOBULARIA** foliis radicalibus crenato-aculeatis, *spinosa*, caulinis integerrimis mucronatis.  
*Globularia spinosa. Tournef. inst.* 476.  
*Bellis cærulea spinosa. Bauh. pin.* 262.  
*Bellis spinosa, flore globofo. Bauh. prodr.* 121.  
*Habitat in Granadæ montibus. 24*
5. **GLOBULARIA** caule subnudo, foliis cuneiformi-cordifolia, bus tricuspидatis: intermedio minimo.  
*Globularia foliis radicalibus cuneiformibus retusis den-*  
*tatis: denticulo intermedio minimo. Hort. cliff.* 491.  
*Roy. lugdb.* 190.

- Scabiosa bellidis folio, humilis, caule nudo, radice repente, folio cordato. *Morif. hist. 3. p. 50. f. 6. t. 15. f. ult.*  
*Bellis cærulea montana frutescens. Bauh. pin. 262.*  
 β. Globularia alpina minima, origani folio. *Tournef. infl. 467.*  
 Scabiosa, bellidis folio, pyrenaica minima. *Morif. hist. 3. p. 51.*  
*Habitat in Pannonia, Austria, Helvetia, Pyrenæis. 24.*  
 audicaulis. 6. GLOBULARIA caule nudo, foliis integerrimis lanceolatis.  
 Globularia pyrenaica, folio oblongo, caule nudo. *Tournef. infl. 467.*  
 Scabiosa, bellidis folio, humilis, caule nudo, radice non repente. *Morif. hist. 3. p. 50. f. 6. t. 15. f. 4.*  
 Bellis cærulea, caule nudo. *Bauh. pin. 262. Raj. hist. 381.*  
*Habitat in Pyrenæis & Austriæ montibus. 24.*  
*Interdum unicum foliolum in scapo.*  
 orientalis. 7. GLOBULARIA caule subnudo, capitulis alternis sessilibus, foliis lanceolato-ovatis integris.  
 Globularia orientalis, floribus per caulem sparsis. *Tournef. cor. 35.*  
*Habitat in Natolia. Bœrting. 24.*  
 Radix perennis. Folia obverse ovata, in petiolos desinentia, acuta, integra, nuda, plurima. Caulis pedalis, herbaceus, simplicissimus: Foliis minimis, lanceolatis, alternis, remotis. Capitula in summitate caulis, 7. ad 10. alterna, sessilia.

## DIPSACUS.

- Nullonum. 1. DIPSACUS foliis sessilibus ferratis.  
 Dipsacus foliis connato-perfoliatis. *Hort. upf. 25. aristis fructus rectis. Sauv. monsp. 156.*  
 Dipsacus capitulis florum conicis. *Hort. cliff. 29. Gron. virg. 15. Roy. lugdb. 188. Dalib. parif. 44.*  
 Dipsacus sylvestris aut Virga Pastoris major. *Bauh. pin. 386.*  
 Dipsacus sylvestris. *Dod. pempt. 735.*  
 β. Dipsacus sativus. *Bauh. pin. 385. aristis fructus hamatis. Sauv. monsp. 156.*  
*Habitat in Gallia, Anglia, Italia, &c.*

2. **DIPSACUS** foliis connatis sinuatis. laciniatus.  
*Dipsacus folio laciniato. Bauh. pin. 385. Morif. hist.*  
*3. p. 158. f. 7. t. 36. f. 4.*  
*Habitat in Alfatia, Azow, Carniola. ♂*
3. **DIPSACUS** foliis petiolatis appendiculatis. *Hort. pilosus.*  
*upf. 25. Roy. lugdb. 188. Dalib. parif. 44.*  
*Dipsacus capitulis florum subglobosis. Hort. cliff. 30.*  
*Roy. lugdb. 188.*  
*Dipsacus sylvestris: capitulo minore f. Virga Pastoris*  
*minor. Bauh. pin. 385.*  
*Dipsacus tertius. Dod. pempt. 735.*  
*Habitat in Anglia, Gallia. ♂*

## SCABIOSA.

\* *Corollulis quadrifidis.*

1. **SCABIOSA** corollulis quadrifidis æqualibus floribus cernuis, foliis pinnatis: foliolis lanceolatis serratis. *Hort. cliff. 30. Hort. upf. 26. Roy. lugdb. 188.* alpina.  
*Scabiosa alpina, foliis centaurii majoris. Bauh. pin. 270.*  
*Scabiosa alpina centauroides. Besl. cyst. æst. 122.*  
*Habitat in alpebus Helveticis, Italicis. 2.*  
*Calyx imbricatus, flore brevior. Fructus semiovatus. Re-*  
*ceptaculum globosum: paleis lanceolatis, canaliculatis,*  
*persistenibus. Semina quadrangularia, coronata den-*  
*tibus 4 majoribus ♂ 4 minoribus alternis. Pappus*  
*radiis 20, patulis.*
2. **SCABIOSA** corollulis quadrifidis æqualibus calycibus aristatis, foliis radicalibus lyratis, caulinis pinnatifidis. *Roy. lugdb. 189. Hort. upf. 26. \** transylvanica.  
*Scabiosa altissima annua, foliis agrimonie nonnihil*  
*similibus. Herm. lugdb. 539. Morif. hist. 3. p. 46. f.*  
*6. t. 13 f. 13.*  
*Habitat in Transylvania. ☉*  
*Calyx ♂ Paleæ mucronato-aristatæ, ut in S. syriaca.*
3. **SCABIOSA** corollulis quadrifidis æqualibus, calycibus aristatis, caule dichotomo, foliis lanceolatis. syriaca.  
*Hort. cliff. 30. Roy. lugdb. 188.*  
*Scabiosa fruticans latifolia alba. Bauh. pin. 269. Morif.*  
*hist. 3. p. 46. f. 6. t. 14. f. 14.*
4. **SCABIOSA** perfoliata folio, flore amethystino. *Vaill.*  
*æst. 1722. p. 233.*  
*Habitat in Syria. ☉*



- leucantha.** 4. SCABIOSA corollulis quadrifidis æqualibus, squamis calycinis ovatis obtusis, foliis pinnatifidis. *Hort. cliff.* 30. *Roy. lugdb.* 188. *Sauv. meth.* 241.  
 Scabiosa flore globoso niveo. *Bauh. pin.* 207.  
 Scabiosa fruticans angustifolia alba. *Bauh. pin.* 270.  
*Habitat in collibus Narbonæ.* 24
- rigida.** 5. SCABIOSA corollulis quadrifidis subradiantibus: squamis calycinis obtusis, foliis lanceolatis serratis. *Amæn. acad.* 6. *af.* 4. \*  
 Scabiosa africana frutescens, foliis rigidis splendentibus & serratis, flore albicante. *Comm. hort.* 1. p. 185. t. 93. *Raj. suppl.* 237.  
*Habitat in Æthiopia.* 6
- succisa.** 6. SCABIOSA corollulis quadrifidis æqualibus, caule simplici, ramis approximatis, foliis lanceolato-ovatis integerrimis. *Hort. cliff.* 30. *Fl. suec.* 112, 119. *Mat. med.* 41. *Roy. lugd.* 188. *Dalib. parif.* 45.  
 Succisa glabra. *Bauh. pin.* 269.  
 Succisa f. Morsus diaboli. *Cam. epit.* 397.  
 β. Succisa hirsuta. *Bauh. pin.* 269.  
*Habitat in Europæ pascuis humidiusculis.* 24
- integrifolia.** 7. SCABIOSA corollulis quadrifidis radiantibus, foliis radicalibus ovatis serratis; rameis lanceolatis, caule herbaceo.  
 Scabiosa corollis quadrifidis, foliis omnibus lanceolatis serratis. *Sauv. monf.* 156.  
 Scabiosa annua integrifolia f. foliis bellidis. *Magn. monsp.* 231. \*  
 Scabiosa montana latifolia non laciniata rubra. *Bauh. pin.* 270.  
 Scabiosa latifolia, purpurascens flore. *Clus. hist.* 2. p. 2. f. 3.  
*Habitat Montpelii & in Helvetia.* ☉  
 Caulis non hispidus; Rami patuli. Calycis squamæ lanceolatæ corollis breviores; Corollæ rubræ. *Sauv.*  
 Folia Radicalia Bellidis, ovata, obtusiuscula, scabra; acutius serrata; caulina pauca; Ramea lanceolata, amplexicaulia, basi ciliata, raro dentata f. pinnatifida, longissima. Calyx 10-phyllus; flore brevior. Corolla radians, purpurascens. Corollulæ bilabiata: Labio interiore indiviso; exteriori tripartito, lineari. Pappus corona denticulata.
- lyrata.** 8. SCABIOSA corollulis quadrifidis æqualibus, foliis omni-

omnibus indivisis ovatis acuminatis subserratis, caule hispido.

Scabiosa corollis quadrifidis, receptaculis pilosis, foliis ovato-lanceolatis serratis hispidis. *Scop. carn.* 351.

Scabiosa foliis ovato-lanceolatis serratis subhirsutis, caule hispido. *Hall. helv.* 671.

Scabiosa latifolia rubra non laciniata. *Bauh. pin.* 270.

Scabiosa latifolia, rubro flore 11. *Clus. hist.* 2. p. 1.

8. Scabiosa montana non laciniata rubra 1. *Bauh. pin.* 270.

*Habitat in Austriae, Helvetiae, Montspelli sylvaticis.*

9. SCABIOSA corollulis quadrifidis radiantibus, caule hispido, foliis lanceolatis pinnatifidis: lobis imbricatis. *tatarica.*

Scabiosa flosculis quadrifidis, foliis pinnatifidis: laciniis lateralibus erectiusculis. *Afl. upf.* 1744. p. 11. 1. 1. \*

Scabiosa altissima segetum. *Triumf. Raj. suppl.* 236?

*Habitat in Tataria.* ♂

*Paleae lineares, seminibus longiores, deciduae. Planta 12-pedalis.*

10. SCABIOSA corollulis quadrifidis radiantibus, caule hispido, foliis pinnatifidis: lobis distantibus. *arvensis.* *Hort. cliff.* 31. *Fl. svec.* 110, 117. *Mat. med.* 40. *Roy. lugdb.* 188. *Dalib. parif.* 44.

Scabiosa pratensis hirsuta. *Bauh. pin.* 269.

Scabiosa arvensis. *Tabern. hist.* 442.

Scabiosa major communior hirsuta, folio non lacinato. *Bauh. hist.* 3. p. 2.

*Habitat in Europa, solo glareoso, iuxta segetes inque pratis.* 24

*Fructus simillimus S. tataricae, sed absque paleis.*

11. SCABIOSA corollulis quadrifidis, foliis caulinis tripinnatis filiformibus. *Sauv. monsp.* 268. †

Scabiosa capitulo globoso, foliis in tenuissimas lacinas divisus. *Bauh. pin.* 271. *Magn. monf.* 231.

*Habitat Montspelli, secus vias autumnno florens.*

\* \* Corollulis quinquefidis.

12. SCABIOSA corollulis quinquefidis radiantibus, columbaria. foliis radicalibus ovatis crenatis, caulinis pinnatis setaceis. *Hort. cliff.* 31. *Fl. svec.* 111, 118. *It. gotl.* 216.

216. 228. \* *Roy. lugdb.* 139. *Dalib. parif.* 45. *Sauv. monsp.* 242.

Scabiosa, capitulo glóbofo, major. *Bauh. pin.* 270.

Scabiosa minor. *Cam. epit.* 711.

β. Scabiosa, capitulo glóbofo, minor. *Bauh. pin.* 270.

*Habitat in Europæ montosis, ficcioribus.*

*Flores pallidi dijco convexo. Pappus setis 5 longitudine corollularum.*

maritima.

13. SCABIOSA corollulis quinquefidis radiantibus calyce brevioribus, foliis pinnatis: summis linearibus integerrimis. *Amæn. acad.* 4. p. 304. \*

Scabiosa maritima parva, *Bauh. hist.* 3. p. 7.

Scabiosa maritima, rutæ caninæ folio. *Bocc. sic.*

*Habitat in Sicilia, Monspeli.* ☉

ucranica.

14. SCABIOSA corollulis quinquefidis radiantibus, foliis radicalibus pinnatifidis; caulinis linearibus basi ciliatis.

Scabiosa foliis planis carnosissimis: inferioribus pinnatis: ramis integerrimis linearibus. *Gmel. sib.* 2. p. 213. t. 87.

*Habitat in Ucraina. P. Forskæhl.*

stellata.

15. SCABIOSA corollulis quinquefidis, foliis dissectis, receptaculis florum subrotundis. *Hort. cliff.* 31. \*

*Hort. upf.* 26. *Roy. lugdb.* 189.

Scabiosa stellata, folio laciniato, major. *Bauh. pin.* 271.

Scabiosa major hispanica. *Clus. hist.* 2. p. 1.

β. Scabiosa stellata, folio laciniato minor. *Bauh. pin.* 271.

γ. Scabiosa stellata minima. *Bauh. pin.* 271. *prodr.* 126. *Habitat in Hispania, præsertim in agro Granatenfi ad versuras.* ☉

*Varietas γ vix palmaris foliis omnibus pinnatis, quam in inferiora lanceolata serrata s. incisa*

*Semina basi lanata, Pappo rotato amplissimo.*

prolifera.

16. SCABIOSA corollulis quinquefidis radiantibus, floribus sessilibus, caule prolifero, foliis indivisis subserratis. †

Scabiosa stellata humilis integrifolia prolifera. *Herm. parad.* 125. t. 125. *Raj. suppl.* 236.

*Habitat in India.*

atropurpurea.

17. SCABIOSA corollulis quinquefidis, foliis dissectis, re-

receptaculis florum subulatis. *Hort. cliff.* 31. *Hort. upf.* 26.

Scabiosa peregrina rubra, capite oblongo. *Bauh. pin.* 270.

Scabiosa VI. Indica. *Clus. hist.* 2. p. 3.

Habitat in India? ☉

Flos atropurpureus Antheris albis. Fructus ovatus. Receptaculum subulatum, paleis setaceis. Semina tetragona, terminata præputio involuto plicato; Pap- po radii 5. setaceis.

18. SCABIOSA corollulis quinquefidis, foliis pinnatis: argentea. laciniis lanceolatis, pedunculis nudis lævibus longissimis. *Roy. lugdb.* 190.

Scabiosa orientalis argentea, foliis inferioribus incis. *Tournef. cor.* 34.

Asterocephalus perennis argenteus, laciniatus, caule tenui eburneo. *Vaill. aët.* 1722. p. 181.

Habitat in Oriente. 24

19. SCABIOSA corollulis quinquefidis, foliis incis, africana. caule fruticoso. *Hort. cliff.* 31. *Roy. lugdb.* 189.

Scabiosa africana frutescens. *Herm. parad.* 219. t. 219.

- β. Scabiosa africana frutescens maxima, foliis rugosis & crenatis, minor. *Herm. parad.* 220.

- γ. Scabiosa africana frutescens maxima, foliis tenuissime incis. *Boerh. lugdb.* 1. p. 123.

- δ. Scabiosa minor æthiopica frutescens, foliis lanuginosis. *Breyn. ic.* 33. t. 26.

Habitat in Africa, Oriente. 24

20. SCABIOSA corollulis quinquefidis, foliis lanceo- cretica. latis confertissimis integerrimis, caule fruticoso.

*Hort. cliff.* 31. *Roy. lugdb.* 169.

Scabiosa stellata, folio non dissecto. *Bauh. pin.* 271.

*Morif. hist.* 3. f. 6. t. 15. f. 31.

- β. Scabiosa cretica frutescens, auriculæ ursi folio.

*Tournef. cor.* 34.

Asterocephalus frutescens, leucoji folio longiore angusto. *Vaill. aët.* 1722. p. 247.

Habitat in Creta & adjacentibus. b

Folia in caule confertissima, ut fere imbricata. Calyces obtusi.

21. SCABIOSA corollulis quinquefidis, foliis lineari- graminifolia. lanceolatis integerrimis, caule herbaceo. *Amæn. acad.* 4. p. 267.

*Scabiosa argentea angustifolia. Bauh. pin. 270. prodr. 127. t. 127.*

*Scabiosa graminea argentea. Bauh. hist. 3. p. 12.*

*Asterocephalus argenteus graminifolius, flore cæruleo. Vaill. parif. 248.*

*Habitat in alpibus Helvetiæ, Baldi, Tridentini. 24*

**Pterocephala. 22. SCABIOSA** corollulis quinquefidis, caule procumbente fruticoso, foliis laciniatis hirsutis, pappo plumoso.

*Pterocephalus perennis humilis laciniatus & incanus. Vaill. act. 1722. p. 184.*

*Habitat in Græcia? b*

*Caulis vix e terra prominens Scapis brevissimis, unifloris.*

**echroleuca. 23. SCABIOSA** corollulis quinquefidis radiantibus, foliis linearibus pinnatis; radicalibus bipinnatis, petiolis perfoliatis.

*Scabiosa corollulis quadrifidis, foliis pinnatis. foliolis lanceolatis superne incis. Sauv. monsp. 241.*

*Scabiosa multifido folio, flore flavescente. Bauh. pin. 270. Moris. hist. 3. p. 48. f. 9. t. 13. f. 23.*

*Scabiosa VII. Clus. hist. 2. p. 3.*

*Habitat in Germaniæ pratis ficis. ♂*

*Folia radicalia pinnata, pinnis pinnatifidis; Caulina pinnata pinnis linearibus. Petiolorum bases utrinque ad caulis latera margine lato exstante unitæ & perfoliatæ. Flores flavescente-albidi: in disco tubulosi & quinquefidi, sed quasi truncati. Pedunculi longi. Fructus ovatus. Caulis genicula purpurascens.*

**papposa. 24. SCABIOSA** corollulis quinquefidis, caule herbaeo, calycibus foliosis florem superantibus, pappo plumoso. *Roy. lugdb. 189.*

*Scabiosa cretica, capitulo pappos mentiente. Tournef. cor. 34. Bærh. lugdb. 1. p. 130.*

*Habitat in Creta.*

## KNAUTIA.

**orientalis. 1. KNAUTIA.** *Hort. cliff. 32. Hort. ups. 27. Roy. lugdb. 190.*

*Lychni-Scabiosa, flore rubro, annua. Bærh. lugdb. 1. p. 131.*

Sc-

*Scabiosa orientalis*, caryophylli flore. *Vaill. act.* 1722.

p. 241.

*Scabiosa orientalis*, foliis superioribus integris, flore parvo purpureo, calyce longo gracili. *Till. pis.* 153.

t. 48. mala.

*Habitat in Oriente.* ☉

*Conferatur Scabiosa cærulea virginiana*, flosculis tetrapetalis caryophyllorum more patulis & lacinia-  
tis. *Pluk. alm.* 335.

## ALLIONIA.

1. *ALLIONIA* foliis cordatis, calycibus quinquefidis violaceo-  
trifloris.

*Allionia. Læfl. it.* 181. \*

*Habitat in Cumana Americæ.*

2. *ALLIONIA* foliis oblique cordatis, calycibus tri- incarnato-  
phyllis trifloris.

*Wedelia. Læfl. it.* 180. \*

*Habitat in Cumanæ arenosis.*

## HEDYOTIS.

1. *HEDYOTIS* foliis lanceolatis petiolatis, corymbis frutesco-  
terminalibus involucretis. *Fl. Zeyl.* 63. \* *Amæn. acad.*

i. p. 392.

*Valerianella* foliis nervosis acutis, flosculis in caulum  
summo quasi involucretis. *Burm. Zeyl.* 227. t. 107.

*Habitat in Zeylona.* b

2. *HEDYOTIS* foliis lanceolato-ovatis, floribus verti- auricularia-  
cillatis. *Fl. Zeyl.* 64. \* *Mat. med.* 47. *Amæn. acad.* i.

p. 391.

*Valerianella palustris*, foliis nervosis oblongis, floscu-  
lis ad caulum nodos inter foliorum sinus collectis.

*Burm. Zeyl.* 227. t. 108. f. 1.

*Muriguti. Rheed. mal.* 10. p. 63. t. 32.

*Habitat in Zeylona.* 24

2. *HEDYOTIS* foliis lineari-lanceolatis, caule herbaceo herbaceo,  
dichotomo, pedunculis geminis. *Fl. Zeyl.* 65. \*

*Habitat in Zeylona.*

## SPERMACOCE.

1. *SPERMACOCE* glabra, foliis lineatis, staminibus tenuioribus  
inclusis.

Spermacoe verticillis tenuioribus. *Dill. elth.* 370. t. 277. f. 359.

Spermacoe annuum, foliis lanceolatis glabris, floribus verticillatis. *Læfl. it.* 202.

Spermacoe. *Roy. lugdb.* 258.

Anonymos americana, foliis parietariæ scabris, floribus albis ad foliorum ortum vix conspicuis. *Pluk. alm.* 33. t. 136. f. 4.

*Habitat in Carolina.* ☉

verticillata. 2. SPERMACOCE glabra, foliis lanceolatis, verticillis globosis.

Spermacoe glabra, staminibus exstantibus. *Sp. pl.* 1. p. 102.

Spermacoe verticillis globosis. *Dill. elth.* 369. t. 277. f. 358. *Hort. cliff.* 33.

Scabiosa jamaicensis hyssopifolia. *Pluk. alm.* 336. t. 58. f. 6?

Pulegium fruticosum erectum, verticillis densissimis. *Sloan. jam.* 64. *hist.* 1. p. 170?

*Habitat in Jamaica, Africa.*

*Variat verticillis globosis amplexicaulibus* ☿ *pedunculatis.*

hirta. 3. SPERMACOCE pubescens, foliis oblongis: summis quaternis, floribus verticillatis.

Spermacoe erecta subhirsuta, foliis oblongis: superioribus approximatis. *Brown. jam.* 141.

*Habitat in Jamaica.*

*Caules herbacei, pubescentes, præsertim ad verticillos oppositi ramosi. Folia elliptica, lineata, subtus pubescentia, sessilia; summa quaterna. Verticilli multiflori.*

spinosa. 4. SPERMACOCE suffruticosa, foliis linearibus spinulis ciliatis.

Spermacoe suffruticosum. *Læfl. it.* 201.

Spermacoe spinosa. *Jacq. amer.* 12.

*Habitat in America.*

*Caules diffusi, duri, ad basin ramosi. Folia linearia rosmarini, articulis longiora, ciliata spinulis minutissimis. Verticilli pauciflori.*

hispida. 5. SPERMACOCE hispida, foliis obovatis. *Fl. zeyl.* 62. \*

*Galeopsis zeylanica*, folio oblongo, flore albo. *Burm.*

*Zeyl.* 163. t. 20. f. 3.

*Habitat in Zeylona.* 24

*Conf.* Tardavel. *Rheed. mal.* 9. p. 149. t. 76.

6. SPERMACOCE procumbens, foliis linearibus, corymbosa  
corymbis lateralibus pedunculatis.

*Habitat in India.*

Caules herbacei, angulati, debiles, diffusi. Folia linearia, reflexa. Pedunculi laterales, oppositi, aphylli. Flores fasciculati, s. umbella simplici, involuero polyphylo umbella longiore. Stamina exsertita, corolla longiora.

### SHERARDIA.

1. SHERARDIA foliis omnibus verticillatis, floribus arvensis terminalibus.

*Sherardia.* *Dill. gen.* 96. *Hort. cliff.* 33. *Fl. suec.* 113,

120. *Roy. lugdb.* 257. *Hall. helv.* 456. *Dalib. parif.* 46.

*Rubeola arvensis repens caerulea.* *Bauh. pin.* 334. *prodr.* 145.

*Rubia parva, flore caeruleo se spargens.* *Bauh. hist.* 2. p. 719.

*Habitat in arvis Scaniae, Germaniae, Angliae.* ☉

2. SHERARDIA foliis floralibus binis oppositis binis muralis floribus.

*Asperula verticillata luteola.* *Bauh. pin.* 334.

*Aspera verticillata muralis minima.* *Column. ecphr.* 302. t. 300.

*Aparine minima.* *Allion. nicaens.* 4.

*Gallium minimum, seminibus oblongis.* *Buxb. cent.* 2. p. 31. t. 30. f. 9.

*Habitat in Italia, Constantinopoli in vetustis muris.* ☉

Caules decumbentes. Folia ovato-lanceolata, inferiora sensa, media quaterna, summa bina. Flores bini propriis pedunculis, pallidi, plani. Fructus oblongi hispidi. Semina subarcuata, vix ac ne vix coronata.

3. SHERARDIA foliis quaternis aequalibus, caule fruticoso.

*Habitat in insula Adscensionis.* *Osbeck.* 5

*Frutex torosus, inaequalis, scaber, Ramis obtuse tetragonis.* Folia quaterna, angusto-lanceolata, integerrima.



*gerrima, glabra, marginibus revoluta, articulis ramorum paulo longiora, interstincta Stipulis brevissimis, subulatis. Flores oppositi, sessiles, axillares. Calyx tetraphyllus, germiini innatus, persistens foliolis lanceolatis. Corolla infundibuliformis, quadrifida, alba, longitudine germinis. Antheræ quatuor, oblongæ. Pericarpium dicoccum.*

# ASPERULA.

odorata.

1. ASPERULA foliis octonis lanceolatis, florum fasciculis pedunculatis. *Fl. suec. 114, 121. Mat. med. 45. Dalib. parif. 46.*  
*Asperula foliis pluribus, floribus pedunculo elevatis. Hort. cliff. 33. Roy. lugdb. 255.*  
*Asperula f. Rubeola montana odorata. Bauh. pin. 334.*

*Habitat in Sveciæ, Germaniæ umbrosis. 2.*

arvensis.

2. ASPERULA foliis senis, floribus sessilibus aggregatis terminalibus. *Hort. upf. 27.*  
*Asperula foliis pluribus, floribus sessilibus. Hort. cliff. 33. Roy. lugdb. 255. Dalib. parif. 47.*  
*Asperula cærulea arvensis. Bauh. pin. 334.*  
*Asperula cærulea. Dod. pempt. 355.*  
*Habitat in Gallia, Flandria, Germania, Anglia. ☉*  
*Folia floralia ciliata, supra hispida. Capitula terminalia sessilia. Fructus glabri.*

taurina.

3. ASPERULA foliis quaternis ovato-lanceolatis, floribus fasciculatis terminalibus.  
*Asperula foliis quaternis ovato-lanceolatis, ramis alternis. Hort. cliff. 33. Virid. cliff. 9. Roy. lugdb. 255. Hall. helv. 457. Sauv. monsp. 164.*  
*Rubia quadrifolia & latifolia lævis. Bauh. pin. 334. Morif. hist. 3. f. 9. t. 21. f. 1.*  
*Habitat in alribus Helvetiæ, Italiæ. 2.*

*Flores, in fasciculum congestos, masculos esse, unico tantum femineo, ex plantis in solo natali inquirendum?*

indictoria.

4. ASPERULA foliis linearibus: inferioribus senis: intermediis quaternis, caule flaccido, floribus plerisque trifidis.  
*Asperula foliis quaternis linearibus, floribus sæpius trifidis. Fl. suec. 115, 122. Sauv. monsp. 163.*  
*Rubeola quadrifolia, caule ramoso flaccido, floribus trifidis. Hall. rupp. 8.*

Gab

**Gallium album tripetalum.** *Morif. prælud.* 267. *Dalib. parif* 49.

**Gallium album 3.** *Tabern. hist.* 433. t. 733. f. 1. *bona.*

*Habitat in Sveciæ, Thuringiæ, Galliæ, Sibiria collibus aridis saxosis.* 24

*Planta, nisi sustentetur, procumbens.*

5. **ASPERULA** foliis quaternis lanceolato-linearibus, pyrenaica, caule erecto, floribus sæpius trifidis.

**Rubia cynanchica saxatilis.** *Bauh. pin.* 333. *prodr.* 146. *Burf.* XIX. 9.

*Habitat in Pyrenæis versus Hispaniam & prope Valentiam Gallorum.* 24

**Caules erecti, spithamæi, quadrangulares. Folia omnia in caule quaterna, linearia, carinata, acuta, levia; inferiora etiam quaterna, sed breviora, lanceolata, obtusiora; folia summa & floralia opposita, latiora, acuta, lanceolata. Flores tubulosi, rubri, sæpius trifidi. Statura Sherardiæ arvensis.**

6. **ASPERULA** foliis quaternis linearibus: superioribus *cynanchica.* oppositis stipulatis, caule erecto, floribus quadrifidis.

**Asperula foliis ternis subulatis triquetris, floribus quadrifidis.** *Sp. pl.* 1. p. 104.

**Asperula foliis linearibus quaternis, summis oppositis.** *Roy. lugdb.* 255. *Sauv. monsp* 163.

**Rubedra foliis quaternis, seminibus glabris, floribus umbellatis.** *Hall. helv.* 458.

**Rubia cynanchica.** *Bauh. pin.* 333. *Bauh. hist.* 3. p. 723.

**Gallium montanum latifolium cruciatum.** *Col. ecphr.* 1. p. 296. t. 297. f. 1.

**Gallium album minus.** *Tabern. hist.* 433. t. 433. f. 2.

*Habitat in Germaniæ, Angliæ, Galliæ, Helvetiæ, Italiæ, Orientis pratis aridis saxosis cretaceis.* 24

*Caulis palmaris, scaber. Germina rubra. Flores extus subrugoso-scabri.*

## DIODIA.

1. **DIODIA.** *Hort. cliff.* 493. \* *Gron. virg.* 71.

*virginiana.*

*Habitat in Virginia aquosis.*

## KNOXIA.

1. **KNOXIA.** *Fl. zeyl.* 400.

K 4

Den-

*zeylanica.*

*Dentellaria zeylanica*, flore minimo, antirrhini folio.

*Raj. suppl.* 246.

*Habitat in Zeylona supra truncos arborum putridarum.*

## HOUSTONIA.

cærulea.

1. *HOUSTONIA* foliis radicalibus ovatis, caule composito, pedunculis primis bifloris. *Gron. virg.* 15.

*Houstonia. Hort. cliff.* 35.

*Rubia parva virginiana*, foliis ad genicula binis, flore cæruleo fistuloso. *Pluk. alm.* 324. *mant.* 164.

*Paronychia* facie planta tetrapetaloides virginiana, flore cæruleo. *Morif. hist.* 3. p. 614. f. 15. t. 4. f. 1.

- β. *Chamejasme inodora*, alines facie, dispermos tetrapetaloides, quadrato caule, virginiana. *Pluk. alm.* 97. *mant.* 45. t. 97. f. 9. *Raj. suppl.* 502.

*Habitat in Virginia.*

purpurea.

2. *HOUSTONIA* foliis ovato-lanceolatis, corymbis terminalibus, floribus superis. *Gron. virg.* 15. †

*Rubia mariana*, alines majoris folio ad caulem binato, flore purpureo-rubente. *Pluk. mant.* 164. *Raj. supp.* 262.

*Habitat in Virginia.*

## GALIUM.

\* *Fructu glabro.*

aristatum.

1. *GALIUM* foliis quaternis lanceolatis æqualibus margine scabris, caule diffuso, petalis caudatis, seminibus glabris.

*Rubia lævis linifolia montis virginis. Bocc. mus.* 75. *Barr. ic.* 356. 583.

*Habitat in monte Ventoso* ☿ *m. Virginis Neapolitano. C. G. Minuart.* 24.

*Caules diffusi, ramosi, leves. Folia quaterna, lanceolata, enervia, utrinque glabra, margine scabra, mucronata. Flores paniculati, albi, pedunculis trifloris. Corollæ quadrifidæ, rarius trifidæ, apice aristatæ s. mucronatæ. Antheræ flavæ; Fructus glabri. Sub autumno novi rami foliis 4, 5, 6. variant.*

rubroides.

2. *GALIUM* foliis quaternis lanceolato-ovatis æqualibus subtus scabris, caule erecto, fructibus glabris.

*Habitat in Europa australi. D. Bæck.*

*Facies Galii borealis, sed folia duplō latiora, nervis subtus scabra, Caulis firmus, erectus. Panicula brevis, conglomerata. Semina levia.*

3. **GALIUM** foliis quaternis obovatis inæqualibus, palustre, caulibus diffusis. *Fl. svec.* 119, 126.

*Gallium caulibus diffusis, foliis quaternis verticillatis. Fl. lapp.* 52.

*Gallium palustre album. Bauh. pin.* 335.

*Habitat in Europæ rivulis limosis.* 24.

4. **GALIUM** foliis quaternis linearibus obtusis, ramis bermudense ramosissimis.

*Aparine foliis quaternis obtusis lævibus. Gron. virg.* 16.

*Rubia tetraphylla glabra, latiore folio, bermudensis, feminibus binis atropurpureis. Pluk. alm.* 224. t. 248. *Raj. suppl.* 261.

*Habitat in Virginia.*

5. **GALIUM** foliis quaternis linearibus, caule procumbente scabro, corollis trifidis.

*Habitat in Canada. Kalm.*

*Caulis procumbens, ramosissimus, Ramis ternis quaternisve, divaricatis. Folia linearia, obtusa, latiuscula, glabra, subtus parum scabra. Pedunculi sæpius terni, tenuissimi, foliorum longitudine, uniflori; Flores minimi, albi, tripartiti. Stamina tria; Fructus didymus, globosus, lævis.*

6. **GALIUM** foliis linearibus: caulinis senis; ramorum tinctorium. quaternis, caule flaccido, pedunculis subbifloris, fructibus glabris.

*Habitat in America septentrionali. Kalm.*

7. **GALIUM** foliis senis lanceolatis retrorsum serrato-uliginosum. aculeatis mucronatis rigidis, corollis fructu majoribus.

*Aparine foliis lineari-lanceolatis acuminatis rigidis, corollis fructu majoribus. Roy. lugdb.* 255. *Fl. svec.* 121, 127.

*Aparine minor palustris parisiensis, flore albo. Fl. lapp.* 58.

*Mollugo montana minor gallio albo similis. Raj. hist.* 482. *angl.* 3. p. 224.

*Gallium album minus. Pet. herb.* 30. f. 6. *Vaill. parif.* 78.

*Rubia quædam minor. Bauh. hist. 3. p. 716.*

*Habitat in Europæ pascuis aquosis sterilibus. 24*

*spuriom.*

8. **GALIUM** foliis senis lanceolatis carinatis scabris retrorsum aculeatis, geniculis simplicibus, fructibus glabris. *Hort. upf. 28. \**

*Aparine semine læviore Raj. hist. 484.*

*Aparine foliis crebrioribus & semine læviore. Moris. hist. 3. p. 332.*

*Habitat in Europæ cultis. ☉*

*Affine G. Aparines, sed minus.*

*saxatile.*

9. **GALIUM** foliis senis obovatis obtusis, caule ramossimo procumbente.

*Galium caule ramossissimo, foliis quinis obverse ovatis. Hort. cliff. 34. Roy. lugdb. 257.*

*Gallium saxatile supinum, molliore folio. Juss. art. 1714. p. 492. t. 15.*

*Habitat in Hispaniæ maritimis lapidosis, inque Saxonia.*

*minutum.*

10. **GALIUM** foliis octonis lanceolatis mucronatis ferrato-aculeatis glabris incurvis, fructibus reflexis.

*Galium foliis senis cuneiformi-lanceolatis mucronatis glabris. Hort. upf. 28. \**

*Galium saxatile minimum supinum & pumilum, flore luteo. Tournef. inst. 1152.*

*Habitat in imperio Ruthenico. 24*

*Simillimum G. vero, sed palmo brevius, decumbens. Caulis lævis, nec margine scaber. Folia utrinque parum convexa, nec margine reflexa. Fructus carnosus, magni, reflexis pedunculis, unde conglomerati.*

*pauciflorum.*

11. **GALIUM** foliis octonis hispidis linearibus acuminatis subimbricatis, pedunculis dichotomis.

*Rubeola saxatilis. Bauh. pin. 334. prodr. 145. Burf. XIX. 17.*

*Aparine minima s. Rubia saxatilis minima. Magn. monf. 291. ex Gerardo.*

*Habitat in montibus Galloprovinciæ. 24*

*Caules numerosi, digitales, angulati; Folia verticillata foliolis 8. s. 6. lanceolato-linearibus vel linearibus, acutis, uti caulis undique pilis patentibus hispidis. Rami rariores, alterni; Foliorum verticilli sæpe ita conferti, ut folia imbricatim digesta conspiciantur.*

*erantur. Panicula varior, fere terminalis ex pedunculis bis dichotomis.*

12. **GALIUM** foliis octonis linearibus sulcatis, ramis *verum.* floriferis brevibus. *Hort. cliff. 34. Fl. fvec. 116, 123. Mat. med. 46. Roy. lugdb. 256.*

*Galium caule erecto, foliis plurimis verticillatis linearibus. Fl. lapp. 61.*

*Gallium luteum. Bauh. pin. 335.*

*Gallium. Dod. pempt. 335.*

*Habitat in Europa frequens. 4*

13. **GALIUM** foliis octonis lævibus subtus scabris; floribus binis, pedunculis capillaribus, caule lævi. *sylvaticum.*  
*Galium caule tereti fulcrato, foliis octonis glaucis & obtusis. Hall. helv. 460.*

*Mollugo montana latifolia ramosa. Bauh. pin. 334.*

*Habitat in Germaniæ & Europæ australis montibus sylvestris.*

*Caules alti, debiles, obsoletissime angulati s. teretiusculi, læves. Folia verticillata, lato-lanceolata, margine & carina s. abra, glaucescentia. Pedunculi capillares, elongati, sæp. ultimi biflori & versus hos foliola duo. Flores minutissimi.*

14. **GALIUM** foliis octonis ovato-linearibus subserratis patentissimis mucronatis, caule flaccido, ramis patentibus. *Mollugo*

*Galium foliis pluribus acutis, caule flaccido, ramis patentissimis. Hort. cliff. 34. Roy. lugdb. 257. Fl. fvec. 117, 125.*

*Mollugo montana angustifolia ramosa, s. Gallium album latifolium. Bauh. pin. 334.*

*Rubia sylvestris lævis. Bauh. pin. 333.*

*Mollugo belgarum. Lob. ic. 802.*

*Habitat in Europæ mediterranea. 4*

15. **GALIUM** foliis subquaternis linearibus lævibus, *montanum.* caule debili scabro, seminibus glabris.

*Galium caule recto, foliis senis inferne canaliculatis. Hall. gert. 189. Zinn. gert. 231.*

*Gallium montanum altissimum, foliis angustis albican- tibus. Kupp. jen. 5.*

*Habitat in Germania. 4*

*Statura inter G. Molluginem & glaucum media. Caulis debilis, scaber. Folia caulina quina, reflexa, sub-*

- sublinearia* ; *Ramea quaterna*. Flores *corymbis trifidis*, *albi*, ante *explicationem* *extus purpurascetes*. *Antheræ fuscae*.
- glaucum. 16. **GALIUM** foliis verticillatis linearibus, pedunculis dichotomis summo caule floriferis, caule lævi. *Roy. lugdb.* 156. *Hort. upf.* 27. *Sauv. monsp.* 161. *Gallium faxatile*, glauco folio. *Bocc. mus.* 2. p. 172. t. 116. *Rubia montana angustifolia*. *Bauh. pin.* 333. *prodr.* 145. *Habitat in* Tataria, Helvetia, Austria, Monspellii. 21.
- purpureum. 17. **GALIUM** foliis verticillatis lineari-setaceis, pedunculis capillaribus folio longioribus. *Hort. cliff.* 34. *Roy. lugdb.* 256. *Gallium nigropurpureum montanum tenuifolium*. *Col. ecphr.* 1. p. 298. *Bauh. pin.* 335. *Habitat in* Italia.
- Hierofolymitanum. 18. **GALIUM** foliis denis lanceolato-linearibus, umbellis fastigiatis, fructibus glabris. *Amæn. acad.* 5. p. 451. *Habitat in* Palæstina. *Statura proxime accedit ad G. rubrum*.
- rubrum. 19. **GALIUM** foliis verticillatis linearibus patulis, pedunculis brevissimis. *Hort. cliff.* 34. *Roy. lugdb.* 256. *Scop. Carn.* 341. *Gallium rubrum*. *Bauh. pin.* 335. *Morif. hist.* 3. p. 332. *Gallium rubro flore*. *Clus. hist.* 2. p. 175. *Habitat in* Italia.
- \* *Fructu hispido.*
- boreale. 20. **GALIUM** foliis quaternis lanceolatis trinerviis glabris, caule erecto, feminibus hispidis. *Fl. lapp.* 60. *Fl. suec.* 118, 124. *Hort. cliff.* 64. *Roy. lugdb.* 257. *Hall. helv.* 450. *Rubia pratensis lævis*, acuto folio. *Bauh. pin.* 333. *prodr.* 145. *Burf. XIX.* 15. *Habitat in* Europæ borealis pratis. 24.
- retundifolium. 21. **GALIUM** foliis quaternis ovatis aculeato-ciliatis, feminibus hispidis. *Rubia semine duplici hispido*, latis & hirsutis foliis. *Bocconi. Morif. hist.* 3. p. 329. f. 9. t. 21. f. 5.

- β. Galium foliis quaternis ovatis lævibus obtusis, panicula dichotoma, seminibus hispidis. Sp. ph. 1. p. 108.*  
*Rubia quadrifolia f. rotundifolia lævis. Bauh. pin. 334 prodr. 145.*  
*Rubia quadrifolia, semine duplici hispido. Bauh. hist. 3. p. 718. Moris. hist. 3. f. 9. t. 21. f. 4.*  
*Aparine Crucjata. Hall. opusc. 275.*  
*Habitat in alpibus Helvetiæ, Styriæ. 2.*
22. **GALIUM** foliis octonis lanceolatis carinatis sca- Aparine.  
 bris retrorsum aculeatis, geniculis villosis, fructu  
 hispido.  
*Aparine foliis lanceolatis acuminatis scaberrimis, corollis fructu minoribus. Roy. lugdb. 455. Fl. sves. 120, 128.*  
*Aparine foliis lanceolatis. Hort. cliff. 34.*  
*Aparine vulgaris. Bauh. pin. 334.*  
*Aparine. Dod. pempt. 353.*  
*Habitat in Europæ cultis & ruderatis. ☉*  
*Caulis æqueter angulis retrorsum aculeatis, basi articulorum villosus. Folia 8. f. 10na, lanceolato-linearia, supra scabra, subtus glabra margine & carina retrorsum aculeatis. Rami oppositi.*
23. **GALIUM** foliis verticillatis linearibus, peduncu- parisense.  
 lis bifidis, fructibus hispidis.  
*Aparine foliis lineari-lanceolatis acuminatis flaccidis, corollis fructu minoribus. Roy. lugdb. 255.*  
*Aparine minima. Ray. angl. 3. p. 225. t. 9. f. 1.*  
*Galium parisiense tenuifolium, flore atropurpureo. Tournef. inst. 664.*  
*Habitat in Anglia, Gallia. ☉*  
*Caules pedales, debiles, tetragoni, retrorsum scabri. Folia septena, lanceolata, mucronata, scabra præsertim margine. Rami floriferi oppositi, breviores. Pedunculi nudi, biflori f. triflori. Corollæ flava, parvæ. Semina hispida.*

## CRUCIANELLA.

1. **CRUCIANELLA** erecta, foliis senis linearibus, flo- angustifolia,  
 ribus spicatis. Hort. upf. 27. ? Sauv. monsp. 164.  
*Crucianella foliis linearibus. Hort. cliff. 32.*  
*Rubia angustifolia spicata. Bauh. pin. 334 prodr. 145.*  
*Habitat Monspelii. ☉*

2. CRU-



- latifolia.* 2. **CRUCIANELLA** procumbens, foliis quaternis lanceolatis, floribus spicatis. *Hort. upf.* 27. \* *Sauv. monsp.* 164.  
*Crucianella* foliis lanceolatis. *Hort. cliff.* 33.  
*Rubia latifolia* spicatis. *Bauh. pin.* 334.  
*Habitat in Creta* ☿ *Monspeli.* ☉
- patula.* 3. **CRUCIANELLA** diffusa, foliis senis, floribus sparsis. *Amzn. acad.* 3. p. 401. \* *Læfl. ii.* 68.  
*Habitat in Hispania* ☉
- maritima.* 4. **CRUCIANELLA** procumbens suffruticosa, foliis quaternis, floribus oppositis quinquefidis.  
*Crucianella erecta*, foliis quaternis, corolla ad solem connivente. *Sauv. monsp.* 164.  
*Rubia maritima.* *Bauh. pin.* 334. *Dod. pempt.* 357. *Clus. hist.* 2. p. 176.  
*Habitat in Creta* ☿ *Monspeli.* *D. Sauvages.* b  
*Caules* provoluti, perennes, lignosi. *Folia* quaterna, lanceolata, rigida, acuta. *Bractæe* ovate. *Corollæ* quinquefidæ, aristatæ; *antheræ* nigrae. *Flores* tristes, lutescentes, die clausi, noctu patentes ambrosiaci.
- monspeliaca* 5. **CRUCIANELLA** procumbens, foliis acutis: caulibus quaternis ovatis; ramis subquinatis linearibus, floribus spicatis.  
*Crucianella repens*, foliis senis, spicis longis. *Sauv. monsp.* 225.  
*Rubia spicata repens.* *Magn. monsp.* 225.  
*Habitat* *Monspeli.*, inque *Palæstina.*  
*Caules* crassiusculi, diffusi. *Rami* alterni, recti, simpliciores. *Folia* inferiora quaterna, ovata, rigida, acuta; *superna* quina s. sena, linearia, acuta. *Pedunculi* nudi. *Spicæ* terminales, similes *C. angustifoliae*.
- RUBIA.**
- tinctorum.* 1. **RUBIA** foliis subsenis. *Hort. cliff.* 35. *Hort. upf.* 28.  
*Mat. med.* 44. *Roy. lugdb.* 254.  
*Rubia perennis*, foliis sæpius quinis. *Sauv. monsp.* 161.  
*Rubia sylvestris aspera.* *Bauh. pin.* 33.
- β. *Rubia tinctorum sativa.* *Bauh. pin.* 33.  
*Habitat* *Monspeli.* ☿ in *Italia.* 2.
- peregrina.* 2. **RUBIA** foliis quaternis. *Roy. lugdb.* 254.

*Rubia quadrifolia asperima lucida peregrina. Herm. lugdb. 523.*

*Habitat - - - 2.*

*Foliorum superior pagina enervis, levis, lucida.*

## FUCHSIA.

### 1. FUCHSIA. Mill. dist. 4.

triphylla.

*Fuchsia triphylla, flore coccineo. Plum. gen. 14. t. 133. f. 1.*

*Habitat in America meridionali.*

*Caulis herbaceus, simplicissimus. Folia terna, lanceolata, sessilia. Racemus terminalis, erectus.*

## SIPHONANTHUS.

### 1. SIPHONANTHUS falicis folio, flore flavescente. indica.

*Amm. ruth. 1736. p. 214. t. 15.*

*Habitat in India. ♀*

*Folia alterna, lanceolata; Pedunculi saepe e regione oppositi folii, umbellati.*

## CATESBÆA.

### 1. CATESBÆA.

spinosa.

*Frutex spinosus, basi foliis plurimis simul nascentibus. flore tetrapetaloide pendulo sordide flavo tubo longissimo, fructu ovali croceo semina parva continente. Catesb. carol. 2. p. 100. t. 100.*

*Habitat in Provincia. ♀*

## IXORA.

### 1. IXORA foliis ovalibus semiamplexicaulibus, floribus fasciculatis. Fl. Zeyl. 22. Osb. it. 220.

*Jasminum flore tetrapetaló, Ixora linnæi, Schetti horti malab. Burm. Zeyl. 125. t. 57.*

*Jasminum indicum, lauri folio, inodorum umbellatum, floribus coccineis. Pluk. alm. 196. t. 59. f. 2.*

*Flamma sylvarum, Kumph. amb. 4. p. 105. t. 46.*

*Arbor indica, lauri amplioribus foliis obtusis. Pluk. mant. 20. t. 364. f. 1.*

*Schetti. Rheed. mal. 2. p. 17. t. 13.*

*Habitat in India. ♀*

alba.

2. *IXORA* foliis ovato-lanceolatis, floribus fasciculatis  
Jasminum indicum, lauri folio, inodorum umbella-  
tum, floribus albicantibus & Schetti album. *Pluk.*  
*alm.* 196. t. 109. f. 2.

Bemischetti. *Rheed. mal.* 2. p. 19. t. 14.

*Habitat in India.* ♀

americana.

3. *IXORA* foliis ternis lanceolato-ovatis, floribus thyr-  
soideis. *Amzn. acad.* 5. p. 393.

*Pavetta* foliis oblongo-ovatis oppositis, stipulis setace-  
is. *Brown. jam.* 142. t. 6. f. 2.

*Habitat in Jamaica.* ♀

## PAVETTA.

indica.

1. *PAVETTA.* *Fl. zeyl.* 56. \* *Amzn. acad.* 1. p.  
388.

Arbor malabariensium, fructulentisci. *Bauh. pin.* 399.

*Pavate. Raj. hist.* 1581.

*Pavetta* f. *Malleomoth.* *Rheed. mal.* 5. p. 19. t. 10.

*Habitat in India.* ♀

Corymbi trichotomi, fastigiati, pedunculo filiformi  
longo insidentes. *Stigma oblongum, bifidum.*

## PETESIA.

stipularis.

1. *PETESIA* foliis lanceolato-ovatis subtus tomentosis,  
thyrsis lateralibus.

*Petesia* fruticosa, foliis ovatis oppositis, stipulis rigidis  
interpositis, racemis minoribus alaribus, calyce  
quinquefido. *Brown. jam.* 143. t. 2. f. 2.

*Habitat in Jamaica.* ♀

*Frutex.* Folia opposita lanceolato-ovata, integerrima.  
Racemi axillares, brachiati, laterales, folio bre-  
viores.

Lygistum.

2. *PETESIA* foliis ovatis nudis lineatis, caule flexu-  
oso.

*Lygistum* flexile fruticosum, foliis ovatis oppositis,  
petiolis pedatis, racemis alaribus. *Brown. jam.* 142.  
t. 3. f. 2.

*Habitat in Jamaica.* ♀

*Frutex caule flexuoso, levi.* Folia ovata venosa a-  
cuta, petiolata. Pedunculi axillares, sæpius gemini,  
uniflori.

3. PETESIA foliis oblongis utrinque tomentosis. Jacq. tomentosa.  
amer. 12. †  
Habitat in America. ‡

## MITCHELLA.

1. MITCHELLA. Amxn. acad. 3. p. 16. repens.  
Chamædaphne. Mitch. gen. 27.  
Lonicera foliis subovatis, germine bifloro, corollis  
interne hirsutis, stylo bifido. Gron. virg. 22.  
Syringa baccifera, myrti subrotundis foliis, floribus  
albis gemellis. Pluk. amalth. 198. t. 444. f. 2. Ca-  
tesb. carol. 1. p. 20. t. 20.  
Baccifera mariana, clematis daphnoides minori folio.  
Pet. gaz. 1. t. 1. f. 13.  
Habitat in Carolina, Terra Mariana, Virginia. ‡

## CALLICARPA.

1. CALLICARPA. Aët. upf. 1741. p. 80. americana.  
Sphondylococcus. Mitch. E. N. C. 8. p. 218.  
Burchardia. Du Ham. arb. 1. p. 111. t. 44.  
Johnsonia. Mill. dict.  
Anonymos baccifera verticillata, folio molli & in-  
cano, ex america. Pluk. alm. 33. t. 136. f. 3.  
Frutex baccifer verticillatus, foliis scabris latis den-  
tatis & conjugatis. Catesb. carol. 2. p. 47. t. 47.  
Frutex foliis amplis subrotundis acuminatis ex ad-  
verso binis, viminibus lentis infirmis quasi leni-  
canitie tectis. Gron. virg. 138.  
Habitat in Virginia, Carolina, Mississippi. ‡

## AQUARTIA.

1. AQUARTIA. Jacq. amer. 12. aculeata.  
Habitat in America meridionali. ‡  
Planta aculeata, superne inermis. Folia alterna, ova-  
ta, obtusa, petiolata. Corolla rotata lacinis linea-  
ribus. Antheræ erectæ, maximæ. Stylus deflexus.  
Jacq. Statura Solani.

## POLYPREMUM.

1. POLYPREMUM. Aët. upf. 1741. p. 78. procumbens.  
Linum carolinianum. Pet. gaz. 9. t. 5. f. 6.  
Habitat in Carolina, Virginia. ○

## PENÆA.

- Sarcocolla. 1. PENÆA foliis ovatis planis.  
Tithymali myrsinitis specie arbuscula æthiopica, subrotundis foliis, stœchadis arabicæ squamato capitulo duro lacrymam fundens. *Pluk. mant.* 183. t. 44.  
*Habitat in Æthiopia.* §  
Folia *obtusiuscula*. Corollæ *obtusæ*. Calyx *intermedio flori nullus*; Stylus *magis subulatus quam in reliquis*.
- mucronata. 2. PENÆA foliis cordatis acuminatis. *Mat. med.* 48.  
Penæa foliis ovatis acuminatis. *Koy. lugdb.* 399.  
Penæa. *Hort. cliff.* 37.  
Erica africana, unedonis flore amplo, foliis cordiformibus in acumen desinentibus. *Raj. dendr.* 97.  
Tithymali myrsinitis specie arbuscula æthiopica, flore parvo e lata basi in acutissimum mucronem subito desinente, capitulis origani. *Pluk. mant.* 183.  
*Habitat in Æthiopia.* §
- squamosa. 3. PENÆA foliis rhombæo cuneiformibus carnosissimis.  
*Habitat in Æthiopia.*  
*Hec reliquis minor.* Corolla (ut in præcedente) *acuta*. Folia parte dimidia superiore patula.

## BLÆRIA.

- ericoides. 1. BLÆRIA. *Hort. cliff.* 49.  
Erica carnea, promontorii bonæ spei, foliis & floribus villosis. *Pet. gaz. t. 2 f. 10.*  
*Habitat ad Cap. b. Spei.* §

## BUDDLEJA.

- americana. 1. BUDDLEJA foliis ovatis. *Amæn. acad.* 5. p. 394. \*  
Buddleja. *Anm. herb.* 577. *Hort. cliff.* 35.  
Buddleja affurgens incana, foliis majoribus, spicis assurgentibus. *Brown. jam.* 144  
Verbasci folio minore arbor, floribus spicatis luteis, seminibus singulis oblongis in singulis vasculis siccis. *Sloan. jam.* 139. *hist.* 2. p. 29. t. 173. f. 1. *Raj. dendr.* 97.  
*Habitat in Caribæis ad ripas & torrentes.* §
- occidentalis. 2. BUDDLEJA foliis lanceolatis.  
Buddleja foliis lanceolatis acuminatis integerrimis oppositis, spicis interruptis. *Mill. dict.*

Ophi-

*Ophioxylon americanum*, foliis oblongis mucronatis  
leviter ferratis bardanæ instar sultus lanuginosis.

*Pluk. alm.* 270 t. 210. f. 1.

*Habitat in America.* b

## EXACUM.

1. EXACUM floribus sessilibus. *Fl. zeyl.* 61. \*

sessile.

*Exacum.* *Amén. acad.* 1. p. 391. \*

*Centaurium minus aureum*, floribus numerosis, æthio-  
picum. *Pluk. alm.* 57. t. 275. f. 3?

*Habitat in Asia & Africa.*

2. EXACUM floribus pedunculatis.

peduncula-

*Centaurium minus hypericoides*, flore luteo, lini capi-  
tulis. *Pluk. mant.* 43. t. 343. f. 3.

*Habitat in India.* c

## PLANTAGO.

\* *Scapo nudo.*

1. PLANTAGO foliis ovatis glabris, scapo tereti, spi-  
ca floribus imbricatis. *major.*

*Plantago foliis ovatis glabris.* *Hort. cliff.* 43. *Fl. svec.*

122, 129. *Mat. med.* 49. *Roy. lugdb.* 401. *Hall. helv.*

470. *Dalb. parif.* 50.

*Plantago scapo spicato*, foliis ovatis. *Fl. lapp.* 62.

*Plantago latifolia sinuata.* *Bauh. pin.* 189.

*Plantago major.* *Cam. epit.* 261.

- β. *Plantago latifolia glabra minor.* *Bauh. pin.* 189.

- γ. *Plantago latifolia rosea*, floribus quasi in spica dispo-  
sitis. *Bauh. pin.* 189. *Hort. upf.* 29.

- δ. *Plantago latifolia*, spica multiplici sparsa. *Bauh. pin.* 189.

- ε. *Plantago latifolia rosea*, flore expanso. *Bauh. pin.* 189.

*Habitat in Europa & Japonia ad vias.* 24

2. PLANTAGO foliis ovatis glabris, scapo angulato, asiatica.  
spica floribus distinctis.

*Habitat in China, Sibiria.*

*Planta ita refert P. majorem, ut quisvis facile eandem  
diceret. At differt Spica longiore, floribus remotis  
& distantibus, Foliis sæpius basi subdentatis, Scapo  
angulato.*

3. PLANTAGO foliis ovato-lanceolatis pubescenti-  
bus, spica cylindrica, scapo tereti. *Hort. cliff.* 36.

L. 2

Fl.

*Fl. suec.* 123, 130. *Roy. lugdb.* 401. *Hall. helv.* 472.  
*Dalib. parif.* 50.

*Plantago latifolia incana. Bauh. pin.* 189.

*Plantago major incana. Clus. hist.* 2. p. 109.

β. *Plantago latifolia hirsuta minor. Bauh. pin.* 189.

*Plantago latifolia incana, spicis variis. Bauh. pin.*  
 189. *Morif. hist.* 3. f. 8. t. 15. f. 7.

*Habitat in Europæ pascuis sterilibus apricis argillosis. 24*  
*Folia non dentata, Calyces glabri, Filamenta purpur-*  
*scientia.*

virginica.

4. **PLANTAGO** foliis lanceolato-ovatis pubescentibus  
 subdenticulatis, spicis cylindricis pubescentibus,  
 scapo angulato. *Gron. virg.* 16.

*Plantago media incana virginiana, ferratis foliis, an-*  
*nua. Morif. hist.* 3. p. 259. f. 8. t. 15. f. 8.

*Plantago virginiana, pilosellæ foliis angustis, radice*  
*turbinata. Pluk. alm.* 298.

*Plantago mariana, spica prætenui. Pet. garz.* 1. t. 1.  
 f. 10.

*Plantago myosotis f. trinervia hirsuta carolina. Raj.*  
*hist.* 2. p. 188.

*Habitat in Virginia. ☉*

altissima.

5. **PLANTAGO** foliis lanceolatis quinquenerviis den-  
 tatis glabris, scapo subangulato, spica oblongo-  
 cylindrica.

*Plantago montana, crasso glabro canaliculato tenui-*  
*oreque folio, profundissime radicata. Till. pif.* 146.

*Habitat in Italia. 24*

*Scapus scaber, tripedalis. Folia sesquipedalia, obtuse*  
*dentata, crassa, glabra; dentibus remotis, obtusis. Spi-*  
*ca cylindrica, longa.*

lanceolata.

6. **PLANTAGO** foliis lanceolatis, spica subovata nu-  
 da, scapo angulato. *Vir. cliff.* 9. *Hort. cliff.* 36.  
*Fl. suec.* 124, 131. *Roy. lugdb.* 401. *Hall. helv.* 471.  
*Dalib. parif.* 50.

*Plantago angustifolia major. Bauh. pin.* 189.

*Plantago minor. Dod. pempt.* 102.

β. *Plantago trinervia, folio angustissimo. Bauh. pin.*  
 189. *prodr.* 98.

γ. *Plantago angustifolia major, caulium summitate fo-*  
*liosâ. Bauh. pin.* 189.

γ. Plan-

6. *Plantago argentea angustifolia e rupe victoriæ. Tournef. infl.* 127.  
*Habitat in Europæ campis sterilibus.* 24
7. **PLANTAGO** foliis lanceolatis obsolete denticulatis, spica subrotunda hirsuta, scapo tereti. lagopus.  
*Plantago angustifolia, paniculis lagopi. Bauh. pin.* 189.  
*prodr.* 98. *Morif. hist.* 3. f. 8. t. 16. f. 13.  
*Plantago Catanance. Rauw. itin.* 4. t. 6.  
*Habitat in G. Narbonensi, Hispania, Lusitania.* 24  
*Folia enervia, subtus pubescentia, apice denticulisque-obsolete fusca. Scapus teres. Spica subglobosa Calycibus hirsutis, Corollis limbo lævibus.*
8. **PLANTAGO** foliis lanceolatis obliquis villosis, spica cylindrica erecta, scapo tereti foliis longiore. albleans.  
*Plantago foliis lanceolato-linearibus, scapo longitudine foliorum, spica oblonga. Hort. cliff.* 37.  
*Roy. lugdb.* 401.  
*Plantago foliis lanceolato-linearibus, scapo foliis duplo longiore, spica oblonga. Sauv. monsp.* 26.  
*Holosteum hirsutum albicans majus. Bauh. pin.* 190.  
*Holosteum salmanticense majus. Clus. hist.* 2. p. 110.
9. *Holosteum hirsutum albicans minus. Bauh. pin.* 190.  
*Habitat in Hispaniæ & Narbonæ aridis.* 24
9. **PLANTAGO** foliis linearibus, planis hirsutis, spica oblonga erecta. alpina.  
*Plantago foliis linearibus, spica oblonga. Hall. helv.* 471. *Scop. carn.* 281.  
*Holosteum hirsutum nigricans. Bauh. pin.* 190.  
*Habitat in Helvetiæ, Austriæ Alpibus.* 24  
*Scapus hirsutus. Spica ante florescentiam cernua. Folia subtus convexa*
10. **PLANTAGO** foliis linearibus, scapo tereti brevissimo limbo lanato, spica subrotunda nutante. cretica.  
*36. Roy. lugdb.* 401.  
*Plantago cretica minima tomentosa, caule adunco. Tournef. cor.* 5.  
*Holosteum f. Leontopodium creticum. Bauh. pin.* 190.  
*Leontopodium creticum. Clus. hist.* 2. p. 11.  
*Habitat in Creta.*
11. **PLANTAGO** foliis semicylindræis integerrimis: basiflanatis, scapo tereti. maritima.  
*Fl. sic.* 127, 133.



- Plantago scapo spicato, foliis linearibus subtus convexis. Fl. lapp. 63. Lob. obs. 163.*  
*Coronopus maritimus major. Bauh. pin. 190.*  
*Habitat in litoribus maritimis Europæ, Americæ borealis. 24.*  
*Basis foliorum pilosa est, unde affinitas summa specierum 7. 8. 11. 12. sed quæ vis naturæ efficeret tam varias? an locus; an copula?*
- subulata.** 19. **PLANTAGO** foliis subulatis triquetris striatis scabris, scapo tereti. *Roy. lugdb. 403*  
*Plantago foliis triangulari-prismaticis. Guett. stamp. 2 p. 428.*  
*Holosteum, strictissimo folio, minus. Bauh. pin. 191.*  
*Serpentina omnium minima. Lob. ic. 439.*  
 β. *Holosteum massiliense. Bauh. pin. 190.*  
*Habitat in maritimis Mediterranei arenosis. 24*  
*Folia conferta, filiformia, acuta, supra plana, margine scabra aculeis vix conspicuis; ad radicem lana.*  
*Scapus teres, foliis longior, minime glaber. Spica ovato-oblonga, non villosa.*
- ferraria.** 13. **PLANTAGO** foliis lanceolatis quinquenerviis dentato-ferratis, scapo tereti.  
*Plantago apula laciniata bulbosa. Col. cephr. 1. p. 258. t. 259. Morif. hist. 3. p. 260. f. 8. t. 16. f. 19.*  
*Habitat in Apulia, Mauritania Brander. 24*  
*Folia lanceolata, quinquenervia, striata, subpubescentia, subpetiolata, dentibus argutis serrata. Scapus teres, pubescens.*
- Coronopus.** 14. **PLANTAGO** foliis linearibus dentatis, scapo tereti.  
*Plantago foliis linearibus pinnato dentatis. Hort. cliff. 37. Fl. svec. 126, 134. Roy. lugdb. 401. Dalib. parif. 50.*  
*Coronopus sylvestris hirsutior. Bauh. pin. 190.*  
*Herba stella f. Cornu Cervinum. Dod. pempt. 109.*  
 β. *Coronopus hortensis. Bauh. pin. 190.*  
*Habitat in Europæ glareosis.*
- lasingii.** 15. **PLANTAGO** foliis linearibus subdentatis, scapo tereti, spica ovata: bracteis carinatis membranaceis.  
*Last it. 124. \**  
*Plantago marina. Pet. herb. 3. f. 9. Raj. angl. 3. p. 315.*  
*Habitat in Hispaniæ collibus agrorumque marginibus. ☉*

*Differt a P. Coronopo quod sit minor & præcior:*  
*Spica ovata; ejus vero longa. Floribus rarius imbricatis: bracteis glabris navicularibus; ejus vero subulatis angustissimis pubescentibus. Stamina brevioribus. Calycis foliolis acutioribus. Colore corollæ magis fusco. Foliis linearibus, nec lanceolatis, parce dentatis. Scapus pilosus est Lxfl.*

16. **PLANTAGO** scapo unifloro. *Fl. lapp. 64. Fl. svec. uniflora. 128, 135. Dalib. parif. 50.*

*Plantago floribus femineis sessilibus ad exortum scapi uniflori maris. Juss. act. 1742. p. 131. t. 7. Guett. stamp. 2. p. 26.*

*Gramen junceum, capitulis quatuor longissimis filamentis donatis. Pluk. alm. 180. t. 35. f. 2. Moris. hist. 3. f. 8 & 9. f. 30.*

*Habitat ad Europæ littora lacuum.*

\* \* *Caule ramoso.*

17. **PLANTAGO** caule ramoso herbaceo; foliis subdentatis recurvatis, capitulis aphyllis. *Hort. upf. 28. Mat. med. 30.*

*Plantago caule ramoso. Virid. cliff. 9. Roy. lugdb. 401.*

*Psyllium majus erectum. Bauh. pin. 191.*

*Psyllium. Dod. pempt. 115.*

*Habitat in Europa australi, inter segetes. ☉*

*Caules diffusi, villosi. Folia inferiora opposita; superiora terna s. quaterna; linearia, recurva, dentata, villosa-viscida. Pedunculi folio longiores. Capitula aphylla Bracteis & calycinis foliolis linearibus.*

18. **PLANTAGO** caule ramoso herbaceo, foliis integerrimis reflexis, capitulis foliosis. *Hort. upf. 29.*

*Kali 111. Alp. ægypt. 128?*

*Habitat in Ægypto. ☉*

*Caulis altus, villosus, diffusus. Folia linearia, integerima, deflexa, longissima, nuda, basi multum villosa, opposita; superiora sæpe terna. Pedunculi longi. Capituli squamæ quatuor infimæ enatæ in foliola lanceolata, capitulo sæpe longiora. Bractææ inter flosculos cuneiformes, retusæ, concavæ, lateribus dilatatæ.*

19. **PLANTAGO** caule ramoso fruticoso, foliis filiformibus integerrimis strictis, capitulis subfoliatis. *cynops.*

*Plantago perennis*, foliis integerrimis, caule ramoso diffuso. *Hort. cliff.* 47.

*Psyllium sempervirens*. *Morif. hist.* 3. p. 262. f. 8. t. 17. f. 1.

*Psyllium majus supinum*. *Bauh. pin.* 191. *Bauh. hist.* 3. p. 513.

*Habitat in Gallo-provincia, Italia.* ☿

*Caulis palmaris, durus, lignosus, fruticosus, purpureascens, nudus. Folia opposita, subulata f. filiformia, canaliculato-trigona, erecta, integerrima, subvillosa. Pedunculi axillares, longitudine foliorum. Capituli squamæ infimæ mucronatæ, patulæ, mediocres. Bractææ, inter flosculos, ovate, concavæ.*

afra.

20. *PLANAAGO* caule ramoso fruticoso, foliis lanceolatis dentatis, capitulis aphyllis.

*Psyllium, foliis crenatis, indicum.* *Bauh. pin.* 191. *prodr.* 99. *Morif. hist.* 3. p. 262. f. 8. t. 19. f. 4.

*Bauh. hist.* 3. p. 513.

*Psyllium foliis laciniatis.* *Bocc. sic.*

*Habitat in Sicilia, Barbaria.* ☿

*Caulis pedalis, frutescens, erectus, pubescens. Folia lanceolata, dentata, subvillosa, striata. Spicæ aphyllæ, plures ad apices ramorum.*

## SCOPARIA.

dulcis.

1. *SCOPARIA* foliis ternis, floribus pedunculatis.

*Capraria foliis ternis, corollis quadripartitis.* *Hort. cliff.* 320. *Roy. lugdb.* 301.

*Samoloides.* *Berh. lugdb.* 2. p. 265. \*

*Veronica americana erecta frutescens & ramosa.* *Herm. parad.* 241. t. 241.

*Veronica fruticosa erecta dulcis, hexangulari caule.* *Sloan. jam.* 81. *hist.* 1. p. 195. t. 108. f. 2.

*phytenoides americanum, flore albo tetrapetalo.* *Fluk alm.* 296. t. 215. f. 1.

β. *Tupelcava.* *Pis. bras.* 246.

*Habitat in Jamaica, Curassao.* ☉

*Planta Pisonis in herbario subvillosa erat.*

procumbens 2. *SCOPARIA* foliis quaternis, floribus sessilibus. *Jacq. amer.* 12.

*Habitat in America calidiorr.* ☉

## RHACOMA.

1. RHACOMA. *Amoen. acad.* 5. p. 393. \* Crossopetalum.  
 Crossopetalum fruticosum tenue, foliis ovatis tenuissimis denticulatis. *Brown. jam.* 145. t. 17. f. 1.  
*Habitat in Jamaica.* §

## CENTUNCULUS.

1. CENTUNCULUS. *Dill. giff.* 161. gen. III. *E. N.* minimus.  
*C. 5. app.* 92. t. 10. f. 31.  
 Centunculus foliis alternis ovatis. *Fl. suec.* 129, 136.  
*Dalib. parif.* 52.  
 Anagallidistrum exiguum, foliis lanceolatis alternis.  
*Mich. gen.* 14. t. 18.  
 Anagallis paludosa minima. *Vaill. parif.* 12. t. 4. f. 2.  
 Alsine palustris minima, flosculis albis. *Mentz. pug.*  
*t. 4. f. 5.*  
*Habitat in Italiæ, Galliæ, Germaniæ, Scaniæ arenosis subudis.* ©

## SANGUISORBA.

1. SANGUISORBA spicis ovatis. *Hort. cliff.* 39. *Fl. officinalis.*  
*suec.* 130, 137. *Mat. med.* 51. *Roy. lugdb.* 240. *Dalib. parif.* 51.  
 Pimpinella sanguisorba major. *Bauh. pin.* 160.  
 Pimpinella sylvestris s. sanguisorba major. *Dod. pempt.*  
 105.  
 3. Pimpinella major rigida præalta auriculata subnuda.  
*Bocc. mus.* 2. p. 19. t. 7.  
*Habitat in Europæ pratis siccioribus.* 24  
 2. SANGUISORBA spicis cylindricis. † media.  
 Sanguisorba tetraſtemon, ſtaminibus tubo longioribus, ſpicis cylindricis. *Zinn. gzt.* 239.  
 Pimpinella minor de canada. *Zanon. hiſt.* t. 138.  
 Pimpinella canadensis major, ſpica brevior rubra, foliis lævibus. *Morif. hiſt.* 3. p. 264. f. 8. t. 18. f. 2.  
*Habitat in Canada.* 24  
*Spicæ cylindricæ longiores, & ſtamina longiora, quam in officinali*  
 2. SANGUISORBA ſpicis longiſſimis. *Hort. cliff.* 39. canadensis  
*Roy. lugdb.* 240.  
 Pimpinella sanguisorba canadensis major, ſpica longiore alba. *Morif. hiſt.* 3. p. 264. f. 8. t. 18. f. 12.

*Pimpinella maxima canadensis longius spicata.* *Corn. canad.* 175. t. 174. *Barr. rar.* 18. f. 739.  
*Habitat in Canada.* 24

### CISSUS.

- vitiginea.* 1. *CISSUS* foliis cordatis subquinelobis tomentosis.  
*Cissus.* *Fl. zeyl.* 60. \* *Amæn. acad.* 1. p. 390. \*  
*Arbuscula baccifera circumplicatilis*, vitigineo folio  
subtus lanato, fructu racemoso fervidiore odore.  
*Pluk. mant.* 27. t. 337. f. 2.  
*Funis crepitans.* *Rumph. amb.* 5. p. 446. t. 164.  
*Schunambu-valli.* *Rhed. mal.* 7. p. t. 11.  
*Habitat in India.* 5
- Cordifolia.* 2. *CISSUS* foliis cordatis integerrimis.  
*Vitis folio subrotundo, uva corymbosa cærulea.*  
*Plum. gen.* 18. ic. 159. f. 3.  
*Habitat in America.*
- Sicyoides* 3. *CISSUS* foliis ovatis nudis fetaceo-ferratis.  
*Irisiola scandens*, foliis oblongo-ovatis ad margines  
denticulis fetaceis. *Brown. jam.* 147. t. 4 f. 1, 2.  
*Bryonia alba geniculata, violæ foliis, baccis e viridi-purpurascens.* *Sloan. jam.* 106. *hist.* 1. p. 233.  
t. 144. f. 1. *Raj. suppl.* 347.  
*Habitat in Jamaica.*
- acida.* 4. *CISSUS* foliis ternatis oblongis carnosissimis incisissimis.  
*Sicyos trifoliata.* *Sp. pl.* 1. p. 1013.  
*Irisiola triphylla scandens & claviculata, foliis crassissimis ferratis.* *Brown. jam.* 147.  
*Bryonia alba triphylla geniculata, foliis crassissimis acidissimis.* *Sloan. jam.* 106. *hist.* 1. p. 233. t. 142. f. 6. *Raj. suppl.* 347.  
*Bryonioides trifoliatum indicum, foliis succulentissimis crassissimis & crenatis.* *Pluk. alm.* 71. t. 152. f. 2.  
*Vitis trifolia minor corymbosa, acinis nigrioribus turbinatis.* *Flum. spec.* 18. t. 259. f. 5.  
*Habitat in America.*
- trifoliata.* 5. *CISSUS* foliis ternatis subrotundis subdentatis.  
*Irisiola triphylla scandens, foliis ovatis subdentatis petiolo communi marginato, calycibus majoribus.*  
*Brown. jam.* 147.

Bry-

*Bryonia alba triphylla maxima.* Sloan. *jam* 106. *hist.* 4.  
p. 233. t. 144. f. 2. *Raj. suppl.* 347.  
*Habitat in Jamaica.*

## EPIMEDIUM.

1. EPIMEDIUM. *Hort. cliff.* 37. *Hort. upf.* 29. *Roy. alpinum.*  
*lugdb.* 402. *Dod. pempt.* 599. *Lob. hist.* 176.  
*Habitat in Alpium Euganeorum, Ligurinarum, Gret-*  
*tanensium, Ponteborum umbrosis.* 24.

## CORNUS.

1. CORNUS arborea, involucro maximo: foliolis ob- *florida*  
cordalis. *Hort. cliff.* 38. *Hort. upf.* 29. *Roy.*  
*lugdb.* 249. *Gron. virg.* 17. *Kalm. it.* 2. p. 341. 3.  
3. p. 104.  
Cornus mas virginiana, flosculis in corymbo digestis  
a perianthio tetrapetalo albo radiatim cinctis. *Pluk.*  
*alm.* 120. t. 20. f. 3. *Cuesb. carol.* 27. t. 27.  
*Habitat in Virginiae sylvis.* 5
2. CORNUS arborea, umbellis involucrum æquantibus. *mas.*  
*Hort. cliff.* 38. *Hort. upf.* 29. *Roy. lugdb.* 249.  
*Dalib. parif.* 52.  
Cornus sylvestris mas. *Bauh. pin.* 447.  
Cornus mas pumillo. *Cus. hist.* 1. p. 13.  
3. Cornus hortensis mas. *Bauh. pin.* 447.  
*Habitat in sepibus Austriae.* 5
3. CORNUS arborea, cymis nudis. *it. W-goth.* 32. *sanguinea.*  
Cornus umbellis involucro multoties longioribus.  
*Hort. cliff.* 38. *Roy. lugdb.* 249. *Hall. helv.* 463.  
*Dalib. parif.* 52. *Fl. suec.* 131.  
Cornus femina. *Bauh. pin.* 447.  
Virga sanguinea. *Dod. pempt.* 782.  
*Habitat in Europæ, Asiæ, Americæ borealis dume-*  
*tis.* 5
4. CORNUS herbacea, ramis binis. *Fl. lapp.* 65. t. 5. *suecica.*  
f. 3. *Fl. suec.* 132, 138. *Roy. lugdb.* 249. *Ord. dan.* t. 5.  
Cornus pumila herbacea, Chamæpericlymenum dicta.  
*Dill. elik.* 108. t. 91.  
Periclymenum humile. *Bauh. pin.* 302. *norvegicum.*  
*Bauh. pin.* 302. *Buxb. ast.* 3. p. 268.  
*Habitat in Svecia, Norvegia, Russia.* 24

- canadensis. 5. CORNUS herbacea, ramis nullis. *Aman. acad.* 1. p. 157.  
 Pyrola; albes flore, brasiliiana. *Bauh. prodr.* 100. pin. 191. *Burf. X.* 106.  
*Habitat in Canada.* 2.  
*Involucrum, quam in priore, longius.*

## FAGARA.

- Pterota. 1. FAGARA foliolis emarginatis. *Aman. acad.* 5. p. 393. \*  
 Schinus foliis pinnatis: foliolis oblongis, petiolo marginato articulato inermi. *Mat. med.* 533. *Sp. pl.* 1. p. 389.  
 Pterota subspinosa, foliis minoribus per pinas marginato-alatas dispositis, spicis geminatis alaribus. *Brown. jam.* 146. t. 5. f. 1.  
 Lauro affinis, jasmini alato folio, costa media membranulis utrinque exstantibus alata, ligno duritie ferro vix cedens. *Sloan. jam.* 137. *hist.* 2. p. 25. t. 162. f. 1. *Raj. dendr.* 86.  
*Habitat in Jamaica.* §
- Tragodes. 2. FAGARA articulis pinnarum subtus aculeatis. *Jacq. amer.* 13.  
 Schinus tragodes. *Sp. pl.* 1. p. 369.  
 Schinoides petiolis subtus aculeatis. *Hort. cliff.* 489. \*  
 Rhus obsoniorum similis leptophyllos tragodes americana spinosa, rachi medio appendiculis aucto. *Pluk. alm.* 319. t. 107. f. 4. ?  
*Habitat in America.* §  
*Præcedenti nimis affinis.*
- Piperita. 3. FAGARA foliolis crenatis, †  
 Piper japonicum Sio & Sanfio. *Kempf. amon.* 892. t. 893.  
*Habitat in Japonia.* §

## TOMEX.

- tomentosa. 1. TOMEX.  
 Tomex tomentosa. *Fl. zeyl.* 59. \* *Aman. acad.* 1. p. 389.  
 Arbor malabarica Illa dicta. *Burm. zeyl.* 26.  
*Habitat in India.* §

## PTELEA.

1. PTELEA foliis ternatis.
- Mill. dict.*
- 211.

trifoliata.

*Ptelea. Hort. cliff.* 36.*Ptelea* similis. *Zinn. gætt.* 207. \**Frutex virginianus trifolius, ulmi samaris. Pluk. alm.*159. *Dill. elth.* 147. f. 122. f. 148.*Habitat in Virginia.* ☿

2. PTELEA foliis simplicibus.

viscosa.

*Dodonæa. Hort. cliff.* 144. *Fl. zeyl.* 141. \* *Roy. lugdb.*206. *Jacq. amer.* 19.*Triopteris erecta fruticosa, foliis oblongis acuminatis, ramulis gracilibus. Brown. jam.* 191. t. 18. f. 1.*Aceri* f. *Paliuro* affinis, angusto oblongo ligustri folio, flore tetrapetalo herbaceo. *Sloan. jam.* 138. *hist.* 2.p. 27. t. 162. f. 3. *Raj. dendr.* 94.*Caryophyllaster litoreus. Rumph. amb.* 4. p. 110. t. 50.*Carpinus viscosa, salicis folio integro oblongo. Burm.**zeyl.* 55. t. 23.*Arbustula viscosa, æleagni foliis late virentibus, americana triccocos. Pluk. phyt.* 142. f. 1.*Habitat in Indiis, locis arenosis.* ☿*Hanc arborem idem genus cum Ptelea agnoscere observavit B. Jussæus, at Florem trifidum hexandrum Browni-  
us, & quadrifidum octandrum Jacquin tribuit, mihi non floruit.*

## LUDVIGIA.

1. LUDVIGIA foliis alternis lanceolatis.

alternifolia

*Ludvigia capsulis cubicis apice perforatis. Hort.  
ups.* 30.*Ludvigia capsulis subrotundis. Hort. cliff.* 491. *Roy.  
lugdb.* 252.*Ludvigia. Gron. virg.* 17.*Lysimachia non papposa, flore luteo majore, filiqua  
caryophylloide minore, ex virginia. Pluk. alm.* 235.  
t. 203. f. 2.*Frutex saligneis foliis, caule purpureo, capsularis.**Pluk. amab.* 99. t. 412. f. 1.*Habitat in Virginia.* ☉

2. LUDVIGIA foliis oppositis lanceolatis, capsulis pendunculatis.

*Ludvigia caule diffuso, foliis lanceolatis capsulis pe-  
dunculatis folio dimidio brevioribus. Fl. zeyl.* 66. \*

Ca-



*cliff.* 483. *Fl. jvec.* 134, 140. *Mat. med.* 52. *Dalib. parif.* 52.

*Tribulus aquaticus.* *Bauh. pin.* 194. *Cam. epit.* 715.

*Panover-Tierraua.* *Rheed. mal.* 11. p. 65. t. 33.

*Habitat in Europæ australis, Asiæque stagnis limosis.* ☉

## DORSTENIA.

*Contrajerva.* 1. **DORSTENIA** acaulis, foliis pinnatifido-palmatis, ferratis, floribus quadrangulis.

*Dorstenia scapis radicatis.* *Hort. cliff.* 32. *Mat. med.* 53.

*Dorstenia dentariæ radice, sphondylii folio, placenta ovali.* *Houst. aët. n.* 421. f. 1.

*Dorstenia sphondylii folio, dentariæ radice.* *Plum. gen.* 29. ic. 119.

*Drakena radix.* *Clus. exot.* 83.

*Cyperus longus odoratus peruanus.* *Bauh. pin.* 14.

*Tuzpatlis.* *Hern. mex.* 147.

*Habitat in nova Hispania, Mexico, Peru, Tabago, insula Vicentii.* 2.

*Drakena.* 2. **DORSTENIA** acaulis, foliis pinnatifido-palmatis integerrimis floribus ovalibus.

*Dorstenia dentariæ radice, sphondylii folio, placenta ovali.* *Houst. aët. n.* 421. f. 1.

*Habitat in Vera Cruce.* 2.

*houstoni.* 3. **DORSTENIA** acaulis, foliis cordatis angulatis acutis, floribus quadrangulis.

*Dorstenia dentariæ radice, folio minus laciniato, placenta quadrangulari & undulata.* *Houst. aët. angl.* 421. f. 2.

*Habitat in Campechia.* 2.

*caulescens.* 4. **DORSTENIA** pedunculis culinis. *Hort. cliff.* 32.

*Parietaria racemosa, foliis ad oras villosis.* *Plum. spec.* 10. ic. 120. f. 1.

*Habitat in America meridionali.* 2.

## ELÆAGNUS.

*angustifolia.* 1. **ELÆAGNUS** foliis lanceolatis. *Roy. lugdb.* 250. *Hort. upf.* 31.

*Elæagnus.* *Cam. epit.* 106. *Hort. cliff.* 33.

*Olea sylvestris, folio molli incano.* *Bauh. pin.* 472.

*Ha-*

## TETRANDRIA MONOGYNIA.

177

*Habitat in Bohemia, Hispania, Syria, Cappadocia,  
inque Aquis sextiis locis subhumidis.*

2. ELÆAGNUS foliis ellipticis. *Amæn. acad.* 4. p. spinosa.  
305. \*

*Elæagnus matthioli incolis Seisefum. Rauw. it.* 112,  
276.

*Habitat in Ægypto.* ‡

3. ELÆAGNUS foliis ovatis. *Roy. lugdb.* 250. *Fl. latifolia.*  
*zeyl.* 58.

*Eleagnus foliis rotundis maculatis. Burm. zeyl.* 92.  
t. 39. f. 2.

*Habitat in Zeylona.* ‡

## BRABEJUM.

1. BRABEJUM. *Hort. cliff.* 36. *Roy. lugdb.* 400. *stellatifo-*  
*Amygdalus æthiopica, fructu holosericeo. Breyn. cent.* lium.

I. D. I.

*Arbor æthiopica hexaphylla, foliis circa caulem ad  
intervalla senis. Pluk. alm.* 47. t. 265. f. 3.

*Habitat in Æthiopia.* ‡

## KRAMERIA.

1. KRAMERIA. *Læfl. it.* 195. \* *Ixina.*

*Habitat in Cumana America.* ‡

*Folia æterna, lanceolata. Racemus terminalis floribus  
alternis.*

## RIVINA.

1. RIVINA floribus tetrandris. *humilis,*  
*Rivina. Hort. cliff.* 35. *Roy. lugdb.* 207.

2. Rivina humilis racemosa, baccis puniceis. *Plum. canescens.*  
*gen.* 48.

*Solanoides americana, circææ foliis canescentibus.*  
*Tournef. art.* 1706.

*Solanum barbadense racemosum minus tinctorium.*

*Pluk. alm.* 353. t. 112. f. 2.

*Amarantus baccifer, circææ foliis. Comm. hort.* 1. p.  
127. t. 66.

*Habitat in Caribæis, Jamaica, Barbados.* ‡

2. RIVINA floribus octandris dodecandrisve. *Amæn. octandra.*  
*acad.* 4. p. 305. \* *Læfl. it.* 207.

*Linnei Spec. plant. pars I.*

M

Ri-

*Rivina scandens racemosa*, amplioribus foliis, baccis violaceis. *Plum. gen.* 48. *ic.* 241.

*Rivina farmentosa*, sarmentis crassioribus, foliis ovatis, floribus spicatis dodecandris. *Brown, jam.* 149, *t.* 23. *f.* 2.

*Habitat in America calidiore.* ‡

### SALVADORA.

perfica.

1. *SALVADORA*. *Garc. act. angl.* 1749. n. 491. *Amæn. acad.* 3. p. 21.

*Rivina paniculata*. *Syst. nat.* 10. p. 899.

*Habitat ad Sinum Perficum, in India orientali.* ‡

*Frutex foliis oppositis, Racemo terminali composito.*

### CAMPHOROSMA.

monspeliaca.

1. *CAMPHOROSMA* foliis hirsutis linearibus. *Amæn. acad.* 1. p. 392. *Mat. med.* 43.

*Camphorosma*. *Sauv. monsp.* 45.

*Selago caule procumbente, foliis sparsis.* *Hort. cliff.* 321. *Roy. lugdb.* 300.

*Camphorata hirsuta.* *Bauh. pin.* 486. *Tournef. act.* 1705. p. 313. *Buxb. cent.* 1. p. 18. *t.* 28. *f.* 1.

*Camphorata monspeliensium.* *Bauh. hist.* 3. p. 379. *Lob. adv.* 174.

*Habitat in Hispaniæ, Narbonæ, Tartariæ arenosis.*

acuta.

2. *CAMPHOROSMA* foliis subulatis rigidis glabris. *Amæn. acad.* 1. p. 393.

*Camphora altera.* *Tabern. h. st.* 57.

*Camphoratae congener.* *Bauh. pin.* 486. (*variæ*)

*Anthyllis altera italorum.* *Lob. ic.* 404. *Dalech. hist.* 1150.

*Habitat in Italia, Tartaria.* 24

glabra.

3. *CAMPHOROSMA* foliis subtriquetris glabris inermibus. *Amæn. acad.* 1. p. 393.

*Camphorata foliis confertissimis inermibus lævibus.* *Hall. helv.* 193.

*Camphorata glabra.* *Bauh. pin.* 286. *Dalech. hist.* 1179.

*Habitat in Helvetia.* 24

### ALCHEMILLA.

vulgaris.

1. *ALCHEMILLA* foliis lobatis. *Fl. fvec.* 135, 141. *Mat. med.* 54.

Al-

- Alchemilla foliis palmatis. Hort. cliff. 38. Roy. lugdb. 231.*  
*Alchemilla foliis simplicibus. Fl. lapp. 66.*  
 ̢. *Alchemilla vulgaris. Bauh. pin. 319.*  
*Alchemilla minor. Tournef. infl. 502.*  
*Habitat in Europæ pascuis. 24*  
 2. **ALCHEMILLA** foliis digitatis ferratis. *Fl. lapp. 61. alpina.*  
*Fl. fvec. 136, 142. \* Hort. cliff. 39. Roy. lugdb. 232.*  
*Alchemilla perennis incana argentea, f. sericea satinum*  
*provocans. Moris. hist. 2. p. 195. f. 2. t. 20. f. 3.*  
*Tormentilla alpina, foliis sericeis. Bauh. pin. 326.*  
 ̢. *Alchemilla alpina pubescens minor. Tournef. infl. 508. hybrida.*  
*Pluk. phyt. 240. f. 1.*  
*Habitat in Alpibus Europæ. 24*  
 3. **ALCHEMILLA** foliis quinatis multifidis glabris. *pentaphyl.*  
*Alchemilla alpina quinquefolia. Bauh. pin. 320. pro-* *lea.*  
*dr. 138.*  
*Alchemilla alpina pentaphyllea minima, lobis fimbri-*  
*atis. Bocc. mus. 1. p. 18. t. 1.*  
*Habitat in monte Canisio, Furca, Gotthardo Rheti-*  
*orum. 24*

## DIGYNIA.

### APHANES.

1. **APHANES.** *Hort. cliff. 39. Fl. fvec. 137, 143. Roy. arvensis.*  
*lugdb. 231 Gron. v rg. 17. Dalb. parif. 53.*  
*Chærophylo nonnihil similis. Bauh. pin. 182.*  
*Alchemilla minima montana. Col. ceph. 145. t. 146.*  
*Habitat in Europæ & Orientis arvis. ☉*

### CRUCITA.

1. **CRUCITA.** *Læfl. it. 203. \** *hispanica,*  
*Habitat in Cumana America.*  
*Caulis altus. Folia opposita lanceolata integerrima. Flores.*  
*spicati, in paniculam collecti.*

### BUFONIA.

1. **BUFONIA.** *Sauv. monsp. 141. Aman. acad. 1. p. 386. tenuifolia.*  
*Læfl. it. 44.*  
*Alfincoides Raj. angl. 3. p. 346.*

M 2

Her-

*Herniaria*, angustissimo gramineo folio, erecta. *Magn. hort.* 97. t. 97.

*Alfina polygonoides tenuifolia*, flosculis ad longitudinem caulis velut in spicam dispositis *Pluk. alm.* 22. t. 75. f. 3.

*Habitat* in Anglia, Gallia, Hispania. 21

*Stamina* 4 numeravit in flore *Læfing*, confirmavit *C. Alstræmer*.

## HAMAMELIS.

*virginiana*. 1. *HAMAMELIS*. *Gron. virg.* 139. *Cold. noveb.* 18. *Catesb. car.* 3. p. 2. t. 2. *Duham. arb.* 1. p. 287. t. 114.

*Trilopus*. *Mitch. gen.* 22.

*Pistacia virginiana nigra*, coryli foliis. *Pluk. alm.* 298.

*Habitat* in Virginia.

## CUSCUTA.

*europæa*. 1. *CUSCUTA* floribus sessilibus. *Cuscuta nuda repens filiformis*. *Fl. svec.* 138. 144. *Mat. med.* 55. *Dalib. parif.* 53.

*Cuscuta nuda repens funiformis*. *Sauv. monsp.* 11.

*Cuscuta major*. *Bauh. pin.* 219.

*Cuscuta*. *Cam. epit.* 984.

*epithymum*. 2. *Cuscuta nuda repens filiformis*. *Sauv. monsp.* 11.

*Epithymum* f. *Cuscuta minor*. *Bauh. pin.* 219.

*Habitat* in Plantis Europæ, parasitica. ☉

*Parasitica planta ecotyledonis, dehiscente semine absque cotyledonibus, extendit spirale corpusculum, nec terram petit radicanda, sed absque omni radice contra solem volubilis ascendit plantas, quasi emissis vasis exhaurit; nec Folia gerit præter squamam, hinc inde minimam, membranaceam, arctam sub ramo. Flos apud nos sæpius quadrifidus; in Europa australi sæpius quinquefidus.*

*americana*. 2. *CUSCUTA* floribus pedunculatis. *Læf. it.* 216.

*Cuscuta caule aphylo volubili repente*. *Gron. virg.* 18.

*Cuscuta* inter majorem & minorem media, filamentis longis & floribus late super arbores & campos se extendens. *Sloan. jam.* 85. *hist.* 1. p. 201. t. 128. f. 4.

*Habitat* in Virginia.

## HYPECOUM.

1. **HYPECOUM** filiquis arcuatis compressis articulatis. procumbens.  
*Hort. upf.* 31.  
*Hypecoum.* *Bauh. pin.* 172. *Dod. pempt.* 449. *Hort. cliff.* 38. *Roy. lugdb.* 402.  
*Habitat inter Archipelagi, Narbonæ & Salmanticensis segetes.* ☉
2. **HYPECOUM** filiquis cernuis teretibus cylindricis. pendulum.  
*Hort. upf.* 31. *Sauv. monsp.* 263.  
*Hypecoum* filiquis pendentibus non articulatis bivalvibus incurvis. *Morif. hist.* 2. p. 280.  
*Hypecoi altera species.* *Bauh. pin.* 172.  
*Cuminum sylvestre filiquatum pone.* *Dalech. hist.* 698.  
*Habitat in Galloprovincia.* ☉
3. **HYPECOUM** filiquis erectis teretibus torulosis. erectum.  
*Hort. upf.* 32.  
*Hypecoum tenuifolium, filiquis erectis teretibus.* *Amm. ruth.* 58. t. 9. *Hall. gætt.* 229.  
*Habitat in Dauria.*

## TETRAGYNIA.

## I L E X.

\* *Flores quadrifidi.*

1. **ILEX** foliis ovatis acutis spinosis. *Hort. cliff.* 40. *Aquifolium*  
*Hort. upf.* 32. *Roy. lugdb.* 400. *Gros. virg.* 12. *Da- lib. parif.* 54.  
*Ilex aculeata baccifera.* *Bauh. pin.* 425.  
*Habitat in Europa australiori, Japonia, Virginia.* ☿
2. **ILEX** foliis ovato-lanceolatis ferratis. *Hort. cliff.* 40. *Cassine.*  
*Mat. med.* 56. *Roy. lugdb.* 400.  
*Aquifolium carolinense, foliis dentatis, baccis rubris.*  
*Catesb. car.* 1. p. 31. t. 31.  
*Habitat in Carolina.* ☿
3. **ILEX** foliis lato-lanceolatis obtusis integerrimis. asiatica.  
*Habitat in India Asia.* ☿

\* \* *Flores trifidi.*

4. **ILEX** foliis cuneiformibus tricuspidatis. cuneifolia.  
*Ilex aculeata, foliis tricuspidi.* *Plum. ic.* 118. f. 2.  
*Habitat in America meridionali.* ☿

Dodonæa.

## 5. ILEX foliis pinnatis †

Dodonæa aquifolii folio tricuspidato. *Plum. gen. 20. ic. 118. f. 1.*

β. Dodonæa aquifolii folio anguloso non aculeato. *Plum. gen. 20.*

*Habitat in America meridionali. ‡*

*Folia pinnata: foliolis 21, sessilibus, rigidis, ovatis, acuminatis, in medio utrinque denticulo angularis. Flores racemosi ut in Rhoe; forte diversi generis?*

## C O L D E N I A.

procumbens

1. COLDENIA. *Fl. zeyl. 69. Mill. dict. \**

*Teucri facie bisnagarica tetracoccus rostrata. Pluk. alm. 363. t. 64. f. 6. Raj. suppl. 281. Morif. hist. 3. p. 423.*

*Habitat in India. ○*

*Frustrificationem ulterius examinent alii in vivente planta, ego perfectam non vidi.*

## POTAMOGETON.

natans.

## 1. POTAMOGETON foliis oblongo-ovatis petiolatis

*natantibus. Hort. cliff. 40. Fl. succ. 139. 145. Roy. lugdb. 212. Hall. helv. 199. Dalib. parif. 54.*

*Potamogeton foliis ovato-oblongis natantibus. Fl. lapp. 68.*

*Potamogeton rotundifolium. Bauh. pin. 193.*

β. Potamogeton foliis lanceolato-oblongis, petiolis longis. *Gron. virg. 139.*

*Habitat in Europæ lacubus & fluviis. 2*

perfoliatum.

## 2. POTAMOGETON foliis cordatis amplexicaulibus.

*Fl. lapp. 69. Fl. succ. 140. 145. Roy. lugdb. 212. Hall. helv. 199. Dalib. parif. 54.*

*Potamogeton perfoliatum. Raj. hist. 188.*

*Potamogeton foliis latis splendentibus. Bauh. pin. 195.*

*Potamogeton rotundifolium alterum. Læs. pruss. 205. t. 65.*

*Habitat in Europæ lacubus fluviisque argillofis. 2*

densum.

3. POTAMOGETON foliis ovatis acuminatis oppositis confertis, caulibus dichotomis, spica quadriflora, *Crust. stamp. 1. p. 102. \**

*Potamogeton minus*, foliis densis mucronatis non serratis. *Magn. monsp.* 303.

*Fontinalis media lucens*. *Bauh. hist.* 3. p. 769. *Raj. hist.* 189.

*Tribulus aquaticus minor alter*. *Clus. hist.* 2. p. 252.

*Habitat* in Gallia, Italia.

*Pedunculus e dichotomia caulis brevissimus*. *Spica 4-flora*.

4. **POTAMOGETON** foliis lanceolatis planis in petiolos definitibus. *Hort. cliff.* 40. *Fl. succ.* 141. 147.

*Roy. lugdb.* 212. *Dalib. parif.* 54.

*Potamogeton foliis angustis splendentibus*. *Bauh. pin.* 193.

*Fontinalis lucens major*. *Bauh. hist.* 3. p. 769.

*Habitat* in Europæ lacubus, stagnis, fluviis argillofis. 2

5. **POTAMOGETON** foliis lanceolatis alternis undulatis serratis. *Roy. lugdb.* 212. *Fl. succ.* 142. 148. *It. sc. an.* 142. \*

*Potamogeton foliis crispis* f. *Lactuca ranarum* sarmenis planis. *Bauh. pin.* 193.

*Habitat* in Europæ fossis & rivulis.

6. **POTAMOGETON** foliis lanceolatis oppositis subundulatis. *ferratum.*

*Potamogeton foliis lanceolatis obscure undulatis, caulibus longe ramosis*. *Gurt. stamp.* 1. p. 102. *Dalib. parif.* 55.

*Potamogeton longo ferrato folio*. *Bauh. pin.* 193.

*Habitat* in Europæ rivulis.

7. **POTAMOGETON** foliis linearibus obtusis, caule compresso. *Hort. cliff.* 40. *Fl. succ.* 143. 149. *Roy. lugdb.* 213. *Dalib. parif.* 55.

*Potamogeton caule plano, foliis gramineis undulatis, spica exigua*. *Hall. helv.* 201.

*Potamogeton caule compresso, folio graminis canini*. *Raj. hist.* 189.

*Habitat* in Europæ fossis paludosis.

8. **POTAMOGETON** foliis setaceis parallelis approximatis distichis. *Fl. succ.* 145, 150. *Dalib. parif.* 55.

*Potamogeton foliis linearibus acutis longissimis alternis confertis*. *Roy. lugdb.* 213.



- Potamogeton gramineum ramosum. *Bauh. pin.* 193.  
*prodr.* 101.  
*Habitat in Europæ fossis & paludibus.*
- setaceum. 9. POTAMOGETON foliis lanceolatis oppositis acuminatis. *Roy. lugdb.* 213. *Dalib. parif.* 55.  
 Potamogeton ramosum angustifolium. *Bauh. pin.* 193.  
*prodr.* 101.  
*Habitat in Europæ fossis paludosis.*
- gramineum. 10. POTAMOGETON foliis linearilanceolatis alternis sessilibus stipula latioribus. *Fl. succ.* 144. 151.  
 Potamogeton gramineum latiusculum, foliis & ramificationibus dense stipatis. *Raj. angl.* 3. p. 149. t. 4. f. 3. *Fl. lapp.* 70.  
 Potamogeton gramineum latifolium. *Laf. pruss.* 206. t. 66.  
*Habitat in Europæ fossis & paludibus. ☉*
- marinum. 11. POTAMOGETON foliis linearibus alternis distinctis inferne vaginantibus. *Fl. succ.* 146, 152.  
 Potamogeton pusillum fluitans. *Bocc. sic.* 42. t. 20. f. 5. *It. goth.* 221:  
 Potamogeton maritimum ramosissimum, grandiusculis capitulis, capillaceo folio. *Pluk. alm.* 305. t. 216. f. 5.  
*Habitat in Europa ad maris littora. ☉*
- pusillum. 12. POTAMOGETON foliis linearibus oppositis alternisque distinctis: basi patentibus, caule tereti. *Fl. succ.* 147, 153. *Dalib. parif.* 56.  
 Potamogeton foliis linearibus alternis remotis. *Roy. lugdb.* 213.  
 Potamogeton minimum, capillaceo folio. *Bauh. pin.* 193. *prodr.* 101.  
 Potamogeton gramineum tenuifolium. *Laf. pruss.* 206. t. 67.  
 Potamogeton pusillum, gramineo folio brevior. *Vaill. parif.* t. 32. f. 4.  
*Habitat in Europæ paludibus. ☉*

## R U P P I A.

- maritima. 1. RUPPIA. *Hort. cliff.* 436. *It. wgoth.* 186. \* *Fl. succ.* 2. n. 154. *Guett. flamp.* 2. p. 416.  
 Buccaferrea maritima, foliis acutissimis. *Mich. gen.* 72. t. 35.

*Potamogeton maritimum*, gramineis longioribus foliis, fructu fere umbellato. *Raj. angl.* 3. p. 134. t. 6. f. 1.

*Gramen maritimum fluitans cornutum.* *Bauh. pin.* 3. prodr. 7.

*Fucus folliculaceus*, fœniculi folio longiore. *Bauh. pin.* 365.

*Fucus ferulaceus* *Lob. ic.* 255.

*Habitat in Europæ maritimis.* ☉

## SAGINA.

1. *SAGINA* ramis procumbentibus, petalis brevissimis. *procumbens*  
*Fl. lapp.* 157. *Fl. suec.* 148. 155. *Roy. lugdb.* 452.

*Sagina scapis & ramis unifloris.* *Guett. stamp.* 2. p. 277.

*Alfinella muscoso flore repens.* *Dill. giff.* 81. *Raj. angl.* 4. p. 90.

*Alfine pusilla graminea, flore tetrapetalo.* *Segu. veron.* 421. t. 5. f. 3.

*Alfine minima, flore fugaci.* *Raj. suppl.* 501.

*Alfine tetrapetala, foliis angustis in origine latescens.* *Hall. helv.* 390.

- β. *Saxifraga graminea pusilla, foliis brevioribus crassioribus & succulentis.* *Raj. angl.* 3. p. 345.

*Habitat in Europæ pascuis sterilibus uliginosis aridis.*

2. *SAGINA* caule erecto unifloro triflorove. *erecta.*

*Sagina scapis unifloris.* *Guett. stamp.* 2. p. 276. *Dalib. parif.* 56.

*Alfine foliis caryophyllæis.* *Raj. angl.* 3. p. 344. t. 15. f. 4.

*Alfine verna glabra.* *Magn. monsp.* 14. *Vaill. parif.* 6. t. 3. f. 2.

*Habitat in Galliæ, Angliæ, Germaniæ sterilibus glareosis.*

3. *SAGINA* caule erecto, floribus oppositis. *virginica.*

*Sagina affinis planta minima, floribus albis.* *Clayt. mss.* 649.

*Habitat in Virginia inter muscos ad margines fontium.* *D. Gronovius.*

*Caulis digitalis, filiformis, erectus; Folia opposita, subulata, remota, minima. Flos terminalis & oppositi nonnulli propriis pedunculis. Perianthium 4partitum, erectum. Petala 4. oblonga. Stamina 4. longitudine calycis. Germen conicum, longitudo-*

*ne calycis. Stylus nullus. Stigma obtusum. Capsula pyramidalis, rostrata, unilocularis. Semina numerosa. An proprii generis?*

## TILLÆA.

- aquatica.* 1. **TILLÆA** erecta dichotoma, foliis acutis. *Fl. suec.* 2. n. 156.  
*Crassula caule dichotomo, floribus linearibus, floribus quadrifidis. Fl. suec.* 259. *Dalib. paris.* 98.  
*Crassula floribus quadrifidis. Hort. cliff.* 497.  
*Sedum minimum annuum, flore roseo tetrapetalo. Vaill. paris.* 181. t. 10. f. 2.  
*Habitat in Europæ inundatis. ☉*
- muscosa.* 2. **TILLÆA** procumbens. *Hort. ups.* 24. *Saxv. monsp.* 129.  
*Tillæa. Dalib. paris.* 43.  
*Tillæa muscosa annua, perfoliata. Mich. gen.* 22. t. 26.  
*Crassula foliis sessilibus connatis, floribus aggregatis in foliorum alis. Guett. stamp.* 1. p. 97.  
*Polygonum muscosum minimum. Bocc. sic.* 56. t. 29.  
*Sempervivum omnium minimum repens muscosum, polygóni faci. Bocc. mus.* 2. p. 36. t. 22.  
*Habitat in Italiæ, Siciliæ, Galliæ muscosis.*  
*Flores hujus sapius trifidi, antecedentium quadrifidi; Phil. Bot.* 178.



## Classis .V.

# PENTANDRIA

## MONOGYNIA.

### HELIOTROPIUM.

1. **HELIOTROPIUM** foliis lanceolato-ovatis , caule **peruvianum**  
fruticoso, spicis numerosis aggregato-corymbosis.  
**Heliotropium** foliis ovato-lanceolatis, spicis plurimis  
confertis, caule fruticoso. *Mill. dict. 1. 143.*  
*Habitat in Peru. b*
2. **HELIOTROPIUM** foliis cordato ovatis acutis sca- **indicum.**  
briusculis, spicis solitariis, fructibus bifidis. *Fl.*  
*zeyl. 70.*  
**Heliotropium** foliis ovatis acutis, spicis solitariis. *Hort.*  
*cliff. 45. Roy. lugdb. 405.*  
**Heliotropium americanum cæruleum.** *Dod. mem. 83.*  
*Fluk. phyt. 245. f. 4.*
3. **Heliotropium americanum cæruleum**, foliis hormini  
angustioribus. *Herm. lugdb. 307. Sloan. jam. 98.*  
*Habitat in India utraque ☉*  
*Semina unilocularia, per paria approximata.*
3. **HELIOTROPIUM** foliis ovatis integerrimis to- **europæum.**  
mentosus rugosis, spicis conjugatis. *Hort. upf. 33.*  
*Sauv. monsp. 305.*  
**Heliotropium** foliis ovatis integerrimis, spicis con-  
junctis. *Hort. cliff. 45. Roy. lugdb. 404.*  
**Heliotropium majus dioscoridis.** *Bauh. pin. 253.*  
*Habitat in Europa australi. ☉*
4. **HELIOTROPIUM** foliis ovatis integerrimis to- **supinum.**  
mentosus pl'catis, spicis solitariis.  
**Heliotropium minus supinum.** *Bauh. pin. 253.*  
**Heliotropium supinum** *Clus. hist. 2. p. 47.*  
*Habitat Salmanticæ juxta agros, Monspelii in litto-*  
*re. ☉*
5. **HELIOTROPIUM** foliis lineari-lanceolatis pilosis, fruticosum.  
spicis solitariis sessilibus. *Amæn. acad. 4. p. 394. \**  
**Heliotropium fruticosum hirsutum**, foliis lanceolatis  
mi-

minoribus, spicis singularibus terminalibus. *Brown. jam.* 151.

*Heliotropium minus*, lithospermi foliis. *Sloan. jam.* 95. *hist.* 1. p. 214. t. 139. f. 4.

*Habitat in Jamaica.* b

curassavi-  
cum.

6. *HELIOTROPIUM* foliis lanceolato-linearibus glabris aveniis, spicis conjugatis. *Hort. cliff.* 45. *Hort. upf.* 33. *Roy. lugdb.* 405. *Dalib. parif.* 57.

*Heliotropium indicum procumbens* glaucophyllon, floribus albis. *Pluk. alm.* 189. t. 36. f. 3.

*Heliotropium maritimum minus*, folio glauco, flore albo. *Sloan. hist.* 1. p. 213. t. 139. f. 3.

*Heliotropium curassavicum*, foliis lini umbilicatis. *Morif. hist.* 3. p. 452. f. 11. t. 31. f. 12.

*Heliotropium americanum procumbens*, facie lini umbilicatis. *Herm. par.* 183. t. 183.

*Habitat in Americæ calidioris maritimis.* ☉

orientale.

7. *HELIOTROPIUM* foliis linearibus glabris aveniis floribus sparsis lateralibus.

*Habitat in Asia.* ☉

*Planta parva, procumbens, repens. Folia linearia, alterna. Flores subsessiles, alterni, solitarii, inter folia sparsi.*

gnaphalodes

8. *HELIOTROPIUM* foliis linearibus obtusis tomentosis, pedunculis dichotomis, spicarum floribus quaternis, caule frutescente.

*Heliotropium arboreum maritimum tomentosum, gnaphalii americani foliis.* *Sloan. jam.* 93. *hist.* 1. p. 213.

*Heliotropium gnaphalodes littoreum frutescens americanum.* *Pluk. alm.* 182. t. 193. f. 5.

*Habitat in Barbados, Jamaica maritimis.* b

## MYOSOTIS.

scorpioides. 1. *MYOSOTIS* seminibus nudis: foliorum apicibus callosis. *Vir. cliff.* 149. *Fl. succ.* 149, 157. *Roy. lugdb.* 404. *Hort. cliff.* 45.

arvensis.

a. *Myosotis* foliis hirsutis. *Hort. cliff.* 46.

*Myosotis hirsuta arvensis major.* *Dill. giss.* 55. *Fl. lapp.* 74.

*Echium scorpioides arvense.* *Bauh. pin.* 254.

palustris.

b. *Myosotis* foliis glabris. *Hort. cliff.* 46. *Dalib. parif.* 58.

Myo-

*Myosotis glabra pratensis. Dill. giff. 67. Fl. lapp. 75.*

*Echium scorpioides palustre. Bauh. pin. 254.*

7. *Echium scorpioides minus, flosculis luteis. Bauh. pin. 254. prodr. 119. Burf. XIV. 6.*

*Habitat in Europæ a campis aridis &  $\beta$  aquosis scaturiginosis. ☉*

2. **MYOSOTIS** feminibus aculeatis glochidibus, foliis virginica, ovato-oblongis, ramis divaricatis.

*Myosotis feminibus hispidis, foliis lanceolato-ovatis. Gron. virg. 19.*

*Cynoglossum virginianum, flore & fructu minimo.*

*Pluk. alm. 126. Morif. hist. 3. p. 449. f. 11. t. 30. f. 9.*

*Habitat in Virginia. ☉*

*Semina interiori latere nuda, angulata, erecta, basi affixa, ut in sequenti, fructus nutans.*

3. **MYOSOTIS** feminibus aculeis glochidibus, foliis lanceolatis pilosis. *Fl. svec. 150, 158. Dalib. parif. 57. Lappula.*

*Lithospermum feminibus echinatis. Hort. cliff. 46. Roy. lugdb. 405.*

*Cynoglossum foliis lanceolatis, floribus in foliorum alis subsessilibus. Guett. stamp. 1. p. 92.*

*Cynoglossum minus. Bauh. pin. 257.*

*Habitat in Europæ argillofis, nudis, ruderatis. ☉*

4. **MYOSOTIS** feminibus nudis, foliis hispidis, racemis foliosis.

*Lithospermum feminibus lævibus, corollis vix calycem superantibus, foliis lanceolatis. Roy. lugdb. 405.*

*Sauv. monsp. 62.*

*Echium luteum minimum. Bauh. pin. 254.*

*Echioides lutea minima apula campestris. Col. ecphr. 1. p. 184. t. 185.*

*Anchusa lutea minima. Lob. ic. 312.*

*Habitat in Italia, Hispania, Narbona. ☉*

## LITHOSPERMUM.

1. **LITHOSPERMUM** feminibus lævibus, corollis officinale, calycem vix superantibus, foliis lanceolatis. *Hort.*

*cliff. 46. Fl. svec. 151. 159. Mat. Med. 57. Roy.*

*lugdb. 505. Dalib. parif. 58.*

*Lithospermum majus erectum. Bauh. pin. 258.*

*Habitat in Europæ ruderatis. 24*

- arvense.* 2. **LITHOSPERMUM** seminibus rugosis, corollis vix calycem superantibus. *Hort. cliff.* 46. *Fl. suec* 152. 160. *Roy. lugdb.* 405. *Hall. helv.* 518. *Dalib. parif.* 59.  
*Lithospermum arvense*, radice rubra. *Bauh. pin.* 258. *Fl. lapp.* 73.  
*Habitat in Europæ agris* ♂ *arvis.* ☉
- virginianum* 3. **LITHOSPERMUM** foliis subovalibus nervosis, corollis acuminatis,  
*Lithospermum corollarum* laciniis acuminatis hirsutis. *Gron. virg.* 140.  
*Lithospermum latifolium virginianum*, flore albido longiore. *Morif. hist.* 3. p. 447. f. 11. t. 28. f. 3. *Raj. suppl.* 272.  
*Habitat in Virginia.*  
*Puncta adpersa foliis sunt excavata, horum latiora seta egrediente notata,*
- purpureo-ceruleum.* 4. **LITHOSPERMUM** seminibus lævibus, corollis calycem multoties superantibus. *Hort. cliff.* 46. *Roy. lugdb.* 405.  
*Lithospermum* flore magno, caule ex apice radices agente. *Hall. helv.* 517.  
*Lithospermum minus repens latifolium.* *Bauh. pin.* 258.  
*Lithospermum repens majus.* *Clus. hist.* 2. p. 163.  
*Habitat in Ungariæ, Angliæ, Galliæ, Italiæ nemoribus* ♂ *secus vias.* 4  
*Caules steriles repentes radicantes: floriferi erecti sunt.*
- fruticosum.* 5. **LITHOSPERMUM** fruticosum, foliis linearibus hispidis, staminibus corollam subæquantibus.  
*Lithospermum fruticosum*, corollis calyce majoribus, foliis linearibus hispidis. *Sauv. monsp.* 50, 63.  
*Anchusa lignosior monspeliensium*, flore violaceo. *Barr. ic.* 1168.  
*Anchusa angustifolia.* *Bauh. pin.* 255.  
*Buglossum fruticosum, rosmarini follo.* *T. Garid. aix.* 68. t. 15.  
 β. *Anchusa arborea.* *Alp. exot.* 67. t. 68.  
*Buglossum samium frutescens, foliis rosmarini obscure virentibus lucidis & hirsutis.* *Tournef. cor.* 6.  
*Habitat in Gallia, Samo & Europa australi.* ♂  
*Stamina longitudine tubi f. limbi expansi.*

6. **LITHOSPERMUM** seminibus duobus, calycibus pa-  
tentibus *Lin. decad. 1. t. 7. \**

*Habitat in Hispania. ☉*

## ANCHUSA.

1. **ANCHUSA** foliis lanceolatis, spicis imbricatis se-  
cundis. *Hort. cliff. 46. Fl. jvec. 152, 161. Mat.*

*med. 59. Roy. lugdb. 406. Dalib. parif. 59.*

*Buglossum angustifolium majus. Bauh. pin. 256.*

*Buglossum sylvestre majus nigrum. Bauh. pin. 256.*

*Segu. ver. 1. p. 214.*

*Habitat ad Europæ rudrata, vias, agros. 2.*

2. **ANCHUSA** racemis subnudis conjugatis. *Roy. lugdb. angustifolia.*  
408.

*Borago sylvestris perennis, flore rufo kermesino. Zan.*  
*hist. 49. t. 20.*

*Buglossum angustifolium minus. Bauh. pin. 256. Mo-*  
*rif. hist. 3. p. 438. f. 11. t. 26. f. 4.*

*Echii facie Buglossum minimum, flore rubente. Lob.*  
*ic. 576.*

*Habitat in Italia, Germania. 2.*

*Simillima præcedenti, sed folia caulina margine subdentata,*  
*spinulis aspera & basi attenuata; nec ovato-lanceolata.*

3. **ANCHUSA** strigosa, foliis linearibus dentatis, pe-  
dicellis bractea minoribus, calycibus fructiferis in-  
flatis. *Laf. Mill. dict. t. 29.*

*Buglossum lusitanicum, echii folio undulato. Tournef.*  
*infl. 134.*

*Anchusa, angustis dentatis foliis, hispanica. Barr. ic.*  
578.

*Habitat in Hispaniæ, Lusitanæ pratis.*

4. **ANCHUSA** villosa-tomentosa, ramis floribusque al-  
ternis axillaribus, bracteis ovatis. *orientalis.*

*Anchusa foliis aveniis, floribus solitariis ex alis ramo-*  
*rum, uti rami e caule. Hort. cliff. 47. Roy. lugdb.*  
406.

*Buglossum orientale, flore luteo. Tournef. cor. 6.*  
*Buxb. cent. 3. p. 17. t. 29. Dill. elih. 60. t. 52. f. 60.*

*Habitat in Oriente. 2.*

5. **ANCHUSA** floribus sparsis, caule glabro.

*virginiana.*

*Anchusa lutea minor, quam alii Paccoon vocant.*  
*Gron. virg. 19.*

**AN-**



*Anchusa minor lutea virginiana*, Puccoon indigenis dicta, qua se pingunt americani. *Pluk. alm.* 30.

*Lithospermum virginianum*, flore luteo duplici ampliori. *Morif. hist.* 3. p. 447. f. 11. t. 28. f. 4. *Kaj. suppl.* 273.

*Habitat in Virginia.* 24

lanata.

6. *ANCHUSA tomentosa*, foliis lanceolatis, staminibus corolla longioribus.

*Pulmonaria mitis*, fragariæ odore. *Bocc. mus.* 2. p. 148. t. 105?

*Habitat Algeriæ.* *Brander.*

*Caulis simplex, pedalis. Folia alterna, lanceolata, villosa, obtusiuscula. Racemi ex alis superioribus, foliis recurvatis, multum villosi. Calyces quinquepartiti, lanati, albido-flavescentes. Corolla cærulea: Nectarium unguibus quinque, brevissimis. Stamina fere corolla longiora. Antheræ incumbentes. Stylus longior.*

tinctoria.

7. *ANCHUSA tomentosa*, foliis lanceolatis obtusis, staminibus corolla brevioribus.

*Lithospermum villosum*; caulibus procumbentibus simplicibus. *Mat. med.* 53. *Sp. pl.* 1. p. 132.

*Anchusa puniceis floribus.* *Bauh. pin.* 255.

*Anchusa monspeliaca.* *Bauh. hist.* 3. p. 584.

*Habitat monspeli.* 24

*Simillima præcedenti* ♂ tota albo villo induta. *Pali staminum longitudine corollæ.*

sempervirens.

8. *ANCHUSA pedunculis diphyllis capitatis.*

*Anchusa scapis diphyllis.* *Hort. cliff.* 47. *Roy. lugdb.* 407.

*Buglossum latifolium sempervirens.* *Bauh. pin.* 256.

*Morif. hist.* 3. p. 437. f. 11. t. 26. f. 2.

*Habitat in Anglia, Hispania.* 24

## CYNOGLOSSUM.

officinale.

1. *CYNOGLOSSUM* staminibus corolla brevioribus, foliis lato-lanceolatis tomentosis sessilibus.

*Cynoglossum foliis lanceolato-ovatis, corollis calycem æquantibus.* *Roy. lugdb.* 406. *Fl. suec.* 154, 162. *Mat. med.* 154.

*Cynoglossum foliis ovato-lanceolatis.* *Hort. cliff.* 47.

Cy-

*Cynoglossum majus vulgare.* Bauh. pin. 257.

β. *Cynoglossum maximum belgicum.* Bauh. pin. 257.

γ. *Cynoglossum sempervirens.* Bauh. pin. 257.

*Habitat in Europæ ruderalis.* ☉

2. **CYNOGLOSSUM** foliis amplexicaulibus ovatis. virginianum.

*Cynoglossum foliis amplexicaulibus.* Gron. virg. 19.

*Morif. hist. 3. f. 11. t. 30. f. 9.*

*Habitat in Virginia.* ☉

3. **CYNOGLOSSUM** corollis calyce duplo longioribus, foliis lanceolatis. cheirifolium.

*Cynoglossum creticum, argenteo angusto folio.* Bauh.

pin. 257. Hort. cliff. 47.

*Cynoglossum creticum, 1. Clus. hist. 2. p. 162.*

*Habitat in Creta, Hispania, Oriente.*

*Corollæ albæ venis sanguineis.*

4. **CYNOGLOSSUM** staminibus corollam æquantibus. apenninum.

*Hort. upf. 33. Sauv. monsp. 64.*

*Cynoglossa montana maxima frigidarum regionum.*

*Col. ecphr. 1. p. 168. t. 170.*

*Habitat in alpinis Apenninis, Campoclarenfibus, locis umbrosis.* ☉

5. **CYNOGLOSSUM** foliis lineari-lanceolatis glabris. linifolium.

*Hort. cliff. 47. Hort. upf. 33. Roy. lugdb. 406.*

*Cynoglossum minus album, lini foliis glaucis, semine umbilicato.* Morif. hist. 3. p. 449. f. 11. t. 30. f. 11.

*Habitat in Lusitania.* ☉

6. **CYNOGLOSSUM** foliis lineari-lanceolatis scabris. lusitanicum.

*Cynoglossum caule erecto ramoso, foliis lanceolatis scabris sessilibus, spicis florum longissimis.* Mill. dict. n. 6.

*Omphaloides lusitanica elatior, cynoglossi folio.* Tournef. inst. 140.

*Habitat in Lusitania.* ☉

*Affinis nimium præcedenti, sed major & asperior.*

7. **CYNOGLOSSUM** repens, foliis radicalibus cordatis. Omphalodes.

*Hort. cliff. 47. Roy. lugdb. 406.*

*Borago minor verna repens, folio lævi.* Morif. hist. 3. p. 437. f. 11. t. 26. f. 3.

*Symphytum minus, boraginis facie.* Bauh. pin. 209.

*Linnaei Spec. plant. pars L.*

N

Ha-

*Habitat in Lusitaniæ, Carniolæ nemorosis ad radices montium.* 24

## PULMONARIA.

\* *Perianthium longitudine tubi corollæ.*

angustifolia. 1. **PULMONARIA** foliis radicalibus lanceolatis. *Hort. cliff.* 44. *Fl. suec.* 2. 164. *Roy. lugd.* 404. *Dalib. parif.* 60. *Hill. plan.* 255. t. 7.

*Pulmonaria* foliis radicalibus ovato-lanceolatis inferius decurrentibus. *Bæhm. lips.* 14.

*Pulmonaria* 5 *pannonica*. *Clus. hist.* 2. p. 170.

*Habitat in Pannonia, Helvetia, Svecia.* 24

*Sequenti nimis affinis, solum foliis angustis distinguenda.*

officinalia. 2. **PULMONARIA** foliis radicalibus ovato-cordatis scabris. *Hort. cliff.* 44. *Fl. suec.* 156, 163. *Mut. med.* 62. *Roy. lugd.* 403. *Hall. helv.* 516. *Dalib. parif.* 60. *Symphytum maculosum* f. *Pulmonaria latifolia*. *Bauh. pin.* 259.

β. *Pulmonaria vulgaris latifolia*, flore albo. *Tournef. inst.* 136.

γ. *Pulmonaria non maculoso folio*. *Clus. hist.* 2. p. 168. *Bauh. pin.* 259.

*Habitat in Europæ nemoribus.* 24

\*\* *Calyces tubo corollæ dimidio breviores.*

virginica. 3. **PULMONARIA** calycibus abbreviatis, foliis lanceolatis obtusiusculis. *Mill. dict.* t. 212.

*Pulmonaria* calyce tubo corollæ brevior, perianthiis quinquepartitis. *Gron. virg.* 20.

*Pulmonaria* foliis ovatis glabris, scapo laxo. *Trew. Ehret.* 11. t. 12.

*Symphytum* f. *Pulmonaria non maculata*, foliis glabris acuminatis, flore patulo cæruleo. *Pluk. alm.* 359. t. 227. f. 6. *Morif. hist.* 3. 444.

*Habitat in Virginia.* 24

sibirica. 4. **PULMONARIA** calycibus abbreviatis, foliis radicalibus cordatis.

*Habitat in Sibiria.* 24

*Radix perennis. Folia glabra, glauca: radicalia cordata; caulina ovata. Corymbi ramosi, nutantes. Media inter antecedentem & sequentem.*

5. PUL-

5. **PULMONARIA** calycibus abbreviatis, foliis ovatis, maritima, caule ramoso procumbente.

*Cerithe foliis ovatis petiolatis. Hort. cliff. 48. Roy. lugdb. 408.*

*Cerithe maritima procumbens. Dill. elh. 75. t. 65. f. 75.*

*Cynoglossum procumbens glaucophyllum maritimum. Pluk. alm. 126. t. 172 f. 3.*

*Cynoglossum perenne maritimum procumbens. Moris. hist. 3. p. 450. f. 11. t. 28. f. 12.*

*Echium marinum. Raj. angl. 3. p. 270. Oed. dan. t. 25.*

*Habitat in Angliæ, Norvegiæ, Islandiæ littoribus silicis. ☉ 24*

*Tres ultime nimis affines : 3. Caule erecto ; Corollis tubo longiore quam limbus. 4. Foliis radicalibus cordatis. 5. Caule ramosissimo ; qui easdem pro varietatibus habeat , vix errat.*

## SYMPHYTUM.

1. **SYMPHYTUM** foliis ovato-lanceolatis decurrentibus. officinale. *Hort. cliff. 47. Fl. svec. 155, 165. Mus. med. 61. Roy. lugdb. 407.*

*Symphytum Consolida major. Bauh. pin. 259.*

*Habitat in Europæ umbrosis subhumidis. 24*

2. **SYMPHYTUM** foliis summis oppositis. tuberosum.

*Symphytum majus, tuberosa radice. Bauh. pin. 259.*

*Symphytum radice tuberosa. Cam. epit. 701.*

*Habitat in Germania australi, Monspelii, Hispania. 24*

*Affine præcedenti, sed folia angustiora : summa magis ovata, nec Folia decurrunt totaliter ad inferiora folia.*

3. **SYMPHYTUM** foliis ovatis subpetiolatis. orientale.

*Symphytum orientale, folio subrotundo aspero, flore cæruleo. Tournef. cor. 7.*

*Symphytum constantinopolitanum, boraginis folio & facie, flore albo. Tournef. cor. 7. Buxb. cen. 5. p. 36. t. 68.*

*Habitat juxta Constantinopolis rivulos, primo vere. 24*

## CERINTHE.

1. **CERINTHE** foliis amplexicaulibus, fructibus geminis, corollis obtusiusculis patulis. major.

N 2

Co

*Cerinthe* foliis cordatis sessilibus. *Hort. cliff.* 48. *Hort. upf.* 35. *Roy. lugd.* 408.

*Cerinthe* foliis caulem amplexantibus, dentibus floris revolutis brevissimis. *Hall. helv.* 515.

*Cerinthe* flore rubro purpurascente. *Bauh. pin.* 258.

β. *Cerinthe*, flavo flore, asperior. *Bauh. pin.* 258.

*Habitat in Europa australi.* ☉

minor.

α. *CERINTHE* foliis amplexicaulibus integris, fructibus geminis, corollis acutis clausis.

*Cerinthe* foliis caulem amplexantibus, floribus profunde quinquefidis. *Hall. helv.* 516.

*Cerinthe* minor. *Bauh. pin.* 258.

*Cerinthe* minor s. quarta. *Clus. hist.* 2. p. 168.

β. *Cerinthe* foliis amplexicaulibus emarginatis, corollis acutis clausis. *Sp. pl.* 1. p. 137.

*Habitat in Austria, Styriae agris.* 24

*Præcedenti maxime affinis, at Corolla ½ quinquefida, præcedentis vero tantum margine quinqueloba.*

## ONOSMA.

simplicissima.

1. *ONOSMA* foliis confertissimis lanceolato-linearibus pilosis.

*Echium creticum.* *Alp. exot.* 130. t. 129. *Morif. hist.* 3. p. 439. f. 11. t. 27. f. 3?

*Habitat in Sibiria.* *Gmelin.* 24

*Caules spithamei, simplices, lignosi, interdum ad basin uno alterove ramo. Folia confertissima ut fere basi imbricata, digiti fere longitudine, angustissime lanceolata & fere linearia, basi attenuata. Racemi sæpius duo, caules terminantes, vix pedunculati. Flores Symphyti absque palis. Symphytum creticum Tournef. cor. esse nequit, cum folia pilis brevissimis adpersa.*

orientalis.

2. *ONOSMA* foliis lanceolatis hispidis, fructibus pendulis.

*Cerinthe orientalis.* *Amæn. acad.* 4. p. 267. \*

*Habitat in Oriente.* β

echioides.

3. *ONOSMA* foliis lanceolatis hispidis, fructibus erectis.

*Cerinthe* foliis lanceolato-linearibus hispidis, femibus quaternis distinctis, fructibus erectis. *Hort. cliff.* 48. *Roy. lugd.* 408.

An-

- Anchusa lutea minor.* *Bauh. pin.* 255.  
*Anchusa echioides lutea*, cerinthe flore, montana. *Col. ceph.* 1. p. 182. t. 183.  
 2. *Anchusa lutea major.* *Bauh. pin.* 255.  
*Anchusa tertia.* *Cam. epit.* 736.  
*Anchusa lutea.* *Dalech. hist.* 1102.  
*Habitat in* Austriæ, Pannoniæ, Helvetiæ, Galliæ, Italiæ *rupibus.* 24

## B O R A G O.

1. BORAGO foliis omnibus alternis, calycibus patentibus. *Hort. upf.* 34. *Mat. med.* 63.  
*Borago calycibus patentibus.* *Hort. cliff.* 44. *Roy. lugdb.* 403. *Dalib. parif.* 61.  
*Buglossum latifolium*, *Borago.* *Bauh. pin.* 256.  
*Habitat hodie in* Normannia ad Colbeck & alibi in Europa; venit olim ex Aleppo. ☉  
 2. BORAGO foliis ramificationum oppositis amplexicaulibus, calycinis foliolis sagittatis.  
*Borago calycinis foliis sagittatis erectis.* *Hort. cliff.* 45.  
*Fl. zeyl.* 71. *Roy. lugdb.* 403.  
*Cynoglossoides folio caulem amplexante.* *Isnard. act.* 1718. p. 325. t. 10.  
*Anchusæ degeneris facie*, indiæ orientalis herba, quadricapsularis. *Pluk. alm.* 30. t. 76. f. 3.  
*Habitat in* India Orientali. ☉  
 3. BORAGO foliis ramificationum oppositis petiolatis, calycinis foliolis ovatis acutis erectis. *Hort. cliff.* 55. *Roy. lugdb.* 403. *africana.*  
*Cynoglossoides africana verrucosa & hispida.* *Isnard. act.* 1718. p. 325. t. 11.  
*Cynoglossum*, boraginis folio & facie, æthiopicum. *Pluk. alm.* 34.  
*Habitat in* Æthiopia. ☉  
*Affinis est admodum antecedenti.*  
 4. BORAGO calycibus tubo corollæ brevioribus, foliis cordatis. *Hort. cliff.* 45. *Roy. lugdb.* 403. *orientalis.*  
*Borago constantinopolitana*, flore reflexo, cæruleo, calyce vesicario. *Tournef. cor.* 6. *itin.* 523. t. 523. *Bauh. cent.* 5. p. 16. t. 30. *Mill. dict.* t. 68.  
*Habitat circum* Constantinopolin. 24

## ASPERUGO.

procumbens. 1. **ASPERUGO** calycibus fructus compressis. *Fl. lapp.* 76 *Fl. rec.* 159. 166. *Hort. cliff.* 44. *Roy. lugdb.* 404. *Hall helv.* 522. *Dalib. parif.* 61.

*Buglossum sylvestre*, caulibus procumbentibus. *Bauh.* p. n. 257. *Morif. hist.* 3. p. 439. f. 11. t. 26. f. 13.

*Habitat in Europæ ruderatis pinguibus* ☉

egyptiaca. 2. **ASPERUGO** calycibus fructus ventricosus.

*Lycopsis ægyptiaca* foliis calloso-verrucosis, caule diffuso, corollis regularibus. *Sp. pl.* 138.

*Habitat in Ægypto.* ☉

*Caules diffusi. Folia petiolata, lato-lanceolata, verrucis callosis hispida, apicibus purpurascentia. Perianthium erectum. Corolla regularis, calyce brevior: limbo albo plano; fauce fornicata lutea. Calyx in fructu ventricosus, parum compressus laciniis inæqualibus, unde Asperuginis genus, præterquam quod Semina compressa, per paria distantia.*

## LYCOPSIS.

vesicaria. 1. **LYCOPSIS** foliis integerrimis, caule prostrato, calycibus fructescentibus inflatis pendulis. *Hort. cliff.* 44. *Hort. upf.* 35. *Roy. lugdb.* 403.

*Echioides flore pullo. Riv. mor.* 11.

*Buglossum procumbens annuum, pullo minimo flore.*

*Morif. hist.* 3. p. 439. f. 11. t. 26. f. 11.

*Habitat in Europa australi.* ☉

pulla. 2. **LYCOPSIS** foliis integerrimis, caule erecto, calycibus fructescentibus inflatis pendulis.

*Echium sylvestre lanuginosum. Bauh. pin.* 254.

*Echium pullo flore. Clus. hist.* 2. p. 164.

*Buglossum sylvestre majus nigrum. Bauh. pin.* 256.

*Habitat in Tartaria, Germania.* 24

*Differt a L. vesicaria caule erecto, radice repente.*

variegata. 3. **LYCOPSIS** foliis repandis dentatis callofis, caule decumbente, corollis cernuis. *Hort. upf.* 34.

*Lycopsis foliis repando-dentatis. Hort. cliff.* 44. *Roy. lug b.* 404.

*Buglossum creticum verrucosum. Stiff. bot.* 57. t. 57.

*Buglossum annuum humile, bullatis foliis, flore cæru-*

*ru-*

ruleo elegant.r variegato. *Morif. hist. 3. p. 439. f. 11.*

t. 26. f. 10.

*Habitat in Creta.* ☉

4. **LYCOPSIS** foliis lanceolatis hispidis, calycibus flo- arvensis.  
rescentibus erectis. *Fl. lapp. 77. Fl. svec. 157, 167.*

*Hort. cliff. 44. Roy. lugdb. 404. Gron. virg. 20.*

*Dalib. parif. 61.*

*Buglossum minus sylvestre. Bauh. pin. 257.*

*Habitat in Europæ agris.* ☉

5. **LYCOPSIS** foliis lanceolatis hirsutis, caule ramo- Echioides.  
sissimo erecto, floribus secundis sessilibus. †

*Echioides orientale, buglossi folio, flore luteo macu-  
lis at. opurpureis notato. Tournef. cor. 47. Buxb.*

*crut. 1. p. 1. t. 1.*

*Habitat in America.* 24

6. **LYCOPSIS** foliis ovatis integerrimis scabris, calyci- orientalis.  
bus erectis.

*Buglossum orientale angustifolium, flore parvo cæru-  
leo Tournef. cor. 6.*

*Habitat in Oriente.* ☉

7. **LYCOPSIS** foliis lineari-lanceolatis confertis tomen- virginica.  
tosis mollibus, caule erecto. *Gron. virg. 140.*

*Habitat in Virginia ad vias.* 24

## ECHIUM.

1. **ECHIUM** caule fruticoso. *Hort. cliff. 43. Roy. fruticosum,  
lugdb. 407.*

*Echium africanum fruticans, foliis pilosis. Comm. hort.*

2. pag. 107. t. 54.

*Habitat in Æthiopia.* 6

2. **ECHIUM** caule lævi, foliis lanceolatis nudis: mar- lævigatum.  
gine carina apiceque scabris.

*Echium africanum perenne, lycopsis facie. Old. afr. 27.*

*Echium africanum minus, foliis oblongis glabris, flo-  
ribus cæruleis. Herm. afr. 2.*

*Buglossum africanum, echii folio, flore purpureo.*

*Fluk. mant. 33. t. 341. f. 7.*

*Habitat ad Cap. b. Spei.* 24

*Caules basi suffruticosi, læves, pedales. Folia lanceo-  
lata, glabra, margine & carina scabra punctis cal-  
losis mucronatis, ut & apice superius. Racemi ex*



*alis foliorum supremorum, secundi, læves. Calyces læves. Corolla parum irregularis. Stamina declinata. Semina muricata.*

italicum.

3. **ECHIUM** caule erecto piloso, spicis hirsutis, corollis subæqualibus, staminibus longissimis. *Echium corollis vix calycem excedentibus margine villosis. Hort. ups. 35. Sp. pl. 1. p. 139. Echium majus & asperius, flore albo. Bauh. pin. 254. Echium flore albo. Cam. epit. 738.*

β. **Lycopsis.** *Bauh. pin. 255.*

*Habitat in Anglia, Italia, Monspeli, in collibus ficcis.*

*Lycopsis β. Corollis minoribus, extus præsertim superne pilosis, & magis regularibus ab E. italico videtur differre, nisi hæc omnia a loco.*

vulgare.

4. **ECHIUM** caule tuberculato-hispido, foliis caulinis lanceolatis hispidis, floribus spicatis lateralibus. *Hort. cliff. 43. Fl. fvec. 158. 168. Roy. lugdb. 407. Dalib. parif. 61. Echium vulgare. Bauh. pin. 254. Clus. hist. 2. p. 143. Habitat in Europa ad vias & agros. ♂*

creticum.

5. **ECHIUM** calycibus fructescentibus distantibus, caule procumbente. *Hort. ups. 35. Echium caule simplici, foliis caulinis linearibus, floribus spicatis ex alis. Hort. cliff. 43. Echium creticum angustifolium rubrum. Bauh. pin. 254. Echium creticum latifolium rubrum. Bauh. pin. 254. Echium creticum. 1. 2. Clus. hist. 2. p. 165. Habitat in Creta, oriente. ○ Stamina non longiora labio brevior corollæ.*

orientale.

6. **ECHIUM** caule ramofo, foliis caulinis ovatis, floribus solitariis lateralibus. *Hort. cliff. 43. Roy. lugdb. 407. Echium orientale, verbasci folio, flore maximo campanulato. Tournef. cor. 6. itin. 3. p. 94. Habitat in Oriente.*

lufitanicum.

7. **ECHIUM** corollis stamine longioribus. *Echium caule simplici, foliis caulinis lanceolatis sericeis, floribus spicatis lateralibus. Roy. lugdb. 407. Echi-*

*Echium*, amplissimo folio, lusitanicum. *Tournef. inst.*

135?

*Habitat in Europa australi.* 2

## TOURNEFORTIA.

1. **TOURNEFORTIA** foliis ovatis serratis, petiolis serratis.  
spinescentibus, spicis terminalibus recurvis.  
*Pittonia arborecens chamædrifolia major. Plum. gen.*  
5. ic. 228. f. 1.
- β. *Pittonia arborecens chamædrifolia minor. Plum. gen.*  
5. ic. 228. f. 2.  
*Habitat in America calidiore.* 5
2. **TOURNEFORTIA** foliis ovatis petiolatis, caule hirsutissima  
hirsuto, spicis ramosissimis terminalibus.  
*Pittonia hirsutissima & ramosissima, baccis albis. Plum.*  
*gen. 5. ic. 209.*  
*Habitat in America calidiore.* 5
3. **TOURNEFORTIA** foliis ovatis acuminatis glaberrimis, petiolis reflexis, caule volubili.  
*Tournefortia foliis ovatis acuminatis. Hort. cliff. 48.*  
*Bryonia nigra fruticosa, racemi ramulis varie implicatis. Sloan. hist. 1. p. 234. t. 143. f. 2.*  
*Virga aurea americana frutescens glabra, foliis subtus cæliis. Pluk. alm. 389. t. 235. f. 6. Raj. suppl. 348.*  
*Habitat in Jamaica, Mexico, arbores scandens.* 5
4. **TOURNEFORTIA** foliis ovato-lanceolatis hirtis, foetidissima.  
pedunculis ramosis, spicis pendulis.  
*Tournefortia foliis ovato-lanceolatis. Hort. cliff. 48.*  
*Roy. lugdb. 408.*  
*Tournefortia scandens, foliis hirtis rugosis ovatis, spicis ramosis. Brown. jam. 169.*  
*Pittonia racemosa, nicotianæ foliis foetidissimis. Plum.*  
*gen 5. ic. 230.*  
*Heliotropium maximum jamaicense, limonii mali folio supra scabro subtus lanugine ferruginea molli. Pluk.*  
*alm. 182. Moris. hist. 3. p. 452.*  
*Heliotropii flore frutex baccifer ramosus, folio rugoso foetido maximo subrotundo hirsuto, fructu albo.*  
*Sloan. jam. 173. hist. 2. p. 108. t. 212. f. 1. Raj.*  
*dend. 75.*  
*Habitat in Mexico, Jamaica.* 5

- cymosa*. 5. **TOURNEFORTIA** foliis ovatis integerrimis nudis, spicis cymosis.  
*Tournefortia frutescens humilis, foliis maximis oblongo-ovatis rugosis, spicis pendulis rarioribus, ramulis crassis sulcatis. Brown. jam. 169.*  
*Pittonia scandens, baccis niveis nigris maculis notatis. Plum. gen. 5?*  
*Heliotropii flore frutex, folio maximo oblongo acuminato glabro. Sloan. jam. 173. hist. 2. p. 109. t. 212. f. 2. Raj. dendr. 76.*  
*Habitat in Jamaica, precedenti nimis affinis. b*
- humilis*. 6. **TOURNEFORTIA** foliis lanceolatis sessilibus, spicis simplicibus recurvis lateralibus.  
*Tournefortia reclinata diffusa & hirsuta, foliis ovatis, ramulis rectis validis. Brown. jam. 169?*  
*Pittonia humilis, anchusæ folio. Plum. gen. 5. ic. 227. f. 2.*  
*Habitat in America calida. b*
- æbirica*. 7. **TOURNEFORTIA** foliis lanceolatis, floribus corymbosis, caule herbaceo.  
*Messerschmidia. Hort. upf. 36.*  
*Arguzia montana. Amm. ruth. 38.*  
*Habitat in Dauriæ apricis glareosis aridis. 24.*  
*Planta dubii etiamnum generis, interim huc relata,*
- suffruticosa*. 8. **TOURNEFORTIA** foliis sublanœolatis incanis, caule suffruticoso.  
*Tournefortia suffruticosa, foliis incanis oblongis, fronde comosa. Brown. jam. 170.*  
*Thymelæa facie frutex maritimus tetraspermos, flore tetrapetalo. Sloan. jam. 138. hist. 2. p. 29. t. 162. f. 4. Raj. dendr. 96.*  
*Habitat in Jamaica. b*

## NOLANA.

- prostrata*. 1. **NOLANA.** *Limn. dec. 1. t. 2. \**  
*Habitat - - - - Dav. Royen. ☉*  
*Plantæ figuram & descriptionem dedit Filius in De- curia plant. 1762.*

## DIAPENSIA.

- lapponica*. 1. **DIAPENSIA** floribus pedunculatis.

Di-

*Diapensia Fl. lapp.* 88. t. 1. f. 1. \* *Fl. svec.* 172, 169.

*Oed. dan.* t. 47.

*Habitat in alpinis Lapponicis.* 2.

2. *DIAPENSIA* floribus subsessilibus.

helvetica!

*Aretia caulibus teretibus, foliis imbricatis, floribus sessilibus. Hall. helv.* 486. t. 8. f. 4.

*Habitat in alpinis Helveticis Gemmi & Pyndnerberg.* 2.

*Habitus antecedentis; fructum non vidi.*

## ARETIA.

1. *ARETIA* villosa, scapis unifloris. *Hall. helv.* 486. t. alpina. 8. f. 1.

*Androsace caulescens, foliis alternis, pedunculis unifloris. Amxn. acad.* 1. p. 164.

*Habitat in Vallesiae monte Loch dicto.* 2.

*Fructus anatome nobis deest in hac planta.*

## ANDROSACE.

5. *ANDROSACE* perianthiis fructuum maximis. *Hort. maxima.* upf. 36.

*Androsace calycibus fructuum maximis. Hort. cliff.*

50. *Roy. lugdb.* 414.

*Androsace foliis lanceolato-ovatis dentatis, calycibus corolla longioribus. Wachend. ultr.* 28.

*Androsace altera. Cam. epit.* 630. *Clus. hist.* 2. p. 134.

*Alpine affinis Androsace dicta major. Bauh. pin.* 251.

*Habitat inter Austriae segetes.* ☉

2. *ANDROSACE* foliis lanceolatis dentatis glabris, septentrio-  
perianthiis angulatis corolla brevioribus. *Fl. svec.* nalis. 160, 170.

*Androsace calyce fructus parvo. Fl. lapp.* 78.

*Androsace montana, flore minore. Buxb. act.* 2. p. 369. t. 23. f. 2.

*Alpine verna, androsaces capitulis. Bauh. pin.* 251. *prodr.* 118. *Oed. dan.* t. 7.

*Alpine minor, androsaces alterius facie. Bauh. pin.* 251. *prodr.* 118. *Burf. XIV.* 75.

*Alpine adfinis Androsace dicta minor. Bauh. pin.* 251.

*Habitat in alpinis Lapponiae, Russiae apricis glareosis.* ☉

3. *ANDROSACE* foliis pilosis, perianthiis hirsutis. villosa.  
Are-

- Aretia villosa*, floribus umbellatis. *Hall. helv.* 486.  
*Sedum alpinum hirsutum*, lacteo flore. *Bauh. pin.* 284.  
*Sedum minus* X, alpinum IV. *Clus. hist.* 2. p. 62.  
*Habitat in alpidibus* Rheticis, Pyrenæis, Carniolicis. 24.  
 lactea. 4. **ANDROSACE** foliis linearibus glabris, umbella involucris multoties longiore.  
*Aretia glabra obtusifolia umbellifera.* *Allion. pedem.* 1. p. 22. t. 4. f. 2.  
*Androsace alpina perennis angustifolia glabra*, flore singulari. *Tournef. inst.* 123.  
*Sedum alpinum*, gramineo folio, lacteo flore. *Bauh. pin.* 284. *Raj. hist.* 1042. *extr.* 291.  
*Sedum minus* XI. *Clus. hist.* 2. p. 61.  
*Habitat in Austriae alpidibus.* 24.  
*Foliorum margo scaber est.*  
 carnea. 5. **ANDROSACE** foliis subulatis glabris, umbella involucrum æquante.  
*Androsace alpina perennis angustifolia glabra & multiflora.* *Tournef. inst.* 123.  
*Sedum alpinum*, angustissimo folio, flore carneo. *Bauh. pin.* 284.  
*Sedum alpinum ejusdem.* *Column. ecphr.* 2. p. 64. t. 65. f. 2.  
*Sanicula alpina*, angustissimo folio. *Pluk. alm.* 332. t. 108. f. 5.  
 6. *Aretia halleri* foliis ciliatis. *Sp. pl.* 1. p. 142.  
*Aretia foliis ciliatis linearibus*, floribus umbellatis. *Hall. helv.* 486. t. 7.  
*Habitat in alpidibus* Pyrenæis, Helveticis. 24.

## P R I M U L A.

- veris. 1. **PRIMULA** foliis dentatis rugosis. *Hort. cliff.* 51. *Fl. suec.* 161, 171. *Roy. lugdb.* 415. *Dalib. parif.* 62.  
*limbo corollarum concavo.* *Mat. med.* 64.  
 officinalis. *Primula veris* odorata, flore luteo simplici. *Bauh. hist.* 3. p. 425.  
*Verbasculum pratense odoratum.* *Bauh. pin.* 241.  
 elatior. 2. *Primula* - - - *limbo corollarum plano.*  
*Verbasculum pratense vel sylvaticum inodorum.* *Bauh. pin.* 241.  
*Primula veris*, pallido flore, elatior. *Clus. hist.* 1. p. 301.

γ. Pri-

- γ. *Primula acaulis*. - - - *scapo nullo.* *acaulis.*  
*Primula veris*, pallido flore, humilis. *Clus. hist.* 1.  
*p.* 302.  
*Verbasculum sylvestre majus*, singulari flore. *Bauh.*  
*pin.* 241.  
*Habitat in Europæ pratis.* 24  
*Varietates licet constantes β. γ. non specie distinguo,*  
*uti nec Maurum ab Europæo.*
- β. PRIMULA foliis crenatis glabris, florum limbo farinosa.  
*plano. Hort. cliff.* 50. *Fl. svec.* 162, 172.  
*Primula foliis crenatis glabris. Vir. cliff.* 12. *Roy.*  
*lugdb.* 415.  
*Primula foliis erectis fastigiatis. Fl. lapp.* 79.  
*Primula veris rubro flore. Clus. dann.* 340.  
*Verbasculum umbellatum alpinum minus. Bauh. pin.*  
*242.*  
*Habitat in Alpibus frigidisque Europæ pratis uliginosis.* 24
- β. PRIMULA foliis ferratis glabris. *Vir. cliff.* 12. *Auricula.*  
*Hort. upf.* 36. *Roy. lugdb.* 415.  
*Primula foliis ferratis carnosis glabris. Hort. cliff.* 50.  
*Sanicula alpina lutea. Bauh. pin.* 242.
- β. *Sanicula alpina purpurea. Bauh. pin.* 242.  
 γ. *Sanicula alpina, foliis rotundis. Bauh. pin.* 242.  
 δ. *Sanicula alpina, flore variegato. Bauh. pin.* 242.  
 ε. *Sanicula alpina angustifolia. Bauh. pin.* 243.  
 ζ. *Sanicula alpina, foliis quasi farina adpersis. Bauh.*  
*pin.* 243.  
*Habitat in alpibus Helveticis, Styriacis.* 24
- 4 PRIMULA foliis integerrimis glabris oblongis, ca- integrifolia.  
 lycibus tubulosis obtusis.  
*Primula foliis glabris carnosis integerrimis. Hall.*  
*helv.* 485.  
*Sanicula alpina rubescens, folio non ferrato. Bauh.*  
*pin.* 243.  
*Auricula urfi IV. Clus. hist.* 1. *p.* 303. t. 304.  
*Habitat in alpibus Helveticis, Styriacis, Pyrenæ-*  
*cis.* 24
5. PRIMULA foliis cuneiformibus dentatis minimis minima.  
 corolla longe minoribus.  
*Sanicula alpina minima carnea. Bauh. pin.* 243.  
*Auricula urfi VIII minima. Clus. hist.* 1. *p.* 305.

*Habitat in Scheeberg. Tauro, Judenberg inque altissimo monte prope salinas Austriæ superioris, qua tenditur in Styriam.*

**Vitaliana.** 6. PRIMULA foliis linearibus integerrimis, floribus sessilibus.

Primula flore subsessili, foliis linearibus. *Anæn. acad.* 1. p. 160.

Auricula urti alpina, gramineo folio, jasmini lutei flore. *Tournef. inst.* 122.

Vitaliana. *Sesl. epist.* 69 t. 10. f. 1. \*

Sanicula alpina, angustissimo folio, flore carneo.

*Pluk. alm.* 332. t. 108. f. 6.

Sedum alpinum, exiguis foliis. *Brüh. pin.* 284.

Sedum alpinum F. gregorii regiensis 1. *Column. ceph.* 2. p. 63. t. 65. f. 1. *Raj. hist.* 1044.

*Habitat in alpinis Pyrenæis & Italicis.* 2.

Calyces longitudine tubi corollæ, ultra medium quinquefidi; corollæ limbus quinquefidus, lobis ovato-oblongis.

Si Semina tantum duo capsulam replentia, videtur Vitaliana genus proprium constituere, quod inquirendum.

**montifolides.** 7. PRIMULA foliis petiolatis cordatis sublobatis crenatis.

*Habitat in Sibiria D. Gmelin.*

*Herba foliis sequentis, floribus antecedentis.*

## CORTUSA.

**matthioli.** 1. CORTUSA calycibus corolla brevioribus.

Cortusa foliis cordatis petiolatis. *Hort. cliff.* 50. *Roy. lugdb.* 414.

Cortusa matthioli. *Clus. hist.* 1. p. 307.

Sanicula montana latifolia laciniata. *Bauh. pin.* 243.

*Habitat in alpinis Austriæ, Sibiriae.*

**Gmelin.** 2. CORTUSA calycibus corollam excedentibus. *Anæn. acad.* 2. p. 340. \*

*Habitat in Sibiria. D. Gmelin.* 2.

## SOLDANELLA.

**alpina.** 1. SOLDANELLA. *Hort. cliff.* 49. *Roy. lugdb.* 414. *Hall. herb.* 488.

Soldanella alpina rotundifolia. *Bauh. pin.* 295. *Morif. hist.* 2. p. 285. f. 3. t. 15. f. 8.

Sol.

*Soldanella alpina. Cam. epit. 254.*

*Habitat in alpinis Helvetiæ, Austriæ, Pyrenæorum. 24*

*Scapus variat uniflorus & multiflorus.*

## DODECATHEON.

### 1. DODECATHEON. *Amen. acad. 3. p. 16. \**

*Meadia.*

*Meadia. Catesb. car. 3. p. 1. t. 1. Trew. Ehret. t. 12.*

*Mill. dict. t. 174.*

*Auricula urfi virginiana floribus boraginis instar rostratis, cyclaminum more reflexis. Pluk. alm. 62. t. 79. f. 6.*

*Habitat in Virginia. 24*

## CYCLAMEN.

### 1. CYCLAMEN corolla retroflexa.

*europæum.*

*Cyclamen foliis cordatis, corolla reflexa. Hort. upf. 47. Mat med. 65.*

*Cyclamen. Hort. cliff. 49. Roy. lugdb. 414.*

*Cyclamina omnia t. 13. Bauh. pin. 307.*

*Cyclaminus. Cam. epit. 35.*

*Habitat in Austriæ, Tartariæ, Europæ australis fœcis umbræ nemorosis 24*

*Varietates sic conjungunt plantas folio anguloso & rotundo, ut vix ac ne vix limites dentur.*

### 2. CYCLAMEN corollæ limbo nutante. *Fl. zeyl. 401. indicum.*

*Habitat in Zeylona. 24*

## MENYANTHES.

### 1. MENYANTHES foliis cordatis integerrimis, corollis ciliatis. *Nymphæides.*

*Menyanthes foliis orbiculatis, corollis margine laceris.*

*Hort. cliff. 53. Roy. lugdb. 430. Dalib. parif. 62.*

*Nymphæa lutea minor, flore imbricato. Bauh. pin. 194.*

*Habitat in Belgii, Angliæ, Dantisci fossis majoribus. 24*

*Umbella simplex, sessilis e latere petioli, ut in sequenti.*

### 2. MENYANTHES foliis cordatis subcrenatis, petio. *indica.*

*lis floriferis, corollis interne pilosis. Fl. zeyl. 72.*

*Nymphæa ceramica. Rumph. amb. 6. p. 173. t. 71. f. 3.*

*Nedel-ambel. Rheed. mal. 11. p. 55. t. 28.*

*Habitat in Malabaræ, Zeylonæ fossis.*



trifoliata.

3. **MENYANTHES** foliis ternatis. *Fl. lapp.* 80. *Fl. svec.* 163, 113. *Hort. cliff.* 52. *Mat. med.* 66. *Roy. lugdb.* 430. *Hall helv.* 487. *Dalib. parif.* 62.  
*Trifolium palustre. Bauh. pin.* 327.  
*Habitat in Europæ paludibus.* 24

## HOTTONIA.

palustris.

1. **HOTTONIA** pedunculo verticillato - multifloro.  
*Boerh. lugdb.* 1. 207. *Hort. cliff.* 51. *Fl. svec.* 164.  
 174. *Roy. lugdb.* 41. *Hall. helv.* 487. *Dalib. pa-*  
*ris.* 93.  
*Millefolium aquaticum* f. *Viola aquatica*, caule nudo.  
*Bauh. pin.* 141.  
*Millefolium aquaticum equisetifolium*, caule nudo.  
*Bauh. pin.* 141.  
*Habitat in fossis & paludibus Europæ borealioris.* 24

indica.

2. **HOTTONIA** pedunculis axillaribus unifloris.  
*Hottonia flore solitario ex foliorum alis. Burm. zeyl.*  
*121. t. 55. f. 1.*  
*Tsiunda-Tsiara. Rheed. mal.* 12. p. 71. t. 36. •  
*Habitat in India. Burmannus.*  
*Caulis digitalis, simplicissimus. Folia verticillata, se-*  
*pe octona, linearia, trifida f. tertrifida. Pedunculi*  
*laterales, axillares foliorum, solitarii, uniflori, fo-*  
*liis longiores. Florem non extricavi, adeoque de*  
*genere non confirmatus.*

## HYDROPHYLLUM.

virginta-  
sum.

1. **HYDROPHYLLUM** foliis pinnatifidis.  
*Hydrophyllum. Hort. cliff.* 44. *Roy. lugdb.* 413. *Gron.*  
*virg.* 21.  
*Dentariæ facie planta monopetalos, fructu rotundo*  
*monopyreno. Morif. hist.* 3. p. 599. f. 51. t. 1. f. 1.  
*Habitat in Virginia.* 24

canadense.

2. **HYDROPHYLLUM** foliis lobato - angulatis.  
*Habitat in Canada.* 24  
*Planta tota Structura antecedentis, ac si genuina filia,*  
*sed diversa folis Foliis, quæ Aceris forma, semi-*  
*quingueloba, glabra lobis acutis, subdentatis, ad pe-*  
*tiolium sinu ut in Aceris foliis.*

LY,

## LYSIMACHIA.

\* *Pedunculis multifloris.*

1. **LYSIMACHIA** paniculata, racemis terminalibus. vulgaris.  
*Lyfimachia* foliis lanceolatis, racemo composito terminali. *Virid. cliff.* 415. *Fl. svec.* 166, 175. *Roy. lugdb.* 416.  
*Lyfimachia* foliis lanceolatis, corymbo terminali. *Fl. lapp.* 81. *Hort. cliff.* 50.  
*Lyfimachia* lutea major. *Bauh. pin.* 245.  
*Habitat in Europa ad ripas* ☿ *paludes.* 2
2. **LYSIMACHIA** racemis simplicibus terminalibus, Ephemere-  
rum.  
petalis obtusis, staminibus corolla brevioribus.  
*Lyfimachia* racemo simplici terminali, foliis lanceolato-linearibus. *Vir. cliff.* 13. *Hort. cliff.* 3. *Roy. lugdb.* 415.  
*Lyfimachia* spicata purpurea minor. *Buxb. cent.* 1. p. 22. t. 33.  
*Lyfimachia* orientalis minor, foliis glaucis. *Till. pis.* 106. t. 40. f. 2.  
*Habitat in Media.* ☉  
*Conf.* *Ephemerum matthioli.* *Bauh. pin.* 244. *Bauh. hist.* 2. p. 905. *Dod. pent.* 203.  
*Convenit cum L. atropurpurea habitu, structura, glauco colore. Differt Foliis subtus fusco-punctatis, quod non sequentis. Petalis obtusis, patulis; staminibus corolla brevioribus; floribus pedicello insidentibus; cum illa caret punctis sub foliis* ☿ *Petala gerit acuminata, calyce paulo longiora, conniventia, saturate rubra; stamina corolla longiora, antheris fuscis; Flores in spicam sessiles; interim maxime affines sunt.*
3. **LYSIMACHIA** spicis terminalibus, petalis lanceo- atropurpu-  
rea.  
latis, staminibus corolla longioribus.  
*Lyfimachia* foliis lanceolato-linearibus caulem & ramos spica ovata terminante. *Hort. cliff.* 51.  
*Lyfimachia* foliis lanceolato-linearibus, spicis terminalibus. *Roy. lugdb.* 415. *Hort. upf.* 37.  
*Lyfimachia* orientalis angustifolia. *Comm. rar.* 33. t. 33.  
*Habitat in Oriente.* ☉  
*Affinitas nimia cum præcedente, tamen distincta.*
4. **LYSIMACHIA** racemis lateralibus pedunculatis. thyrsiflora.

*Lyfimachia racemis simplicibus lateralibus. Virid. cliff.*  
13. *Fl. fvec.* 167, 177. *Roy. lugdb.* 416.

*Lyfimachia ex alis foliorum thyrsifera. Fl. lapp.* 82.  
*Hort. cliff.* 52.

*Lyfimachia bifolia, flore globofo luteo, Bauh. pin.*  
242.

*Lyfimachia lutea. Cluf. hift.* 2. 53. f. 12.

*Habitat in Europa in paludibus.* 24

\* \* *Pedunculis unifloris.*

quadrifolia. 5. **LYSIMACHIA** foliis quaternis subseffilibus, pedunculis quaternis unifloris.

*Lyfimachia foliis ovato-acutis quaternis. Gron. virg.* 20.

*Anagallis marina lutea, foliis latis stellatis. Pet. gaz.*  
5. t. 2. f. 5.

*Anagallis lutea, foliis & floribus ex eodem exortu quatuor ex æqualibus furculorum intervallis cruciatim positis. Pluk. mant.* 12. t. 333. f. 1.

*Habitat in Virginia.*

punctata. 6. **LYSIMACHIA** foliis subquaternis, pedunculis verticillatis unifloris.

*Lyfimachia foliis lanceolatis, floribus solitariis. Hort. cliff.* 52. *Roy. lugdb.* 416. *Gron. virg.* 20.

*Lyfimachia lutea minor, foliis nigris punctis notatis. Bauh. pin.* 245.

*Lyfimachia lutea. 11. Cluf. hift.* 2. p. 52. f. 2.

*Blattariæ affinis planta minor, flore luteo, foliis nigris punctis notatis. Morif. hift.* 2. p. 491. f. 5. t. 10. f. 15.

*Habitat in Hollandia inter arundines.*

*Calyces pubescentes. Petala maculis fulvis punctata.*

ciliata. 7. **LYSIMACHIA** petiolis ciliatis, floribus cernuis.

*Lyfimachia foliis ovato-lanceolatis subcordatis, petiolis margine utrinque lanatis, flore solitario. Wach. ultr.* 390.

*Lyfimachia canadensis. jalappæ foliis. Sarac. canad. Walth. hort.* 32. t. 12.

*Habitat in Virginia, Canada.*

*Caulis erectus. Folia opposita, oblongo-ovata, integerrima, acuta, glabra; nervis lateralibus; petioli canaliculati, ciliati. Pedunculi solitarii, uniflori, nudi.*

8. *Ly-*

8. **LYSIMACHIA** calycibus corollam superantibus, *Linum stel-*  
caule erecto ramosissimo. *Roy. lugdb. 416. Hort. latum.*  
*upf. 37. Sauv. monsp. 130.*  
*Linum minimum stellatum. Bauh. pin. 214. prodr.*  
*107. Magn. monsp. 163. t. 162.*  
*Habitat in Galliæ, Italiæ collibus. ☉*
9. **LYSIMACHIA** foliis ovatis acutis, floribus solita- *nemorum.*  
riis, caule procumbente. *Hort. cliff. 52. Roy. lugdb.*  
*416.*  
*Anagallis lutea nemorum. Bauh. pin. 252. Morif. hist.*  
*2. p. 569. f. 5. t. 26. f. 5.*  
*Anagallis. Clus. hist. 2. p. 182.*  
*Habitat in Germaniæ, Galliæ, Angliæ nemoribus gla-*  
*reosis roridis.*
10. **LYSIMACHIA** foliis subcordatis, floribus solita- *Nummula-*  
riis, caule repente. *Vir. cliff. 13. Hort. cliff. 52. ria.*  
*Fl. svec. 168. Mat. med. 67. Roy. lugdb. 416. Hall.*  
*helv. 481.*  
*Nummularia major lutea. Bauh. pin. 309.*  
*Anagallis mas. Cam. epit. 394.*  
*Habitat in Europa juxta agros ☿ scrobes. 24*
11. **LYSIMACHIA** foliis ovatis acutiusculis, pedun- *tenella.*  
culis folio longioribus, caule repente.  
*Lysimachia foliis subrotundis, floribus solitariis longe*  
*pedunculatis. Guett. stamp. 2. p. 119.*  
*Lysimachia foliis orbiculatis planis, floribus solitariis*  
*rubellis, caule repente. Sauv. monsp. 135.*  
*Nummularia minor, purpurascens flore. Bauh. pin.*  
*310. prodr. 139. Morif. hist. 2. p. 567. f. 5. t. 26. f. 2.*  
*Habitat in Galliæ, Angliæ, Italiæ Ericetis humidis.*

## ANAGALLIS

1. **ANAGALLIS** foliis indivisis, caule procumbente. *arvensis.*  
*Anagallis foliis indivisis. Roy. lugdb. 416. Fl. svec. 169.*  
*178.*  
*Anagallis foliis ovatis. Hort. cliff. 52.*  
*Anagallis phœniceo flore. Bauh. pin. 252.*  
*Habitat in Europæ arvis. ☉*
2. **ANAGALLIS** foliis indivisis, caule erecto. *monelli.*  
*Anagallis foliis lanceolatis. Hort. cliff. 52. Hort. upf.*  
*38.*

*Anagallis cærulea*, foliis binis ternisve ex adverso nascentibus. *Bauh. pin.* 252.

*Anagallis tenuifolia monelli. Chuf. app. alt.*

*Habitat Veronæ.* ☉

latifolia.

3. *ANAGALLIS* foliis cordatis amplexicaulibus, caulibus compressis.

*Anagallis hispanica latifolia*, maximo flore. *Tournef. infl.* 143.

*Cruciata montana minor*, flore cæruleo. *Barr. ic.* 584.

*Habitat in Hispania. Læfl.* ☉

Caulis erecto patens, brachiatus, ramosus, quadrangularis: angulis oppositis triplo latioribus, angustioribus vero margine acuto utrinque septis. Folia subcordata maxime sessilia & amplexicaulia, basi deorsum spectantia, obtusiuscula, nervosa, glabra, subtus pallide punctata. Pedunculi axillares, solitarii, uniflori, nudi, teretes, corolla cærulea, fundo purpurascens. Filamenta purpurea: Antheris oblongis flavis. Notabilis foliis magnis & latis, caule compresso, *A. arvensi* attamen valde assinis.

linifolia.

4. *ANAGALLIS* foliis linearibus, caule erecto.

*Anagallis lusitanica*, linariæ folio angustiore. *Tournef. infl.* 143.

*Habitat in Lusitania, Hispania. Claud. Alstræmer.* ☉

Caulis tetragonus, erectus, palmaris, rigidulus; Ramis alternis, erectis. Folia opposita, sessilia, linearia, acutiuscula, levia, internodiis longiora. Pedunculi filiformes, foliis duplo longiores. Fructus cernui, calyce breviores.

## THEOPHRASTA.

americana.

1. *THEOPHRASTA. Roy. lugdb.* 528.

*Eresia foliis aquifolii longissimis. Plum. gen.* 8. *ic.* 126.

*Habitat in America æquinoctiali.* ‡

Caudex nudus, simplicissimus, apice frondosus, ut in *Palmis*. Folia lanceolata, longissima, dentata nervis undulatis; omnia ex apice caudicis undique diffusa. Fructificationes ramosæ, diffusæ, inter folia ad apicem caulis.

## PATAGONULA.

americana.

1. *PATAGONULA.* †

## PENTANDRIA MONOGYNIA.

213

*Patagonica foliis partim serratis partim integris. Dill.*

*elk. 306. t. 226. f. 293.*

*Habitat in Patagonia Americæ australis. ♀*

*Ignota nobis, videtur Cordiæ affinis.*

## SPIGELIA.

1. SPIGELIA. *Amæn. acad. 5. p. 133. t. 2. \**

*Anthelmia.*

*Spigelia quadrifolia, spicis terminalibus. Brown. jam.*

*156. t. 37. f. 3.*

*Arapabaca quadrifolia, fructu testiculato. Plum. gen.*

*11. Barr. æquin. 15.*

*Brazeel-parilis. Pet. gaz. t. 59. f. 10.*

*Arapabaca brasiliensis dicta planta. Marcgr. bras. 46.*

*Habitat in Cajenna, Brasilia. ☉*

*Caulis simplex, herbaceus. Folia terminalia, quatuor, lanceolata. Folia caulina alia duo opposita. Ramus ex ala singula folii caulini solitarius, cauli smillimus, terminatus foliis quatuor cruciatim positis, ut in caule. Racemi terminales solitarii in caule & ramis.*

## OPHIORRHIZA.

1. OPHIORRHIZA foliis lanceolato-ovatis.

*Mungos.*

*Ophiorrhiza. Fl. zeyh. 402. Mat. med. 79. t. 1. Amæn.*

*acad. 2. p. 118.*

*Habitat in India orientali. 24*

*Conf. Limonium mauritanicum, flore minimo, caule*

*folioso. Pet. gaz. 66. t. 41. f. 12.*

2. OPHIORRHIZA foliis ovatis.

*Mitreola.*

*Mitreola. Hort. cliff. 492. Gron. virg. 28.*

*Mitra. Houst. mss. t. bona.*

*Habitat in America meridionali.*

## RANDIA.

1. RANDIA subinermis.

*mitis.*

*Randia. Hort. cliff. 485.*

*Cacao affinis frutex spinosus, lycii facie, jasmini flo-*

*re albo. Sloan. jam. 135. hist. 2. p. 18. t. 161. f. 1.*

*Raj. dendr. 82.*

*Habitat in America. ♀*

*Hortensis spinis fere destituitur; an sola cultura a sequenti diversa?*

aculeata.

2. *RANDIA* ramis bispinosis.

*Randia* foliis subrotundis confertis, summis ramulis bispiniferis, floribus solitariis. *Brown. jam.* 143. t. 8. f. 1.

*Lycium majus americanum*, jasmini flore, foliis subrotundis lucidis. *Pluk. alm.* 234. t. 97. f. 6.

*Habitat in jamaica.* ♀

## A Z A L E A.

indica.

1. *AZALEA* floribus subsolitariis, calycibus pilosis.

*Chamærhododendron exoticum*, amplissimis floribus liliaceis. *Breyn. prodr.* 1. p. 24.

*Cistus indicus*, ledi alpini foliis, floribus ampliis. *Herm. lugdb.* 152. t. 153. *Raj. hist.* 1895.

*Tsfutiusi. Kämpf. amæn.* 845. t. 846.

*Habitat in India.* ♀

nudiflora.

2. *AZALEA* foliis ovatis, corollis pilosis, staminibus longissimis. *Kalm. it.* 3. p. 110. *Duham. arb.* 1. p. 85. t. 3.

*Azalea* ramis infra flores nudis. *Gron. virg.* 21.

*Azalea* scapo nudo, floribus confertis terminalibus, staminibus declinatis. *Hort. cliff.* 69. *Trew. chret.* 48.

*Azalea erecta*, foliis ovatis integris alternis, flore luteo piloso præcoci. *Cold. eb.* 25.

*Cistus virginiana*, periclymeni flore ampliori minus odorato. *Pluk. mant.* 49.

*Habitat in Virginie siccis.* ♀

*Floret ante folia vel sub ipsa vernatione* ♂ *stamina gerit duplo corollis longiora.*

viscosa.

3. *AZALEA* foliis margine scabris, corollis piloso-glutinosis. *Kalm. it.* 3. p. 110.

*Azalea* ramis infra flores foliosis. *Gron. virg.* 21. *Cold. eb.* 24.

*Cistus virginiana*, flore & odore periclymeni. *Pluk. alm.* 106. t. 161. f. 4. *Catesb. car.* 1. p. 57. t. 57.

*Habitat in Virginia.* ♀

*Affinis nimis præcedenti, sed floret postquam omnia adulta sunt folia, ♂ stamina vix corollis longiora gerit.*

japonica.

4. *AZALEA* foliis adspersis punctis excavatis.

*Azalea* ramis compositis suberectis. *Fl. succ.* 171, 180.

AZA-

*Azalea* maculis ferrugineis subtus adspersa. *Fl. lapp.* 89.

t. 6. f. 1.

*Habitat in Alpibus Lapponiæ.* †

5. *AZALEA* ramis diffuso-procumbentibus. *Fl. lapp.* 90. procumbens

t. 6. f. 2. *Fl. suec.* 170, 179. *Hall. helv.* 416.

*Chamærhododendros* supina ferruginea, thymi folio,  
alpina. *Bocc. mus.* 2. p. 64. t. 53.

*Chamæcistus* serpyllifolia, floribus carneis. *Bauh. pin.*  
466.

*Chamæcistus* VII. *Oel. dan.* t. 9.

*Habitat in Alpibus Europæ.* †

## PLUMBAGO.

1. *PLUMBAGO* foliis amplexicaulibus lanceolatis sca- europæa.

bris. *Hort. cliff.* 53. *Mat. med.* 75. *Roy. lugdb.* 417.

*Sauv. monsp.* 63.

*Lepidium* Dentillaria dictum. *Bauh. pin.* 97.

*Tripolium* dioscoridis. *Col. ecphr.* 1. p. 160. t. 161.

*Habitat in Europa australi.* †

2. *PLUMBAGO* foliis petiolatis ovatis glabris, caule zeylanica.

filiformi. *Hort. cliff.* 53. *Hort. ups.* 43. *Fl. zeyl.* 73.

*Roy. lugdb.* 417.

*Lychnis* indica spicata, ocymastri foliis, fructibus lap-  
pazeis oblongis, radice urente. *Comm. hort.* 2. p.  
169. t. 85.

*Tumba* codiveli. *Rheed. mal.* 10. p. 15. t. 8. *Raj. suppl.*  
550.

*Habitat in India.* †

3. *PLUMBAGO* foliis petiolatis ovatis glabris, caule rosea.

geniculis gibbosis.

*Plumbago* zeylanica, folio splendente ocymastri, flo-  
re rubro. *Burm. zeyl.* 195.

*Radix* vesicatoria. *Rumph. amb.* 5. p. 453. t. 168.

*Schetti* codiveli. *Rheed. mal.* 10. p. 17. t. 9. *Raj. suppl.*  
550.

*Habitat in India.* †

4. *PLUMBAGO* foliis petiolatis ovatis glabris, caule scandens.

flexuoso-scandente.

*Plumbago* americana, viticulis longioribus semper vi-  
rentibus. *Morif. hist.* 3. p. 599.

*Dentellaria* lychnoides sylvatica scandens, flore albo.

*Sloan. jam. hist.* 1. p. 211. t. 133. f. 1.



*Caapomonga. Marcgr. bras. 28. Pif. bras. 200.*

*Habitat in America calidiorē. 5*

*Calyx a D. Jacquino missus distinctam speciem demonstravit.*

## PHLOX.

- paniculata.** 1. PHLOX foliis lanceolatis margine scabris, corymbis paniculatis. *Mill. dict. t. 205. f. 2.*  
*Lychnidea folio salicino. Dill. elth. 205. t. 166. f. 203.*  
*Lysimachia virginiana umbellata maxima, lysimachiae luteae floribus amplioribus. Pluk. mant. 121?*  
*Habitat in America septentrionali. Collinson. 24.*  
*Planta magna, altā. Caulis laevis, obtusus. Folia lato-lanceolata: margine aculeis minutissimis sparsis exasperata. Corymbi conferti in umbellam Floribus numerosissimis, tubo pubescentibus.*
- maculata,** 2. PHLOX foliis lanceolatis laevibus, racemo opposito corymbofo. *Kalm. it. 3. p. 153.*  
*Lychnoides marilandica, foliis binis oppositis basi & auriculis caulem utrinque amplexantibus. Raj. suppl. 489?*  
*Habitat in Virginia. Kalm. 24.*  
*Plantae caulis teretiusculus, subscaber, rubro viridique undique tenuissime maculatus, erectus. Folia laevia, e basi amplexicauli cordata, elongata in lanceolata sensim acutiora. Flores in Racemum digesti, ex pedunculis brevissimis corymbosis.*
- pilosa.** 3. PHLOX foliis lanceolatis villosis, caule erecto, corymbo terminali.  
*Lychnoides marilandica, calycibus lanuginosis, foliis angustis acutis. Raj. suppl. 490.*  
*Lychnidea umbellifera, blattariae accedens virginiana, major repens, pseudomelanthii foliis pilosis. flore pentapetaloidae fistuloso. Pluk. alm. 133. t. 98. f. 1.*  
*Habitat in Virginia. 24.*
- carolina.** 4. PHLOX foliis lanceolatis laevibus, caule scabro, corymbis subfastigiatis.  
*Phlox foliis lanceolatis sessilibus glabris crassis, caule erecto, floribus verticillatis terminalibus. Mill. dict.*  
*Lychnidea caroliniana, floribus quasi umbellatim dispositis, foliis lucidis crassis acutis. Mart. cent. 10. t. 10.*  
*Habitat in Carolina. 24.*

*Similis Ph. glaberrimæ, sed Caulis triplo altior, nonnihil scaber. Folia latiora, ex ovato-lanceolata. Corymbus floribus numerosis, pluribus pedunculis ex alis supremis foliorum, erectis, quasi in corymbum fastigiatis. Corollarum tubus lævis est.*

5. PHLOX foliis lineari-lanceolatis glabris, caule erecto, corymbo terminali. *Horr. cliff. 63. Roy. lugdb. 433. Gron. virg. 21. Kalm. it. 3. p. 153.*

*Lychnidea folio melampyri. Dill. elth. 203. t. 166. f. 202.*

*Habitat in Virginia. 24*

6. PHLOX foliis lato-lanceolatis: superioribus alternis, caule bifido, pedunculis geminis. *Mill. dict. t. 205. f. 1.*

*Lychnidea virginiana, alines aquaticæ foliis, floribus in ramulis divaricatis. Pluk. mant. 121 ?*

*Habitat in Virginia.*

*Caules erectiusculi, simplices, dein divisi in Ramos duos. Folia opposita in caule simplici per 5 oppositiones, ovato-lanceolata, molliuscula, scabra, superiora alterna. Flores ex divaricatione caulis 3 foliorum alternorum alis, bini, propriis pedunculis. Calyx 5-partitus. Corollæ pallide cærulescens tubo curvato.*

7. PHLOX foliis ovatis, floribus solitariis.

*ovata.*

*Lychnidea fistulosa marilandica, clinopodii vulgaris folio, flore amplo singulari. Pluk. mant. 122. t. 348.*

*f. 4.*

*Habitat in Virginia. 24*

8. PHLOX foliis subulatis hirsutis, floribus oppositis.

*subulata.*

*Lychnidea, blattariæ accedens virginiana, minor repens, hirsutis camphoratæ foliis. Pluk. alm. 233. t. 98. f. 2.*

*Habitat in Virginia.*

*Corollæ lobi emarginati 3 tubus calyce duplo longior est.*

9. PHLOX foliis linearibus villosis, pedunculis ternis.

*sibirica.*

*Amæn. acad. 2. p. 342.*

*Habitat in Asia boreali. 24*

*Lobi Corollæ rotundati, integri 3 calyces longitudine tubi corollæ.*

10. PHLOX foliis fetaceis glabris, floribus solitariis.

*setacea.*

*Lychnidea*, *blattariæ accedens virginiana*, *camphoratæ* congeneris, glabris foliis. *Pluk. alm.* 233. t. 98. f. 3.  
*Habitat in Virginia.*

### CONVOLVULUS.

\* *Caule volubili.*

arvensis.

1. **CONVOLVULUS** foliis sagittatis utrinque acutis, pedunculis unifloris. *Fl. fvec.* 173, 181. *Dalib. parif.* 65.

*Convolvulus foliis sagittatis utrinque acutis. Hort. cliff.* 66. *Roy. lugdb.* 427.

*Convolvulus minor arvensis. Bauh. pin.* 294.

- β. *Convolvulus minimus*, angusto auriculato folio. *Bocc. mus. t.* 33.

*Habitat in Europæ agris.* 21

sepium.

2. **CONVOLVULUS** foliis sagittatis postice truncatis, pedunculis tetragonis unifloris. *Roy. lugdb.* 427. *Fl. fvec.* 174, 182. *Dalib. Parif.* 65.

*Convolvulus foliis sagittatis postice truncatis. Hort. cliff.* 66. involucris cordatis. *Virid. cliff.* 18.

*Convolvulus major albus. Bauh. pin.* 294.

*Habitat in Europæ sepibus.* 21

*Foliorum margo fuscus est. Corollæ albæ, at in America septentrionali, incarnatæ.*

Scammonia.

3. **CONVOLVULUS** foliis sagittatis postice truncatis, pedunculis teretibus subtrifloris. *Roy. lugdb.* 427. *Mat. med.* 82.

*Convolvulus syriacus* f. *Scammonia syriaca. Morif. hist.* 2. p. 12. f. 1. t. 3. f. 5. *Mill. dict. t.* 102.

*Scammonia syriaca. Bauh. pin.* 294.

*Habitat in Syria, Mysia, Cappadocia.* 21

*Simillima præcedenti, sed Folia non margine fusca;*

*Pedunculi teretes, biflori f. triflori, nec tetragoni uniflori, folio longiores, nec breviores. Involucrum minimum, patulum, remotum, nec calyce majus erectum approximatum. Calyx lateribus dilatatus, emarginatus, nec teres acuminatus.*

Medium.

4. **CONVOLVULUS** foliis linearibus hastato-acuminatis: auriculis dentatis, pedunculis subunifloris, calycibus sagittatis, caule volubili.

*Tala-neli. Rheed. mal.* 11. p. 113. t. 55.

*Habitat in India.*

Cau-

*Caulis volubilis, non radicans. Folia lanceolata, glabra, acuta, integra basi dilatata in 2 auriculas recurvas, extrorsum subdentatas. Pedunculi uniflori, rarius biflori. Bractææ in medio pedunculi, duæ, oppositæ, lanceolatæ, acuminatæ. Calycis foliola sagittata, ut in Campanula Medio.*

5. CONVULVULUS foliis cordatis integris panduriformibus, calycibus lævibus. panduratus.

*Convolvulus foliis inferioribus cordatis, superioribus trilobis, calycibus pedunculis petiolisque glabris, caule rubescente. Gron. virg. 141.*

*Convolvulus megalorhizus, flore amplo lacteo; fundo purpureo. Dill. elth. 101. t. 85. f. 99.*

*Habitat in Virginiae arenosis.*

6. CONVULVULUS foliis cordatis integris trilobisque villosis, calycibus lævibus, capsulis hirsutis, pedunculis subbifloris. carolinus.

*Convolvulus, folio hederaceo, arvensis, flore dilute purpureo. Dill. elth. 100. t. 84. f. 98.*

*Habitat in Carolina. 24*

*Pedunculi uniflori aut biflori.*

7. CONVULVULUS foliis cordatis integris trilobisque, corollis indivisis, fructibus erectis. hederaceus.

*Convolvulus annuus, foliis cordatis rarius trilobis, calycibus tuberculato-pilosis. Hort. cliff. 67.*

*Convolvulus flore purpureo, calyce punctato. Dill. elth. 99. t. 83. f. 96. infer. 97. t. 81. f. 93.*

*Habitat in India Asiæ, Africæ, Americæ. ☉*

*Pedunculi erecti, etiam fructus, longitudine petiolorum.*

8. CONVULVULUS foliis cordatis trilobis, corollis Nil. semiquinquefidis, pedunculis petiolo brevioribus.

*Convolvulus cærulens, hederaceo anguloso folio. Bauh. pin. 295. Bauh. hist. 2. p. 164. Dill. elth. 96.*

*t. 80. f. 91. 92.*

*Habitat in America. ☉*

*Pedunculi brevissimi, Fructus erecti.*

9. CONVULVULUS foliis cordatis indivisis, fructibus cernuis, pedicellis incrassatis. purpureus.

*Convolvulus calycibus tuberculatis pilosis. Virid. cliff. 18. Hort. upf. 38. Gron. virg. 141.*

Con-

*Convolvulus purpureus*, folio subrotundo. *Bauh. pin.* 295. *Ehret. pict.* 7. f. 2.

β. *Convolvulus cæruleus minor*, folio subrotundo. *Dill. elth.* 97. t. 82. f. 94.

γ. *Convolvulus folio cordato glabro.* *Dill. elth.* 100. t. 84. f. 97.

*Habitat in America.* ○

*Pedunculi erecti, at Pedicelli in fructu incrassati cernui, calyce punctato-scabro & piloso.*

obscurus.

10. **CONVOLVULUS** foliis cordatis indivisis, caule subpubescente, pedunculis incrassatis unifloris, calycibus glabris.

*Convolvulus flore minore lacteo, fundo atro rubente.* *Dill. elth.* 98. t. 83. f. 95.

*Habitat in China, Batavia, Zeylona, Surinamo.* ○

*Caulis teres. Folia exacte cordata, subtus subvillosa, margine scabra petiolis subpubescentibus. Pedunculi uniflori, petiolis longiores, subpubescentes, apice crassiores & glabri: bracteis duabus obsoletis. Corolla C. hederacei, sed minor, colore diversa.*

Batatas.

11. **CONVOLVULUS** foliis cordatis hastatis quinquenerviis, caule repente hispido tuberifero.

*Convolvulus foliis cordatis angulatis, radice tuberosa.* *Hort. cliff.* 67. *Roy. lugdb.* 427.

*Convolvulus radice tuberosa esculenta &c.* *Catesb. car.* p. 60. t. 60.

*Convolvulus indicus orientalis Inhamé f. Batatas.* *Morif. hist.* 2. p. 11. f. 1. t. 3. f. 4.

*Convolvulus indicus vulgo Patates dictus.* *Raj. hist.* 728.

*Batatas.* *Bauh. pin.* 91. *Rumph. amb.* 5. p. 367. t. 130. *Kalm. it.* 2. p. 300.

*Kappa kelengu.* *Rheed. mal.* 7. p. 95. t. 50.

*Habitat in India utraque.* b

*Conf. Convolvulus indicus vulgo Patates dictum.*

*Few. per.* 3. p. 16. t. 11. foliis palmatis.

verticillatus. 12. **CONVOLVULUS** foliis cordatis oblongis nudis, pedunculis umbellatis bifidis multifloris.

*Convolvulus minor scandens, floribus plurimis alaribus, calycibus glabris, foliis oblongo-cordatis.* *Brown. jam.* 153.

*Ipomœa verticillata.* *Amæn. acad.* 5. p. 394.

Qua-

Quamoclit purpurea folanifolia minor. *Plum. ic.* 94. f. 2.

*Habitat in America.*

Corollæ figura est *Convolvuli campanulata*, nec secta ut *Plumieri*. Folia cordata vel ovata, margine subrepando scabriusculo. Flores cæruleſcentes, parvi. Pedunculi folio breviores, umbellati, at pedicelli laterales sæpe ſubdiviſi.

13. CONVOLVULUS foliis cordatis, pedunculis umbellatus, umbellatis, caule volubili.

*Convolvulus americanus*, vulgaris folio, capsulistriquetris numerosis. *Pluk. alm.* 114. t. 167. f. 1.

*Convolvulus luteus polyanthos.* *Plum. amer.* 88. t. 102.

*Convolvulus polyanthos*, folio subrotundo, flore luteo. *Sloan. jam.* 53.

*Habitat in Martinica, Domingo, Jamaica.* 2.

14. CONVOLVULUS foliis cordatis glabris, caule malabari- perenni villoſo. cus.

*Convolvulus foliis cordatis acuminatis*, caule arboreo ſcandente. *Roy. lugdb.* 429.

*Kattu-kelengu.* *Rheed. mal.* 11. p. 105. t. 51. *Raj. ſuppl.* 376.

*Habitat in Malabriæ arenofis.* 6

15. CONVOLVULUS foliis cordatis pubeſcentibus, canarienſis: caule perenni villoſo, pedunculis multifloris.

*Convolvulus foliis cordatis*, caule fruticoſo villoſo.

*Vr. cliff.* 18. *Hort. cliff.* 67. *Roy. lugdb.* 428.

*Convolvulus canarienſis ſempervirens*, foliis mollibus & incanis. *Comm. hort.* 2. p. 101. t. 51.

*Convolvulus canarienſis*, foliis longioribus mollibus incanis. *Pluk. alm.* 114. t. 325. f. 1.

*Habitat in Canarienſibus inſulis.* 6

16. CONVOLVULUS foliis peltatis, pedunculis mul- peltatus, tifiſloris.

*Convolvulus lævis indicus major.* *Rumph. amb.* 6. p. 428. t. 159.

*Habitat in Amboina.*

17. CONVOLVULUS foliis cordatis angulatis, caule Turpethum membranaceo-quadrangulari, pedunculis multifloris.

*Fl. zeyl.* 74. *Mat. med.* 84.

Con-

*Convolvulus zeylanicus alatus maximus*, foliis ibisci nonnihil similibus angulosis. *Herm. lugdb.* 177. t. 178. 179.

*Turpethum repens*, foliis althææ vel indicum. *Bauh. pin.* 149.

*Habitat in Zeylona.* 24

tomentosus. 18. *CONVOLVULUS* foliis trilobis tomentosis, caule lanuginoso.

*Convolvulus folio lanato in tres lacinias diviso*, flore oblongo purpureo. *Sloan. jam.* 55. *hist.* 1. p. 154. t. 98. f. 2. *Raj. suppl.* 381.

*Convolvulus jamaicensis* trilobus, foliis tomentosa lanugine densiori ac veluti instar utrinque obductis. *Pluk. alm.* 115. t. 167. f. 4.

*Habitat in Jamaica.*

althæoides. 19. *CONVOLVULUS* foliis cordatis palmatis sericeis: lobis repandis, pedunculis subbifloris.

*Convolvulus foliis cordato-digitatis sericeis: foliolis linearibus: intermediis duplo longioribus.* *Vir. cliff.* 18. *Roy. lugdb.* 429. *Sauv. monsp.* 202.

*Convolvulus foliis ovatis divisis basi truncatis: laciniiis intermediis duplo longioribus.* *Hort. cliff.* 97.

*Convolvulus argenteus*, folio althææ. *Bauh. pin.* 295.

*Convolvulus*, betonicæ althææque foliis, repens argenteus. *Barr. rar.* 5. t. 312.

β. *Convolvulus argenteus elegantissimus*, foliis tenuiter incis. *Tournef. inst.* 85.

*Habitat in Europæ meridionalis, Orientis, Africæ collibus.* 24

macrocarpos. 20. *CONVOLVULUS* foliis palmato-pedatis quinquepartitis, pedunculis unifloris.

*Convolvulus polyphyllus*, flore & fructu purpureis maximis. *Plum. spec.* 1. ic. 91. f. 1.

*Habitat in America.*

*Pedunculi articulati, incrassati, plicati.*

viridicus. 21. *CONVOLVULUS* foliis palmato-pinnatifidis integerrimis, pedunculis paniculatis.

*Convolvulus foliis laciniatis & quinquefoliis.* *Bauh. pin.* 295.

*Convolvulus quinquefolius.* *Bauh. prodr.* 134.

Con-

*Convolvulus ægyptius. Vegl. ægypt. 73. t. 74.*

*Habitat in Ægypto.*

22. CONVOLVULUS foliis digitatis septenis integer- maerorhi-  
zos.  
rimis, caule glabro, pedunculis trifloris.

*Convolvulus coccineus heptaphyllus, radice crassissima. Plum. spec. 1. ic. 90. f. 1.*

*Habitat in America.*

23. CONVOLVULUS foliis palmatis: lobis septem paniculatus.  
ovatis acutis integerrimis, pedunculis paniculatis.

*Pal-modecca. Rheed. mal. 11, p. 101. t. 49. Raj. suppl.*

374.

*Habitat in Malabariz arenosis.*

24. CONVOLVULUS foliis digitatis quinis pilosis in- pentaphyl-  
lus.  
tegerrimis, caule piloso. *Roy. lugdb. 429.*

*Ipomœa foliis digitatis supra glabris, caule piloso, pedunculis multifloris. Hort. upf. 39. Sp. pl. 162.*

*Convolvulus americanus pentaphyllus & heptaphyllus major. Herm. lugd. 183. t. 185. Raj. hyst. 1881.*

*Convolvulus pentaphyllus, flore pallide flavescente, caule hirsuto pungente. Sloan. jam. 55. hist. 1. p. 153.*

- β. *Convolvulus pentaphyllos, folio glabro dentato, viticulis hirsutis. Plum. amer. 1. ic. 91. f. 2.* serpens.

*Habitat in America. ☉*

*Flos albus. Stylus bicapitatus. Bractea 2. magnæ calycem includunt.*

25. CONVOLVULUS foliis digitatis glabris, foliis quinque-  
dentatis, caule lævi. *Roy. lugdb. 429.* litus.

*Convolvulus quinquefolius glaber americanus. Pluk. alm. 116. t. 167. f. 6.*

*Habitat in America. ☉*

\* \* *Caule prostrato.*

26. CONVOLVULUS foliis cordato-ovatis, peduncu- ficulus.  
lis unifloris: bracteis lanceolatis, flore sessili.

*Convolvulus foliis ovatis acutis, floribus duplici bractea succinctis. Roy. lugdb. 428.*

*Convolvulus foliis ovatis acutis. Hort. cliff. 67.*

*Convolvulus ficulus minor, flore parvo auriculato. Bocc. sic. 89. t. 48. Moris. hist. 2. f. 1. t. 7. f. 5.*

*Habitat in Sicilia. ☉*

*Caulis vix vel admodum parum, sæpius minime volubilis.*



- terrestris. 27. CONVULVULUS foliis linearibus sericeis, pedunculis subtrifloris, calycibus hirsutis mucronatis.  
*Volvulus terrestris. Dalech. hist. 1425. Bauh. hist. 2. p. 160.*  
*Habitat in Europa australi & Africa. 24*
- Dorycnium. 28. CONVULVULUS foliis sublinearibus sericeis, caule paniculato, calycibus obtusis nudiusculis.  
*Habitat in Oriente. 5*  
*Caulis paniculatus. Folia sub lanceolata, sericea, in summis ramis minora rariora. Rami rigidi, rarius dichotomi, terminati floribus subsolitariis.*
- Linearis. 29. CONVULVULUS foliis lanceolatis sericeis lineatis petiolatis, pedunculis bifloris, calycibus sericeis subfoliaceis.  
*Convolvulus minor repens rupellensis, flore rubro. Moris. hist. 2. p. 17. f. 1. t. 4. f. 2. mala.*  
*Convolvulus marinus repens, angusto & oblongo folio, flore purpureo. Barr. rar. 31. t. 1132.*  
*Convolvulus serpens maritimus, spicæ foliis. Triumph. obs. 91. t. 91. f. 2.*  
*Habitat in Hispaniæ, Siciliæ, Mediterranei maritimis. 24*  
*Caules semipedales, prostrati, flexuosi. Folia remota, lanceolata, basi angustiora, sericea, acuta, ad angulum acutum lineata. Pedunculi folio breviores, Bracteis 2, calyce longioribus. Calyces subfoliacei, non pilosi, sed obsolete sericei. Corollæ extus hirtæ.*
- Cneorum. 30. CONVULVULUS foliis lanceolatis tomentosis, floribus capitatis, calycibus hirsutis, caule erectiusculo.  
*Convolvulus argenteus umbellatus erectus. Tournef. inst. 84.*  
*Convolvulus major erectus creticus argenteus. Moris. hist. 2. p. 11. f. 1. t. 3. f. 1.*  
*Convolvulus saxatilis erectus villosus perennis. Barr. rar. 4. t. 470. Bocc. mus. 2. p. 79. t. 70.*  
*Dorycnium. Clus. hist. 2. p. 254.*  
*Cneorum album, folio argenteo molli. Bauh. pin. 463.*  
*Habitat in Hispania, Creta, Syria. 24*  
*Calyces obrusi & valde hirsuti.*

21. CONVULVULUS foliis linearibus acutis, caule cantabrica, ramoſo ſubdichotomo calycibus piloſis. *Hort. cliff.* 68. *Sauv. Monſp.* 56.  
 Convolvulus minimus ſpicæfolius. *Moriſ. hiſt.* 2. p. 17. f. 1. t. 4. f. 3.  
 Convolvulus linariæ folio. *Bauh. pin.* 295.  
 Cantabrica quorundam. *Cluſ. hiſt.* 2. p. 49.  
*Habitat in Italia, Sicilia, Narbona, Verona.*
22. CONVULVULUS foliis cordatis pubeſcentibus, ſpithameus, caule recto, pedunculis uniſloris.  
 Convolvulus ſpithameus, caule recto, foliis ſubrotundis. *Gron. virg.* 141.  
*Habitat in Virginia.*
23. CONVULVULUS foliis cordatis, pedunculis um- corymbosus bellatis, caule repente.  
 Convolvulus niveus polyanthus, folio ſubrotundo majore. *Plum. ſpec.* 1. ic. 89. f. 2.  
*Habitat in America.*
24. CONVULVULUS foliis ovalibus tomentofis, pe- perſicus. dunculis uniſloris. *Amæn. acad.* 2. p. 341. \*  
*Habitat in Perſia, ad maris Caſpici litus.* 21
25. CONVULVULUS foliis lanceolato-ovatis glabris, tricolor. caule declinato, floribus ſolitariis. *Vir. cliff.* 19.  
*Roy. lugdb.* 428.  
 Convolvulus foliis lanceolato-ovatis nudis, caule recto, floribus ſolitariis. *Hort. cliff.* 68. *Hort. wſp.* 38.  
 Convolvulus peregrinus cæruleus, folio oblongo. *Bauh. pin.* 295. flore triplici colore inſignito. *Moriſ. hiſt.* 2. p. 17. f. 1. t. 4. f. 4.  
*Habitat in Africa, Mauritania, Hiſpania, Sicilia.* ☉
26. CONVULVULUS foliis ſagittatis poſtice obtuſis, repens. caule repente, pedunculis uniſloris.  
 Convolvulus lacteſcens, foliis ſagittatis, radice longa alba perenni. *Gron. virg.* 141.  
 Convolvulus marinus catharticus, foliis acetofæ, flore niveo, *Plum. amer.* 89. t. 105.  
*Ballel. Rheed. mal.* 11. p. 107. t. 52.  
*Habitat in America maritimis.* 24
27. CONVULVULUS foliis haſtato-lanceolatis: auri- reptans. culis rotundatis, caule repente, pedunculis ſubbiſloris.  
 Olus vagum. *Rumph. amb.* 3. p. 419. t. 155. f. 1.

*Habitat in India.*

*Planta ex minoribus. Caulis filiformis, nec volubilis, levis. Folia cordata, hastata, sagittata, basi auricula utrinque recurvata, interdum dentata, glabra, acuminata. Pedunculi biflori, breves, pedicellis petiolo longioribus. Calyx rotundatus.*

hirtus.

38. CONVOLVULUS foliis cordatis subhastatisque villosis, caule petiolisque pilosis, pedunculis multifloris.

*Convolvulus foliis cordato-hastatis glabris acuminatis angustis rotundatis, caule petiolisque villosis.*

*Hort. cliff. 496? Roy. lugdb. 429.*

*Habitat in India. Osbeck.*

*Planta non magna, minus volubilis. Pedunculi 3*

*Caulis pilis flavescentibus exstantibus hirti. Folia parum villosa: primaria cordato-subrotunda; secundaria proprie cordata; floribus proxima saepe hastata. Pedunculi nudi, foliis longiores, floribus alternis pedicellatis, involucris ad singulum pedicellum minimis lanceolatis.*

Soldanella.

39. CONVOLVULUS foliis reniformibus, pedunculis unifloris. *Hort. cliff. 67. Mat. med. 81. Roy. lugdb. 428.*

*Soldanella maritima minor. Bauh. pin. 295.*

*Brassica marina. Cord. hist. 205.*

*Habitat in Angliæ, Frisiæ littoribus maris.*

Pes capræ.

40. CONVOLVULUS foliis bilobis, pedunculis unifloris.

*Convolvulus foliis bilobis. Fl. zeyl. 75.*

*Convolvulus foliis subrotundis apice emarginatis basi integris. Roy. lugdb. 428.*

*Convolvulus maritimus zeylanicus, folio crasso cordiformi. Herm. lugdb. 174. t. 175.*

*Convolvulus marinus. Rumph. amb. 5. p. 433. t. 159. f. 1.*

*Schovanna-adamboe. Rheed. mal. 11. p. 117. t. 57.*

*Habitat in India.*

brasiliensis.

41. CONVOLVULUS foliis emarginatis basi biglandulosus, pedunculis trifloris.

*Convolvulus maritimus, foliis nitidis subrotundis emarginatis, petiolis biglandulosus. Brown. jam. 152.*

Con-

*Convolvulus marinus catharticus*, folio rotundo, flore purpureo. *Plum. amer.* 89. t. 104.

*Convolvulus marinus* f. *Soldanella brasiliensis*. *Marcgr. bras.* 51. *Pisf. bras.* 258.

*Habitat in Brasiliæ, Domingo maritimis.*

42. *CONVOLVULUS* foliis lobato - palmatis oblongis, pedunculis unifloris, caule repente.

*Convolvulus albus*, folio laciniato, maritimus. *Plum. spec.* 1. ic. 90. f. 2.

*Habitat in America.*

## IPOMOE A.

\* *Floribus sparsis.*

1. *IPOMOE A* foliis pinnatifidis linearibus, floribus subsolitariis. *Hort. cliff.* 60. *Hort. upf.* 39. *Fl. zeyl.* 77.

*Roy. lugdb.* 430. *Mill. dict.* t. 214.

*Quamoclit* f. *Jasminum americanum*. *Clus. poshk.* 9.

*Convolvulus pennatus exoticus rarior*. *Col. aquat.* 73. t. 72.

*Flos cardinalis*. *Rumph. amb.* 5. p. 421. t. 155. f. 2.

*Jasminum millefolii* folio. *Bauh. pin.* 398.

*Tsiuria - cranti*. *Rheed. mal.* 11. p. 123. t. 60.

*Habitat in India, innotuit 1580. per Cæsarpinum.* ☉

2. *IPOMOE A* foliis digitatis septenis, pedunculis umbellatis brevissimis.

*Quamoclit heptaphyllus*, flore coccineo umbellato.

*Plum. spec.* 3. ic. 92. f. 2.

*Habitat in America.*

3. *IPOMOE A* foliis digitatis: foliolis petiolatis, pedunculis unifloris, carolina.

*Convolvulus minor pentaphyllos*, flore purpureo minore. *Catesb. car.* 2. p. 9. t. 19.

*Habitat in Carolina.*

4. *IPOMOE A* foliis palmatis: lobis septenis lanceolatis acuminatis integerrimis, pedunculis trifloris. *Hort. upf.* 39.

*Convolvulus major heptaphyllus*, flore sulphureo odorato. *Stoan. jam.* 55. *hist.* 1. p. 152. t. 96. f. 2.

*Convolvulus americanus*, mandjuccæ multifido folio, heptaphyllos, flore albo fundo purpureo, radice tuberosa, cortice albo. *Pluk. alm.* 116. t. 276. f. 5.

*Habitat in Jamaica.* b

- digitata.* 5. *IPOMOEA* foliis palmatis: lobis septenis lanceolatis obtusis, pedunculis trifloris.  
*Quamoclit* foliis digitatis, flore coccineo. *Plum. spec.* 3. ic. 92. f. 1.  
*Habitat in America.*
- coccinea.* 6. *IPOMOEA* foliis cordatis acuminatis: basi angulatis, pedunculis multifloris. *Hort. ups.* 39.  
*Ipomœa* foliis cordatis acuminatis vix dentatis. *Hort. cliff.* 66. *Roy. lugdb.* 489.  
*Ipomœa* foliis cordatis integerrimis, limbo floris quinquefido. *Sauv. méth.* 114.  
*Quamoclit americana*, folio hederæ, flore coccineo. *Comm. rar.* 21. t. 21.  
*Convolvulus coccineus*, folio anguloso. *Plum. amer.* 89. t. 103. *Raj suppl.* 380.  
*Habitat in Domingo.* ☉
- lacunosa.* 7. *IPOMOEA* foliis cordatis acuminatis scrobiculatis basi angulatis, pedunculis subunifloris flore brevioribus.  
*Convolvulus stellatus*, periplocæ rotundioris folio. *Dill. elth.* 103. t. 87. f. 102.  
*Habitat in Virginia, Carolina.* ☉  
*Simillima I. coccineæ, sed Pedunculi uniflori f. biflori; Flores tubo crassiore albi, oris purpurascetes, nec coccinei.*
- campanulata.* 8. *IPOMOEA* foliis cordatis, pedunculis multifloris, perianthio exteriore orbiculari, corollis campanulatis lobatis.  
*Adamboe. Rheed. mal.* 11, p. 115. t. 56.  
*Habitat in India.*  
*Corolla crassior, quam in reliquis; sub Perianthio quinquepartito, aliud perianthium minus orbiculatum integerrimum.*
- bana nox.* 9. *IPOMOEA* foliis cordatis acutis integerrimis, caule aculeato, floribus ternis, corollis indivisis.  
*Convolvulus* foliis cordatis, caule aculeato. *Hort. cliff.* 496.  
*Ipomœa sylvestris*, foliis & floribus amplissimis: tubis florum subteretibus. *Brown. jam.* 155.  
*Convolvulus maximus*, caule spinulis obtusis obliquo. *Sloan. jam.* 55. *hist.* 1. p. 151. t. 96. f. 1.

*Convolvulus americanus*, subrotundis foliis, viticulis spinosis. *Pluk. alm.* 115. t. 275. f. 3.

*Smilax aspera* indiæ occidentalis. *Bauh. pin.* 296.

*Munda-valli*. *Rheed. mal.* 11. p. 103. t. 50.

*Habitat in Indiæ arenosis, arborescens.*

*Plantæ non florentes incertos de genere reliquissent Botanicos, nisi Brownius figuram variare docuisset; præsertim cum aliæ caule tantum aculeato & foliis cordatis, aliæ. ex alio loco, foliis panduratis, caule & petiolis aculeatis prodire.*

10. *IPOMOEA* foliis cordatis integerrimis, floribus con- violacea.  
fertis, corollis indivisis. *Sauv. monsp.* 114. †

*Quamoclit* foliis amplissimis cordiformibus. *Plum. spec.*

3. ic. 93. f. 1.

*Habitat in America meridionali.*

11. *IPOMOEA* foliis cordatis acutis integerrimis, flo- solanifolia.  
ribus solitariis. *Roy. lugdb.* 430.

*Quamoclit* solanifolio, flore roseo. *Plum. spec.* 3. ic.

94. f. 1.

*Habitat in America.*

12. *IPOMOEA* foliis sagittatis postice truncatis, pedun- glaucifolia.  
culis bifloris.

*Convolvulus* foliis cordato-lanceolatis. *Hort. cliff.* 67.

*Convolvulus stellatus arvensis*, folio glauco. *Dill.*

*elh.* 103. t. 87. f. 101.

*Habitat in Mexicæ arvis.*

13. *IPOMOEA* foliis trilobis cordatis, pedunculis tri- triloba.  
floris.

*Convolvulus pentaphyllos minor*, flore purpureo?

*Sloan. jam.* 55. hist. 1. p. 153. t. 97. f. 1. *Raj. suppl.*

381.

*Habitat in America.* ☉

*Caulis volubilis, angulatus. Folia cordata, profunde triloba: intermedio ovato; lateratibus semicordatis, integerrima, glabra. Pedunculi triflori. Corollæ cylindricæ, violacæ. Stigmata duo, globosa. Capsulæ pilosæ.*

14. *IPOMOEA* foliis trilobis cordatis, pedunculis multi- hesjerifolia.  
floris racemosis.

*Quamoclit hederæ folio trifido. Plum. spec.* 3. t. 92.

f. 2.

*Habitat in America.*

*Corollæ quam in J. triloba quadruplo longiores & pedunculi plurimis floribus racemosis.*

\* \* *Floribus aggregatis.*

- hepaticifolia. 15. IPOMOEA foliis trilobis, floribus aggregatis. *Fl. zeyl.* 79.  
*Convolvulus indicus villosus, hederæ folio tripartito, flore cæruleo. Herm. lugdb.* 182.  
*Convolvulus zeylanicus hirsutus, foliis hepaticæ. Herm. prodr.* 327.  
*Habitat in Zeylona.*
- tamniifolia. 16. IPOMOEA foliis cordatis acuminatis pilosis, floribus aggregatis.  
*Volubilis carolinienfis, tamni folio subhirsuto. Dell. elth.* 428. t. 318. f. 414.  
*Habitat in Carolina.*
- Pes tigridis. 17. IPOMOEA foliis palmatis, floribus aggregatis. *Fl. zeyl.* 79.  
*Convolvulus zeylanicus villosus pentaphyllos & heptaphyllos minor, Pes tigrinus. Herm. lugdb.* 184. t. 187.  
*Afl. bonon.* 11. p. 2. p. 362. t. 23. f. 2.  
*Volubilis zeylanica, Pes tigrinus dicta. Dill. elth.* 429. t. 318. f. 411.  
*Pulli-schovadi. Rheed. mal.* 11. p. 121. t. 59.  
*Habitat in India.*

## POLEMONIUM.

- cæruleum. 1. POLEMONIUM foliis pinnatis, floribus erectis, calycibus corollæ tubo longioribus.  
*Polemonium. Fl. lapp.* 86. 183. *Fl. suec.* 175. *Hort. cliff.* 163. *Roy. lugdb.* 432. *Hall. helv.* 410.  
*Valeriana cærulea. Bauh. pin.* 164.
- α. Polemonium calycibus lanatis. *Hort. ups.* 40.  
*Habitat in Europæ, Asiæ, septentrionalibus.* 24.
- reptans. 2. POLEMONIUM foliis pinnatis (septenis, floribus nutantibus.  
*Polemonium foliis pinnatis, radicibus reptatricibus. Gron. virg.* 22. *Mill. dict.* t. 209.  
*Habitat in Virginia.* 24.

2. **POLEMONIUM** foliis pinnatifidis linearibus, flo rubrum.  
ribus racemosis pendulis.  
Polemonium calycibus tubo corollæ brevioribus. *Sp.*  
*pl. 1. p. 163.*  
Quamoclit pinnatum erectum, floribus in thyrsum dig-  
gestis. *Dill. elth. 321. t. 241. f. 312.*  
*Habitat in Carolina citerioris arenosis. B. Jussieu.*
4. **POLEMONIUM** foliis inferioribus hastatis, supe-  
rioribus lanceolatis. *Gron. virg. 142. †*  
*Habitat in Virginia.*  
*Facies aliena a Polemoniis, genus mihi dubium.*
3. **POLEMONIUM?** foliis pinnatifidis acutis dentatis, *Nyctelea.*  
caule diffuso.  
*Ipomœa Nyctelea. Sp. pl. 1. p. 160.*  
*Scorpiurus humilis virginianus, foliis rutaceis. Morif.*  
*hist. 3. p. 451. f. 11. t. 28. f. 3.*  
*Habitat in Virginia.*  
Caulis herbaceus, dichotomus, angulatus, diffusus. Fo-  
lia alterna, petiolata, pinnatifida, fere *Hydrophylli*:  
laciniis acutis, utrinque dente angulatis. Pe-  
dunculis oppositifolii, uniflori, pubescentes, elongati.  
Calyx monophyllus, quinquesidus ultra medium, a-  
cutus, corolla longior patens. Corolla monopetala,  
campanulata, quinquesida. Stamina 5. Pistillum 1.  
Fructus sphericus. Facies *Solani peruviani* s. radi-  
cantis, sed corolla campanulata; forte proprii gene-  
ris planta.

## CAMPANULA.

a. Foliis levioribus angustioribus.

1. **CAMPANULA** caule unifloro, calyce corollam uniflora.  
æquante. *Fl. lapp. 85. t. 9. f. 5. 6. Fl. suec. 177, 185.*  
*Habitat in alpinis Lapponicis. 2.*
2. **CAMPANULA** caulibus unifloris, foliis caulinis pulla.  
ovatis crenatis, calycibus cernuis.  
*Campanula alpina latifolia, flore pullo. Bauh. pin. 93.*  
*prodr. 33. Burf. IV. 21.*
3. **Campanula** foliis subrotundis. *Bauh. prodr. 34. t. 35.*  
*Habitat in Austria.*  
*Radix filiformis, repens. Caules spithamæi, erecti,*  
*flexuosi, filiformes, raro ramo uno alterove, eoque*  
*flo-*



*florifero. Folia radicalia & caulina ovata, obtusa, subcrenata crenis distantibus, nuda, petiolata. Pedunculus terminalis. Flos cernuus, magnitudine C. rotundifoliae, calyce levi. β. floribus pluribus in summitate caulis variat.*

- rotundifolia 3. CAMPANULA foliis radicalibus reniformibus, caulinis linearibus. *Fl. lapp. 83. Fl. svec. 176, 184. Hort. cliff. 65. Roy. lugdb. 247. Dalib. parif. 66.*

*Campanula minor rotundifolia vulgaris. Bauh. pin. 93.*

- β. Campanula minor rotundifolia alpina. *Bauh. pin. 93. prodr. 34. t. 34.*

- γ. Campanula alpina linifolia caerulea. *Bauh. pin. 93. Fl. lapp. 84. it. gotl. 229. Magn. monsp. 47. t. 46.*

*Habitat in Europæ pascuis. 2.*

patula.

4. CAMPANULA foliis striatis: radicalibus lanceolato-ovalibus, panicula patula. *Fl. svec. 2. n. 186.*

*Campanula minor rotundifolia, flore in summis caulis. Bauh. pin. 93.*

*Campanula esculentæ facie, ramis & floribus patulis. Dill. elth. 68. t. 58. f. 68.*

- β. Campanula decurrens, foliis caulinis lanceolatis serratis decurrentibus. *Fl. svec. 1. n. 178.*

*Habitat in Angliæ, Sveciæ arvis. ♂*

*Calyx ad basin laciniarum utrinque denticulo livido.*

- Rapunculus 5. CAMPANULA foliis undulatis: radicalibus lanceolato-ovalibus. panicula coarctata. *Hort. ups. 40.*

*Campanula foliis radicalibus lanceolato-ovalibus, caule ramosissimo patulo. Hort. cliff. 65. Dalib. parif. 68.*

*Rapunculus esculentus. Bauh. pin. 92.*

*Rapunculum. Dod. pempt. 165.*

*Habitat in Helvetia, Anglia, Gallia. ♂*

*Folia lanceolata, Racemus terminalis. Rami similes breviores, erecti. Pedunculi terni sæpius, intermedio longiore. Caulis angulatus, scaber.*

- perfoliata 6. CAMPANULA foliis radicalibus obovatis; caulinis lanceolato-linearibus subferratis sessilibus remotis. *Fl. svec. 179. 187. Dalib. parif. 67.*

*Campanula foliis lineari-lanceolatis crenatis, caule longissimo simplicissimo, floribus raris. Vir. cliff. 17. Hort. cliff. 65. Roy. lugdb. 246.*

- Rapunculus persicifolius*, magno flore. *Bauh. pin.* 93.  
 6. *Rapunculus nemorosus angustifolius*, magno flore,  
 major. *Bauh. pin.* 93.  
*Habitat in Europæ septentrionalis asperis.* 24.  
 7. CAMPANULA foliis ovatis glabris subferratis, cau- pyramidalis  
 le erecto paniculato: ramulis brevibus.  
*Campanula foliis ovatis glabris obsolete ferratis, ra-*  
*mis brevissimis cauli approximatis. Vir. cliff.* 17.  
*Roy. lugdb.* 246.  
*Campanula foliis ovatis margine cartilagineo crenatis,*  
*caule ramosissimo angustato. Hort. cliff.* 64.  
*Rapunculus hortensis*, latiore folio, f. *Pyramidalis.*  
*Bauh. pin.* 93.  
*Habitat* - - - - ♂  
 8. CAMPANULA foliis cordatis lanceolatisque, peti- americana.  
 olis ciliatis, floribus secundis, corollis quinque-  
 partitis planis.  
*Campanula caule ramoso, foliis linguiformibus crenu-*  
*latis margine cartilagineo. Roy. lugdb.* 246.  
*Campanula minor americana, foliis rigidis, flore cæ-*  
*ruleo patulo. Herm. lugdb.* 107.  
*Trachelium americanum minus, flore cæruleo patulo.*  
*Dodart. mem.* 4. p. III. t. III.  
*Habitat in Pennsylvania.* ♂  
 9. CAMPANULA foliis lanceolatis: caulinis acute liliifolia.  
 ferratis, floribus paniculatis nutantibus, *Hort. upf.* 41.  
*Campanula urticæ foliis glabra, floribus minoribus*  
*pendulis. Amm. ruth.* 11. n. 17.  
*Habitat in Tataria, Sibiria.* ♂  
 10. CAMPANULA foliis rhomboidibus ferratis, spica rhomboida-  
 secunda. lis.  
*Campanula foliis rhomboidibus ferratis, spica unilate-*  
*rali subnuda. Hall. helv.* 491.  
*Campanula drabæ minoris foliis. Bauh. pin.* 94.  
*prodr.* 36.  
*Rapunculus teucris folio. Barr. rar.* 78. t. 367.  
*Rapunculus alpinus, (teucris folio) rhomboidalis.*  
*Bocc. musf.* 75. t. 61.  
*Habitat in alpinis Helvetiæ, Italiæ.*

β *Foliis scabris latioribus.*

11. CAMPANULA foliis ovato-lanceolatis, caule liliifolia.  
 P 5 flus-

simplicissimo tereti, floribus solitariis pedunculatis, fructibus cernuis. *Vir. cliff.* 17. *Hort. cliff.* 65. *Fl. fvec.* 180, 188. *Roy. lugdb.* 245.

*Campanula maxima*, foliis latissimis. *Bauh. pin.* 94.

*Habitat in Angliæ, Sueciæ montosis, sepibus.* 24.

*Caulis simplicissimus, teres. Folia lanceolato-ovata, serrata. Flores ex alis superioribus, solitarii, pedunculati. Calyces glabri. Fructus cernui.*

rapuncu-  
lodes.

18. *CAMPANULA* foliis cordato-lanceolatis, caule ramoso, floribus secundis sparsis, calycibus reflexis. *Vir. cliff.* 17. *Roy. lugdb.* 246. *Dalib. parisi.* 66.

*Campanula foliis ovatis crenatis, floribus per caulem uno versu sparsis. Hort. cliff.* 64.

*Campanula hortensis, rapunculi radice. Bauh. pin.* 94. *Morif. hist.* 2. p. 460. f. 5. t. 3. f. 32.

*Campanula urticæ foliis oblongis minus asperis. Bauh. pin.* 94. *Hallero.*

*Habitat in Helvetia, Gallia, Austriæ siccissimis.*

bononiensis.

13. *CAMPANULA* foliis ovato-lanceolatis subtus scabris sessilibus, caule paniculato. †

*Campanula* f. *Cervicaria bononiensis*, flore parvo. *Bauh. hist.* 2. p. 804. *Morif. hist.* 2. p. 461. f. 5. t. 4. f. 38.

*Campanula foliis lanceolatis leviter ferratis alternis, floris thyrso spicato & ad basin ramoso. Segu. ver.* 1. p. 176.

*Habitat in Baldi, Lessinensium jugis, Bononiæ.*

graminifolia.

14. *CAMPANULA* foliis lineari-subulatis, capitulo terminali. †

*Campanula alpina, tragopogi folio. Bauh. pin.* 94.

*Trachelium, tragopogi folio, montanum. Col. phyt.* 2. p. 25. *Bauh. hist.* 2. p. 802.

*Trachelium minus gramineum cæruleo-violaceum. Barr. ic.* 332.

*Habitat in Italiæ montibus, Aprutii Salmone vicinis.*

spicata.

15. *CAMPANULA* hispida, spica laxa floribus alternis, foliis linearibus.

*Campanula hirsuto folio angusto leviter serrato, floribus parvis spicatis. Pluk. alm.* 76. t. 153. f. 3.

*Habitat in Vallesia.*

*Radix crassitie digiti. Folia radicalia lineari-lanceolata, caulina similia, sed breviora. Caulis pedalis; ramis al-*

*alternis, terminatis spica longissima ex Floribus sessilibus, alternis, remotis, plurimis. Corollæ subcylindricæ. Planta caule, foliis, calycibus, pilis albis hispida.*

16. CAMPANULA caule angulato, foliis petiolatis, *Trachelium* calycibus ciliatis, pedunculis trifidis. *Vir. clif.* 16. *Fl. svec.* 181, 189. *Roy. lugdb.* 245. *Dalib. parif.* 66. Campanula foliis radicalibus cordatis, calycibus ciliatis. *Hort. clif.* 64.

Campanula vulgatio, foliis urticæ, vel major & asperior. *Bauh. pin.* 94.

*Habitat in Europæ sepibus.* 24.

Folia ovata, subcordata, petiolata, dentata. Pedunculi trifidi, triflori.

17. CAMPANULA caule angulato fissili, floribus *glomerata.* sessilibus, capitulo terminali. *Vir. clif.* 16. *Fl. svec.* 182, 190. *it. gotl.* 196. *it. scan.* 306. *Roy. lugdb.* 245. *Dalib. parif.* 67.

Campanula foliis lanceolato-ovatis crenatis, ramis capitulo florali terminatis. *Hort. clif.* 64.

Campanula pratensis, flore conglomerato. *Bauh. pin.* 94.

*Trachelium alpinum, floribus conglomeratis, foliis asarinæ rigidis & hirsutis. Herm. par.* 235. t. 235.

*Habitat in Angliæ, Galliæ, Austriæ, Sveciæ pratis aridis.* 24.

Folia ovato-oblonga, sessilia, obtusiuscula. Flores 3, sessiles in alis foliorum; capitulum terminale.

18. CAMPANULA hispida, floribus sessilibus, capitulo terminali, foliis lanceolato-linearibus undulatis. *cervicaria.* *it. aland.* 115. *Fl. svec.* 183, 191. \*

Campanula foliis echii. *Bauh. prodr.* 36. (non pinac.).

*Trachelium altissimum, foliis asperis angustis, floribus parvis. Bauh. hist.* 2. p. 801. t. 801.

*Habitat in Helvetiæ, Sveciæ, Germaniæ asperis sylvaticis.*

Corollæ anguli exteriores longitudinaliter ciliati.

19. CAMPANULA hispida, racemo ovato-oblongo *thyrsoides.* terminali, caule simplicissimo, foliis lanceolato-linearibus.

Campanula foliis echii. *Bauh. pin.* 94.

Als-

- Alopecurus alpinus* quibusdam, *Echium montanum* dalechampii. *Bauh. hist.* 2. p. 809.  
*Cervicaria major tenuifolia*. *Thal. harc.* 32. t. 4. \*  
*Habitat in alpinis* Harcyniæ, *Carniolæ*. 24.
- petraea. 20. **CAMPANULA** caule angulato simplici, floribus sessilibus capitato-glomeratis, foliis subtus tomentosis.  
*Campanula* foliis lanceolato-ovatis serratis hispidis, floribus per caulem dense congestis. *Segu. ver.* 1. p. 179.  
*Campanula alpina sphaerocephalos*. *Bauh. pin.* 94.  
*Trachelium majus petræum*. *Pon. bald.* 333. *Bauh. hist.* 2. p. 301.  
*Habitat in Baldo*.  
*Folia ovato-oblonga, obtusiuscula, serrata, supra scabra, subtus tomentosa nivea*.
- Medium. 21. **CAMPANULA** capsulis quinquelocularibus obte-ctis, caule simplici erecto folioso floribus erectis.  
*Vir. cliff.* 16. *Roy. lugdb.* 245.  
*Campanula* calycibus a tergo lamellis quinque notatis. *Hort. cliff.* 64.  
*Campanula hortensis*, folio & flore oblongo. *Bauh. pin.* 94.  
*Viola mariana*. *Dod. pempt.* 163.  
*Habitat in Germaniæ, Italiæ sylvis apricis.* ♂
- barbata. 22. **CAMPANULA** capsulis quinquelocularibus obte-ctis, caule simplicissimo unifolio, foliis lanceola-tis, corollis barbatis.  
*Campanula* foliis echii, floribus villosis. *Bauh. pin.* 94. *prodr.* 36. t. 36.  
*Campanula alpina asperior*, foliis echii, flore magno villoso. *Fluk. Phyt.* 153. f. 6.  
*Rapunculus montanus*. *Bauh. hist.* 2. p. 808.  
*Habitat in Alpibus Austriæ, Helvetiæ, Pedemontii.* Allioni.  
*Caulis simplicissimus, spithameus, in medio Folio unico, radicalibus simili. Folia radicalia lanceolata, scabra, obtusiuscula, integerrima. Flores ad apicem caulis 3 f. 4, sicundi, intus valde barbati. Capsulæ 5-locula-ris; obtecta sinibus reflexis.*
- sibirica. 23. **CAMPANULA** capsulis trilocularibus obte-ctis, caule paniculato.

*Habitat in Sibiria. Gmelin.*

*Plantæ* Caulis pedalis, angulatus parum hispidus, rectus, indivisus. Panicula rarior, laxa. Folia linearia, semiamplexicaulia: Caulina scabra. Flores oblongi, parvi. Calyces hispidi. Varietas ludit caule subdiviso, foliis latioribus.

24. CAMPANULA capsulis quinquelocularibus obdichotoma, tectis, caule dichotomo, floribus cernuis. *Amæn. acad. 4. p. 306. \**

*Campanula* hirsuta, ocymi folio caulem ambiente, flore pendulo. *Bocc. sic. 83. t. 45. f. 1. Morif. hist. 2. p. 459. f. 5. t. 3. f. 26. Raj. hist. 736.*

*Habitat in Syria, Sicilia.*

25. CAMPANULA capsulis quinquelocularibus obmollis, tectis pedunculatis, caule prostrato, foliis suborbiculatis.

*Campanula* rotundifolia hirsuta saxatilis, folio molli. *Bocc. mus. app.*

*Habitat in Oriente, Hispania. Alströmer.*

Caules decumbentes, subramosi, rigidiusculi, villosi. Folia parva, sessilia, suborbiculata, subintegerrima, tomentosa, pubescentia. Flores ex alis foliorum, solitarii, longius pedunculati. Calyces magni, nudi sinibus reflexis. Flores foliis sextuplo majores. Verbo: *Herba Antirrhini mollis, Flores Campanulæ Medii, cæterum statura Camp. heterophyllæ.*

26. CAMPANULA foliis obovatis crenatis, floribus saxatilibus alternis nutantibus, capsulis quinquecarinatis. †

*Campanula* cretica saxatilis, bellidis folio, magno flore. *Tournef. inst. 111. Barr. rar. 79. t. 813.*

*Trachelium* saxatile, bellidis folio, cæruleum creticum. *Bocc. mus. 2. p. 76. t. 64.*

*Habitat in Cretæ scopulis saxosis.*

27. CAMPANULA capsulis obtectis pedunculatis, laciniatis. foliis serratis: radicalibus lyratis; caulinis lanceolatis.

*Campanula* foliis longis laciniatis. *Sp. pl. 1. p. 165.*

*Campanula* græca saxatilis, jacobææ folio. *Tournef. it. 1. p. 99. t. 99. cor. 1.*

*Campanula* mariana, laciniatis foliis peregrina. *Bauh. pin. 94. Morif. hist. 2. p. 460. f. 5. t. 3. f. 31.*

Cam-

*Campanula* foliis radicalibus dentato-pinnatis, caulinis cordatis sessilibus. *Gron. orient.* 19.

Medium dioecorisidis. *Raw. it.* 284.

*Habitat in Græcia, Libano. Burmannus.*

*Planta leviuscula.* Folia radicalia pinnatifida, lyrata, serrata; Caulina lanceolata, subcuneiformia, serrata. Calyces sinibus reflexis, germen obtegentibus.

- frida. 28. *CAMPANULA* capsulis obtectis, foliis hirtis: caulinis lanceolatis ferratis, caule simplicissimo, floribus sessilibus.

*Campanula orientalis*, folio oblongo rigido aspero, flore sursum spectante. *Tournef. cor.* 3.

*Habitat in Syria, Palæstina.*

Caulis sesquipedalis, simplicissimus, hirtus. Folia Radicalia cordata f. ovata, oblongiuscula, serrata, utrinque hirta, petiolata. Caulina alterna, remota, sessilia, lanceolata, obtusa, serrato-dentata, hirta, basi angustiora. Flores axillares, solitarii, sessiles, erecti. Calyces hirti, germen sinibus tegentes.

\* Caule subdiviso.

- canariensis. 29. *CAMPANULA* capsulis quinquelocularibus, foliis hastatis dentatis oppositis petiolatis.

*Campanula* foliis hastatis dentatis, caule determinate folioso. *Hort. cliff.* 65. t. 8. *Roy. lugdb.* 247.

*Campanula canariensis* regia f. Medium radice tuberosa, foliis sinuatis cæsius atriplicis ænulis ternis circum caulem ambientibus, flore amplo pendulo: colore flammeo rutilante. *Pluk. alm.* 76. t. 276. f. 1.

*Habitat in insulis Canariis.* 24

- fruticosa. 30. *CAMPANULA* capsulis quinquelocularibus columnaribus, caule fruticoso, foliis lineari-subulatis, pedunculis longissimis.

*Campanula africana*, ericæ folio, flore cæruleo patulo. *Herm. afr.* 5.

*Habitat ad Caput b. Spel.* 6

- Speculum, ♀ 31. *CAMPANULA* caule ramosissimo diffuso, foliis oblongis subcrenatis, calycibus solitariis corolla longioribus, capsulis prismaticis. *Hort. ups.* 41.

*Campanula* caule ramoso, foliis ovato-oblongis crenatis, *Hort. cliff.* 65.

Cam-

*Campanula calycibus corollam superantibus, capsulis columnaribus. Vir. cliff. 17. Roy. lugdb. 247. Dalib. parif. 68.*

*Onobrychis arvensis* f. *Campanula arvensis erecta. Bauh. pin. 215.*

*Habitat inter segetes Europæ australis. ☉*

32. **CAMPANULA** caule basi subramoso stricto, foliis oblongis crenatis, calycibus aggregatis corolla longioribus, capsulis prismaticis. hybrida.

*Campanula arvensis minima erecta. Morif. hist. 2. p. 457. f. 5. t. 2. f. 22.*

*Speculum veneris minus. Raj. hist. 743.*

*Habitat in Anglia, Gallia, inter segetes.*

*Differt a Speculo ♀ f. matre: Caule basi tantum ramoso, erecto, superne vix ramoso, stricto; Floribus sessilibus, 3. f. 4. simul; Corollis sæpius in rudimento latentibus nec explicatis. Quin hæc ex C. Spec. ♀ olim orta sit, dubium vix videtur.*

33. **CAMPANULA** ramis patentibus indivisis, foliis limonifolia radicalibus ellipticis lævibus integerrimis, floribus sessilibus ternis.

*Campanula orientalis, limonli minimi facie, flore patulo. Tournef. cor. 3.*

*Habitat in Oriente. Burmannus.*

*Folia radicalia ovali-oblonga, integerrima, glabra, petiolata, non tamen rigida. Caulis ramis simplicissimis, virgatis, foliis linearibus f. subulatis. Flores remoti, axillares, sessiles, sæpius terni, bracteis longitudine germinis.*

34. **CAMPANULA** caule subdiviso ramosissimo, foliis linearibus acuminatis, *Pentagonia. Hort. cliff. 66. Roy. lugdb. 247.*

*Campanula pentagonia, flore amplissimo, thracica. Tournef. inst. 112.*

*Speculum veneris, flore amplissimo, thracicum. Raj. hist. 742.*

*Habitat in Thracia. ☉*

35. **CAMPANULA** caule simplici, foliis cordatis dentatis amplexicaulibus, floribus sessilibus aggregatis. *Hort. upf. 40.*

*Campanula caule simplicissimo, foliis amplexicaulibus. Hort. cliff. 65. Roy. lugdb. 247. Grev. virg. 22.*

Cam-



*Campanula pentagonia perfoliata. Moris. hist. 2. p. 457. f. 5. t. 2. f. 23. Barr. rar. 83. t. 1133.*

*Habitat in Virginia. ☉*

*Flores sæpius mutilati, excepto terminali.*

capensis.

35. **CAMPANULA** foliis lanceolatis dentatis hispidis, pedunculis longissimis, capsulis strigosis.

*Campanula africana annua hirsuta, latis serratisque foliis, flore magno violaceo. Comm. hort. 2. p. 69.*

*t. 35.*

*Habitat ad Caput b. Spei. ☉*

elatines.

37. **CAMPANULA** foliis cordatis dentatis pubescentibus petiolatis, caulibus prostratis, pedunculis capillaribus multifloris.

*Habitat in alpinis Europæ australis. Allioni.*

*Specimen: Caulis prostratus, teres, pubescens, varius ramosus. Folia alterna, cordata, dentato-serrata, longius petiolata, utrinque subpubescentia. Pedunculi axillares, capillares s. filiformes, aphylli, laxi, ramosi, sæpius triflori, pubescentes.*

hederacea.

38. **CAMPANULA** foliis cordatis quinquelobis petiolatis glabris, caule laxo. *Dalib. parif. 68.*

*Campanula foliis subrotundis quinquangularibus basi emarginatis glabris, floribus solitariis. Læst. it. 127.*

*Campanula cymbalaræ vel hederæ folio. Baub. pin. 93. prodr. 84. Moris. hist. 2. p. 456. f. 5. t. 4 f. 18.*

*Pluk. phyt. 23. f. 1.*

*Habitat in Anglia, Gallia, Hispania, locis umbrosis humidiusculis.*

heterophylla.

39. **CAMPANULA** foliis subovatis glabris intertremis, caulibus diffusis.

*Campanula saxatilis, foliis inferioribus bellidis, cæteris nummulariæ. Tournef. cor. 2. itin. 1. p. 243. t. 243.*

*Habitat in Oriente. 24*

Krinus.

40. **CAMPANULA** caule dichotomo, foliis sessilibus utrinque dentatis, floralibus oppositis. *Hort. t. ff. 65. Roy. lugdb. 247. Guett. stamp. 429.*

*Campanula caule quadrangulo patulo scabro, calycibus sessilibus axillaribus corollæ tubulosæ æqualibus. Læst. it. 127. \**

*Erini s. Rapunculi minimum genus. Column. phytob. 122. t. 28.*

## PENTANDRIA MONOGYNIA.

241

- Rapunculus minor*, foliis incis. *Bauh. pin.* 92.  
*Alfne oblongo folio serrato, flore cæruleo. Bauh. hist.*  
 3. p. 367.  
*Habitat in Italia, Hispania, G. Narbonensi. ☉*  
*Genus dubium monent Corolla inæqualis, Capsula non*  
*extus dehiscens, Ramificatio opposita, Stigma sim-*  
*plex.*

## ROELLA.

1. *ROELLA foliis ciliatis: mucrone recto.* *ciliata.*  
*Roella. Hort. cliff. 492. t. 35.*  
*Campanula africana frutescens aculeosa, flore viola-*  
*ceo. Comm. hort. 2. p. 77. t. 30.*  
*Campanula africana humilis pilosa, flore exalbido lan-*  
*guide purpureo. Seb. thes. 1. p. 25. t. 16. f. 5.*  
*Aculeosa mauritanica, ericæ foliis hirsutis rigidis infe-*  
*sto mucrone pungentibus. Pluk. alm. 8. t. 252. f. 4.*  
*Habitat in Mauritania, Aethiopia. 24*  
 2. *ROELLA foliis ciliatis: mucrone reflexo.* *reticulata.*  
*Roella foliis imbricatis. Roy. lugdb. 248.*  
*Campanula capitis bonæ spei, foliis reticulatis spinos-*  
*is. Pet. mus. 21. f. 157.*  
*Habitat ad Caput. b. Spei. 24*

## PHYTEUMA.

1. *PHYTEUMA capitulo subfolioso, foliis omnibus pauciflora,*  
*lanceolatis.*  
*Rapunculus foliis obtusis, spica pauciflora. Hall. helv.*  
*497.*  
*Rapunculus alpinus parvus comosus. Bauh. hist. 2. p.*  
*811.*  
*Habitat in alpihus Helveticis, Styriacis.*  
*Capsula bi-f. trilocularis.*  
 2. *PHYTEUMA capitulo subrotundo, foliis linearibus hemisphæri-*  
*integerrimis.* *ca.*  
*Phyteuma foliis linearibus, floribus capitatis. Roy.*  
*lugdb. 248.*  
*Rapunculus umbellatus, folio gramineo. Bauh. pin. 92.*  
*Rapunculus sylvestris cæruleus umbellatus. Thal. herc.*  
*94. t. 3.*  
*Rapuntium alterum angustifolium alpinum. Column.*  
*ecphr. 2. p. 23. t. 26.*  
*Habitat in alpihus Helvetiæ, Italiæ, Pyrenæis. 24*  
*Linnaei Spec. plant. P. 1. Q 2. PHY-*

- orbicularis. 3. PHYTEUMA capitulo subrotundo, foliis serratis : radicalibus cordatis.  
*Phyteuma foliis oblongis, spica orbiculari. Guett. stamp. 1. p. 34. Dalib. parif. 69.*  
*Rapunculus folio oblongo, spica orbiculari. Bauh. pin. 92. Morif. hist. 2. p. 463. f. 5. t. 5. f. 47.*  
*Rapuntium montanum rarius corniculatum. Column. ecphr. 1. p. 222. t. 224.*
- β. *Rapunculus umbellatus latifolius. Bauh. pin. 92.*  
 γ. *Rapunculus umbellatus angustifolius, Bauh. pin. 92.*  
*Rapunculus sylvestris umbellatus. Thal. herc. 1. 8. f. 1. 2.*  
*Habitat in alpibus Italiæ, Helvetiæ, Veronæ, Suffesinæ.*
- spicata. 4. PHYTEUMA spica oblonga, capsulis bilocularibus, foliis radicalibus cordatis.  
*Phyteuma spica oblonga nuda, foliis caulinis lanceolatis serratis. Amæn. acad. 1. p. 149.*  
*Phyteuma foliis cordato-lanceolatis, spica oblonga. Guett. stamp. 1. p. 34. Dalib. parif. 69.*  
*Rapunculus spicatus. Bauh. pin. 92. cæruleus. prodr. 32. t. 32.*  
*Rapunculus foliis cordatis, spica florum oblonga. Monnier. obs. 130.*  
*Rapunculus corniculatus spicatus f. alopecuroides, flore albo & cæruleo. Morif. hist. 2. p. 463. f. 5. t. 5. f. 46.*  
*Habitat in Alpestribus Helvetiæ, Austriæ, Baldi, Angliæ, Galliæ, Italiæ.*  
*Capfula non trilocularis, sed bilocularis. Dill. giff. 65.*
- comosa. 5. PHYTEUMA fasciculo terminali sessili, foliis dentatis : radicalibus cordatis.  
*Rapunculus alpinus corniculatus. Bauh. pin. 113. prodr. 33. t. 33.*  
*Trachelium petræum minus. Pon. bald. 326. t. 326.*  
*Campanula sphærocephalus pervenusta, foliorum ad oras insigniter denticulata. Pluk. alm. 77. t. 152. f. 6.*  
*Habitat in Baldi & Tyrolensibus montibus. ♂*
- pinnata. 6. PHYTEUMA floribus sparsis, foliis pinnatis. Roy. lugdb. 248.  
*Rapunculus creticus f. Pyramidalis altera. Bauh. pin. 93.*

*Rapunculus creticus*, *Petromarula*. *Bauh. hist.* 2. p. 811. t. 712.

*Habitat in Creta.*

## TRACHELIUM.

1. *TRACHELIUM*. *Hort. upf.* 41.

caeruleum;

*Trachelium foliis ovatis ferratis, corymbis compositis.*

*Vir. cliff.* 17. *Roy. lugdb.* 248.

*Trachelium foliis ovatis ferratis, caule umbella terminato.* *Hort. cliff.* 66.

*Cervaria valerianoides caerulea.* *Bauh. pin.* 95.

*Habitat in Italiæ & Orientis umbrosis.* ♂

## SAMOLUS.

1. *SAMOLUS*. *Hort. cliff.* 51. *Hort. upf.* 42. *Fl. fvec.* valerandi. 165, 192. *Roy. lugdb.* 249. *Gron. virg.* 23. *Dalib. parif.* 69.

*Anagallis aquatica, rotundo folio non crenato.* *Bauh. pin.* 252.

*Alsine aquatica, foliis rotundis beccabungæ.* *Morif. hist.* 2. p. 323. f. 3. t. 24. f. 28.

2. *Samolus africanus, folio rotundiore.* *Walsh. hort.* 162. *africanus.* t. 23.

*Habitat in maritimis Europæ, Asiæ & Americæ borealis, ad litora, fontes.* ♂

*Differt africana planta Caule magis ramoso & firmiore; sed vix sufficiunt notæ pro distincta specie.*

## NAUCLEA.

1. *NAUCLEA*.

orientalis.

*Cephalanthus foliis oppositis.* *Fl. zeyl.* 53. *Sp. pl.* 1. p. 95.

*Platanocephalos citri foliis bijugis, capite majore.* *Vaill. art.* 1722. p. 259.

*Arbor indica, fructu aggregato globoso.* *Raj. hist.* 1441.

*Katu-Tsiacca.* *Rheed. mal.* 3. p. 29. t. 33.

*Bancalus.* *Rumph. amb.* 3. p. 84. t. 55?

*Habitat in India Afratica.* ♀

## RONDELETIA.

1. *RONDELETIA foliis sessilibus, panicula dichotoma.* americana.

Q 2

Ron-

*Rondeletia arborescens*, tini facio. *Plum. gen.* 15. ic. 142. f. 1.

*Habitat in America.* §

*Arbustula* Folia opposita, sessilia, lanceolata. Pedunculi communes solitarii, longissimi, nudi, apice formantes corymbum dichotomum; in cujus singula dichotomia Flos sessilis, cum involucro diphylo.

asiatica.

2. RONDELETIA foliis petiolatis oblongis acutis. *Fl. Zeyl.* 80.

Cupi. *Rheed. mal.* 2. p. 37. t. 23. *Raj. hist.* 1494.

*Habitat in Malabar, Zeylona.* §

### MACROCNEUMUM.

jamaicense. 1. MACROCNEUMUM. *Aman. acad.* 5. p. 413. \*  
*Macrocnemum arborescens*, foliis ovatis oppositis, racemis sustentaculis longis insidentibus. *Brown. jam.* 165.

*Habitat in Jamaica.*

### BELLONIA.

aspera.

1. BELLONIA.

*Bellonia frutescens*, folio melissæ aspero. *Plum. gen.* 19. ic. 47.

*Habitat in America.* §

*Arbustula* Folia opposita, ovata, serrata, subius aspera: petiolis brevissimis Flores in corymbum.

### PORTLANDIA.

grandiflora.

1. PORTLANDIA.

*Portlandia floribus pentandris.* *Jacq. amer.* 16.

*Portlandia* foliis majoribus nitidis ovatis oppositis, floribus amplissimis. *Brown. jam.* 164. t. 11. f. 1.

*Habitat in Jamaica.* §

*Corolla Daturæ speciosa.*

### CINCHONA.

officinalis.

1. CINCHONA panicula brachiata.

*Cinchona.* *Mat. med.* 71.

*Quinquina.* *Cond. aët. parif.* 1738. *Godofr. mat.* 2. p. 180.

*Arbor febrifuga peruviana.* *Raj. hist.* 1796.

*Habitat in Loxa Peruvia.* §

2. CIN.

2. CINCHONA pedunculis unifloris. *Jacq. amer. caribæa.*

16. †

*Habitat in Caribæis. ‡*

## PSYCHOTRIA.

1. PSYCHOTRIA stipulis emarginatis, foliis lanceola-asiatica.  
to-ovatis. *Aman. acad. 5. p. 395.*

*Psychotrophum fruticosum, foliis amplioribus ovatis,  
stipulis rigidis interpositis. Brown. jam. 160. 9. 17.*

f. 2.

*Habitat in Jamaica & India Orientali. ‡*

2. PSYCHOTRIA caule repente, foliis cordatis petio-herbacea.  
latis.

*Psychotrophum herbaceum repens sylvaticum, foliis  
subrotundo-cordatis oppositis, floribus paucioribus  
alaribus, laciniis corollæ erecto-patentibus. Brown.  
jam. 161.*

*Jasminum inodorum repens, violæ foliis, fructu pu-  
niceo. Barr. æqu. 63.*

*Violæ folio baccifera repens, flore albo pentapetaloi-  
de, fructu rubro dispermo. Sloan. jam. 115. hist. 1.  
p. 243.*

*Periclymeno accedens planta indiæ, foliis periclymeni  
rotundioribus, fructu bipyreno. Morif. hist. 2. p. 535.*

*Karinta-kali. Rheed. mal. 10. p. 41. t. 21. Raj. suppl.  
358.*

*Habitat in Indiæ utriusque umbrosis roridis.*

## COFFEA.

1. COFFEA floribus quinquefidis dispermis. *Aman. arabica.  
acad. 6. Coffea. \**

*Coffea. Hort. cliff. 59. Hort. upf. 41. Mat. med. 70.  
Roy. lugdb. 239.*

*Jasminum arabicum, lauri folio, cujus semen apud nos  
Coffe dicitur. Juss. ast. 1713. p. 388. t. 7.*

*Jasminum arabicum, castaneæ folio, flore albo odo-  
ratissimo. Till. pis. 87. t. 32.*

*Evonymo similis ægyptica, fructu baccis lauri simili,  
Bauh. pin. 498. Pluk. phyt. 272. f. 1.*

*Bon. Alp. ægypt. 36. t. 36.*

*Habitat in Arabia felici, Æthiopia.*

- occidentalis. 2. *COFFEA* floribus quadrifidis, baccis monospermis.  
*Jacq. amer.* 16.  
 Pavetta foliis oblongo-ovatis oppositis, stipulis setaceis. *Brown. jam.* 142. t. 6. f. 1.  
*Jasminum* arborescens, lauri foliis, flore albo odoratissimo. *Plum. spec.* 17. ic. 156. f. 2.  
*Habitat in America calidiore.* ‡  
*Flores in panicula brachiata.*

## CHIOCOCCA.

- racemosa. 1. *CHIOCOCCA* foliis oppositis. *Jacq. amer.* 16.  
*Chiococca* fermentosa, foliis myrtinis, spicis plurimis tenuissimis & terminalibus & ex alis supremis. *Brown. jam.* 164.  
*Lonicera* racemis lateralibus simplicibus laxis, floribus oppositis pendulis, geniculis compressis. *Sp. pl.* 1. p. 175. n. 12. *Hort. cliff.* 496.  
*Periclymenum* racemosum, flore flavescente, fructu niveo. *Plum. ic.* 211. t. 217. f. 2, *Dill. elth.* 306. t. 228. f. 295.  
*Jasminum* folio myrtino acuminato, flore albicante racemoso. *Sloan. jam.* 196. *hist.* 2. p. 97. t. 188. f. 3.  
*Raj. dendr.* 64.  
*Habitat in Jamaica, Barbados, locis confragosis.* ‡  
 nocturna. 2. *CHIOCOCCA* foliis alternis, *Jacq. amer.* 16. †.  
*Habitat in America.* ‡

## HAMELLIA.

- patens. 1. *HAMELLIA* racemis patentibus. *Jacq. amer.* 16. †  
*Periclymenum* arborescens, ramulis inæxatis, flore luteo. *Plum. ic.* 218 ?  
*Habitat in America meridionali.* ‡

## LonICERA.

\* *Periclymena caule volubili.*

- Caprifolium 1. *LonICERA* floribus verticillatis terminalibus sessilibus, foliis summis connato-perfoliatis. *Hort. cliff.* 45. *Hort. upf.* 42. *Virid. cliff.* 14. *Roy. lugdb.* 237. *Sauv. monsp.* 139.  
*Periclymenum* perfoliatum. *Bauh. pin.* 302.

*Caprifolium Italicum.* *Dod. pempt. 411. Hortul. angl. t. 5. f. 1, 2, 3.*

*Habitat in Europa australi. h*

2. LONICERA spicis nudis verticillatis terminalibus, foliis summis connato-perfoliatis. *sempervi-rens.*  
*Lonicera floribus capitatis terminalibus, foliis summis connatis, inferioribus perfoliatis. Hort. cliff. 58.*  
*Roy., lugdb. 237.*

*Periclymenum perfoliatum virginianum sempervirens & florens. Herm. lugdb. 484. t. 483.*

*Periclymenum virginianum. Riv. mon. 116.*

*Habitat in Virginia, Mexico. h*

3. LONICERA capitulis ovatis imbricatis terminalibus, foliis omnibus distinctis. *Hort. cliff. 58. Hort. upf. 42. \* Fl. fvec. 191, 193. Roy. lugdb. 237. Dalib. paris. 69. Sauv. monsp. 139. it. scan. 305. Mat. med. 68.* *Peric'yme-aurum.*

*Periclymenum non perfoliatum germanicum. Bauh. pin. 302.*

- β. *Caprifolium non perfoliatum. Hort. angl. t. 6. c. 8. f. 1, 2.*

*Caprifolium germanicum, flore rubelló, ferotinum, Tournef. instit. 226.*

*Habitat in Europa media. h*

*Caulis & folia pubescentia in α, at in β glabra.*

\* \* *Chamæcerasa pedunculis bifloris.*

4. LONICERA pedunculis bifloris, baccis distinctis, nigra. *foliis ellipticis integerrimis. Roy. lugdb. 238. Sauv. monsp. 140.*

*Chamæcerasus alpina, fructu nigro gemino. Bauh. pin. 451.*

*Periclymenum alpinum nigrum. Gesn. fasc. 37. t. 8. f. 48.*

*Periclymenum rectum, folio serrato. Bauh. hist. 2. p. 107.*

*Periclymenum rectum 2. Clus. hist. 1. p. 58.*

*Habitat in Delphinatu, Gallia, Helvetia.*

*Involucrum germinibus brevius. Folia integerrima, sæpe leguntur serrata, sed hæc a Papilione Camilla sunt erosa. Schmiedel.*

5. LONICERA pedunculis bifloris, baccis distinctis, tatarica. *foliis cordatis obtusis. Hort. upf. 42.*



*Chamæcerasus fructu gemino rubro, foliis glabris cordatis. Amm. ruth.* 184.

*Habitat in Tataria.* §

*Folia cordata glabra, brevius petiolata. Pedunculi axillares, biflori. involucrum lanceolatum; involuella obovata. Corolla pallide incarnata: labio superiore tripartito: lacinia media bifida; inferiore simplici.*

*Xylosteum.* 6. *LONICERA* pedunculis bifloris, baccis distinctis, foliis integerrimis pubescentibus. *Roy. lugdb.* 238. *Fl. svéc.* 192, 194. *Dalib. parif.* 69. *Sauv. monsp.* 140.

*Lonicera pedunculis bifloris, foliis ovatis obtusis integris. Hort. cliff.* 58.

*Chamæcerasus dumetorum, fructu gemino rubro. Bauh. pin.* 451.

*Xylosteum. Dod. pempt.* 412.

*Habitat in Europæ frigidioris sepibus.* §

*pyrenaica.* 7. *LONICERA* pedunculis bifloris, baccis distinctis, foliis oblongis glabris.

*Lonicera pedunculis bifloris, baccis distinctis, floribus infundibuliformibus, ramis divaricatis. Roy. lugdb.* 238.

*Xylosteum pyrenaicum. Tournes. inst.* 609. *Magn. hort.* 209. t. 209. *Raj. dendr.* 29. *Duham. arb.* 2. p. 374. t. 110.

*Habitat in Pyrenæis.* §

*Differt a præcedente Corollis regularibus, Ramis divaricatis, Foliis glabris.*

*alpigena.* 8. *LONICERA* pedunculis bifloris, baccis coadunatis didymis. *Mill. dist.* § 167. f. 2.

*Lonicera pedunculis bifloris, foliis ovatis acutis integris. Hort. cliff.* 58. *Sauv. monsp.* 140.

*Lonicera baccis bifloris concretis, floribus bilabiatis. Roy. lugdb.* 238.

*Chamæcerasus. Gesn. fasc.* 33. t. 14. f. 44.

*Chamæcerasus alpina, fructu rubro gemino: duobus punctis notato. Bauh. pin.* 451.

*Periclymenum rectum quartum. Clus. hist.* 1. p. 59.

*Habitat in Alpibus Helveticis, Pyrenæicis, Allobrogicis.* §

9. **LONICERA** pedunculis bifloris, baccis coadunato-  
globois, styliis indivisis. *carulea.*  
Lonicera pedunculis bifloris bilabiatis, bacca singulari  
globosa integerrima. *Roy. lugdb. 239.*  
Chamæcerasus montana, fructu singulari cæruleo.  
*Bauh. pin. 451.*  
Periclymenum rectum, fructu cæruleo. *Clus. pon. 85.*  
*Bauh. hist. 2. p. 108. Raj. hist. 1492.*  
*Habitat in Helvetia. ♀*  
Rami teretes, rigidi, rubri. Folia ovata, integra.  
Gemmæ. 2. f. 3, altera supra alteram, tamquam due  
spicæ, subulatæ, compressæ, horizontales. Pedunculi  
ex infimis oppositionibus foliorum, cernui, brevissi-  
mi. Germen unicum, absque visibili perianthio,  
cum Corollis duabus, 5-fidis, regularibus, albis.  
\* \* \* *Caule erecto pedunculis multifloris.*
10. **LONICERA** spicis terminalibus, foliis ovato-ob-  
longis acuminatis distinctis sessilibus. *marilandica.*  
*Gron. virg. 142.*  
Periclymeni virginiani flore coccineo planta marilan-  
dica, spica erecta, foliis conjugatis. *Raj. dendr. 32.*  
*Catesb. carol. 2. p. 78. t. 78.*  
*Habitat in Virginia, Marilandia, Carolina. ♀. 24*  
*Generis dubij planta a nobis non viso fructu.*
11. **LONICERA** capitulis lateralibus pedunculatis, fo-  
liis petiolatis. *Symphoricarpos.*  
Lonicera pedunculis axillaribus capitatis, foliis petio-  
latis. *Hort. cliff. 58. Roy. lugdb. 237. Gron. virg. 23.*  
Symphoricarpos foliis alatis. *Dill. elh. 371. t. 278.*  
*f. 360.*  
Vitis idæa caroliniana, foliis subrotundis hirsutis. *Hor-  
tul. angl. 85. t. 20.*  
*Habitat in Virginia, Carolina. ♀*
12. **LONICERA** racemis terminalibus, foliis serratis. *Diervilla.*  
*Diervilla. Hort. cliff. 63. t. 7.*  
Diervilla acadensis fruticosa, flore luteo. *Tournef.  
art. 1706. t. 7. f. 1. Duham. arb. 1. p. 209. t. 87.*  
*Habitat in Acadia, Noveboraco. ♀*
13. **LONICERA** corymbis terminalibus, foliis ovatis corymbosa.  
acutis.  
Peryclimenum foliis acutis, floribus profunde dissectis.  
*Fewill. peruv. 1. p. 760. t. 45.*  
*Habitat in Peru. ♀*

## TRIOSTEUM.

- perfoliatum. 1. **TRIOSTEUM** floribus verticillatis sessilibus.  
*Triosteospermum*, latiore folio, flore rutilo. *Dill.*  
*elth.* 394. t. 293. f. 378.  
*Habitat in America septentrionali.* b
- angustifolium. 2. **TRIOSTEUM** floribus oppositis pedunculatis.  
*Lonicera humilis* hirsuta, caule obsolete rubente quadrato, foliis lanceolatis adversis, flore luteo ad alas unico. *Gron. virg.* 143.  
*Periclymenum herbaceum rectum virginianum.* *Pluk. alm.* 287. t. 104. f. 2.  
*Habitat in Virginia.* 2

## MORINDA.

- umbellata. 1. **MORINDA** erecta, foliis lanceolato-ovatis, pedunculis confertis. *Fl. zeyl.* 81. \*  
*Bancudus angustifolia.* *Rumph. amb.* 3. p. 157. t. 98.  
*Habitat in India.* b
- clitrofolia. 2. **MORINDA** arborea, pedunculis solitariis. *Fl. zeyl.* 82.  
*Bancudus latifolia.* *Rumph. amb.* 2. p. 158. t. 99.  
*Coda-pilava.* *Rheed. mal.* 1. p. 97. t. 52. *Raj. hist.* 1442.  
*Habitat in India.* b
- Royoc. 3. **MORINDA** procumbens. *Hort. cliff.* 73. *Roy. lugdb.* 187.  
*Morinda americana humifusa laurifolia.* *Vaill. act.* 1722. p. 275.  
*Royoc humifusum*, fructu cupressino. *Plum. spec.* 11.  
*Periclymenum americanum*, e cujus radice fit atramentum. *Pluk. alm.* 287. t. 212. f. 4.  
*Habitat in America calidiore.* b

## CONOCARPUS.

- erecta. 1. **CONOCARPUS** erecta, foliis lanceolatis.  
*Conocarpus.* *Hort. cliff.* 485.  
*Conocarpus* foliis oblongis, petiolis brevibus, floribus in caput conicum. *Brown. jam.* 159.  
*Alnus maritima myrtifolia coriariorum.* *Pluk. alm.* 18. t. 240. f. 3.

Alut

*Alni fructu laurifolia arbor maritima. Sloan. jam. 135.*

*hist. 2. p. 18. t. 161. f. 2. Raj. dendr. 11.*

*Rudbeckia laurifolia maritima. Amm. herb. 581.*

*Innominata. Plum. ic. 135. t. 144. f. 2.*

*Habitat in Jamaica, Bermudensium, Brasiliæ maritimis. ‡*

2. **CONOCARPUS** foliis lanceolato-ovatis obtusiusculis, fructibus sejunctis. *racemosa*

*Conocarpus foliis elliptico-ovatis, petiolis biglandulosis, racemis laxis, fructibus sejunctis. Brown. jam. 159.*

*Mangle julifera, foliis ellipticis ex adverso nascentibus. Sloan. jam. 156. hist. 2. p. 66. t. 187. f. 1. Raj. dendr. 115.*

*Habitat in Jamaica maritimis. ‡*

3. **CONOCARPUS** procumbens, foliis subrotundis. *procumbens.*

*Rudbeckia supina, foliis subrotundis. Amm. herb. 581.*

*Habitat in Cuba.*

## ERITHALIS.

1. **ERITHALIS.**

*Erithalis fruticulosa, foliis obovatis crassius nitidis oppositis, pedunculis ramosis ad alas superiores. Brown. jam. 165. t. 17. f. 3.*

*Habitat in Jamaica. ‡*

*fruticulosa.*

## MENAIIS.

1. **MENAIIS.** *Last. it. 306.*

*Habitat in America meridionali. ‡*

*Caules teretes, subvillosi. Folia alterna, ovata, integra, aspera.*

*teretia.*

## GENIPA.

1. **GENIPA.**

*Genipa fructu ovato. Plum. spec. 20. ic. 136.*

*Janipaba. Marcgr. bras. 92. Pis. bras. 158.*

*Habitat in America meridionali. Montier. ‡*

*americana.*

## MUSSÆNDA.

1. **MUSSÆNDA.** *Herm. zeyl. 36. Fl. zeyl. 84.*

*Mussænda zeylanica, flore rubro, folio ex florum thyrsis prodeunte albo. Burn. zeyl. 168. t. 76.*

*Fo-*

*frondosa.*

Folium principissæ. *Rumph. amb. 4. p. 111. t. 51.*

Belilla. *Rheed. mal. 2. p. 27. t. 18. Raj. hist. 1493. dendr. 126.*

*Habitat in India. 5*

### MIRABILIS.

Jalapa.

1. **MIRABILIS** floribus congestis terminalibus erectis. *Mirabilis. Hort. cliff. 53. Hort. upf. 43. Fl. zeyl. 85. Mat. med. 76. Rumph. amb. 5. p. 253. t. 89. Osb. it. 225. Nyctage. Roy. lugdb. 417. Cold. noveb. 29. Solanum mexicanum, flore magnâ. Bauh. pin. 168. Admirabilis peruviana. Clus. hist. 2. p. 87.*

*Habitat in India utraque. 24*

*Flores terminales, subseni, congesti absque interjectis foliis, nec folio longiores.*

dichotoma.

2. **MIRABILIS** floribus sessilibus axillaribus erectis solitariis. *Amæn. acad. 4. p. 267. \**

*Admirabilis Jasmini rosa. Clus. hist. 2. p. 90.*

*Jalapa officinarum. Mart. cent. 1. t. 1.*

*Solanum mexicanum, flore parvo. Bauh. pin. 168. prodr. 91.*

*Habitat in Mexico. 24*

*Simillima M. Jalapæ. Flores noctu fragrantissimi.*

longiflora.

3. **MIRABILIS** floribus congestis terminalibus longissimis nutantibus, foliis subvillois. *act. Holmensch. 1756. p. 176. t. 6. f. 1. Amæn. acad. 4. p. 268.*

*Alzoyatl, Mirabilis mexicana. Hern. mex. 170. f. 2.*

*Habitat in Mexicæ frigidis montibus. Monnier. 24*

### CORIS.

mons-pelien-  
sis.

1. **CORIS.** *Hort. cliff. 68. Hort. upf. 46. Mat. med. 80. Roy. lugdb. 304. Sauv. monsp. 46.*

*Coris cærulea maritima. Bauh. pin. 280.*

*Symphytum petræum. Cam. epit. 699.*

*Habitat in Europæ australis arenosis maritimis. ☉*

*Caulis ruber. Folia alterna, linearia, crassiuscula, patentia. Flores spicati.*

### VERBASCUM.

Thapsus.

1. **VERBASCUM** foliis decurrentibus utrinque tomentosis. *Vir. cliff. 13. Fl. svec. 186, 195. Mat. med. 90. Roy. lugdb. 421. Dalib. parif. 71.*

Ver-

*Verbascum* caule simplici superne floribus sessilibus clavato, foliis utrinque lanigeris. *Hort. cliff.* 55.

*Verbascum* mas latifolium luteum. *Bauh. pin.* 239.

*Verbascum* latius. *Dod. pemp.* 153.

*Habitat in Europæ glareosis sterilibus.* ♂

2. *VERBASCUM* foliis cuneiformi-oblongis. *Hort. lychnitis.*  
*upf.* 45. *Fl. svéc.* 2. p. 196.

*Verbascum* foliis ovato-acutis subtus villosis crenatis, spicis laxis lateralibus & terminalibus. *Hort. cliff.*

54 *Roy. lugdb.* 421. *Dalib. paris.* 71.

*Verbascum* pulverulentum, flore luteo parvo. *Bauh. hist.* 3. p. 812. *Raj. hist.* 1094.

*Verbascum* mas, angustioribus foliis, floribus pallidis. *Bauh. pin.* 239.

- β. *Verbascum* lichnitis, flore albo parvo. *Bauh. pin.* 240.

*Verbascum* candidum femina. *Fuch. hist.* 847.

- γ. *Phlomis* mas alter. *Lob. ic.* 562.

*Habitat in Europæ ruderalis cultis.* ♂

3. *VERBASCUM* foliis ovatis utrinque tomentosis: phlomoides. inferioribus petiolatis.

*Verbascum* femina, flore luteo magno. *Bauh. pin.* 239.

*Verbascum* montanum tomentosum & incanum, folio subrotundo; caule non alato, staminibus purpureis. *Till. pis.* 171.

*Habitat in Italia.* ♂

*Caulis erectus, spicatus, valde tomentosus. Folia ovata, minime cordata, crenata: inferiora petiolata petiolo supra plano; superiora sessilia, semiamplexicaulia, non decurrentia. Racemus spiciformis ex Bracteis sparsis, lanceolatis. Flores intra singulas bractæas quatuor, quorum intermedius primum, tum infimus, demum laterales florent. Capsulæ oblongæ.*

4. *VERBASCUM* foliis oblongo - cordatis petiolatis. algrum. *Mat. med.* 91.

*Verbascum* foliis ovatis crenatis subtus tomentosis, caule angulato. *Vir. cliff.* 13. *Fl. svéc.* 187, 197.

*Roy. lugdb.* 421. *Dalib. paris.* 91.

*Verbascum* foliis ovatis subtus incanis crenatis, spica laxa rarius ramosa. *Hort. cliff.* 54.

*Verbascum* nigrum, flore ex luteo purpurascente. *Bauh. pin.* 240.

*Habitat in Europa ad pagos, vias.* 24

5. VER-

- phoeniceum* 5. VERBASCUM foliis ovatis crenatis radicalibus, caule subnudo racemoso.  
Verbascum foliis ovatis crenatis nudis scabris, caule ramoso. *Hort. cliff.* 55.  
Blattaria perennis, flore violaceo. *Morif. hist.* 2. p. 497.  
Blattaria purpurea. *Bauh. pin.* 241. *Raj. hist.* 1096.  
*Habitat in Europa orientali.* ♂  
*Blattaria nimis affinis.*
- Blattaria.* 6. VERBASCUM foliis amplexicaulibus oblongis glabris, pedunculis solitariis. *Hort. upf.* 46.  
Verbascum annuum, foliis oblongis sinuatis obtusis. *Hort. cliff.* 15. *Guett. stamp.* 2. p. 309.  
Verbascum foliis glabris serrato-dentatis, radicalibus sinuatis. *Roy. lugdb.* 421. *Dalib. paris.* 72. *Sauv. monsp.* 276.  
Blattaria lutea, folio longo laciniato. *Bauh. pin.* 240.
- ♂. Blattaria alba. *Bauh. pin.* 241.  
*Habitat in Europæ australioris argillaceis.* ☉
- Arcturus.* 7. VERBASCUM foliis lyrato-pinnatis.  
Verbascum humile creticum laciniatum. *Bauh. pin.* 240.  
Verbascum brassicæ folio. *Column. ecphr.* 2. p. 81. t. 82.  
Verbascum sylvestre creticum. *Alp. exot.* 123. t. 122.  
*Habitat in Creta.* ♀
- spinosum.* 8. VERBASCUM -caule folioso spinoso frutescente.  
*Amæn. acad.* 4. p. 307.  
Verbascum creticum spinosum frutescens. *Tournef. cor.* 8.  
Verbascum spinosum creticum. *Lob. illust.* 113.  
Leucojum creticum spinosum incanum luteum. *Bauh. pin.* 201.  
Leucojum spinosum. *Alp. exot.* 37. t. 36.  
Glastivida 1. cretica. *Pon. ital.* 114.  
*Habitat in Creta.* ♀
- sinuatum.* 9. VERBASCUM foliis radicalibus pinnatifido-repandis tomentosis, caulinis amplexicaulibus nudiusculis, rameis primis oppositis.  
Verbascum foliis pinnato-sinuatis hirsutis: subtus tomentosis. *Roy. lugdb.* 421. *Sauv. monsp.* 276.  
Verbascum nigrum, foliis papaveris corniculati. *Bauh. pin.* 240. *Camer. hist.* 403. t. 403.

8. *Verbascum græcum* fruticosum, folio sinuato candidissimo. *Tournef. cor. 8. itin. 1. p. 335. t. 335.*  
*Habitat* Monspeli, Florentiæ. ♂  
*Folia radicalia repanda f. obtuse sinuato-pinnata; Caulina oblonga, undulata, basi utrinque parum decurrentia. Ramea ovata f. cordata, parum decurrentia: horum prima opposita; reliqua alterna.*
10. *VERBASCUM* foliis incisus nudis, caule folioso, Osbeckii calycibus lanatis, pedunculis bifloris. *Ofb. it. 52.*  
*Blattaria orientalis, bugulæ folio, flore maximo virescente lituris luteis in semicirculum striato. Tournef. cor. 8. it. 2. p. 83. t. 83. Buxb. cent. 5. p. 17. t. 32.*  
*Habitat in Hispania; circa Bosphorum.*
11. *VERBASCUM* foliis lanatis radicalibus, scapo numyconi.  
*Cortusa foliis ovatis sessilibus. Hort. cliff. 50. Roy. lugdb. 417.*  
*Sanicula alpina, foliis boraginis, villosa. Bauh. pin. 243.*  
*Auricula urfi myconi. Dalech. hist. 837. Bauh. hist. 3. p. 869.*  
*Habitat in Pyrenæorum nemorosis. 24*

## D A T U R A.

1. *DATURA* pericarpis spinosis erectis ovatis: spinis ferox, supremis maximis convergentibus. *Amæn. acad. 3. p. 403. \**  
*Datura cochiniensis spinosissima. Zanon. hist. 1. p. 76.*  
*Stramonium f. Datura ferox, pomo crassioribus aculeis robustioribus. Herm. lugdb. 583. Moris. hist. 3. p. 607. f. 15 t. 2. f. 4. Raj. hist. 748.*  
*Stramonium ferox Bocc. sic. 50.*  
*Stramonium longioribus aculeis. Barr. rar. 109. t. 1112.*  
*Habitat in China. ☉*  
*Similis D. Stramonio, sed minus glabra & Capsule 4 spinis supremis maximis: in mea erectæ erant.*
2. *DATURA* pericarpis spinosis erectis ovatis, foliis ovatis glabris. *Hort. cliff. 55. Hort. upf. 43. Fl. sv. c. 185, 198. Gron. virg. 23. Roy. lugdb. 422. Dalib. parif. 70.* *Stramonium*



*Solanum foetidum*, pomo spinoso oblongo, flore albo. *Bauh. pin.* 168.

*Tatula. Cam. epit.* 176.

*Habitat in America, nunc vulgaris per Europam.* ☉

*Tatula.* 3. *DATURA* pericarpis spinosis erectis ovatis, foliis cordatis glabris dentatis.

*Solanum sativum*, pomo spinoso oblongo, flore albo.

*Bauh. pin.* 168?

*Stramonium majus purpureum.* *Raj. hist.* 748.

*Habitat* . . . . . ☉

*Similis D. Stramonio, sed duplo major. Caulis purpurascens; punctis albis adpersus, ad angulum acutum divisus, laevis. Corollae pallide caeruleae. Folia argutius dentata, si complanentur cordata.*

*Metel.* 4. *DATURA* pericarpis spinosis nutantibus globosis, foliis cordatis subintegris pubescentibus. *Hort. cliff.* 55. *Hort. ups.* 44. *Fl. zeyl.* 86. *Mat. med.* 85. *Roy. lugdb.* 422.

*Datura alba.* *Rumph. amb.* 5. p. 242. t. 87.

*Solanum pomo spinoso rotundo, longo flore.* *Bauh. pin.* 168.

*Humuatu.* *Rheed. mal.* 2. p. 47. t. 28.

*Habitat in Asia, Africa.* ☉

*fastuosa.* 5. *DATURA* pericarpis tuberculatis nutantibus globosis, foliis laevibus.

*Datura rubra.* *Rumph. amb.* 5. p. 243.

*Solanum foetidum*, fructu spinoso rotundo, semine pallido. *Bauh. pin.* 168.

*Nux Metella.* *Cam. epit.* 175.

β. *Solanum aegyptiacum*, flore pleno. *Bauh. pin.* 168.

*Stramonium* fructu spinoso rotundo, flore duplicitriplicive. *Tournef. inst.* 119.

*Habitat in Aegypto.* ☉

*Corolla extus violacea, intus alba, saepe gemina, una intra alteram s. duplicata.*

*arborea.* 6. *DATURA* pericarpis inermibus nutantibus, caule arboreo.

*Stramonoides arboreum*, oblongo & integro folio, fructu laevi. *Fewil. peruv.* 2. p. 761. t. 46.

*Habitat in Peru.* ☿

## HYOSCYAMUS.

1. **HYOSCYAMUS** foliis amplexicaulibus sinuatis, niger. floribus sessilibus. *Hort. cliff.* 56. *Fl. suec.* 184, 199. *Mat. med.* 86. *Roy. lugdb.* 422. *Dalib. parif.* 70. *Hall. helv.* 512.  
*Hyoscyamus vulgaris & niger. Bauh. pin.* 169.  
*Habitat in Europæ ruderatis pinguibus. ♂*
2. **HYOSCYAMUS** foliis caulinis petiolatis cordatis reticulatus. sinuatis acutis; floralibus integerrimis, corollis ventricosis.  
*Hyoscyamus foliis caulinis lanceolatis subdentatis, radicalibus sinuato-dentatis. Gron. orient.* 51.  
*Hyoscyamus, cauliculis spinosissimis, ægyptiacus. Bauh. pin.* 169.  
*Hyoscyamus ægyptius. Raj. hist.* 713.  
*Hyoscyamus rubello flore. Bauh. pin.* 169.  
*Hyoscyamus peregrinus. Clus. p. n.* 502.  
*Hyoscyamus peculiaris Cam. hort.* 77. t. 22.  
*Habitat in Creta, Syria, Ægypto. ☉*  
*Similis H. nigro, sed Folia caulina ovata, repanda, supra magis glabra; Floralia ovata, sessilia, integra. Flores pedunculo brevissimo. Corollæ campanulata, rubra, venis obscuris pulchre reticulata.*
3. **HYOSCYAMUS** foliis petiolatis sinuatis obtusis, albus. floribus subsessilibus. *Hort. cliff.* 56. *Roy. lugdb.* 422. *Sauv. monsp.* 275.  
*Hyoscyamus albus major. Bauh. pin.* 169.  
*Hyoscyamus albus minor. Bauh. pin.* 169.  
*Hyoscyamus albus vulgaris. Clus. hist.* 2. p. 118.  
*Habitat in Europa australi. ☉*  
*Variat Corollæ fauce atropurpurea & viridi.*
4. **HYOSCYAMUS** foliis petiolatis eroso-dentatis acutis, floribus pedunculatis, fructibus pendulis. *Hort. cliff.* 56. *Roy. lugdb.* 422.  
*Hyoscyamus creticus luteus major. Bauh. pin.* 169. *prodr.* 92.  
*Hyoscyamus albus creticus. Clus. hist.* 2. p. 84.  
*Hyoscyamus creticus luteus minor. Bauh. pin.* 169.  
*Hyoscyamus aureus. Alp. exot.* 99. t. 98.  
*Habitat in Creta, Oriente. ♂*  
*Caulis biennis, pilosus pedalis. Petioli valde pilosi. Folia lobata, denticulata. Pedunculi laterales petioli-  
*Limæi Spec. plant. pars I. R tib-**

tiolorum, erectis sub flore, omnino deflexi sub fructu. Corolla flavissima, aurèa, fauce atropurpurea striata, limbo semiquinquefido: lobo exteriori majore: sinu inferiore ultra limbum secto, ut in *Teucriis*, ad quod flamine & pistillum longissimum declinata.

- pufillus*. 5. *HYOSCYAMUS* foliis lanceolatis dentatis: floralibus inferioribus binis, calycibus spinosis. *Hort. upf.* 44. *Hyoscyamus* foliis lanceolatis. *Hort. cliff.* 56. *Roy. lugdb.* 422.

*Hyoscyamus* *pufillus* aureus americanus, antirrhini foliis glabris. *Pluk. alm.* 188. t. 37. f. 5.

*Habitat in Persia.* ☉

- physalodes*. 6. *HYOSCYAMUS* foliis ovatis integerrimis, calycibus inflatis subglobosis. *Hort. upf.* 44. \*  
- *Habitat in Sibiria.* 24

## NICOTIANA.

- pufilla*. 1. *NICOTIANA* foliis oblongo-ovalibus radicalibus, floribus racemosis acutis.

*Nicotiana* foliis ovato-lanceolatis obtusis rugosis, calycibus brevissimis. *Mill. ic.* 185. f. 2.

*Habitat in Vera cruce.*

- fruticosa*. 2. *NICOTIANA* foliis lanceolatis subpetiolatis amplexicaulibus, floribus acutis, caule fruticoso.

*Nicotiana* foliis lineari-lanceolatis acuminatis semiamplexicaulibus, caule fruticoso. *Mill. dict.*

*Habitat ad Cap. b. spei, China.* ☿

*Simillima N. Tabaco, ut fere varietas basi persistente.*

- Tabacum*. 3. *NICOTIANA* foliis lanceolato-ovatis sessilibus decurrentibus, floribus acutis.

*Nicotiana* foliis lanceolatis. *Hort. cliff.* 56. *Hort. upf.* 45. *Mat. med.* 87. *Roy. lugdb.* 423. *Sp. pl.* 1. p. 180.

*Nicotiana* major latifolia. *Bauh. pin.* 169.

*Blennochoes. Rencal. spec.* 37. t. 38.

*Habitat in America, nota Europæis ab 1560.* ☉

- rustica*. 4. *NICOTIANA* foliis petiolatis ovatis integerrimis, floribus obtusis.

*Nicotiana* foliis ovatis. *Hort. cliff.* 56. *Hort. upf.* 45. *Roy. lugdb.* 423. *Sp. pl.* 1. p. 180.

*Nicotiana* minor. *Bauh. pin.* 170.

*Pachyphylla Remeal. Spec. 42.*

*Habitat in America, nunc in Europa. ☉*

5. NICOTIANA foliis petiolatis cordatis integerrimis, paniculata. floribus paniculatis obtusis clavatis.

*Nicotiana foliis cordatis, floribus paniculatis: tubis clavatis. Sp. pl. 1. p. 180.*

*Nicotiana minor, folio cordiformi, tubo floris prælongo. Fowill. peruv. 1. p. 717. t. 10.*

*Habitat in Peru. ☉*

*Differt a N. rustica Caule teneviore, altiore, paniculato.*

*Corollis angustioribus: limbo brevissimo, obtusissimo, tubo longo clavato; calycibus capsulisque acutis.*

6. NICOTIANA foliis petiolatis cordatis integerrimis, glutinosa. floribus racemosis secundis ringentibus.

*Nicotiana foliis cordatis, corollis racemosis subringentibus, calycibus inæqualibus. Sp. pl. 1. p. 181.*

*Nicotiana militaris. Ait. hobb. 1753. p. 41. t. 2.*

*Habitat in Peru. B. Jussieu. ☉*

*Folia magis cordata, glutinosa. Flores inracemos paucos, longos. Corollæ ferme speciei primæ, sed magis inæquales & fere ringentes, fauce hiantes. Stamina ad latus superius inflexa. Calycis una lacinia reliquis duplo major.*

7. NICOTIANA foliis petiolatis cordatis crenatis, racemis recurvatis caule aculeato pruriginoso frutescente.

urena.

*Nicotiana arborescens spinosissima, flore exalbido.*

*Plum. Spec. 3. ic. 211.*

*Habitat in America meridionali. ‡*

*Frustrificatio Racemis secundis revolutis, Corollis campanulatis, Foliaque corollata præcedentis, sed crenata & tota arbor aculeata.*

## ATROPA.

1. ATROPA acaulis, scapis unifloris.

*Mandragora. Hort. cliff. 51. Roy. lugdb. 423. Mat. med. 88. Sp. pl. 1. p. 181.*

Mandragora

*Mandragora fructu rotundo. Bauh. pin. 169.*

*Mandragoras. Dod. pempt. 457.*

*Habitat in Hispaniæ, Italiæ, Cretæ, Cycladum apri-  
cis.*

- Belladonna.** 2. **ATROPA** caule herbaceo, foliis ovatis integris.  
*Atropa. Hort. cliff. 57. Hort. upf. 45. Roy. lugdb. 423.*  
*Solanum melanocerasus. Bauh. pin. 166.*  
*Solanum lethale. Clus. hist. 2. p. 86.*  
*Habitat in Austria, Angliæ, Italia montibus sylvo-*  
*sis. 24.*
- physalodes.** 2. **ATROPA** caule herbaceo, foliis sinuato-angulatis,  
 calycibus clausis acutangulis.  
*Alkekengi amplo flore violaceo. Few. peruv. 724. t. 16.*  
*Habitat in Peru. B. Jussieu. ☉*  
*Radix fibrosa, annua. Caulis bipedalis, patulus, ere-*  
*ctus, ramis angulatis. Folia alterna, glabra, oblonga,*  
*per petiolum decurrentia, obtuse sinuata (Daturæ).*  
*Pedunculi ad latus petiolorum, solitarii, nudi, uni-*  
*fiori. Calyx ovatus, basi excisus, angulis quinque*  
*compressis, profunde 5-partitus: foliolis sagittato-ova-*  
*tis. Corolla campanulata, parum quinqueloba, obtu-*  
*sa: Limbo erecto, subplicato, cærulescente. Stamina*  
*5, subulata, basi dilatata, subvillosa, corolla longe bre-*  
*viora, superne divergentia. Antheræ erectæ. Ger-*  
*men intra basin flaminum. Stylus filiformis longitudi-*  
*ne flaminum. Stigma capitatum. Fructus cernuus,*  
*tectus calyce majore, quinquangulati angulis compressis*  
*arcte clausis, basi mucronatis. Bacca exsueca, 3. 4. 5-*  
*locularis. Semina subrotunda, compressa. Media*  
*inter Atropam & Physalidem; a Physalide differt figu-*  
*ra corollæ, flaminibus distantibus, calyce quinqueparti-*  
*to, capsula loculamentis pluribus.*
- frutescens.** 4. **ATROPA** caule fruticoso, pedunculis confertis,  
 foliis cordato-ovatis obtusis.  
*Bella-donna frutescens rotundifolia hispanica. Tour-*  
*nef. inst. 77. Barr. ic. 1173.*  
*Habitat in Hispania. §*  
*Habitus & omnia Physalidis somniferæ, ut limites ge-*  
*nerum in solo calyce quinquepartito.*
- arborescens.** 5. **ATROPA** caule frutescente, pedunculis confertis,  
 corollis revolutis, foliis oblongis. *Amzn. acad. 4.*  
*p. 307.*  
*Bella-donna frutescens, flore albo, nicotianæ foliis.*  
*Plum. spec. 1. ic. 46. f. 1.*  
*Habitat in America meridionali. §*

## PHYSALIS.

\* *perennes.*

1. **PHYSALIS** caule fruticoso, ramis rectis, floribus somnifera. confertis.

*Physalis* caule fruticoso tereti, foliis ovatis integerrimis, floribus confertis. *Hort. cliff.* 62. *Roy. lugdb.* 426.

*Solanum somniferum verticillatum.* *Bauh. pin.* 166.

*Solanum somniferum.* *Clus. hist.* 2. p. 85.

*Habitat in Mexico, Creta, Hispania.* *h*

*Fructus* *Physalidis*; *Flos* *Atropæ*; *Herba præcedentis*, cui ita similis, ut vix genere separari queat.

2. **PHYSALIS** caule fruticoso, ramis flexuosis, floribus confertis. flexuosa.

*Baccifera indica*, floribus ad foliorum exortus, fructu sulcato decapyreno. *Raj. hist.* 1632.

*Pevetti.* *Rheed. mal.* 4. p. 113. t. 55.

*Habitat in India.* *h*

*Habitus* & *statura* *P. somnifera*; Flores itidem sparsi ad axillas foliorum; Calyces etiam nascuntur & involvunt baccam; manifeste vero differt floribus minoribus, ramis bifariam valde flexuosis, hinc folia quasi distiche disposita, flexurae angulis exterioribus inserta.

3. **PHYSALIS** caule fruticoso, foliis ovatis pilosis, floribus subsolitariis, corollis revolutis. arborefcens.

*Physalis* foliis ovato-lanceolatis integerrimis oppositis, caule fruticoso. *Mill. dict.* t. 206. f. 2.

*Habitat in Campechia.* *h*

4. **PHYSALIS** caule fruticoso, foliis ovatis tomentosis. curassavica.

*Vir. cliff.* 16. *Roy. lugdb.* 426.

*Physalis* caule perenni, foliis ovatis solitariis, ramis annuinis simplicissimis. *Hort. cliff.* 62. \*

*Solanum vesicarium curassavicum*, solano antiquorum simile, foliis origani subincanis. *Morif. hist.* 3. p. 527. *Pluk. alm.* 352. t. III. f. 5.

*Habitat in Curassao.* *h*

5. **PHYSALIS** foliis geminis repandis obtusis subto-mentosis, caule herbaceo superne paniculato. viscosa.

*Physalis* foliis cordatis integerrimis obtusis scabris, corollis glabris. *Sp. pl.* 1. p. 183.

*Physalis radice perenni, foliis cordatis obtusis. Hort. cliff. 496. Roy. lugdb. 426. Gron. virg. 24.*

*Alkekengi bonariense repens, bacca turbinata viscosa. Dill. elth. 11. t. 10. f. 10.*

*Habitat in Virginia, Bonaria. 24*

**Alkekengi. 6. PHYSALIS** foliis geminis integris acutis, caule herbaceo inferne subramoso.

*Physalis caule simplici annuo, foliis integerrimis geminatis, floribus solitariis. Hort. cliff. 62. Hort. upf. 49. Mat. med. 93. Dalib. parif. 73.*

*Physalis radice perenni, foliis lanceolato-cordatis. Vir. cliff. 16. Roy. lugdb. 426.*

*Solanum vesicarium. Bauh. pin. 166. Dod. pempt. 454.*

*Habitat in Italia, Germania, Japonia scrobibus. 24*

*Calyces maturi fructus saturatissime sanguinei.*

\* \* *Annua*

**angulata. 7. PHYSALIS** ramosissima, ramis angulatis glabris, foliis ovatis dentatis.

*Physalis annua ramosissima, ramis angulosis glabris, foliis dentato-ferratis. Hort. cliff. 62. Hort. upf. 50.*

*Fl. zeyl. 97. Roy. lugdb. 427.*

*Halicacabum f. Solanum indicum. Cam. hort. 70. t. 17.*

*Solanum vesicarium indicum. Bauh. pin. 166.*

*Alkekengi indicum glabrum, chenopodii folio. Dill. elth. 13. t. 12. f. 12.*

**β. Alkekengi indicum glabrum, capsicii folio. Dill. elth. 12. t. 11. f. 11.**

*Habitat in India utraque. ☉*

**pubescens. 8. PHYSALIS** ramosissima, foliis villoso-viscosis, floribus pendulis.

*Physalis annua ramosissima, ramis teretibus pubescentibus: geniculis nodosis. Hort. cliff. 62.*

*Alkekengi virginianum, fructu luteo. Fervill. peruv. 3. p. 5. t. 1.*

*Solanum vesicarium virginianum procumbens annuum, folio lanuginoso. Morif. hist. 3. p. 527. f. 13. t. 3. f. 24.*

*Habitat in India utraque, imprimis in Virginia. ☉*

*Fructus villoso-viscosus, nec glaber; maculae corollae fuscae, nec pallidae. Pedunculi nutantes, nec cernui.*

9. PHY-

9. **PHYSALIS** *ramosissima*, pedunculis fructiferis folio villoso longioribus. *Hort. cliff.* 62. *Roy. lugdb.* 427. *Solanum vesicarium indicum minimum.* *Herm. lugdb.* 569. t. 571.  
*Pee-inota-inodien.* *Rheed. mal.* 10. t. 140. f. 71.  
*Habitat in Indiæ aridis sordidis.* ☉
10. **PHYSALIS** *ramosissima*, foliis villosis, pedunculis strictis. *Alkekengi indicum glabrum, chenopodii folio.* *Dill. elkh.* 12. t. 12. f. 12.  
*Habitat in America.* ☉  
*Facies antecedentium* 7, 8, 9, *sed antheræ flavae, nec caeruleæ; rami teretes: supra plani villosi; Folia villoso-viscosa; Alæ ramificationum viridi quasi prulina germinant. Pedunculi floriferi longi, erecti, stricti. Flores cernui. Calyces fructus maximi, quinquanguli, pedunculo stricto penduli.*

## SOLANUM.

\* *inermia.*

1. **SOLANUM** caule inermi fruticoso, foliis ovatis tomentosis integerrimis, umbellis compositis. *Solanum arborescens, verbasci folio.* *Plum. spec.* 4. *Solanum maxime tomentosum, spinis carens, virginianum.* *Pluk. alm.* 351. t. 216. f. 1.  
*Habitat in America.* ☿
2. **SOLANUM** caule inermi fruticoso, foliis ovatis integerrimis; pedunculis lateralibus filiformibus. *Solanum caule inermi perenni, foliis ovatis integris, floribus ad basin ramulorum confertis.* *Roy. lugdb.* 424.  
*Solanum africanum lignosum, folio atroviridi angusto oblongo obtuso.* *Bærh. lugdb.* 2. p. 68.  
*Habitat in Guinea.* ☿
3. **SOLANUM** caule inermi fruticoso, foliis lanceolatis repandis, umbellis sessilibus. *Solanum caule inermi fruticoso, foliis ovato-lanceolatis integris, floribus solitariis.* *Hort. cliff.* 61. *Hort. ups.* 48. *Roy. lugdb.* 424.  
*Solanum fruticosum bacciferum.* *Bauh. pin.* 61.



*Pseudocapsicum. Dod. pempt. 718.*

*Habitat in Madera. §*

- diphyllum. 4. **SOLANUM** caule inermi fruticoso, foliis geminis: altero minore, floribus cymosis.

*Solanum caule inermi perenni, foliis ovato-lanceolatis geminis: altero minimo. Vir. cliff. 15. Hort. cliff. 61. Roy. lugdb. 424.*

*Solanum americanum strychnodendro accedens, fructu medio quasi fisso. Pluk. alm. 549. t. 111. f. 4. mlla.*

*Habitat in America? §*

- Dulcamara. 5. **SOLANUM** caule inermi frutescente flexuoso, foliis superioribus hastatis, racemis cymosis. *Hort. cliff. 60. Fl. spec. 189, 201. Mat. med. 95. Roy. lugdb. 424. Dalib. parif. 73.*

*Solanum scandens f. Dulcamara. Bauh. pin. 167.*

- β. **Solanum dulcamarum africanum**, foliis crassis hirsutis. *Dill. elth. 365. t. 273. f. 352.*

*Habitat in Europæ sepibus humentibus. §*

- quercifolium. 6. **SOLANUM** caule inermi herbaceo angulato flexuoso scabro, foliis pinnatifidis, racemis cymosis.

*Solanum foliis quernis. Few. peruv. 722. t. 15.*

*Habitat in Peru. D. Jusseu. 24*

*Habitus S. Dulcamara. Caules angulati, erecti, scabri. Folia oblonga, in petiolum decurrentia, venosa, glabra, subtus scabra, pinnatifida. Pedunculi ramosi. Corollæ violacæ, obtusiores quam Dulcamara: maculis duabus viridibus ad basin singuli lobi. Antheræ flavæ, stylo breviores. Baccæ ovatæ.*

- radicans. 7. **SOLANUM** caule inermi herbaceo lævi teretiusculæ prostrato radicante, foliis pinnatifidis, racemis cymosis. *Linn. decad. 1. t. 10. \**

*Habitat in Peru. 24*

- bonariensis. 8. **SOLANUM** caule subfrutescente subinermi, foliis cuneiformibus sinuato-repandis.

*Solanum bonariense arborescens, papas floribus. Dill. elth 364 t. 272 f. 351.*

*Solanum arborescens, foliis angulatis, fructu aureo.*

*Plum. spec. 4. ic. 224. f. 2.*

*Habitat in agro Bonariensi. §*

- Folia* ☿ *Flores maximi*; *Aculei aliquot in tenella planta, non vero in adulta. Flores plures steriles. Fructus globosi, grandes, lutei.*
9. SOLANUM caule inermi herbaceo, foliis ovatis æthiopicum  
repando-angulatis, pedunculis fertilibus unifloris  
cernuis. *Amæn. acad.* 4. p. 367. \*  
*Solanum pomiferum, fructu rotundo striato duro.*  
*Bauh. pin.* 167. *Bauh. hist.* 3. p. 620.  
*Solanum pomiferum herbariorum. Lob. ic.* 265.  
*Habitat in Æthiopia. Burserus. China.* ☉  
*Caulis rigidulus. Folia S. bonariensis, palmaria, obtusa. Fructus ruber, magnus, depressus, profunde torulosus* ☿ *quasi in lobos sectus, durus.*
10. SOLANUM caule inermi herbaceo, foliis pinna- tuberosum.  
tis integerrimis, pedunculis subdivisis. *Vir. cliff.*  
15. *Hort. cliff.* 60. *Hort. ups.* 48. *Roy. lugdb.* 423.  
*Dalib. parif.* 73.  
*Solanum tuberosum esculentum. Bauh. pin.* 167. *prodr.*  
89. t. 89.  
*Habitat in Peru: innotuit 1590. C. Bauhino.* ☉. 24
11. SOLANUM caule inermi herbaceo, foliis pinna- Lycopersi-  
cam.  
tis incis, racemis simplicibus. *Vir. cliff.* 15. *Hort.*  
*cliff.* 60. *Roy. lugdb.* 423.  
*Solanum pomiferum, fructu rotundo striato molli.*  
*Bauh. pin.* 167.  
*Pomum amoris. Cam. epit.* 821. *Rumph. amb.* 3. p.  
416. t. 154. f. 1.
- β. *Solanum racemosum cerasorum forma. Bauh. pin.*  
167. *prodr.* 90.  
*Habitat in America calidiore.* ☉. β. 24
12. SOLANUM caule inermi herbaceo, foliis pinna- pimpinelli-  
folium.  
tis integerrimis, racemis simplicibus. *Amæn. acad.*  
4. p. 268. \*  
*Habitat in Peru.* ‡
13. SOLANUM caule inermi herbaceo, foliis pinna- peruvianum  
tis tomentosis incis, racemis bipartitis foliosis,  
baccis subpilosis.  
*Lycopersicon pimpinellæ folio. Ferv. peruv.* 3. p. 37.  
t. 25.  
*Habitat in Peru. B. Jussieu.* 24  
*Radix perennis, vix vel parum supra radicem fruti-*  
*caus. Folia pinnata, tomentosa, incana, pinnata*  
abs.

- absque foliolis alternis minutis. Racemi bipartiti foliolo ad quemeumque pedicellum ovato. Corollæ late flavæ, multo majores quam in Lycopersico.*
- montanum. 14. SOLANUM caule inermi herbaceo, foliis subcordatis repandis.  
*Solanum tuberosum minus, atriplicis folio, vulgo Papa montana. Few. peruv. 3. p. t. 46.*  
*Habitat in Peru. 2.*
- nigrum. 15. SOLANUM caule inermi herbaceo, foliis ovatis dentato-angulatis, umbellis nutantibus.  
*vulgatum. Solanum caule inermi herbaceo, foliis ovatis angulatis. Vir. cliff. 15. Hort. cliff. 60. Hort. upf. 48. Fl. fvec. 188, 200. Mat. med. 94. Gron. virg. 24. Roy. lugdb. 433. Dalib. parif. 72.*  
*Solanum officinarum. Bauh. pin. 166.*
- patulum. β. — *ramis teretibus glabris, foliis integerrimis glabris.*  
*Solanum procerius patulum vulgaris fructu. Dill. elth. 367. t. 275. f. 355.*
- villosum. γ. — *ramis teretibus villosis, foliis angularis subvillosis.*  
*Solanum annuum hirsutius, baccis luteis. Dill. elth. 366. t. 274. f. 353.*
- guineense. δ. — *ramis angulatis dentatis, foliis integerrimis glabris.*  
*Solanum guineense, fructu magno instar cerasi. Dill. elth. 360. t. 274. f. 354.*
- virginicum. ε. — *ramis angulatis dentatis, foliis repandis glabris.*  
*Solanum nigrum vulgari simile, caulibus exasperatis. Dill. elth. 368. t. 275. f. 256.*
- judaicum. ζ. — *ramis aculeis incurvis, foliis repandis nudis.*  
*Habitat in Orbis totius cultis. ☉*  
*Tot varietates β. γ. δ. ε. ζ. videntur esse hybride proles.*
- \* \* Spinosæ.
- Melongena. 16. SOLANUM caule inermi herbaceo, foliis ovatis tomentosis integris, calycibus aculeatis, fructu pendulo.  
*Solanum calycibus aculeatis, foliis ovatis integerrimis tomentosis. Hort. cliff. 61. Hort. upf. 49. Fl. zeyl. 93. Roy. lugdb. 426.*  
*Solanum pomiferum, fructu oblongo. Bauh. pin. 167. Mala insana. Dod. pempt. 458.*  
*Trongum hortense. Rumph. amb. 5. p. 238. t. 85.*  
*Habitat in Asia, Africa, America. ☉*

17. SOLANUM caule aculeato herbaceo, foliis cordatis angulatis tomentosis aculeatis, baccis hirtis calyce obtectis. ferox.

*Habitat in Malabar.*

Statura *S. Melongena*. Caulis herbaceus. Folia cordata, angulata, ampla, petiolata, venis aculeis erectis adpersa, supra villosa, subtus tomentosa. Aculeati sunt petioli, pedunculi, calyces. Pedunculi axillares, simplices, breves. Calyces campanulati, tomentosi, valide aculeati, obtegentes totas baccas. Baccæ admodum hirtæ.

18. SOLANUM caule aculeato herbaceo hirtio, foliis cordatis sinuatis subtomentosis, calycibus aculeatis. Campechiense.

*Vir. cliff.* 16. *Hort. cliff.* 61. Solanum campechiense, calycibus echinatis. *Dill. elth.* 361. t. 268. f. 347.

*Habitat in America ad finem Campechiensem.*

*Folia utrinque testâ setis stellatis divergentibus.*

19. SOLANUM caule aculeato herbaceo, foliis cordatis angulato-lobatis: utrinque villosis aculeatis. mammosum

*Vir. cliff.* 15. *Hort. cliff.* 61. *Hort. upf.* 49. *Roy. lugdb.* 425.

Solanum barbādense spinosum annuum, fructu aureo rotundiore pyri parvi inversi forma & magnitudine.

*Pluk alm.* 350. t. 225. f. 1. *Merian. surin.* 27. t. 27.

*Habitat in Virginia, Barbados.* ☉

*Folia longitudine & latitudine æqualia, obtusiuscula.*

20. SOLANUM caule petiolisque aculeatis; foliis sinuato-angulatis supra glabris, floribus paniculatis. paniculatum.

*Jurepeba. Pis. bras.* 181.

*Habitat in Brasilia.*

Caules & Petioli albo-tomentosi aculeis rectis, rarioribus, nudis. Folia tomentosa, lata, cordata, profunde angulata, acutiuscula, inermia, supra glabra. Panicula terminalis, composita, tomentosa, uti Calyces inermis.

21. SOLANUM caule aculeato herbaceo, foliis pinatifidis utrinque aculeatis: lacinis sinuatis obtusis, calycibus aculeatis. virginianum.

*Roy. lugdb.* 425. \* Solanum americanum laciniatum spinosissimum. *Dill. elth.* 360. t. 267. f. 346.

*Solanum* annuum virginianum nigricans spinosissimum, flore cæruleo. *Pluk. alm.* 351. t. 62. f. 3.

*Habitat in America.* ☉

Indicum.

22. *SOLANUM* caule aculeato fruticoso, foliis cuneiformibus angulatis subvillosis integerrimis: aculeis utrinque rectis. *Fl. zeyl.* 94.\*

*Solanum* caule aculeato fruticoso, foliis ovatis sinuatis integerrimis: spinis utrinque erectis. *Hort. cliff.* 61. *Roy. lugdb.* 424.

*Solanum indicum* spinosum, flore boraginis. *Robert. ic.* 28. *Dill. elth.* 362. t. 270. f. 349.

*Solanum americanum* perenne subincanum, fructu pyriformi longiore, spinis plurimis armatum. *Pluk. alm.* 350. t. 225. f. 6.

*Habitat in India utraque.* ☿

fuscatum.

23. *SOLANUM* caule aculeato fruticoso, foliis ovatis obtuse lobatis: aculeis utrinque rectis: superioribus coloratis.

*Solanum americanum*, caule & pedunculo nigro, foliis acanthi spinosis. *Barrh. lugdb.* 2. p. 68.

*Habitat in America.* ☿ ☉

Rami varie spinosi, latere solari fuscescentes. Folia repanda, sinuata, obtusa, utrinque viridia: spinis in superiore pagina late purpureis, sæpe fascia pallida cinctis. Pedunculi spinis validioribus. Flores violacei.

carolinense.

24. *SOLANUM* caule aculeato annuo, foliis hastatoangulatis: aculeis utrinque rectis, racemis laxis.

*Solanum* caule aculeato fruticoso, foliis ovatis lateribus laciniatis: aculeis utrinque rectis. *Hort. cliff.* 61.\* *Vir. cliff.* 15. *Roy. lugdb.* 424.

*Solanum carolinense* spinosum, boraginis floribus spicatis. *Dill. elth.* 362. t. 269. f. 348.

*Habitat in Carolina.* ☉

Caulis quasi fruticosus, sed annuus. Folia basi integra, sinuata angulis acutiusculis, spinis in costa nec lateribus. Racemi foliis longiores laxi, simplices.

sodomeum.

25. *SOLANUM* caule aculeato fruticoso, foliis obovatis pinnatifido-sinuatis obtusis sparfe aculeatis nudis, calycibus aculeatis.

*Solanum* caule aculeis recurvis, foliis sinuatis subtus to-

tomentosis undique aculeatis, pedunculis aculeatis.

*Fl. zeyl.* 95. \* *Hort. ups.* 49.

*Solanum* caule aculeato fruticoso, foliis oblongis pinato-laciniatis: spinis supinis erectis. pronis recurvis. *Hort. cliff.* 61. *Vir. cliff.* 15. *Roy. lugdb.* 424.

*Solanum* pomiferum frutescens africanum spinosum nigricans, boraginis flore, foliis profunde laciniatis. *Herm. lugdb.* 573. t. 575. *Morif. hist.* 3. p. 521. f. 13. t. 1. f. 15.

*Solanum* spinosum, profunde laciniatis foliis subtus lanuginosis, maderaspatanum. *Pluk. alm.* 351. t. 316. f. 4.

*Habitat in Africa.* §

26. *SOLANUM*. caule aculeato fruticoso, aculeis tomentosis foliisque oblique ovatis repandis. sanctum.

*Solanum* caule aculeato fruticoso, foliis repandis, calycibus aculeatis. *Roy. lugdb.* 425.

*Solanum* spinosum, fructu rotundo. *Bauh. pin.* 167. a quo *Burserus*.

*Habitat in Palæstina.* §

Caulis tomentosus, cinereus, aculeis crassis, rectis, brevibus, flavescentibus, excepto apice tomentosus. Folia ovata, altera basi breviora, tomentosa, crassa, obtusa: tenellis pinnatim sinuatis margine albidis; aculeis in costa tribus. Petioli subtus aculeati. Pedunculi ex latere caulis; pedicello primario cum calyce spinoso, reliquis masculis inermibus. Corolla Boraginis, caruleo-purpurascens.

27. *SOLANUM* caule aculeato fruticoso, aculeis aceto-mentosis, foliis cordatis inermibus subrepandis: tenellis purpureo-pulverulentis.

*Solanum* caule aculeato fruticoso, foliis cordatis villosis repandis, calycibus inermibus. *Hort. cliff.* 61. *Roy. lugdb.* 425.

*Solanum* africanum spinosum, folio canescente undulato. *Triumf. præl.* 46. t. 6. *Raj. suppl.* 355.

*Solanum* foliis & caule spinosis. *Morif. blæs.* 310.

*Solanum* spinosum maxime tomentosum. *Bocc. fr.* 8. t. 5.

*Habitat in Æthiopia.* §

Caulis & tota planta tomentosa aculeis angustissimis, nudis. Folia cordata, obtusa, repanda, fere sola cysta spinosa: tenella prodeuntia margine pulvere viol.

*olaceo ex pilis stellatis. Baccæ fulvæ, magnitudine Grossulariæ.*

- bahamense. 28. SOLANUM caule aculeato fruticoso, foliis lanceolatis repandis obtusis margine reflexis, racemis simplicibus. *Hort. cliff. 61. Roy. lugdb. 424.*

*Solanum bahamense spinosum, petalis angustis reflexis. Dill. elth. 263. t. 271. f. 250.*

*Solanum fruticosum bacciferum spinosum, flore cæruleo. Sloan. jam. 108. hist. 1. p. 38. t. 11. f. 3.*

*Habitat in Americæ insula Providentia. b*

- igneum. 29. SOLANUM caule aculeato fruticoso, foliis lanceolatis acuminatis basi utrinque revolutis, racemis simplicibus.

*Solanum spiniferum frutescens, spinis igneis, americanum. Pluk. alm. 350. t. 225. f. 5.*

*Habitat in America. b*

*Simillimum S. bahamensi, sed Flores albi, nec violacei; Folia acuminata, nec obtusifuscula. Spinæ validiores, copiosiores, rubro-ignæ.*

- trilobatum. 30. SOLANUM caule aculeato fruticoso, foliis cuneiformibus subtrilobis glabris obtusis inermibus.

*Solanum spinosum jamaicense glabrum, foliis parvis minus profunde lacinlatis. Pluk. alm. 351. t. 316. f. 5.*

*Habitat in Jamaica, ad Cap. b. Spei. b*

*Caulis subfrutescens, alius, angustus. Folia parva, glabra, obtusa, subtri-f. quinqueloba, raro etiam ad basin lobata. Aculei reflexi in caule, petiolis, foliis, pedunculis. Pedunculi uniflori; Calyces minimi, inermes aut aculeati. Stylus adscendens. Baccæ parvæ Sambuci: An duæ species, cum una floribus parvis, albis; altera magnis, violaceis?*

## CAPSICUM.

- annuum. 1. CAPSICUM caule herbaceo, pedunculis solitariis. *Roy. lugdb. 426.*

*Capsicum annuum. Hort. cliff. 59. Hort. ups. 47. Fl. zeyl. 92. Mat. med. 96.*

*Capsicum filiquis longis propendentibus. Tournes. inst. 152.*

*Piper indicum vulgatissimum. Bauh. pin. 102.*

*Vallis-capo-molago. Rheed. mal. 2. p. t. 35.*

*Ha-*

*Habitat in America meridionali.* ☉

*Fructus hujus & sequentis speciei mire variat figura.*

1. CAPSICUM caule fruticoso, pedunculis geminis. frutescens.

*Roy. lugdb.* 426.

*Capsicum frutescens. Hort. cliff.* 59.

*Capsicum brasiliannum Clus. exot.* 340. f. 2.

*Capsicum indicum. Rumph. amb.* 5. p. 247. t. 88. f.

1, 3, 4.

- β. *Capsicum fructu minimo conico rubro.* • *Brown.*

*jam.* 176.

*Capsicum minus, fructu parvo pyramidali erecto.*

*Sloan. jam.* 112. *hist.* 1. p. 240. t. 146. f. 2.

*Piper filiquosum magnitudinis baccarum asparagi.*

*Bauh. hist.* 2. p. 944.

*Piper brasiliannum Petita. Clus. eur.* 55. *Rumph. amb.*

5. p. 247. t. 88. f. 2.

*Habitat in Indiis.* ☿

## STRYCHNOS.

1. STRYCHNOS foliis ovatis, caule inermi. *Fl. Zeyl.* Nux vomica.

91. *Mat. med.* 77.

*Strychnos foliis quinquenerviis. Wach. ultraj.* 32.

*Nux Vomica officinarum. Bauh. pin.* 511. *Raj. hist.*

1814.

*Colubrina ligni tertium genus in malabar. Bauh. pin.*

301.

*Caniram. Rheed. mal.* 1. p. 67. t. 37. *Raj. hist.* 1661.

*Habitat in India.* ☿

2. STRYCHNOS foliis ovatis acutis, cirrhis simplicibus. colubrina.

*bus. Mat. med.* 78. *Amæn. acad.* 2. p. 119.

*Strychnos foliis trinerviis ovatis binatis. Wach. ultraj.* 32.

*Clematis indica spinosa, foliis luteis. Bauh. pin.* 301.

*Fructus orbicularis major fuscus striatus. Bauh. pin.* 405.

*Modira-caniram. Rheed. mal.* 7. p. 10. t. 5. *Raj. hist.*

1661.

*Habitat in India.* ☿

## JAQUINIA.

1. JAQUINIA foliis lanceolatis acuminatis. *Jacq.* 4- ruscifolia.

*mer.* 15.

*Medeola aculeata foliis verticillatis, ramis aculeatis.*

*Sp. pl.* 1. p. 329.

Fru-



*Fruticulus foliis rufci stellatis. Dill. clth. 148. t. 123.  
f. 149.*

*Habitat in America meridionali. b*

armillaris. 2. JAQVINIA foliis obtusis cum acumine. *Jacq. amer.  
15. †*

*Chrysophyllum Barbasco. Læfl. it. 204. 277.*

*Habitat in America. b*

linearis. 3. JAQVINIA foliis linearibus acuminatis. *Jacq. amer.  
15. t. 40.*

*Habitat in America. b*

## CHIRONIA.

trinervia. 1. CHIRONIA herbacea, calycinis foliolis membrana-  
ceo carinatis. *Fl. zeyl. 90. \**

*Chironia foliis lanceolatis trinerviis. Hort. cliff. 54.*

*Lyfimachia folio sinuato acuminato trinervio, flore  
cæruleo amplo, calyce carinato alato. Burm. zeyl.  
145. t. 67.*

*Habitat ad Cap. b. Spei. O*

Usoides. 2. CHIRONIA herbacea, foliis linearibus. *Hort. cliff.  
54. Roy. lugdb. 433.*

*Rapuntio affinis, lini facie, capitis b. spei. Breyn.  
cent. 175. t. 90.*

*Habitat in Capitis b. Spei berbidis. 24*

jasminoides. 3. CHIRONIA herbacea, foliis lanceolatis. *Amxr. acad.  
6. afr. 5. \**

*Habitat ad Cap. b. spei.*

campanula-  
ta. 4. CHIRONIA herbacea, foliis sublinearibus, calyci-  
bus longitudine corollæ.

*Habitat in Canada. Kalm.*

*Caulis pedalis, teres: Ramis longis. Folia lanceolato li-  
nearia, lævia. Flores terminales, solitarii, rotati,  
purpurei, pedunculo longo insidentes; Calycis foliola  
longitudine corollæ subulata. Antheræ spirales.*

angularis. 5. CHIRONIA herbacea, caule acutangulo, foliis ova-  
tis amplexicaulibus.

*Habitat in Virginia. Kalm.*

*Facies Centaurii minoris. Caulis pedalis, tetragonus;  
alis membranaceis. Folia ovata, hypericæ, opposita,  
ses-*

*Jessilia* & *semiamplexicaulia*. Flores congesti, ut in *Hyperico*; Antheræ spirales; Stylus bifidus.

6. **CHIRONIA** floribus duodecimfidis. dodecandra.  
*Gentiana* floribus duodecim-petalis, foliis distinctis.

*Gron. virg.* 27.

*Habitat in Virginia.*

7. **CHIRONIA** frutescens baccifera. baccifera.  
*Centaureum minus* arborescens pulpiferum. *Comm. rar.*  
*9. t. 9.*

*Centaureum minus africanum* arborescens angustifolium.

*Old. afr.* 26.

*Habitat in Æthiopia.* §

8. **CHIRONIA** fruticosa, foliis linearibus subtomentosis, calycibus campanulatis. frutescens.  
*Anan. acad.* 4. p. 308.  
*n. 130.*

*Chironia frutescens capsulifera.* *Sp. pl.* 1. p. 190. *Mill. dist.* 1. 97.

*Centaureum minus africanum* arborescens latifolium, flore ruberrimo. *Comm. rar.* 3. t. 8. *Old. afr.* 26.

*Centaureum* foliis binis oppositis angustis linearibus, flore magno rubente. *Burm. afric.* 205. t. 74. f. 1.

*Habitat in Æthiopia.* §

*Calyx tubo corollæ multo amplior. Stylus deflexus.*

## CORDIA.

1. **CORDIA** foliis ovatis tomentosis, corymbis latera- Myxa.  
libus, calycibus decemstriatis.

*Cordia* foliis subovatis serrato-dentatis. *Hort. cliff.* 53.

*Mat. med.* 171. *Sp. pl.* 1. p. 190.

2. *Sebestena sylvestris* & domestica. *Bauh. pin.* 445. *Alp. ægypt.* 30.

*Sebestena domestica* f. *Myxa.* *Comm. hort.* 1. p. 139.  
t. 72.

*Vidi-maram.* *Rheed. mal.* 4. p. 77. t. 37.

*Habitat in Ægypto, Malabar.* §

3. **CORDIA** foliis lanceolato-ovatis scabris, panicula Geraschan-  
terminali calycibus decemstriatis. thus.

*Geraschanthus* foliis ovato-oblongis utrinque productis, racemis terminalibus. *Brown. jam.* 170. t. 29. f. 3.

*Habitat in Jamaica.* §

- Sebestena.* 3. *CORDIA* foliis oblongo - ovatis repandis scabris.  
*Hasselq. ii. 458.*  
*Cordia nucis juglandis folio. Plum. gen. 13. ic. 105.*  
*Cordia foliis amplioribus hirtis, tubo floris subæquali.*  
*Brown. jam. 202.*  
*Sebestena scabra, flore miniato crispo. Dill. elh. 341.*  
*t. 255. f. 331.*  
*Caryophyllus spurius inodorus, folio subrotundo scabro, flore racemoso hexapetaloide coccineo. Sloan. jam. 136. hist. 2. p. 20. t. 64. Raj. suppl. 86. Catesb. car. 2. p. 91. t. 91.*  
*Novella nigra. Rumph. amb. 2. p. 226. t. 75.*  
*Habitat in Indiis. h*  
*Folia juniora serrata, adulta vix repanda, suprema certe integerrima, ovato - oblonga, scabra. Petioli teretes. Stipulæ nullæ. Calyx cylindricus, oblongus ore trifido. Corolla figura Mirabilis, fulvo-lutea, æstivatione rugosissima Papaveris. Stamina 5. Stylus bis bifidus.*
- macrophylla.* 4. *CORDIA* foliis ovatis villosis sesquipedalibus.  
*Collococcus platyphyllos major, racemis umbellatis.*  
*Brown. jam. 168.*  
*Prunus racemosa, foliis oblongis hirtutis maximis, fructu rubro. Sloan. jam. 184. hist. 2. p. 130. t. 221. f. 1. Raj. dendr. 43.*  
*Habitat in Jamaica. h*
- Collococa.* 5. *CORDIA* foliis oblongo-ovatis integerrimis, floribus subcorymbosis, calycibus interne tomentosis.  
*Cordia glabra. Sp. pl. 1. p. 191.*  
*Collococcus foliis rugosis venosis oblongo-ovatis, floribus laxè racemosis. Brown. jam. 167.*  
*Ceraso affinis arbor baccifera racemosa, flore pentapetalo herbaceo guttato, fructu coccineo monopyreno viscido, semine rugoso. Sloan. jam. 169. hist. 2. p. 95. t. 203. f. 2.*  
*Habitat in Jamaica. h*

## EHRETIA.

- clinifolia.* 1. *EHRETIA* foliis oblongo-ovatis integerrimis glabris, floribus paniculatis. *Amæn. acad. 5. p. 395.*  
*Ehretia foliis alternis oblongis acuminatis, spica florum sparsa, petalis reflexis albis. Trew. ehret. t. 24.*  
*Ehre-*

*Ehretia arborea*, foliis oblongo-ovatis alternis, racemis terminalibus. *Brown. jam.* 168. t. 16. f. 1.

*Cerafo* affinis arbor baccifera racemosa, flore albo pentapetalo, fructu flavo monoppyreno eduli dulci. *Sloan. jam.* 2. p. 94. t. 203. f. 1. *Raj. dendr.* 45.

*Habitat in Jamaica.* ♀

2. *EHRETIA spinosa.* *Jacq. amer.* 14. †

*spinosa.*

*Habitat in America.* ♀

2. *EHRETIA* foliis ovatis integerrimis lævibus, floribus subcorymbosis, calycibus glabris. *Bourreria.*

*Cordia Bourreria.* *Amæn. acad.* 5. p. 395. \*

*Bourreria* fructibus succulentis. *Jacq. amer.* 14.

*Bourreria arborea* foliis ovatis alternis, racemis rarioribus terminalibus. *Brown. jam.* 168. t. 15. f. 2.

*Mespilus americana* laurifolia glabra, fructu rubro mucaginoso. *Com. hort.* 1. p. 153. t. 79.

*Pittonia* similis laureolæ foliis, floribus albis, baccis rubris. *Catesb. car.* 2. p. 79. t. 79.

*Jasminum periclymeni* folio, flore albo, fructu flavo rotundo tetrapyreno. *Sloan. jam.* 169. *hist.* 2. p. 96. t. 204. f. 1. *Raj. dendr.* 63.

*Habitat in Jamaica.* ♀

*Folia exacte ovata*, utraque pagina lævia. *Corollæ* tubus longitudine tantum *Calycis.* *Bourreria* affinitate jungit *Ehretias* cum *Cordiis.*

4. *EHRETIA* foliis cuneiformi-lanceolatis margine exsucca, reflexis. †

*Bourreria exsucca.* *Jacqu. amer.* 14.

*Rhamnus cumanensis.* *Læfl. it.* 182.

*Habitat in America meridionali.* ♀

## VARRONIA.

1. *VARRONIA* foliis lanceolatis lineatis, pedunculis linearibus lateralibus petiolo adnatis, spicis globosis. *Amæn. acad.* 5. p. 394. \*

*Lantana corymbosa*, foliis alternis, floribus corymbosis. *Sp. pl.* 628.

*Ulm* angustifoliæ facie baccifera jamaicensis, foliis superne scabris, subtus villosis, floribus flavis perpusillis, fructu botryoide monospermo. *Pluk. alm.* 393. t. 328. f. 5.

*Habitat in America.* ♀

82

For

*Folia lanceolata, utrinque attenuata. Pedunculi laterales, elongati, filiformes, basi petiolo adnati.*

bullata,

2. VARRONIA foliis ovatis venoso-rugosis, spicis globosis. *Amén. acad.* 5. p. 394.

Varronia mirabiloides, spicis subrotundis inæqualibus, corollis hypocrateriformibus. *Jacqu. amer.* 14.

*Habitat in America.* §

globosa.

3. VARRONIA foliis lanceolato-oblongis, caule dichotomo, pedunculis axillaribus elongatis nudis, spicis globosis.

Varronia globosa spicis æqualibus globosis. *Jacqu. amer.* 14.

*Habitat in America.* §

*Pedunculus e dichotomia nudus. Calycis dentes lineares, longi, recurvati.*

curassavica.

4. VARRONIA foliis lanceolatis, spicis oblongis. *Jacqu. amer.* 14.

Varronia assurgens farmentosa, foliis & capitulis oblongis. *Brown. jam.* 172.

Lantana foliis alternis, spicis oblongis. *Sp. pl.* 1. p. 627.

Periclymenum rectum, salviæ folio rugoso majore oblongo bullato, flore albo, fructu longiore. *Sloan. jam.* 163. *hist.* 2. p. 81. *Raj. dendr.* 31.

*Habitat in America.* §

alba.

5. VARRONIA foliis cordatis, floribus cymosis. *Jacqu. amer.* 14.

Mespilus americana, alni vel coryli foliis, fructu mucaginoso albo. *Comm. hort.* 1. p. 155. t. 80.

*Habitat in America.* §

## LAUGIERIA.

oderata.

1. LAUGIERIA. *Jacqu. amer.* 16. †

Edechia. *Læfl. it.* 306, 271, 259.

*Habitat in America.* §

## BRUNSFELSIA.

americana.

1. BRUNSFELSIA.

Brunsfelsia flore albo, fructu croceo molli. *Plum. gen.* 12. ic. 65.

*Habitat in America meridionali.* §

*Folia oblonga, obtusa, integerrima, petiolata. Flores in apice ramorum plures; Antheræ extra corollæ tubum prominentes.*

## CESTRUM.

1. **CESTRUM** floribus pedunculatis. *Hort. cliff.* 490. nocturnum.  
Jasminoides foliis pishaminis, flore virescente noctu odoratissimo. *Dill. elth.* 183. t. 153. f. 185.

Jasminum laurinis foliis, flore pallido luteo, fructu atrocæruleo polypyreno venenato. *Sloan. jam.* 169. *hist.* 2. p. 96. t. 204. f. 2. *Raj. dendr.* 63. *Brown. jam.* 173.

*Syringa laurifolia jamaicensis, floribus ex flavo palliscentibus. Pluk. alm.* 35. t. 64. f. 3.

*Parxu. Fewill. peruv.* 2. p. 32. t. 32. f. 1.

*Habitat in Jamaica, Chilli. ♀*

2. **CESTRUM** floribus sessilibus. *Hort. cliff.* 491. diurnum.

Jasminoides laureolæ folio, flore candido interdū odorato. *Dill. elth.* 186. t. 154. f. 186.

Laureola sempervirens americana, latioribus foliis, floribus albis odoratis. *Pluk. alm.* 209. t. 95. f. 1. *Raj. dendr.* 53.

*Hediunda jasmini flore. Fewill. peruv.* 2. p. 25. t. 20. f. 3.

*Habitat in Chilli, Havana. ♀*

## LYCIUM.

1. **LYCIUM** foliis linearibus. *Hort. cliff.* 57. *Hort. upf.* afrum.  
47. *Mill. dict.* t. 171. f. 1. *Trew. chret.* 4. t. 24.

Jasminoides africanum, jasmini aculeati foliis & facie. *Niff. act.* 1711. p. 420. t. 12.

Jasminoides aculeatum, salicis folio. *Mish. gen.* 224. t. 105. f. 2.

*Rhamnus alter, foliis salis, flore purpureo. Bauh. pin.* 477.

*Habitat in Africa & in Regno Valentino. ♀*

*Caulis rectus, rigidus. Flores atropurpurei. Baccae globosæ, atrorubæ.*

2. **LYCIUM** foliis lanceolatis crassiusculis, calycibus barbarum  
bifidis.

*Rhamnus peregrinus, rosmarini folio, candidior. Pluk. alm.* 317. *mont.* 160. t. 322. f. 2. mala.

*Jasminoides aculeatum* humile, halimi minoris folio, fructu per maturitatem flavescente. *Mich. gen.* 224. t. 105. f. 1.

*Jasminoides sinense*, halimi folio longiore & angustiore. *Duham. arb.* 1. p. 306. t. 121. f. 4.

*Jasminoides aculeatum*, polygoni folio, floribus parvis albidis. *Schaw. afric.* 349. f. 349?

*Habitat in Asia, Africa, Europa.* ‡

*Calycis ore bilobo, aut interdum trilobo, a reliquis speciebus diversum. Frutex debilis, ni sustentetur nutans & decumbens. Flores intus pallide rubicundi. Baccæ ovata, flavæ. Cortex ramorum exalbidus. Hyemes nostrates fert.*

capfulare.

3. **LYCIUM** foliis lanceolatis tenuibus glabris, pedunculis calycibusque pubescentibus, pericarpiliis capsularibus. *Amzn. acad.* 4. p. 308. \*

*Habitat in Mexico. Miller.* ‡

## CHRY SOPHYLLUM.

Cainito,

1. **CHRY SOPHYLLUM** foliis ovatis parallele striatis subtus tomentoso-nitidis. *Jacq. amer.* 15.

*Chrysophyllum* foliis ovatis superne glabris parallele striatis, subtus tomentoso-nitidis. *Hort. cliff.* 49.

*Anona* foliis subtus ferrugineis, fructu rotundo majore lævi purpureo, semine nigro partim rugoso partim glabro. *Sloan. jam.* 206. *hist.* 2. p. 170. t. 229. *Raj. dendr.* 78.

*Cainito. Laet. amer.* 390.

*Cainito* folio subtus aureo, fructu maliformi. *Plum. gen.* 10.

- β. *Cainito* folio subtus aureo, fructu oliviformi. *Plam. gen.* 10. ic. 69.

*Sideroxylon Pacurero. Læfl. it.* 204.

*Habitat in America calidiore.* ‡

*Folia alterna, petiolata, ovata, integerrima. Pedunculi laterales, uniflori, numerosi, brevissimi.*

glabrum.

2. **CHRY SOPHYLLUM** foliis utrinque glaberrimis. *Jacq. amer.* 15. \*

*Habitat in America.* ‡

## SIDEROXYLON.

inermis

1. **SIDEROXYLON** inermis. *Hort. cliff.* 69.

Side-

*Sideroxylon primum* f. dein *Coriæ indorum* nomine data arbor. *Dill. elth.* 357. t. 265. f. 344.

*Padus foliis oblongis, fructu solitario.* *Burm. afric.* 238. t. 84. f. 2.

*Habitat in Æthiopia.* ‡

1. *SIDEROXYLON spinosum, foliis deciduis.* *Duham.* lycioides. arb. 2. p. 260. t. 68.

*Lycioides.* *Hort. cliff.* 488. \*

Arbor folio salicis viridi alterno splendente, spinis longioribus alternis ad alas foliorum. *Boerh. lugdb.* 2. p. 263.

*Habitat in Canada.* ‡

3. *SIDEROXYLON spinosum, foliis perennantibus.* spinosum. *Hort. cliff.* 69.

Caro - mœll. *Reed. mal.* 5. p. 77. t. 39. *Raj. hist.* 1634.

*Habitat in Malabar.* ‡

## RHAMNUS.

\* *Spinosi.*

1. *RHAMNUS spinis terminalibus, floribus quadrifidis* catharticus. dioicis, foliis ovatis. *Hort. cliff.* 70. *Fl. jvec.* 193.

202. *Mat. med.* 72. *Roy. lugdb.* 224.

*Rhamnus catharticus.* *Bauh. pin.* 478.

*Cervispina.* *Cord. hist.* 175.

*Habitat in Europæ sepibus.* ‡

2. *RHAMNUS spinis terminalibus, foliis linearibus.* lycioides. *Rhamnus tertius flore herbaceo, baccis nigris.* *Bauh.*

*pin.* 477.

*Rhamnus tertius forte niger.* *Clus. hist.* 1. p. 110.

*Habitat in Hispania.* ‡

3. *RHAMNUS spinis terminalibus, foliis oblongis integerrimis.* Oleoides.

*Rhamnus hispanicus, oleæ folio.* *Tournef. inst.* 539.

*Habitat in Hispaniâ.* C. *Alströmer.* ‡

*Similis R. cathartico, sed minor.* Folia ad gemmas 2.

f. 3, *Oleæ similia* & *minora, levia, confidentia, integerrima, quasi venis marginata, brevius petiolata, obtusiuscula, reticulato-venosa. Fructus solitarii, similes R. cathartico.*



\* \* *inermes*.

- alpinus. 4. **RHAMNUS** *inermis*, floribus dioicis, foliis duplicato-crenatis.  
*Rhamnus inermis*, foliis glabris crenatis. *Aët. gatt.* 349. t. 16.  
*Frangula ora folii ferrata*. *Hall. helv.* 164.  
*Frangula rugosiore & ampliore folio*. *Tournef. inst.* 612.  
*Frangula altera polycarpus*. *Bauh. prodr.* 160.  
*Alnus nigra baccifera*, rugosiore folio f. major. *Bauh. hist.* 1. p. 562.  
*Alnus nigra polycarpus*. *Bauh. pin.* 428.  
*Habitat in alpinis Helveticis.* ‡
- Frangula. 5. **RHAMNUS** *inermis*, floribus monogynis hermaphroditis, foliis integerrimis. *Hort. cliff.* 70. *Fl. suec.* 194. 203. *Mat. med.* 73. *Roy. lugdb.* 224.  
*Rhamnus inermis*, foliis annuis. *Fl. lapp.* 60.  
*Alnus nigra baccifera*. *Bauh. pin.* 428.  
*Frangula*. *Dod. pempt.* 784.  
*Habitat in Europæ borealis nemorosis humidiusculis.* ‡
- Sarcophalus. 6. **RHAMNUS** *inermis*, foliis ovalibus coriaceis integerrimis emarginatis. *Amæn. acad.* 5. p. 395. \*  
*Sarcophalus foliis ovatis glabris alternis apice leviter emarginatis*, cortice interiore ferrugineo. *Brown. jam.* 179.  
*Habitat in America.* ‡
- micranthus. 7. **RHAMNUS** *inermis*, foliis ovato-lanceolatis obliquis pubescentibus, stipulis lanceolatis deciduis. *Amæn. acad.* 5. p. 395. \*  
*Rhamnus an Zizyphus arborefcens*, foliis oblongo-ovatis hirsutis & leniter serratis. *Brown. jam.* 173. t. 12. f. 2.  
*Muntingia folio corni*, fructu minore. *Plum. gen.* 41. ic. 205. f. 1.  
*Habitat in America.* ‡
- colubrinus. 8. **RHAMNUS** *inermis*, floribus monogynis hermaphroditis, capsulis trilocularibus, ramis horizontalibus, floribus erectis. *Jacq. amer.* 16.  
*Arbor baccifera indica*, foliis majoribus splendentibus, flore pentapetalo. *Comm. hort.* 1. p. 175. t. 90.  
*Ha-*

*Habitat in America.* ‡

*Folia magna, ovata, integerrima.*

9. RHAMNUS inermis, floribus dioicis, stigmate triplici, foliis ferratis. *Vir. cliff.* 19. *Hort. upf.* 47. *Sauv. monsp.* 95. Alaternus.

Rhamnus inermis, floribus polygamis, stigmate triplici, foliis ferratis. *Hort. cliff.* 70. *Roy. lugdb.* 225.

Phylica elatior. *Bauh. pin.* 477.

Phylica humilior. *Bauh. pin.* 477.

Alaternus 1, 2. *Clus. hist.* 1. p. 50.

*Habitat in Europa australi.* ‡

10. RHAMNUS inermis, floribus hermaphroditis, capsulis trilocularibus, foliis rugosis integerrimis ciliatis. *Jacq. amer.* 16. cubensis.

*Habitat in America.* ‡

*Hic uti R. colubrinus Fructu capsulari est.*

11. RHAMNUS inermis, floribus hermaphroditis, foliis ovatis lineatis repandis subtus reticulatis. *Amæn. acad.* 4. p. 308. \* lineatus.

Rhamnus lineata. *Osborn. it.* 219. t. 7.

*Habitat in Chinæ elevatis.* ‡

\* \* \* *Aculeati.*

12. RHAMNUS aculeis geminatis: inferiore reflexo, *Paliurus.* floribus trigynis. *Hort. cliff.* 69. *Hort. upf.* 47. *Roy. lugdb.* 224. *Sauv. monsp.* 306.

Rhamnus folio subrotundo, fructu compresso. *Bauh. pin.* 477.

*Paliurus, Dod. pempt.* 576.

*Habitat in Europa australi.* ‡

12. RHAMNUS aculeis geminatis: altero recurvo, *Lotus.* foliis ovato-oblongis.

*Zizyphus sylvestris. Tournesf. instit.* 627. *Shaw. afr.* 631. f. 631.

*Habitat in Regno Tunetano.* ‡

Frutex ramulis teretibus. Aculei duo ad exortum foliorum: altero horizontali, altero recurvo ut in *Paliuro*, sed Folia omnia *zizyphi*, figura, nervis, glabritie, magnitudine, at margo obsolete & rarius crenatus.

- Napeca.** 14. **RHAMNUS** aculeis subgeminatis recurvis, pedunculis corymbosis, floribus semidigynis, foliis serratis utrinque lævibus. *Fl. zeyl.* 87. \*
- Jujuba indica spinosa, folio & fructu longiore. *Pluk. alm.* 199. t. 216. f. 2. *Raj. dendr.* 44.
- Vidara littorea. *Rumph. amb.* 2. p. 119. t. 42.
- Habitat in Zeylona.* h
- Jujuba.** 15. **RHAMNUS** aculeis folitariis recurvis, pedunculis aggregatis, floribus semidigynis, foliis retatis subtus tomentosis. *Fl. zeyl.* 89. \*
- Jujuba indica spinosa, folio & fructu rotundo. *Pluk. alm.* 199. *Raj. dendr.* 44.
- Malus indica. *Rumph. amb.* 2. p. 117. t. 36.
- Perim toddal. *Rheed. mal.* 4. p. t. 41.
- Habitat in India.* h
- Oenopli.** 16. **RHAMNUS** aculeis folitariis recurvis, pedunculis aggregatis subseffilibus, foliis semicordatis subtus tomentosis. *Fl. zeyl.* 88. \*
- Jujuba aculeata, nervosis foliis infra sericeis flavis. *Burm. zeyl.* 131. t. 61.
- Habitat in Zeylona.* h
- Iguaneus.** 17. **RHAMNUS** aculeis geminatis altero patente, racemis axillaribus monoicis, foliis nudis. *Jacq. amer.* 16.
- Jujube americana spinosa, loti arboris foliis, & facie, fructu rotundo parvo dulci. *Comm. hort.* 1. p. 141. t. 73.
- Habitat in America.* h
- Zizyphus.** 18. **RHAMNUS** aculeis geminatis rectis, floribus digynis, foliis ovato-oblongis glabris. *Hort. cliff.* 69.
- Mat. med.* 74. *Roy. lugdb.* 224. *Sauv. monsp.* 59.
- Jujuba sylvestris. *Bauh. pin.* 446.
- Zizyphus. *Dod. pempt.* 807.
- β. Jujubæ majores oblongæ. *Bauh. pin.* 446.
- Habitat in Europa australi.* h
- Spina Christi.** 19. **RHAMNUS** aculeis geminatis rectis, foliis ovatis. *Hort. cliff.* 63. \* *Roy. lugdb.* 224.
- Oenoplia spinosa. *Bauh. pin.* 477. *Clus. hist.* 2. p. 313.
- Nabca Paliurus athenæi credita. *Alp. ægypt.* 16. t. 19.
- Jujube f. Zizyphus africana, mucronatis foliis, spina gemella. *Pluk. alm.* 199. t. 197. f. 3.

Ha-

*Habitat in Æthiopia, Palæstina. ‡*  
*Arbustula e seminibus missis sub nomine Spinæ Christi*  
*Hierosolyma a D. Hasselquist.*

PHYLICA.

1. PHYLICA foliis linearibus verticillatis. ericoides.  
*Phylica foliis ovato-linearibus. Horr. cliff. 70. Roy. lugdb. 199.*  
*Alaternoides africana, ericæ foliis, floribus albicantibus & muscosis. Comm. hort. 2. p. 1. t. 1.*  
*Habitat in Æthiopia. ‡*
2. PHYLICA foliis lineari-subulatis: summis hirsutis. plumosa.  
*Roy. lugdb. 199.*  
*Ricinus arborescens africanus, tomentosis capitulis. Sch. thes. 1. p. 38. t. 23. f. 4, 5.*  
*Chamaelea africana tricoccus & tetracoccus taxifolia, villosa cæsarie coronata. Pluk. mant. 45. t. 342. f. 3.*  
*Chamaelea foliis angustis subtus incanis, floribus capitatis muscosis. Burm. afr. 117. t. 43. f. 2.*  
*Alaternoides africana, rorismarini latiore & pilosiore folio. Comm. prælud. 63. t. 13.*  
*Habitat in Æthiopia. ‡*
3. PHYLICA foliis linearibus triquetris sparsis, caule radiata,  
 piloso, bracteis apice glabris coloratisque. *Amæz. acad. 4. p. 938.*  
*Chrysanthemum ericoides coronatum. C. B. S. Breyer. cent. 165. t. 82. Pluk. mant. 47. t. 454. f. 7.*  
*Habitat ad Cap. b. Spei. ‡*
4. PHYLICA foliis cordato-ovatis patentibus, caule cordata,  
 prolifero.  
*Alaternoides africana, chamæmespilio folio. Comm. rar. 62. t. 12.*  
*Habitat ad Cap. b. Spei. ‡*  
*Differt a, PH. buxifolia, quod Folia non ovata, sed cordata exquisite attenuato apice. Caulis prolifer, nec capitula lateralita.*
5. PHYLICA foliis cordatis, floribus dioicis. dioica.  
*Habitat ad Cap. b. Spei. ‡*
6. PHYLICA foliis ovatis sparsis ternisque subtus to- buxifolia.  
 mentosis.

Cha-

- *Chamælea folio subrotundo subtus incano, floribus in capitulum collectis. Burm. afr. 119. t. 44. f. 1.*  
*Chamælea africana, foliis subrotundis. Herm. afr. 6.*  
*Habitat in Æthiopia. ‡*

## CEANOTHUS.

- americanus. 1. **CEANOTHUS** foliis trinerviis. *Mill. dist. t. 86.*

*Duham. arb. 1. p. 138. t. 51.*

*Ceanothus corymbis folio longioribus. Hort. upf. 51.*

*Aët. upf. 1741. p. 77.*

*Celastrus inermis, foliis ovatis ferratis trinerviis, racemis ex summis alis longissimis. Hort. cliff. 73.*  
*Gron. virg. 25. Roy. lugdb. 435.*

*Evonymus novi belgii, corni feminæ foliis. Comm. hort. 1. p. 167. t. 86. Raj. dendr. 69.*

*Evonymus, jujubinis foliis, carolinensis, fructu parvo fere umbellato. Pluk. alm. 139. t. 28. f. 6.*

*Habitat in Virginia, Carolina. ‡.*

- asiaticus. 2. **CEANOTHUS** foliis ovatis enerviis.

*Ceanothus racemis folio brevioribus. Fl. zeyl. 98. \**

*Grossularia spinis vidua, baccis in racemo congestis spadiceis, foliis crenatis ovato-acuminatis. Burm. zeyl. 111. t. 48.*

*Spireæ theophrasti similis frutex indicus, latiori folio.*

*Pluk. alm. 350. t. 63. f. 2?*

*Habitat in Zeylona. ‡*

- africanus. 3. **CEANOTHUS** foliis lanceolatis enerviis, stipulis subrotundis.

*Celastrus inermis, foliis lanceolatis obtuse ferratis petiolatis appendiculatis. Hort. cliff. 73. Roy. lugdb. 435.*

*Alaternoides africana, lauri ferratæ folio. Comm. præl. 61. t. 11.*

*Ricinoides africana arborefcens, folio phillyreæ longifoliæ ferratæ. Seb. thes. 1. p. 35. t. 22. f. 6.*

*Habitat in Æthiopia. ‡*

## BYTTNERIA.

- scabra. 1. **BYTTNERIA**. *Læfl. it. 313.*

*Chætæa aculeata. Jacq. amer. 17.*

*Habitat in America meridionali. ‡*

*Fru-*

*Frutex asuleatus. Folia lanceolata. Flores racemosi, albi.*

## MYRSINE.

1. MYRSINE. *Hort. cliff. 72. \* Zinn. gart. 202.* africana.  
*Vitis idæa æthiopica, myrti tarentinæ folio, flore atropureo. Comm. Hort. 1. p. 123. t. 64.*  
*Buxus africana rotundifolia ferrata. Pluk. alm. 74. t. 80. f. 5.*  
*Frutex æthiopicus baccifera, foliis myrtilli. Breyn. cent. 9. t. 5.*  
*Habitat in Æthiopia. ‡*

## CELASTRUS.

1. CELASTRUS inermis, foliis ovatis integerrimis. bullatus.  
*Evonymoides virginiana, foliis non serratis, fructu coccineo eleganter bullato. Isnard. act. 1716. p. 369.*  
*Evonymus virginianus rotundifolius, capsulis coccineis eleganter bullatis. Pluk. alm. 139. t. 28. f. 5.*  
*Habitat in Virginia. ‡*
2. CELASTRUS inermis, caule volubili, foliis serratis. scandens.  
*Evonymoides canadensis scandens, foliis serratis. Isn. act. 1716. p. 369. Duham. arb. 1. p. 223. t. 95.*  
*Frutex viminibus lentis infirmis, foliis profunde serratis. Gron. virg. 55.*  
*Habitat in Canada. ‡*
3. CELASTRUS inermis, foliis ovatis ferrulatis, floribus racemosis, caule erecto. myrtifolius.  
*Myrtifolia arbor, foliis latis subrotundis, flore albo. Sloan. jam. 162. hist. 2. p. 79. t. 193. f. 1. Raj. dendr. 36.*  
*Habitat in Virginia, Jamaica. ‡*
4. CELASTRUS spinis foliosis, ramis angulatis, foliis obtusis. buxifolius.  
*Lycium portoricense, buxi foliis angustioribus. Pluk. alm. 234. t. 202. f. 3.*  
*Habitat in Æthiopia. ‡*  
*Rami & ramuli angulati. Folia obovata, obtusa, acute serrata. Cymæ pedunculatæ, ex alis.*
5. CELASTRUS spinis nudis, ramis teretibus, foliis pyracanthus acutis. *Hort. cliff. 72. Mill. dict. t. 87.*

Ly-

*Lycium æthiopicum*, pyracanthæ folio. *Comm. hort.* 1. p. 163. t. 84. *Raj. dendr.* 72.

*Evonymo* affinis æthiopica, lycii foliis & aculeis, fructu evonymi. *Pluk. alm.* 139. t. 280. f. 5.

*Habitat in Æthiopia.* †

Rami & ramuli teretiusculi. Folia ovali-lanceolata, utrinque acuta, in petiolos desinentia, superne serraturis acuminatis ut fere spinosis, persistētia; Cymæ pedunculatæ, laterales.

## EVONYMUS.

- europæus. 1. *EVONYMUS* floribus plerisque quadrifidis.  
*Evonymus* foliis oblongo-ovatis. *Hort. cliff.* 38. *Tl. fvec.* 133, 204. *Roy. lugdb.* 436. *Hall. herb.* 523.
- temisfolius. *Evonymus vulgaris*, granis rubentibus. *Bauh. pin.* 428.  
*Evonymus.* 2. *Clus. hist.* 1. p. 57.
- latifolius. β. *Evonymus latifolius.* *Bauh. pin.* 428.  
*Evonymus* 1. f. latifolia. *Clus. hist.* 2. p. 56.  
*Habitat in Europæ sepibus; β in Pannonia.* †  
*Ex flore primario convenit cum congeneribus.* *Phil. Bot.* 178.
- americanus. 2. *EVONYMUS* floribus omnibus quiquefidis.  
*Evonymus* foliis lato-lanceolatis serratis. *Hort. upf.* 30.  
*Evonymus* foliis lanceolatis. *Gron. virg.* 17.  
*Evonymus virginianus*, pyracanthæ foliis, capsula verrucarum instar exasperata. *Pluk. alm.* 139. t. 115. f. 5.  
*Celastrus* foliis oppositis ovatis integerrimis, floribus subfolitariis. *Hort. cliff.* 32.  
*Rhus virginianum*, folio myrti. *Comm. hort.* 1. p. 157. t. 81. *Raj. dendr.* 57.  
*Habitat in Virginia.* †

## DIOSMA.

- oppositifolia. 1. *DIOSMA* foliis subulatis acutis oppositis. *Hort. cliff.* 71. *Roy. lugdb.* 434.  
*Spiræa africana*, foliis cruciatim positis. *Comm. rar.* 1. t. 1.  
*Hypericum africanum vulgare.* *Seb. thes.* 2. p. 41. t. 40. f. 5.  
*Habitat ad Caput b. Spei.* †
- hirsuta. 2. *DIOSMA* foliis linearibus hirsutis. *Hort. cliff.* 71.  
*Roy. lugdb.* 424.

Spi-

*Spiræa africana* odorata, foliis pilosis. *Com. rar.* 3.

t. 3.

*Habitat ad Caput b. Spei.* §

2. DIOSMA foliis linearibus mucronatis glabris carina-  
tis subtus bifariam punctatis.

*Diosma foliis fetaceis acutis.* *Hort. cliff.* 72.

*Spiræa africana* odorata, floribus suave rubentibus

*Comm. rar.* 2. t. 2.

*Erica æthiopica*, rosmarini sylvestris folio eleganter  
punctato, flore tetrapetalo purpureo. *Pluk. mant.* 68.

t. 347. f. 4.

*Habitat in Æthiopia.* §

*Habitus juniperi.* Folia juniperina, glabra, subtri-  
etra, subtus punctis pertusa.

4. DIOSMA foliis lineari-lanceolatis subtus convexis ericoides  
bifariam imbricatis.

*Spiræa africana*, ericæ bacciferæ foliis. *Raj. dendr.* 91.

*Ericaformis*, coridis folio, æthiopica, floribus pen-  
tapetalis in apicibus. *Pluk. amalrh.* 236. t. 279. f. 5.

*Habitat in Æthiopia.* §

*Hæc valde fragrans in Emplastris Hottentottorum.*

3. DIOSMA foliis lanceolatis ciliatis.

ciliata

*Spiræa* forte genus africanum, serpilli hirsutis foliis,  
fruticosum, floribus albis umbellatis. *Pluk. amalrh.*

197. t. 411. f. 3. *Seb. mus.* 2. t. 17. f. 5.

*Habitat in Æthiopia.* §

6. DIOSMA foliis lanceolatis glabris.

lanceolata.

*Spiræa africana*, saturejæ foliis brevioribus. *Raj.*  
*dendr.* 91.

*Habitat in Æthiopia.* §

7. DIOSMA foliis lanceolato-ovalibus glanduloso-cre-  
natis, floribus solitariis. *Amæn. acad.* 4. p. 308. \*

*Habitat in Æthiopia.* §

8. DIOSMA foliis ovato-oblongis, floribus solitariis uniflora,  
terminalibus.

*Cistus humilis æthiopicus*, inferioribus foliis rosmarini  
sylvestris punctatis, cæteris autem serpilli subrotun-  
dis, flore carneo. *Pluk. mant.* 49. t. 342. f. 3.

*Habitat in Æthiopia.* §

Folia rigidiora, & longe majora, quam in præceden-  
tibus.



tibus. Flos etiam sextuplo major, margine ut ♀ sub-  
tus punctatus.

- pulchella. 9. DIOSMA foliis ovatis obtusis glanduloso-crenatis,  
floribus geminis axillaribus.  
*Habitat in Æthiopia. ♂*  
Fruticulus pulcherrimus Ramis virgatis. Folia exquisi-  
ta. Floſculi cærulei, minimi, axillares, gemini pro-  
priis pedunculis.

### HARTOGIA.

- capensis. 1. HARTOGIA.  
*Habitat ad Cap. b. ſpei. ♂*  
*Statura Dioſmæ aut Brunie. Folia oppoſita, ſubulata,*  
*triquetra. Flores albi in Corymbum.*

### BRUNIA.

- nodiflora. 1. BRUNIA foliis quadrifariam imbricatis: *Hort. cliff. 70.*  
*Cupreſſo-Pinulus capitis b. ſpei. Breyn. cent. 22. t. 10.*  
*Erica capitata ſ. nodiflora cupreſſiformis africana. Pluk'*  
*mant. 69. t. 346. f. 4.*  
*Scabiolaæ affinis Arbuſcula africana ericoides ſphæro-*  
*cephalos. Raj. hiſt. 1444.*  
*Habitat in Æthiopia. ♂*
- lanuginosa. 2. BRUNIA foliis linearibus patulis apice calloſis. *Hort.*  
*cliff. 71.*  
*Tamarifcus monomopatensis, ericæ tenuiſſimo folio,*  
*pillulifera, ramulis lanugine araneorum inſtar imple-*  
*xis. Pluk. alm. 361. t. 318. f. 4.*  
*Habitat in Æthiopia. ♂*
- ciliata. 3. BRUNIA foliis ovatis acuminatis ciliatis.  
*Habitat in Æthiopia. ♂*
- abrotanoides. 4. BRUNIA foliis lineari-lanceolatis patentibus trique-  
tris apice calloſis.  
*Brunia foliis creberrimis lanceolatis, floribus conglom-*  
*batis. Burm. afric. 266. t. 100. f. 1.*  
*Leviſanus africanus, ericæ folio, capitulo majore.*  
*Raj. dendr. 130.*  
*Leviſanus capensis, ſerpylli folio. Pet. gaz. 9. t. 5.*  
*f. 7.*  
*Erica capitata ſ. nodiflora, corioſi foliis rectis, æthio-*  
*pica. Pluk. mant. 69. t. 346. f. 7.*  
*Habitat in Æthiopia. ♂*

5. BRU-

5. **BRUNIA** foliis spatulatis acutis carnosiss. Levisianus.  
*Brunia* foliis oblongis incanis, florum capitulo ramu-  
 lum terminante. *Burm. afric.* 267. t. 100. f. 2.  
*Protea fusca* foliis ovatis obtusis imbricatis, capitulis  
 globosis. *Sp. pl.* 1. p. 95.  
*Protea* foliis oblongo-ovatis lanis involutis. *Roy. lugdb.*  
 187.  
*Conocarpodendron* folio subrotundo brevissimo *Boerh.*  
*lugdb.* 2. p. 202. t. 202.  
*Chrysanthemum* fruticosum, polygoni foliis, africa-  
 num, caulibus scabris, flore minore. *Pluk. mant.*  
 47. t. 343. f. 9.  
*Habitat in Æthiopia.* ♀
6. **BRUNIA** floribus solitariis. *Hort. cliff.* 71. uniflora.  
*Ericaformis* æthiopica, cupressi foliis compressiusculis:  
*Pluk. alm.* 136. t. 279. f. 2.  
*Habitat in Æthiopia.* ♀

## I T E A.

1. **ITEA.** *Gron. virg.* 143. *Duham. arb.* 1. p. 219. t. 126. virginica.  
*Diconangia.* *Mitch. gen.* 5.  
*Habitat in Virginia.* ♀  
*Folia Padi integrâ.* *Racemi angustii.*

## G A L A X.

1. **GALAX.** aphylla.  
*Viticella.* *Mitch. gen.* 24.  
*Anonymos* f. *Belvedere.* *Gron. virg.* 25.  
*Habitat in Virginia.* 24.

## C E D R E L A.

1. **CEDRELA** floribus paniculatis. ocorata.  
*Cedrela Cedro.* *Læfl. it.* 183.  
*Cedrela* foliis pinnatis, floribus laxè racemosis, ligno  
 levi odorato. *Brown. jam.* 158. t. 10. f. 1.  
*Cedrus barbadensium*, alatis fraxini foliis. *Pluk. alm.*  
 92. t. 157. f. 1.  
*Pruno* forte affinis arbor maxima materie rubra laxa  
 odorata. *Sloan. jam.* 182. *hist.* 2. p. 128. t. 220 f. 2.  
*Raj. dendr.* 43.  
*Habitat in America meridionali.* ♀

*Linnaei Spec. plant. pars I.*

T

MAN.

## MANGIFERA.

indica.

## 1. MANGIFERA.

*Mangifera arbor.* Bon. jav. 95. *Fl. zeyl.* 471.*Manga domestica.* Rumph. amb. 1. p. 93. t. 25.*Manga indica*, fructu magno reniformi. *Raj. hist.* 1550.*Perficæ* similis, putamine villosa. *Bauh. pin.* 440.*Mao* f. *Mau* f. *Manghos.* *Rheed. mal.* 4. p. 1. t. 1. 2.*Habitat in India.* ‡*Racemus compositus, brachiatus; an flores omnes hermaphroditi? utrum una aut plures species? alius dies determ.net.*

## CUPANIA.

americana.

## 1. CUPANIA.

*Cupania castaneæ folio*, fructu sericeo & ramoso.*Plum. gen.* 45. ic. 110.*Habitat in America calidiore.* ‡*Folia alterna, obovata, oblonga, obtusa, dentato-ser-rata: nervis transversalibus crassis. Flores racemosi, an Monoici?*

## HIRTELLA.

americana.

1. HIRTELLA. *Hort. cliff.* 17. \**Frutex innominatus.* *Maregr. bras.* 78. f. 2, mala.*Habitat in Basilia.* ‡*Facies Royenæ lucida. Racemus simplex, erectus.*

## RIBES.

\* *Ribes inermis.*

rubrum.

## 1. RIBES inermis, racemis glabris pendulis, floribus planiusculis.

*Ribes inermis, floribus planiusculis, racemis pendulis.**Hort. cliff.* 82. *Fl. svec.* 197, 205. *Hort. upf.* 51. *Mat.**med.* 99. *Vir. cliff.* 21. *Roy. lugdb.* 270. *Dalib. parif.* 75.*Ribes vulgare acidum.* *Bauh. hist.* 2. p. 97. *Fl. lapp.* 68.*Grossularia*, multiplici acino, f. non spinosa hortensis rubra. *Bauh. pin.* 455.2. *Grossularia hortensis*, fructu margaritis simili. *Bauh. pin.* 455.*Habitat in Sueciæ borealibus.* ‡

2. RI-

2. RIBES inerme, racemis erectis, bracteis flore longioribus. *alpinum*.  
Ribes inerme, floribus planiusculis, racemis erectis.  
*Hort. cliff. 82. Fl. suec. 198, 206.*  
Ribes alpinum dulce. *Bauh. hist. 2. p. 93. Fl. lapp. 97.*  
Grossularia vulgaris, fructu dulci. *Bauh. pin. 455.*  
Grossularia distinctis baccis. *Bauh. pin. 455.*
3. Ribes montana, oxyacanthæ sapore. *Bauh. prodr. 160.*  
*Amæn acad. 1. p. 170.*  
*Habitat in Sueciæ, Helvetiæ, Angliæ sepibus ficcis. §*
3. RIBES inerme, racemis pilosis, floribus oblongis. *nigrum*.  
Ribes inerme, floribus oblongis. *Hort. cliff. 269. Hort. upf. 51. Fl. suec. 196, 207. Mat. med. 100. Roy. lugdb. 269. Dalib. parif. 74.*  
Ribes vulgaris, fructu nigro. *Rudb. vall. 32. Fl. lapp. 99.*  
Grossularia non spinosa, fructu nigro. *Bauh. pin. 455.*
3. Ribesium nigrum pennsylvanicum, floribus oblongis.  
*Dill. elth. 324. t. 244. f. 315.*  
*Habitat in Suecia, Helvetia, Pennsylvania. §*  
\* \* Grossulariæ aculeatæ.
4. RIBES ramis subaculeatis reclinatis, pedunculi bractea triphylla. *reclinatum*.  
Ribes ramis subaculeatis reclinatis. *Hort. cliff. 82. Hort. upf. 51. Roy. lugdb. 270.*  
Grossularia spinosa sativa altera, foliis latioribus. *Bauh. pin. 455.*  
*Habitat in Germania, Helvetia. §*
5. RIBES ramis aculeatis, petiolorum ciliis pilosis, baccis hirsutis. *Grossularia*.  
Ribes ramis aculeatis, racemis erectis, baccis hirsutis.  
*Hort. cliff. 82.*  
Ribes ramis aculeatis erectis, fructu hispido. *Vir. cliff. 21. Roy. lugdb. 269.*  
Grossularia, fructu maximo hispido margaritarum sere colore. *Raj. hist. 1484.*  
*Habitat in Europa. §*
6. RIBES ramis undique aculeatis. *Hort. upf. 51.* *oxyacanthoides*.  
Grossularia, oxyacanthæ foliis amplioribus, e sinu Hudsonis. *Pluk. amalth. 212. Dill. elth. 166. t. 139. f. 116.*  
*Habitat in Canada. §*

- Uva crispa. 7. **RIBES** ramis aculeatis, baccis glabris, pedicellis bractea monophylla.  
*Ribes* ramis aculeatis erectis, fructu glabro. *Hort. cliff.* 82. *Fl. svec.* 195, 208. *Roy. lugdb.* 273. *Hall. helv.* 346. *Dalib. parif.* 74.  
*Grossularia* simplici acino vel spinosa sylvestris. *Bauh. pin.* 455.  
*Habitat in Europa boreali.* §
- cynosbati. 8. **RIBES** aculeis subaxillaribus, baccis aculeatis racemosis.  
*Habitat in Canada.* *Kalm.* §  
*Facies præcedentium.* Folia parum fissâ. Aculeus infra spinæ sub alis. Pedunculi sæpius triflori. Germen in flore hispidum, nec aculeatum. Corolla *R. Grossulariæ*. Baccæ magnitudine nucis coryli, undique armatæ aculeis robustis.

## GRONOVIA.

- scandens. 1. **GRONOVIA.** *Hort. cliff.* 74.  
*Gronovia* scandens lappacea, pampinea fronde. *Houfl. Mart. cent.* 1. p. 40. t. 40. *Amm. herb.* 346.  
*Habitat in Vera Cruce.*

## HEDERA.

- Helix. 1. **HEDERA** foliis ovatis lobatisque. *Fl. lapp.* 91. *Fl. svec.* 190, 209. *Hort. cliff.* 74. *Mat. med.* 98. *Roy. lugdb.* 223. *Hall. helv.* 164.  
*Hedera* arborea. *Bauh. pin.* 305.  
 β. *Hedera* poetica. *Bauh. pin.* 305.  
 γ. *Hedera* major sterilis. *Bauh. pin.* 305.  
 δ. *Hedera* humi repens. *Bauh. pin.* 305.  
*Habitat in Europæ arboribus putrescentibus, inque sepibus.* §
- quinquefo- 2. **HEDERA** foliis quinatis ovatis ferratis. *Hort. cliff.* 74. *Roy. lugdb.* 223. *Gron. virg.* 24.  
 lia. *Vitis* hederacea indica. *Stapel. theatr.* 364.  
*Edera* quinquefolia canadensis. *Corn. canad.* 99. t. 100.  
*Helix. Mich. gen.* 30.  
*Habitat in Canada.* §  
*Capsula* utrinque turbinata, bilocularis, polysperma.  
*Flores* masculi a femineis distincti. *D. Mitchel.*

## VITIS.

1. VITIS foliis lobatis sinuatis nudis. vinifera.  
*Vitis foliis palmato - angulatis. Hort. cliff. 74. Hort. upl. 50. Mat. med. 97. Gron. virg. 144. Roy. lugdb. 222.*  
*Vitis vinifera. Bauh. pin. 299.*
- β. Vitis corinthiaca f. apyrena. *Bauh. hist. 2. p. 72. apyrena.*  
*Habitat in Orbis quatuor partibus temperatis. §*  
*Cirrhus oppositifolius racemifero - floriferus.*
2. VITIS foliis cordatis dentatis subtus villosis, cirrhis indica. indica.  
*racemiferis. Fl. zeyl. 99. \**  
*Vitis sylvestris indica, acinis rotundis. Raj. dendr. 67.*  
*Scembra Valli. Rheed. mal. 7. p. t. 6.*  
*Habitat in India. §*
3. VITIS foliis cordatis subtrilobis dentatis subtus to- Labrusca.  
*mentosis.*  
*Vitis hederæ folio ferrato. Plum. spec. 18. ic. 259. f. 1.*  
*Vitis sylvestris virginiana. Bauh. pin. 299.*  
*Vitis vinifera sylvestris americana, foliis averſa parte denſa lanugine teſtis. Pluk. phyt. 249. f. 1.*  
*Vitis fructu minore rubro acerbo, folio subrotundo minus laciniato, subtus alba lanugine teſto. Sloan. hiſt. 2. p. 104. t. 210. f. 4.*  
*Habitat in America ſeptentrionali. §*
4. VITIS foliis cordatis dentato-ferratis utrinque nudis. vulpina.  
*Vitis vulpina dicta virginiana nigra, Pluk. alm. 392.*  
*Vitis aceris folio. Raj. dendr. 68.*  
*Habitat in Virginia. §*
5. VITIS foliis ternatis : foliolis subrotundis ferratis. trifolia.  
*Vitis pearme doorica, foliis ternis subrotundis ferratis. Raj. dendr. 68.*  
*Folium Cauffmanis. Rumph. amb. 5. p. 450. t. 166. f. 2.*  
*Habitat in India. §*  
*Flores non vidi, hinc dubius utrum Vitis f. Cissi generis.*
6. VITIS foliis quinatis : foliolis multifidis. *Hort. cliff. laciniola.*  
*74. Roy. lugdb. 223.*  
*Vitis laciniatis foliis. Corn. canad. 182. t. 183.*  
*Vitis apii folio. Bauh. hiſt. 2. p. 73.*  
*Habitat - - - §*

arbores.

7. *VITIS* foliis supradecompositis : foliolis lateralibus pinnatis.

*Vitis caroliniana*, foliis apii, uva corymbosa purpurascens. *Act. bonon.* 2. part. 2. p. 365. t. 24.

Frutex scandens, petroselinifolius, virginianus claviculis donatus. *Pluk. mant.* 85. t. 412. f. 2.

*Habitat in Carolina, Virginia.* ♀

## LAGOECIA.

Cuminoides.

1. *LAGOECIA*. *Hort. cliff.* 73. *Hort. upf.* 52.

*Cuminum sylvestre*, capitulis globosis. *Bauh. pin.* 146.

*Cuminum sylvestre*. *Cam. epit.* 518. *Gesn. epist.* 50.

*Habitat in Creta, Lemno, Lybia, Galatia.* ☉

## SAUVAGESIA.

erecta.

1. *SAUVAGESIA*.

*Ireon herbaceus minor*, foliis oblongis tenuissime crenatis, stipulis, ciliatis, floribus singularibus ad alas. *Brown. jam.* 179. t. 12. f. 3.

*Habitat in Domingo, Jamaica, Surinamo.* ☉

*Habitus Hyperici aut Corchori.* Caulis erectus. Folia alterna, ovato-lanceolata, obtuse serrata, Stipulae acutae, ciliatae.

## CLAYTONIA.

virginica.

1. *CLAYTONIA* foliis linearibus.

*Claytonia*. *Gron. virg.* 25.

*Ornithogalo affinis virginiana*, flore purpureo pentapetaloide. *Pluk. alm.* 272. t. 102. f. 3. *Rudb. elyf.* 2. p. 129. f. 6.

*Habitat in Virginia.* 24

sibirica.

2. *CLAYTONIA* foliis ovatis. *Hort. upf.* 52. \*

*Limnia*. *Act. flokh.* 1746. p. 120. t. 5. \*

*Habitat in Sibiria* 24

## ACHYRANTHES.

sanguinolenta.

1. *ACHYRANTHES* caule erecto simplicissimo, spicis brachiatis laevibus.

*Verbena rubra*. *Rumph. amb.* 7. p. 60. t. 27. f. 2.

*Habitat in India.* 21

Caulis humanae altitudinis, simplicissimus. Folia opposita, ovata, integra, petiolata, subtus sanguinolenta.

- 1a. Spicæ terminales, brachiatae, laeves. Flores parvi, Stamina 5 alterna castrata.
2. ACHYRANTHES caule patulo, floribus spicatis remotis ovatis, calycibus squarroso-patulis, foliis alternis. *muricata.*  
*Blitum frutescens. Rumph. amb. 5. p. 235 t. 83. f. 2. Habitat in India. §*  
*Erutex Caule patulo. Folia alterna, petiolata, ovata, integra, nudiuscula. Spicæ alternae, pedunculatae, longae, A. asperae, at Flores remotiores, magis ovati, non reflectuntur, sed calycis squamæ tres breviores patulae.*
3. ACHYRANTHES caule erecto, calycibus reflexis aspera. spicæ adpressis. *Fl. zeyl. 105.*  
*Achyranthes caule erecto. Hort. cliff. 42. Roy. lugdb. 418.*
- α. Amaranthus ficulus spicatus. *Bocc. sic. 16. t. 9. Pluk. ficula. phyt. 260. f. 2.*
- β. Amaranthus spicatus zeylanicus, foliis obtusis amarantho ficulo bocconi similis. *Burm. zeyl. 16. t. 5. f. 3. indica.*  
*Amaranthus spicatus, dictamni cretici folio, maderaspatis. Pluk. alm. 26. t. 10. f. 4. Mill. dict. t. 11. f. 2.*  
*Habitat in Sicilia, Zeylona, Jamaica. §*  
*Indica β. caule ♂ fructificatione simillima, sed folia obovata obtusa, crassa, undulata quintuplo majora, hybridam indicant.*
4. ACHYRANTHES caule patulo, spicis interruptis; flosculis ternis: lateralibus utrinque fasciculo setarum uncinato. *lappacea.*  
*Achyranthes caule erecto, spica interrupta, floribus externe lanatis. Sp. pl. 1. p. 204.*  
*Achyranthes floribus externe lanatis interrupte spicatis. Fl. zeyl. 103.*  
*Blitum scandens, fructu lappaceo. Burm. zeyl. 47. t. 18. f. 1.*  
*Centaurium ciliare minus, circæ foliis firmioribus, spicatis floribus. Pluk. alm. 93. t. 82. f. 2.*  
*Wellia Codivelli. Rheed. mal. 10. p. t. 59.*  
*Habitat in India.*  
*Spicæ longæ, solitariae, interruptæ: ad singulos dentes pedunculi Flores 3 transverse positi, ad latera florum*



rum lateralium utrinque penicillus purpureus, e setis rigidis uncinatis, sensim se pandentibus in pilulam lappaceam, maturo fructu discedentem & adherentem.

prostrata.

5. *ACHYRANTHES* caulibus prostratis, spicis oblongis: flosculis binatis utrinque fasciculo setarum unginato.

Auris canina femina. *Rumph. amb. 6. p. 26. t. 11.*

*Verbena indica. Bont. jav. 150. t. 150.*

*Habitat in India. 21.*

Caules prostrati, sæpe repentes, sesquipediales, divaricati, bipediales, atropurpurei, pilosi. Folia opposita, obovata, subpetiolata, subpilosa, acuta, margine purpurascens. Spicæ e dichotomia, pedunculatæ, foliariæ, oblongæ, virides, apice purpurascens. Flores minuti, subpedicellati, gemini, calyce communi triphylo mucronato; ad latera alterius flosculi penicillus purpureus setis uncinatis. Fructus reflexi, ovati, obtusi.

lanata.

6. *ACHYRANTHES* caule prostrato, spicis ovatis lateralibus, calycibus tomentosis. *Mill. dict. t. 12. f. 1.*

*Chenopodium incanum* r. cemosum, folio majore minori opposito. *Burm. zeyl. 60. t. 26. f. 1.*

*Amaranthus indicus verticillatus albus*, foliis lanugine incanus. *Pluk. alm. 27. t. 57. f. 8.*

*Scherubala. Rheed. mal. 10. p. 75. t. 29.*

*Habitat in India.*

corymbosa.

7. *ACHYRANTHES* foliis oppositis linearibus, panicula dichotoma. *Fl. zeyl. 100.*

*Paronychia* foliis ad genicula confertis acutis, floribus umbellatis. *Burm. zeyl. 184. t. 65. f. 2.*

*Amaranthus* folio tenuissimo incano mollugini æmulante. *Bocc. mus. 44. t. 39.*

*Elichryso* affinis herba indica polygonoides ramosior, foliis caulium nodos ambientibus. *Pluk. alm. 134. t. 86. f. 6.*

*Habitat in Zeylona.*

## C E L O S I A.

argentea.

1. *CELOSIA* foliis lanceolatis, stipulis subsulcatis, pedunculis angulatis, spicis scariosis.

Ce-

*Celosia* foliis lineari-lanceolatis. *Hort. cliff.* 43. \* *Roy. lugdb.* 419.

*Amaranthus* spica albescente habitiore. *Mart. cent.* 7. t. 7.

*Habitat in China.* ☉

2. *CELOSIA* foliis ovatis, stipulis falcatis, pedunculis angulatis, spicis scariosis. margaritacea

*Amaranthus* simplici panicula. *Bauh. pin.* 121.

*Amaranthus* spica albescente habitiore. *Mart. cent.* 7. t. 7.

*Habitat in America.* ☉

*Simillima C. argentea*, ut fere varietas, sed Folia fere ovata, stamina purpurea.

3. *CELOSIA* foliis lanceolato-ovatis recurvis subundatis, pedunculis angulatis, spicis oblongis cristatis. cristata.

*Celosia* foliis oblongo-ovatis, pedunculis teretibus substriatis, spicis oblongis. *Sp. pl.* 1. p. 205.

*Celosia* foliis lanceolato-ovatis. *Hort. cliff.* 43. *Hort. upf.* 52.

*Amaranthus* panicula conglomerata. *Bauh. pin.* 121.

- β. *Amaranthus* major, paniculis surrectis flavescentibus. *Herm. lugdb.* 30.

*Habitat in Asia.* ☉

4. *CELOSIA* foliis lanceolato-ovatis lineatis acuminatissimis, stipulis falcatis, spicis cristatis. cristatis.

*Amaranthus* cristatus. *Cam. epit.* 792.

*Amaranthus* minor. spica singulari, lunatis circum caulem foliis. *Barr. rar.* 471. t. 1195. *Bocc. mus.* 2. p. 77. t. 66.

*Amaranthus* vulgaris. *Rumph. amb.* 5. p. 236. t. 84.

*Habitat in India.* ☉

Rami laterales ex singulis alis. Folia subtus venis rubris striata: inferiora saepe terna. Stipulae dimidiatae, falcatae.

5. *CELOSIA* foliis ovatis strictis inauriculatis, caule coccineofulcato, spicis multiplicibus cristatis.

*Amaranthus* panicula incurva. *Bauh. pin.* 121.

*Amaranthus* panicula speciosa cristata. *Bauh. hist.* 2. p. 969.

*Habitat in India.*

*Differt ab C. cristata* foliis triplo crassioribus, fragilibus.

Floribus penitus purpureis, nec rubicundis carina purpurea. Stamina corolla brevioribus, nec longioribus.

paniculata. 6. CELOSIA foliis oblongo-ovatis, caule assurgente, spicis terminalibus alternis remotis.

Celosia major farmentosa assurgens, foliis majoribus ovatis. *Brown. jam. 179.*

Blitum album majus scandens. *Sloan. jam. 49. hist. 1. p. 142. t. 91. f. 2.*

Habitat in Jamaica.

Caulis prostratus, teres, ramosus. Folia alterna, ovata, acuminata. Rami terminati panicula e spicis alternis, saepe ramosis. Flores alterni, distincti. Stamina corolla breviora. Stigma trifidum.

lanata. 7. CELOSIA foliis lanceolatis tomentosis obtusis, spicis confertis, staminibus lanatis. *Fl. zeyl. 102. \**

Habitat in Zeylona.

nodiflora. 8. CELOSIA foliis cuneiformibus acutiusculis, spicis globosis lateralibus. *Fl. zeyl. 101. \**

Amaranthoides indicum nodiflorum, capitulis ex albo virescentibus. *Burm. zeyl. 16. t. 5. f. 2.*

Amaranthoides indicum globosum, ad caulium nodos floridum, foliis solitariis summo apice barbatis.

*Pluk. alm. 27. t. 133. f. 2.*

Habitat in Zeylona.

## ILLECEBRUM.

\* *Caulibus prostratis.*

verticillatum. 1. ILLECEBRUM floribus verticillatis nudis, caulibus procumbentibus.

Illecebrum caulibus procumbentibus. *Roy. lugdb. 214. Dalib. parif. 76, 126.*

Illecebrum. *Hort. cliff. 492.*

Paronychia serpillifolia palustris. *Vaill. parif. 157. t. 15. f. 1.*

Polygala repens nivea. *Bauh. pin. 215.*

Habitat in Europæ pascuis udis.

suffruticosum. 2. ILLECEBRUM floribus lateralibus solitariis, caulibus suffruticosis.

Illecebrum caule erecto. *Roy. lugdb. 214.*

*Polygonum herniariae* foliis. *Lob. adv. 404.<sup>p</sup> Dalech. hist. 1124.<sup>p</sup>*

*Paronychia hispanica fruticosa, myrti folio. Tournef. inst. 508.*

*Habitat in Hispania.*

3. **ILLECEBRUM** spicis cymosis secundis, caule dif- cymosum.  
fuso, foliis quaternis.

*Polygonum capitula inter genicula echinata. Bocc. sic.*

*41. t. 2. f. 3. Raj. bist. 214.*

*Polygoni hisp nci genus Clusii. Dalech. hist. 1125?*

*Habitat Montpelii. D. Sauvages. Lusitania, Itva. ☉*

*Planta facie S di aut Polygoni; Caulis filiformis, brachiatus. Folia subquaternata, linearia, crassiuscula. Flores digesti in spicas secundas, Spicae in cymastam laterales quam terminales. Calyces pentaphylli, colorati, apice fornicati aristati; in viva Generis certitudinem inquirant autoptae.*

4. **ILLECEBRUM** floribus bracteis nitidis obvallatis, Paronychia.  
caulibus procumbentibus.

*Herniaria squamis nitidis flores superantibus. Hort. cliff. 41. Hort. upf. 54. Roy. lugdb. 215. Sauv. monsp. 129.*

*Paronychia hispanica. Clus. hist. 2. p. 183.*

*Polygonum minus candicans. Bauh. pin. 281.*

*Habitat in Hispania & Narbona. 24*

5. **ILLECEBRUM** floribus bracteis nitidis occultant- capitulum.  
tibus capitula terminalia, caulibus erectis, foliis ciliatis.

*Herniaria erecta, squamis nitidis flores occultantibus.*

*Sauv. monsp. 129.*

*Paronychia narbonensis erecta. Tournef. inst. 508.*

*Polygonum minus candicans, capitulis surrectis. Magn. monsp. 209.*

*Polygonum montanum niveum minimum. Lob. it. 420.*

*Habitat in G. Narbonensi, Hispania, Oriente.*

*Folia ciliata, subius villosa.*

\* \* *Caulibus repentibus.*

6. **ILLECEBRUM** caulibus repentibus pilosis, foliis Achyrantha.  
ovatis mucronatis opposito minore, capitulis sub-  
globosis subspinosis.

*Achyranthes repens*, caule repente, capitulis laterali-  
bus sessilibus. *Sp. pl.* 1. p. 205.

*Achyrantha repens*, foliis bliti pallidi. *Dill. elth.* 8.  
t. 7 f. 7.

*Habitat in Turcomannia.* ☉

polygono-  
idea.

7. **ILLECEBRUM** caulibus repentibus hirtis, foliis  
lato-lanceolatis petiolatis, capitulis orbiculatis nudis,  
*Gomphrena polygonoides*. *Sp. pl.* 1. p. 225.

*Herniaria hirsuta repens* ad nodos alternos florida.  
*Brown. jam.* 184.

*Amaranthoides humile curassavicum*, foliis polygoni.  
*Herm. prodr.* 17. t. 17. *Sloan. jam.* 48. *hist.* 1. p.  
141. t. 86. f. 2. *Raj. suppl.* 126.

- β. *Amaranthoides marina hirsuta*, halimi folio. *Plum.*  
*ic.* 94. t. 21. f. 2.

*Habitat in Americæ littoribus maris.*

*Simillima sequenti, sed caule, non flore villoso.*

ficoidea.

8. **ILLECEBRUM** caulibus repentibus glabris, foliis  
lato-lanceolatis petiolatis, capitulis orbiculatis pube-  
scentibus.

*Gomphrena ficoidea*. *Syst. nat.* 1. p. 225.

*Gomphrena* caule diffuso repente, capitulis sessilibus.  
*Roy. lugdb.* 418.

*Amaranthoides marina repens*, polygoni folio, capi-  
tulis argenteis. *Plum. spec.* 20.

*Habitat in marinis littoribus Americæ, nunc Hispa-  
niæ.* 24

*Corollæ acutis oculis pubescentes sunt.*

sessile.

9. **ILLECEBRUM** caulibus repentibus glabris, foliis  
lanceolatis subsessilibus, capitulis oblongis glabris.  
*Gomphrena sessilis*. *Sp. pl.* 1. p. 225. *Fl. zeyl.* 116.

*Amaranthoides humile maderaspatanum*, capitulis can-  
dificantibus, folio molli. *Pluk. alm.* 27. t. 133. f. 1.

*Amaranthus humilis*, foliis oppositis, flosculis ex alis  
glomeratis. *Burm. zeyl.* 17. t. 4. f. 2.

*Coluppa. Rheed. mal.* 10. p. 21. t. 9.

*Habitat in India Orientali.*

*Folia angustiora quam in I. ficoideo; petala glabra.*

vermicula-  
tum.

10. **ILLECEBRUM** caulibus repentibus glabris, foliis  
subteretibus carnosissimis, capitulis oblongis glabris ter-  
minalibus.

*Gomphrena vermicularis*. *Sp. pl.* 224.

*Caraxeron humile*, cepeæ foliis, capitulis albis. *Vaill. art.* 1722. p. 264.

*Gomphrena repens rufescens*, foliis linearibus crassiusculis. *Brown. jam.* 184.

*Amaranthoides humile curassavicum*, cepeæ foliis lucidis, capitulis albis. *Herm. par.* 15. t. 15. *Pluk. alm.* 27. t. 75. f. 9.

*Amarantho affinis aizoides* f. *Amaranthoides minor americana procumbens*, sedi teretifolii foliis & facie, flore oblongo niveo. *Breyn. prodr.* 2.

*Trifolii spica Crithmum maritimum non spinosum brasiliense*. *Raj. hist.* 1331.

*Habitat in Brasilia, Curassao.*

## GLAUX.

1. *GLAUX. Hort. cliff.* 43. *Fl. succ.* 199, 210. *Roy. maritima. lugdb.* 417.

*Glaux foliis elliptico-oblongis. Fl. lapp.* 72.

*Glaux maritima. Bauh. pin.* 215.

*Alfene bifolia*, fructu coriandri, radice geniculata.

*Læf. pruss.* 13. t. 3.

*Habitat in Europæ maritimis, salsis.* 24

## THESIUM.

1. *THESIUM panicula foliata*, foliis lineari-lanceolatis. *Linophyllon.*

*Thesium. Hort. cliff.* 41. *Hall. helv.* 183. *Dalib. par.* 75.

*Thesium floribus sparsis. Sauv. monsp.* 48.

*Linaria montana*, flosculis albicantibus. *Bauh. pin.* 213.

*Sesamoides procumbens montanum*, linariæ folio, floribus albicantibus. *Morif. hist.* 3. p. 601. f. 15. t. 1. f. 3.

*Anonymos lini folio. Clus. hist.* 1. p. 324.

β. *Onobrychis IV lutea. Bauh. pin.* 215. *Dalech. hist.* 491.

*Habitat in Europæ siccis montosis, Cretaceis & in Oriente.* ○

2. *THESIUM racemo foliato*, foliis linearibus.

*alpinum,*

*Linophyllum collinum lignosum minus, flore albicante.*

*Pont. anth.* 261. *Segu. ver.* 3. p. 90.

*Alchimilla alpina*, linariæ foliis, calyce florum albo in foliorum alis. *Till. pis.* 6.

Al-

*Alchimilla linearis* folio, floribus & vasculis in foliorum alis sessilibus *Shaw. spec. 14.*

*Habitat in Italix alpidibus, in Sveciæ collibus.* ☉

Differt a precedenti Caulibus non ramosis; Foliis linearibus, nec lanceolatis; confertis, nec remotis. Racemo pedicellis unifloris brevissimis, folio unico terminali, duobusque lateralibus minoribus, nec paniculato floribus ternis quinisve. Floribus sæpius quadrifidis, nec quinquefidis.

capitatum. 3. **THESIUM** capitulis sessilibus terminalibus, foliis triquetris lævibus.

*Thesium floribus capitatis. Roy. lugdb. 114.*

*Habitat in Æthiopia.* ☿

Caules duri: Ramis alternis, remotis, superioribus sensim longioribus. Folia alterna, parva, subulata, mucronata. Capitula terminalia, sub quibus ramus excrefcit. Florum laciniae valde acuminatae & interne secundum totam longitudinem villosae.

scabrum. 4. **THESIUM** capitulis pedunculatis, foliis triquetris margine scaberrimis.

*Habitat ad Cap. b. spei.* ☿

Simillima *T. capitato*, sed Capitula longo pedunculo nudo; Flores obtusiusculi, tantum ad apices interne villosi: Folia tripliei angulo cartilagineis dentibus exasperata.

funale. 5. **THESIUM** floribus spicatis, corollis ciliatis, caule suffruticoso, foliis subulatis brevissimis.

*Habitat ad Cap. b. spei.* ☿

Fruticulus facie *Schæni capensis*, subnudus: Ramis numerosis, alternis, confertis, virgatis, simplicibus, teretibus, filiformibus. Folia subulata, alterna, remota, minutissima. Spicae terminales, 2. f. 3., alternæ.

umbellatum 6. **THESIUM** floribus umbellatis, foliis oblongis. *Centaurium luteum ascyroides virginianum. Pluk. mant. 43. t. 342. f. 1.*

*Habitat in Virginiæ, Pennsylvaniae pascuis siccis. Kalm. 24.*

Radix perennis. Folia alterna, ovali-lanceolata, integerrima. Rami alterni, in summitate caulis: Umbel-

*bellula racemosa, terminales : involucri tetraphyllo ,  
parvo : Flosculi 5. f. 3.*

## RAUVOLFIA.

1. RAUVOLFIA glaberrima nitidissima. *Jacq. amer. 14. nitida,*  
*Rauvolfia. Hort. cliff. 75. t. 9.*  
*Rauvolfia tetraphylla angustifolia. Plum. gen. 19. ic.*  
*236. f. 1.*  
*Habitat in America meridionali. †*
2. RAUVOLFIA subpubescens. canescens.  
*Rauvolfia fruticosa, foliis verticillatis tenuissime vil-*  
*losis. Brown. jam. 180.*  
*Rauvolfia tetraphylla latifolia. Plum. gen. 19. ic. 236. f. 2.*  
*Solan. fructu fruticosa, foliis laurinis oblongis inte-*  
*gris subtus hirsutis. Sloan. jam. 173. hist. 2. p. 107.*  
*t. 211. f. 1. Raj. dendr. 75.*  
*Arbor Sycophora jamaicensis, foliis minoribus. Pluk.*  
*phyt. 266. f. 2.*  
*Habitat in Jamaica. †*  
*Rami subtomentosi. Folia subtus potius tomentosa, quam*  
*hirsuta.*

## CERBERA.

1. CERBERA foliis ovatis. Ahouaij.  
*Thevetia. Hort. cliff. 75. Roy. lugdh. 413.*  
*Ahouai. Thev. antarct. 66. Tournef. inst. 658. Raj.*  
*hist. 16. 6.*  
*Ahouai major. Pif. bras. 49.*  
*Arbor americana, foliis pomi, fructu triangulo. Bauh.*  
*pin. 434.*  
*Habitat in Brasilia. †*
2. CERBERA foliis lanceolatis : nervis transversalibus. Manghas.  
*Fl. zeyl. 106. \* Osb. it. 91.*  
*Manghas lactescens, foliis nerii crassius venosis, jasmī*  
*ni flore, fructu persici simili venenato. Burm. zeyl.*  
*150. t. 70. f. 1.*  
*Manghas fructu venenato. Bauh. pin. 440. Raj. hist.*  
*1552.*  
*Arbor lactaria. Rumph. amb. 2. p. 243. t. 81.*  
*Odollam. Rheed. mal. 1. p. 71. t. 39.*  
*Habitat in Indiis ad aquas. †*  
*Arbor lactescens. Folia ueterna, sparsa ad apices rami-*  
*rum.*



rum, latè-lanceolata, petiolata, glabra, integerrima, pedalia. Racemi terminales, ramosi, inæquales. Perianthium pentaphyllum foliis lanceolatis, patentibus, coloratis, deciduis. Corolla alba, infundibuliformis: Tubo calyce longiore, interne angustato: Limbo quinquepartito, lobis ovatis, tubo brevioribus. Stamina Filamenta 5, brevissima, tubi superiori parti inserta. Anthera ovata, tetragona, testis lanugine tubi. Pist. Germen bifidum. Stylus filiformis, tubo fere brevior. Stigma ovatum, fissum. Pericarpium Drupa ovata, magnitudine ovi anseris; viride punctis albis minutissimis, uno latere compressum futuraque obsoleta; Semina 2, veluti duas castaneas magnas includens, venenatum, vomitorium. Osbeck.

Theretha.

2. CERBERA foliis linearibus longissimis confertis.

Jacq. hist. t. 34.

Ahouai nerii folio, flore luteo. Plum. spec. 20. ic. 18.

Nerio affinis angustifolia lactescens, flore luteo. Pluk.

alm. 253. t. 207. f. 3.

Habitat in America calidiore. b

## V I N C A.

minor.

1. VINCA caulibus procumbentibus, foliis lanceolatis-ovatis, floribus pedunculatis.

Vinca foliis ovatis. Hort. cliff. 77. Hort. upf. 53. Mat. med. 101. Roy. lugdb. 412. Dalib. parif. 75.

Clematis Daphnoides minor. Bauh. pin. 301.

Clematis Daphnoides. Dod. pempt. 405.

Habitat in Germania, Anglia, Gallia. b

Caules procumbentes, Folia obtusiora margine lævia; Pedunculi curvati; Calyces breves.

major.

2. VINCA caulibus erectis, foliis ovatis, floribus pedunculatis.

Clematis Daphnoides major. Bauh. pin. 302. Dod. pempt. 406.

Habitat in Gallia Narbonensi, Hispania. b

Affinis nimis præcedenti, sed Caules erecti; Folia acutiora, margine scabra; pedunculi recti; Calyces longitudine tubi.

lutea.

2. VINCA caule volubili foliis oblongis. Amæn. acad. 4. p. 307.

**Apocynum scandens**, salicis folio, flore amplo plano.

*Catesb. car. 2. p. 53. t. 53.*

*Habitat in Carolina.* ♀

4. **VINCA** caule suffrutescente erecto, floribus geminis sessilibus, foliis oblongis petiolis basi bidentatis. rosea.

**Vinca** foliis oblongo-ovatis integerrimis, tubo floris longissimo, caule ramofo fruticoso. *Mill. dict. t. 186.*

*Habitat in Madagascar, Java.* ♀

**Caulis** rigidulus, teres, levis. **Folia** opposita, lanceolato-ovalia, coriacea rachi alba: **Petiolis** utrinque basi denticulo. **Flores** axillares foliorum, sessiles, gemini. **Calyx** in setas 5, subulatas. **Corolla** hypocrateriformis, rosea, pulcherrima, plana: **Lacinii** semisalcatis, obtusis, mucronatis: umbilico atro-purpureo, obtuso, villoso; **Tubus** longiusculus, clavatus, clava 5 tuberculis.

## GARDENIA.

1. **GARDENIA**. *Ellis act. angl. vol. 51. p. 392. t. 23. Florida.*

**Jasminum** foliis lanceolatis oppositis integerrimis, calycibus acutioribus. *Mill. ic. t. 180.*

*Cotsjopiri. Rumph. amb. 7. p. 26. t. 14. f. 2.*

*Habitat in India orientali: Suratte, Amboina, Cap. b. spei.* ♀

*Misi figuram Wachendorf, iconem Ellis, vivam plantam Gorden, at flore pleno; unde fructus etiamnum ignotus, diversa certe est a Nandi Ervalam Rheed. mal. 2. t. 54. Et Rumph. amb. 4. t. 39. uti ex calyce constat; propius accedere videtur ad Jasminum indicum, mali aurantie foliis, flore albo pleno amplissimo. Breyn. prodr. 3. p. 73. Till. pis. 87. t. 30, nisi flores pedunculatos pingeret; qui in Gardenia sessiles sunt.*

## NERIUM.

1. **NERIUM** foliis lineari-lanceolatis ternis. *Hort. cliff. Olandet.*

*76. Hort. upf. 53. Fl. zeyl. 108. Roy. lugdb. 412.*

**Nerium** floribus rubescentibus. *Bauh. pin. 464. Burm. zeyl. t. 77.*

**Rhododendrum.** *Dod. pempt. 851.*

*Areli. Rheed. mal. 9. p. t. t. 1, 2.*

*Linnei Spec. plant. pars I.*

V

Ha-

*Habitat a Gades in Indiam orientalem locis subhumidis. §*

- divaricatum. 2. **NERIUM** foliis lanceolato-ovatis, ramis divaricatis.  
*Fl. zeyl.* 109. \*  
 Apocynum zeylanicum indicum frutescens, nerii flore candidissimo. *Herm. par.* 40.  
*Habitat in India. §*
- antidysentericum. 3. **NERIUM** foliis ovatis acuminatis petiolatis. *Fl. zeyl.* 107. *Mat. med.* 102.  
 Codaga pala. *Kheed. mal.* 1. p. 85. t. 47.  
*Habitat in India. §*
- zeylanicum. 4. **NERIUM** foliis lanceolatis oppositis, ramis rectis.  
*Amzn. acad.* 4. p. 309. \*  
 Apocynum arborecens, nerii flore, minus. *Burm. zeyl.* 23. t. 12. f. 2.  
*Habitat in India. §*

## PLUMERIA.

- rubra. 1. **PLUMERIA** foliis ovato-oblongis, petiolis biglandulosis. *Hort. cliff.* 76. *Roy. lugdb.* 412.  
 Plumeria arborecens, ramulis crassis, foliis oblongo-ovatis, petiolis biglandulosis, floribus geminatis per spicas terminales. *Brown. jam.* 181.  
 Plumeria flore roseo odoratissimo. *Tournef. inst.* 659.  
*Catesb. car.* 2. p. 92. t. 92. *Ehret. pict.* t. 10. *Trew. Ehret.* t. 41.  
 Nerium arboreum, folio maximo obtusifloro, flore incarnato. *Sloan. jam.* 154. *hist.* 2. p. 61. t. 185. f. 1. *§* t. 186. f. 1.  
 Jasminum indicum. *Mer. surin.* 8. t. 8.  
 Clematis arborea americana, laurinis amplissimis foliis, flore laccæ colore odoratissimo. *Pluk. mant.* 50. *alm.* 109. t. 207. f. 2.  
*Habitat in Jamaica, Surinamo. §*
- alba. 2. **PLUMERIA** foliis lanceolatis revolutis, pedunculis superne tuberosis.  
 Plumeria flore niveo, foliis longis angustis & acuminatis. *Plum. spec.* 20. *ic.* 231.  
 Apocynum americanum frutescens, longissimo folio, flore albo odoratissimo. *Comm. hort.* 2. p. 47. t. 24.  
 Ne-

*Nerium arboreum altissimum, folio angusto, flore albo.*

*Sloan. jam. 154. hist. 2. p. 62. Raj. dendr. 114.*

*Herm. parad. 49.*

*Habitat in Jamaica.*

*Folia lateribus revoluta. Flores spicati.*

3. **PLUMERIA** foliis lanceolatis petiolatis obtusis. obtusa.

*Plumieria flore niveo, foliis brevioribus obtusis. T.*

*Catesb. car. 2. p. 93. t. 93. Plum. ic. 232.*

*Flos convolutus. Rumph. amb. 4. p. 85. t. 387.*

*Habitat in America calidiore. ♀*

*Folia ovali-lanceolata. Pedunculi ramosi.*

## ECHITES.

1. **ECHITES** pedunculis bifloris. *Jacq. amer. 13. biflora*  
*hist. t. 21.*

*Apocynum scandens, flore neri albo. Plum. amer.*

*82. t. 96.*

*Habitat in America. ♀*

2. **ECHITES** pedunculis racemosis, foliis obovatis acuminatis. *Jacq. amer. 13. hist. t. 25.* quinquangularis.

*Habitat in America meridionali. ♀*

3. **ECHITES** pedunculis racemosis, foliis subovatis obtusis mucronatis. *Jacq. amer. 13. hist. t. 26.* suberecta.

*Apocynum erectum fruticolum, flore luteo maximo & speciosissimo. Sloan. jam. 89. hist. 1. p. 206. t. 130. f. 2.*

*Habitat in Jamaica. ♀*

4. **ECHITES** pedunculis subracemosis, foliis lanceolatis acuminatis. *Jacq. amer. 13. hist. t. 27.* torulosa.

*Nerium sarmentosum scandens, ramulis tenuibus, folliculis gracilibus torosis. Brown. jam. 181. t. 16. f. 2.*

*Habitat in Jamaica. ♀*

*Folliculi filiformes, torulosi, longissimi.*

5. **ECHITES** pedunculis umbellatis, foliis ovatis obtusis mucronatis, caule volubili. *Jacq. hist. t. 22.* umbellata.

*Periploca alia, floribus amplis circinatis & crispis, radice Bryoniae tuberosa. Plum. ic. 216. f. 27.*

*Apocynum scandens majus, folio subrotundo. Sloan. jam. 89. hist. 1. p. 207. t. 131. f. 2.*

- Echites scandens*, foliis ovatis nitidis venosis, floribus herbaceis. *Brown. jam.* 182.

*Habitat in Jamaica.*

trifida.

6. *ECHITES* pedunculis trifidis multifloris, foliis ovato-oblongis acuminatis. *Jacq. amer.* 13. *hist.* 1. 24.

*Habitat in America.* ‡

### CAMERARIA.

latifolia.

1. *CAMERARIA* foliis ovatis utrinque acutis transverse striatis. *Hort. cliff.* 76. *Roy. lugdb.* 412.

*Cameraria lato myrti folio.* *Plum. gen.* 18. *ic.* 72. *f.* 1.

*Cameraria arborea*, foliis ovato-acuminatis nitidis rigidis reflexis. *Brown. jam.* 182.

*Habitat in America calidiore.*

angustifolia.

2. *CAMERARIA* foliis linearibus. †

*Cameraria angusto linariæ folio.* *Plum. gen.* 18. *ic.* 72. *f.* 2.

*Habitat in America calidiore.*

### TABERNÆMONTANA.

citrifolia.

1. *TABERNÆMONTANA* foliis oppositis ovatis, floribus lateralibus glomerato-umbellatis.

*Tabernæmontana foliis lanceolatis.* *Hort. cliff.* 76. *Roy. lugdb.* 413.

*Tabernæmontana citri foliis undulatis.* *Plum. gen.* 18. *ic.* 248. *f.* 2.

*Pala. Rheed. mal.* 1. *p.* 81. *t.* 46.

*Habitat in utraque India.* ‡

laurifolia.

2. *TABERNÆMONTANA* foliis oppositis ovalibus obtusiusculis. †

*Tabernæmontana laurifolia*, flore albo, fructu rotundiore. *Amm. herb.* 212.

*Nerium arboreum*, folio latiore obtuso, flore luteo minore. *Sloan. jam.* 154. *hist.* 2. *p.* 62. *t.* 186. *f.* 2.

*Habitat in Jamaica ad ripas fluviorum.* ‡

alternifolia.

3. *TABERNÆMONTANA* foliis alternis, caule arboricente.

*Curutu-pala.* *Rheed. mal.* 4. *p.* 1. 43. *Raj. hist.* 1754.

*Habitat in Malabar.* ‡

Amsonia.

4. *TABERNÆMONTANA* foliis alternis, caulibus subherbaceis.

ANO-

*Anonymus suffrutex*. Gron. virg. 26.

*Habitat in Virginia*. 2.

Affinis *Cameraria* & *Tabernaemontana*. Folia alterna, lato-lanceolata, subsessilia, lœvia. Rami alterni. Racemus compositus, terminalis. Folliculi 2, reflexi, teretes, longi, acuminati. Semina cylindrica, scabra. Antheræ intra corollæ faucem. Stigma capitulum supra orbiculum membranaceum.

### CEROPEGIA.

1. **CEROPEGIA** umbellis pendulis, floribus erectis. *Candelabrum*.  
Niota-niodem-valli. *Rheed. mal.* 9. p. 27. t. 16.  
*Habitat in Malabar*. 2.
2. **CEROPEGIA** pedunculis bifloris. *Fl. zeyl.* 46. \* *biflora*.  
*Habitat in Zeylona*. 2.

## D I G Y N I A.

### PERIPLOCA.

1. **PERIPLOCA** floribus interne hirsutis. *græca*.  
*Periploca* foliis lanceolato-ovatis. *Hort. cliff.* 78. *Roy. lugdb.* 410.  
*Apocynum* folio oblongo. *Bauh. pin.* 303.  
*Apocynum* 2 angustifolium. *Clus. hist.* 1. p. 195.  
*Habitat in Syria*. 2.
2. **PERIPLOCA** spicis imbricatis. *indica*.  
*Periploca* foliis ovatis acutis glabris, spicis imbricatis.  
*Fl. zeyl.* 113. \*  
*Apocynum* caule volubili perenni, foliis ovato-lanceolatis. *Roy. lugdb.* 412.  
*Periploca* foliis angustis acutis glabris. *Burm. zeyl.* 187. t. 83. f. 1.  
*Apocyni* genus scandens ex india orientali. *Pluk. mant.* 17. t. 359. f. 2.  
*Habitat in Zeylona*. 3.
3. **PERIPLOCA** caule hirsuto. *africana*.  
*Cynanchum* caule volubili ramoso, foliis subovatis cum acumine. *Hort. cliff.* 79.  
*Apocynum* scandens africanum, vincæ pervincæ foliis subincanum. *Comm. rar.* 18. t. 18.

*Apocynum scandens africanum, convolvuli minoris folio, & caule hirsuto. Pluk. alm. 37. t. 137. f. 4.*

*Apocynum indicum minus, nummulariæ foliis. Morif. hist. 3. p. 611. f. 15. t. 3. f. 62.*

β. *Cynanchum foliis planis sinuatis, flore pallide viridi, fructu crasso glabro viridi. Burm. afric. 34. t. 14. f. 2.*

*Habitat in Africa. 24*

tennifolia,

4. *PERIPLOCA foliis lineari-lanceolatis.*

*Cynanchum radice grandulosa, foliis angustis sinuatis, floribus urceolatis miniatis. Burm. afric. 36. t. 15.*

β. *Cynanchum linearibus foliis acutis, floribus urceolatis rubris. Burm. afric. 37. t. 16. f. 1.*

*Habitat in Cap. b. Spei dunis.*

## CYNANCHUM.

acutum,

1. *CYNANCHUM caule volubili herbaceo, foliis cordato-oblongis glabris. Hort. cliff. 79. Hort. upf. 54. Roy. lugdb. 409.*

*Scammonia monspeliacæ affinis, foliis acutioribus. Bauh. pin. 294.*

*Apocynum 3 lat. folium. Clus. hist. 1. p. 125.*

*Habitat in Sicilia, Hispania, Astracan. 24*

suberosum.

2. *CYNANCHUM caule volubili inferne suberoso fisso, foliis cordatis acuminatis. Hort. cliff. 79. Roy. lugdb. 409. Gron. virg. 27.*

*Periploca carolinensis, flore minore stellato. Dill. elth. 300. t. 229. f. 296.*

*Habitat in America calidiore.*

hirsutum.

3. *CYNANCHUM caule volubili fruticoso inferne suberoso fisso, foliis ovato-cordatis. Hort. cliff. 79. Roy. lugdb. 410.*

*Periploca scandens, folio citrei, fructu maximo. Plum. spec. 2.*

*Apocynum scandens virginianum rugosum, pullis amplis floribus, capsulis alatis. Morif. hist. 3. p. 611. f. 15. t. 3. f. 61.*

*Apocynum scandens fruticosum, fungoso cortice, surinamenfe. Herm. par. 53.*

*Habitat in America. 5*

4. **CYNANCHUM** caule volubili herbaceo, foliis reniformi-cordatis acutis. *Hort. cliff.* 79. *Roy. lugdb.* 409. *Sauv. monsp.* 133. *monspeliacum.*  
*Scammonea monspeliaca*, foliis rotundioribus. *Bauh. pin.* 294.  
*Apocynum* IV. latifolium: *Scammonea valentina.*  
*Clus. hist.* 1. p. 126.  
*Habitat in Hispania & Narbonæ maritimis.* 24.  
 5. **CYNANCHUM** caule erecto divaricato, foliis cordatis glabris. *Hort. cliff.* 79. *Roy. lugdb.* 409. *Sauv. monsp.* 133. *erectum.*  
*Apocynum folio subrotundo.* *Bauh. pin.* 302.  
*Apocynum* I. latifolium. *Clus. hist.* 1. p. 124.  
*Habitat in Syria.* 24.

**A P O C Y N U M.**

1. **APOCYNUM** caule rectiusculo herbaceo, foliis ovatis utrinque glabris, cymis terminalibus. *androsæmifolium.*  
*Apocynum* caule erecto annuo, foliis ovatis. *Roy. lugdb.* 411.  
*Apocynum foliis ovatis.* *Hort. cliff.* 80.  
*Apocynum canadense*, foliis androsæmi majoris. *Bocc. fcc.* 35. t. 16. f. 3. *Morif. hist.* 3. p. 609. f. 15. t. 3. f. 16.  
*Habitat in Virginia, Canada.* 24.  
 2. **APOCYNUM** caule rectiusculo herbaceo, foliis oblongis, paniculis terminalibus. *cannabinum.*  
*Apocynum* foliis ovatis acutis subtus tomentosis. *Gron. virg.* 28.  
*Apocynum canadense* ramosum, flore e viridi albicante, siliqua tenuissima. *Morif. hist.* 3. p. 609. f. 15. t. 3. f. 14.  
*Apocynum virginianum*, flore herbaceo, siliqua longissima. *Morif. præl.* 232.  
*Apocynum canadense* maximum, flore minino herbaceo. *Pluk. alm.* 35. t. 13. f. 1.  
*Apocynum erectum virginianum* ramosum, caule subrubente, esulæ raræ foliis, siliquis tenuissimis. *Pluk. alm.* 35. t. 260. f. 4.  
*Habitat in Canada, Virginia.* 24.  
*Corollæ virides, nec pictæ sunt.*  
 3. **APOCYNUM** caule rectiusculo herbaceo, foliis ovato-lanceolatis. *venetum.*  
*Roy. lugdb.* 411. *Sauv. monsp.* 133.



*Apocynum foliis ovato-lanceolatis. Hort. cliff. 80.*  
*Tithymalus maritimus, purpurascens floribus. Bauh.*  
*pin. 291.*

*Etula rara e lio venetorum insula. Lob. hist. 201.*

*Habitat in Adriatici maris insulis. 24*

- frutescens. 4. **APOCYNUM** caule erecto frutescente, foliis lanceolato-ovalibus; corollis acutis fauce villosis. *Fl. zeyl. 114.*

*Apocynum caule erecto arboreo, foliis ovatis acutis.*  
*Roy. lugdb. 412.*

*Apocynum floribus fasciculatis. Fl. zeyl. 23. t. 12. f. 1.*  
*Habitat in Zeylona. 5*

- reticulatum. 5. **APOCYNUM** caule volubili perenni, foliis ovatis venosis. *Roy. lugdb. 412.*  
*Habitat in India.*

## ASCLEPIAS.

\* *foliis oppositis planis.*

- undulata. 1. **ASCLEPIAS** foliis sessilibus oblongis lanceolatis, undulatis, petalis ciliatis.

*Apocynum africanum, lapathi folio. Comm. rar. 16.*  
*t. 16.*

*Habitat in Africa. 24*

- gigantea. 2. **ASCLEPIAS** foliis amplexicaulibus oblongo-ovalibus. *Fl. zeyl. 112.*

*Apocynum erectum majus latifolium indicum. Pluk.*  
*alm. 35. t. 175. f. 3.*

*Beidel offar. Alp. aegypt. 85. t. 86.*

*Ericu. Rheed. mal. 2. p. 53. t. 31. Seb. thes. 1. p. 41.*  
*t. 26. f. 1.*

*Habitat in India, Aegypto. 24*

*Nectaris promunt non cornicula subulata, sed laminas solidas conniventes. Occidit inadsueta Pecora, uti Anemone nemorosa. Rumph. amb. 7. p. 25.*

- variegata. 3. **ASCLEPIAS** foliis ovatis rugosis nudis, caule simplici, umbellis subsessilibus; pedicellis tomentosis.

*Asclepias e virginia. Bauh. pin. 215.*

*Apocynum vetus americanum, Wisk. Dill. elth. 32.*

*Apocynum americanum erectum, tuberosa radice non incanum, foliis rigidioribus laevis subrotundis, floribus*

bus albis intus purpureis, summo caule corymbum magnum efformantibus. *Fluk. alm.* 34. t. 77. f. 1.

*Habitat in America boreali.* 24.

*Affinis A. Syriacæ.*

4. ASCLEPIAS foliis ovalibus subtus tomentosis, caule *syriacæ* simplicissimo, umbellis nutantibus.

*Asclepias* caule erecto simplici herbaceo, foliis ovalibus subtus tomentosis, umbellis nutantibus. *Vir. clif.* 20. *Hort. upf.* 53. *Gron. virg.* 27. *Roy. lugdb.* 410.

*Asclepias* caule erecto simplici annuo, foliis ovato-oblongis subtus incanis, umbella nutante. *Hort. clif.* 78.

*Apocynum* majus *syriacum* rectum. *Corn. canad.* 90.

*Apocynum syriacum.* *Clus. hist.* 2. p. 87.

*Beidellfar. Vegl. agypt.* 28. (non vero *Alpini*).

- β. *Asclepias* foliis lanceolato-ellipticis, caule simplici *exaltata*, glabro, nectarii corniculis conniventibus. *Amæn. acad.* 3. p. 404. *Kalm. it.* 3. p. 259.

*Habitat in Virginia.* 24.

5. ASCLEPIAS foliis ovatis subtus villosis, caule simplici, umbellis erectis, nectarialis resupinatis. *purpurascens.*

*Asclepias* floribus obsoleto purpureis, corniculis resupinatis. *Dill. elih.* 32. t. 28. f. 31.

*Apocynum* erectum noveboracense, foliis minus incanis, flore ex obsoleto dilute purpurascente. *Herm. par.* 33.

*Habitat in Carolina.* 24.

*Affinis valde A. syriacæ.*

6. ASCLEPIAS foliis ovatis subtus pilosiusculis, caule *amœna*, simplici, umbellis nectarisque erectis.

*Apocynum* floribus *amœne* purpureis, corniculis furrectis. *Dill. elih.* 31. t. 27. f. 30.

*Habitat in America septentrionali.* 24.

7. ASCLEPIAS foliis lanceolatis glabris, caule simplici, nivea, umbellis erectis lateralibus solitariis.

*Asclepias* caule erecto simplici annuo, foliis lanceolato-ovatis glabris, pedunculis alternis, umbellis erectis. *Gron. virg.* 27.

*Apocynum* *perficariæ* mitis folio, corniculis lacteis. *Dill. elih.* 33. t. 29. f. 32.

*Apocynum americanum*, foliis amygdali longioribus.

*Plum. spec. 2. ic. 30.*

*Habitat in Virginia & America calidiorē. ♂*

incarnata.

8. *ASCLEPIAS* foliis lanceolatis, caule superne diviso, umbellis erectis geminis.

*Asclepias* caule erecto ramoso annuo, foliis lanceolatis, umbellis terminalibus erectis pluribus. *Vir. cliff. 20 Hort. cliff. 78. Roy. lugdb. 411. Gron. virg. 27.*

*Apocynum minus rectum canadense. Corn. canad. 9. t. 93. Barr. rar. 8. t. 72. Raj. hist. 1089.*

*Habitat in Canada, Virginia. 2.*

*Caulis, ubi dividitur, folia e regione tria producit.*

curassavica.

9. *ASCLEPIAS* foliis lanceolatis petiolatis glabris, caule simplici, umbellis erectis solitariis.

*Apocynum* caule erecto simplici annuo, foliis lanceolatis, umbellis alternis erectis. *Hort. cliff. 78. Roy. lugdb. 411.*

*Apocynum* radice fibrosa, petalis coccineis, corniculis croceis. *Dill. elth. 34. t. 30. f. 33.*

*Apocynum curassavicum* f. *americanum*, fibrosa radice, floribus aurantiis, chamænerii foliis latioribus.

*Herm. parad. 36. t. 36.*

*Habitat in Curassao. 4. §*

decumbens.

10. *ASCLEPIAS* foliis villosis, caule decumbente.

*Asclepias* caule decumbente hirsuto, foliis ovatis obtusis villosis subsessilibus. *Gron. virg. 27.*

*Apocynum carolinianum aurantiacum pilosum. Pet. ficc. 90.*

*Habitat in Virginia. 2.*

lactifera.

11. *ASCLEPIAS* foliis ovatis, caule erecto, umbellis proliferis brevissimis.

*Asclepias* caule erecto simplici herbaceo, foliis ovatis acuminatis, corymbis compositis longitudine petiolorum. *Fl. zeyl. 111. \**

*Habitat in Zeylona. 2.*

Vincetoxicum.

12. *ASCLEPIAS* foliis ovatis basi barbatis, caule erecto, umbellis proliferis.

*Asclepias* caule erecto simplicissimo herbaceo, foliis cordato-lanceolatis, racemis conglomeratis alternis. *Fl. succ. 200, 212. Mat. med. 103. Dalib. parif. 76.*

*Ascle-*

*Asclepias* caule erecto annuo, foliis ovato-lanceolatis, floribus confertis. *Hort. cliff.* 78.

*Asclepias* albo flore. *Bauh. pin.* 303.

*Vincetoxicum*. *Dod. pempt.* 407.

*Habitat in Europæ glareosis.* 24

13. *ASCLEPIAS* foliis ovatis basi barbatis, caule subvolubili.

*Asclepias* caule subvolubili herbaceo, foliis ovato-lanceolatis, floribus confertis patentissimis. *Roy. lugdb.*

410. *Hort. ups.* 53. *Sauv. monsp.* 133.

*Asclepias* nigro flore. *Bauh. pin.* 303.

*Vincetoxicum* flore nigro. *Cam. epit.* 560.

*Habitat* Monspelli in collibus. 24

*Nectaria* huic & præcedenti sunt Calli quinque truncati absque auricula & ungue. Hæc in fertiliori solo incipit scandere subvolubili caule.

\* \* foliis latere revolutis.

14. *ASCLEPIAS* foliis revolutis lanceolatis, caule fruticoso.

*Asclepias* caule erecto fruticoso, foliis lanceolatis, umbellis nutantibus, conceptaculis inflatis pilosis.

*Hort. cliff.* 78. *Hort. ups.* 54. *Vir. cliff.* 20. *Roy. lugdb.* 410.

*Apocynum* erectum africanum, villosa fructu, salicis folio. *Herm. parad.* 23. t. 24. *Mill. dict.* t. 45.

*Apocynum* erectum elatius, salicis angusto folio, folliculis pilosis. *Pluk. alm.* 36. t. 138. f. 2.

*Habitat in Æthiopia.* 3

*Nectaria compressa* absque ungue, cujus loco auriculæ duæ, longiores, reflexæ. Folliculi inflati, aculeati, sed mites.

15. *ASCLEPIAS* foliis revolutis lineari-lanceolatis oppositis ternatisque, caule decumbente.

*Asclepias* montana humilis, radice longius proserpente, lini foliis. *Amm. ruth.* 10.

*Habitat in Sibiria.* 24

16. *ASCLEPIAS* foliis revolutis linearibus verticillatis, caule erecto.

*Asclepias* foliis verticillatis lineari-setaceis. *Gron. virg.* 26.

Apo-

*Apocynum marianum* erectum, linearis angustissimis foliis, umbellatum. *Fluk. mant.* 17. t. 336. f. 4.

*Apocynum marianum*, foliis angustissimis stellatis. *Pet. mus.* 609.

*Habitat in Virginia.* 24

*Folia sepe quaterna, verticillata. Pedunculi umbelliferi sunt oppositifolii.*

\* \* \* *foliis alternis.*

rubra

17. *ASCLEPIAS* foliis alternis ovatis, umbellis ex eodem pedunculo communi pluribus.

*Apocynum* caule erecto simplici annuo, foliis ovatis acuminatis alternis, pluribus in pedunculo umbellis.

*Gron. virg.* 27.

*Habitat in Virginia.*

tuberosa.

18. *ASCLEPIAS* foliis alternis lanceolatis, caule divaricato piloso.

*Asclepias* caule erecto divaricato villosa, foliis lanceolatis, umbellis simplicibus subterminalibus. *Hort. cliff.* 78. *Roy. lugdb.* 411.

*Apocynum novæ anglæ* hirsutum, tuberosa radice, floribus aurantiis. *Herm. lugdb.* 646. t. 647. *Dill. elrh.* 35. t. 30. f. 34.

*Habitat in America boreali.* 24

## STAPELIA.

variegata.

1. *STAPELIA* denticulis ramorum patentibus. *Vir. cliff.* 20. *Roy. lugdb.* 409.

*Stapelia* denticulis ramorum extrorsum prominulis. *Hort. cliff.* 77. *Hort. ups.* 53.

*Asclepias aizoides africana.* *Bratl. succ.* 3. p. 3. t. 22.

*Asclepias aizoides aphylla*, flore fritillariæ, siliquis longis angustis erectis. *Morif. hist.* 3. p. 610. f. 15. t. 3. f. 4.

*Apocynum humile aizoides*, siliquis erectis, africanum, *Herm. lugdb.* 52. t. 53. *Striff. bot.* 32.

*Fritillaria crassa* promont. bonæ spei. *Stap. theophr.* 335. *Habitat ad Cap. b. Spei.* 5

hirsuta.

2. *STAPELIA* denticulis ramorum erectis. *Hort. cliff.* 77. *Roy. lugdb.* 409.

*Asclepias africana aizoides*, flore pulchræ fimbriatæ. *Comm.*

*Comm. rar.* 19. t. 19. *Bratl. succ.* 3. p. 5. t. 23.

*Raf. inf. musc.* t. 9.

*Habitat ad Cap. b. Spei.* ‡

## HERNIARIA.

1. **HERNIARIA** glabra herbacea. *Bauh. hist.* 3. p. 378. glabra.

*Herniaria calycibus bractea nudis. Hort. cliff.* 41. *Fl.*

*svet.* 207, 213. *Mat. med.* 104. *Roy. lugdb.* 215.

*Hall. herb.* 182. *Dalib. parif.* 76.

*Polygonum minus* f. *Millagrana major. Bauh. pin.* 281.

*Habitat in Europæ apricis, glareosis, siccis.* ○

2. **HERNIARIA** hirsuta herbacea. *Bauh. hist.* 3. p. 379. hirsuta.

*Habitat in Angliæ, Hispaniæ, Italiæ agris.*

*Affinis nimium præcedenti; forte varietas.*

3. **HERNIARIA** caulibus fruticosis, floribus quadri-fruticosa, fidis. *Amæn. acad.* 4. p. 269. \* *Last. it.* 128. \*

*Herniaria fruticosa, viticulis lignosis. Bauh. pin.* 382.

*Polygonum herniariæ folio & facie, perampla radice.*

*Bauh. hist.* 3. p. 378. *Lob. ic.* 85.

*Habitat in Hispania.* ‡

4. **HERNIARIA** ? quæ *Polygonum lenticulatum*

*lenticulata*

*Polygonum minus tenuifolium. Bauh. pin.* 281.

*Polygonum minus lentifolium. Bauh. prodr.* 131.

*Polygonum maritimum longius radicatum nostras, serpylli folio crasso nitente. Pluk. alm.* 302. t. 53.

*f.* 3. *Pet. herb.* 10. *f.* 6. *Raj. angl.* 3. p. 161.

*Camphorata frutescens, foliis latis & angustis. Buxb. cent.* 1. p. 18. t. 28. *f.* 2. quoad figuram.

*Habitat in Hispaniæ montibus Escorial, Monspelii, Anglia.*

*Flores non vidi; at Parietariæ herniariæ folio. Plum. characterem misit D D. Aymen: cui Perianthium 5-partitum, acutum, vix patens, persistens. Corolla minima, 5-partita. Stamina 5, capillaria. Antheræ simplices. Germen subrotundum. Stylus vix ulius. Stigma obtusum. Capsula parva, unilocularis, calyce obvoluta. Semen 1, subovatum, compressum. Habitus proximus Herniariæ, sed mollior.*

CHE.

## CHENOPODIUM.

\* *foliis angulosis.*

- Bonus Henricus.** 1. **CHENOPODIUM** foliis triangulari - sagittatis integerrimis, spicis compositis aphyllis. *Hort. cliff.* 84. *Fl. fvec.* 208, 214. *Mat. med.* 106. *Roy. lugdb.* 278. *Hall. helv.* 174. *Dalib. parif.* 79.  
*Lapathum unctuosum. Bauh. pin.* 115.  
**Bonus Henricus.** *Bauh. hist.* 2. p. 965.  
*Habitat in Europæ ruderatis.* 2.  
*Spica composita, aphylla: Spiculis alternis sessilibus: floribus glomeratis, sessilibus.*
- urbicum.** 2. **CHENOPODIUM** foliis triangularibus subdentatis, racemis confertis strictissimis cauli approximatis longissimis. *Fl. fvec.* 209, 215. \*  
*Chenopodium latifolium minus ramosum, petiolis longissimis ex foliorum alis confertim nascentibus. Buxb. hall.* 69. t. 1.  
*Habitat in Europæ borealis plateis.* ☉
- rubrum.** 3. **CHENOPODIUM** foliis cordato - triangularibus obtusiusculis dentatis, racemis erectis compositis foliosis caule brevioribus. *Fl. fvec.* 210, 217. *Mat. med.* 107. *Dalib. parif.* 78.  
*Chenopodium erectum, foliis subtriangularibus antice sinuato - dentatis, racemis erectis foliosis lateralibus. Hort. cliff.* 85. *Roy. lugdb.* 219.  
*Atriplex sylvestris latifolia. Bauh. pin.* 119.  
*Pes anserinus. Dalech. hist.* 542. t. 342. *Fuchs. hist.* 653. *Bauh. hist.* 2. p. 975.  
*Habitat in Europæ cultis, rud-ratis.* ☉  
*Spica glomerulis sessilibus, interstinctæ foliolis linearibus; adulta rubet diffusa. Folia nitida, crassa.*
- murale.** 4. **CHENOPODIUM** foliis ovatis nitidis acutis dentatis, racemis ramosis nudis. *Fl. fvec.* 211, 216. *Dalib. parif.* 211.  
*Chenopodium erectum ramosissimum, foliis triangularibus dentatis, racemis ramosis caulem superantibus. Hort. cliff.* 85. *Lech. scan.* 16.  
*Chenopodium 1. Tabern. hist.* 812. f. 1.  
*Atriplex sylvestris latifolia, acutiore folio. Bauh. pin.* 119.  
*Atriplex dicta Pes anserinus. Bauh. hist.* 2. p. 975. c. figur.

Ha-

*Habitat in Europæ muris aggeribusque.* ☉

*Caules erecti. Fructus subpaniculati. Color viridis.*

5. **CHENOPODIUM** foliis deltoideis sinuato-dentatis ferocinum.  
rugosis glabris uniformibus, racemis terminalibus.

*Amzn. acad. 4. p. 309. \**

*Chenopodium hispanicum procerius, folio deltoide.*

*Tournef. inst. 666.*

*Blitum ficus folio. Raj. angl. 3. p. 155.*

*Habitat in Hispania, Gallia, Anglia.* ☉

6. **CHENOPODIUM** foliis rhomboideo-triangulari- album.  
bus erosis postice integris: summis oblongis, race-  
mis erectis. *Fl. fvec. 212, 218. Dalib. parif. 80.*

*Chenopodium foliis inferioribus ovatis acutis antror-  
sum dentatis: summis lanceolatis. Vir. cliff. 22.*

*Hort. cliff. 85. Gron. virg. 145. Roy. lugdb. 219.*

*Hall. herb. 175.*

*Atriplex sylvestris. Fuchs. hist. 1. 119.*

*Atriplex sylvestris, folio sinuato candicante. Bauh.  
pin. 119.*

*Habitat in agris Europæ.* ☉

*Planta matura albicat. Racemi recti, conglobati: spicis  
alternis, sessilibus, confertis.*

7. **CHENOPODIUM** foliis rhomboideis dentato-sinua- viride.  
tis, racemis ramosis subfoliatis. *Fl. fvec. 213, 219.*

*Dalib. parif. 78.*

*Chenopodium sylvestre, opuli folio. Vaill. parif. 36. t.*

*7. f. 1*

*Chenopodium folio oblongo integro. Dill. app. 62?*

*Atriplex sylvestris. Bauh. hist. 2. p. 972.*

*Habitat in Europæ cultis oleraceis, nimium abun-  
dans.* ☉

*Caulis erectus, viridis, angulis purpurascens. Fo-  
lia ramea lanceolata, integerrima, vix uno alte-  
rove dente. Racemi filiformes, divisi, longi, nudi.  
Calyces fructus argute pentagoni. Affine nimium  
precedenti.*

8. **CHENOPODIUM** foliis cordatis angulato-acumi- hybridum.  
natis, racemis ramosis nudis. *Fl. fvec. 214, 220.*

*Dalib. parif. 79.*

*Chenopodium foliis triangulari-sagittatis: infra medi-  
um sinuato-dentatis, racemis longissimis. Hort. cliff.*

*84. Roy. lugdb. 219.*



*Chenopodium stramonii folio. Vaill. parif. 36. t. 7. f. 2.*  
*Raj. angl. 4. p. 125.*

*Chenopodium 2. Tabern. hist. 812. f. 2.*

*Chenopodio affinis, folio lato laciniato in longissimum mucronem procurrens. Raj. suppl. 123.*

*Habitat in Europæ cultis. ☉*

Botrys. 9. **CHENOPODIUM** foliis oblongis sinuatis, racemis nudis multifidis. *Hort. cliff. 84. Hort. upf. 55. Mat. med. 109. Roy. lugdb. 219. Sauv. monsp. 272.*

*Botrys ambrosioides vulgaris. Bauh. pin. 138.*

*Botrys. Dod. pempt. 34.*

*Habitat in Europæ australis arenosis. ☉*

ambrosioides. 10. **CHENOPODIUM** foliis lanceolatis dentatis, racemis foliatis simplicibus. *Hort. cliff. 84. Hort. upf. 56. Roy. lugdb. 219.*

*Botrys ambrosioides mexicana. Bauh. pin. 138. 516.*

*Botrys odorata suaveolens americana mexicanave.*

*Morif. hist. 2. p. 605. f. 5. t. 35. f. 8.*

*Habitat in Mexico, Lusitania. ☉*

multifidum. 11. **CHENOPODIUM** foliis multifidis: segmentis linearibus, floribus axillaribus sessilibus.

*Chenopodium sempervirens, foliis tenuiter laciniatis.*

*Dill. elih. 78. t. 66. f. 77.*

*Habitat in Bonaria. ☿*

anthelminticum. 12. **CHENOPODIUM** foliis ovato-oblongis dentatis, racemis aphyllis. *Kalm. canad. 2. p. 283.*

*Chenopodium, lycopi folio, perenne. Dill. elih. 77. t. 66. f. 76.*

*Habitat in Pennsylvania, Bonaria. 2.*

glaucum. 13. **CHENOPODIUM** foliis ovato-oblongis repandis, racemis nudis simplicibus glomeratis. *Fl. suec. 215, 221. Dalib. parif. 77.*

*Chenopodium foliis ovato-oblongis repande sinuatis, florum racemis glomeratis ex ala solitariis. Vir. cliff. 22. Hort. cliff. 85. Roy. lugdb. 219.*

*Chenopodium angustifolium laciniatum minus. Tournef. inst. 506.*

*Atriplex angustifolia laciniata. Bauh. hist. 2. p. 472. t. 473.*

*Atriplex sylvestris secunda. Tabern. ic. 427.*

*Habitat ad Europæ fineta. ☉*

\* \* *foliis integris.*

14. **CHENOPODIUM** foliis integerrimis rhombeo-ovatis, floribus conglomeratis axillaribus. *Fl. fvec.* 216, 222. *Mat. med.* 108. *Dalib. parif.* 77. *Chenopodium foliis triangulari-ovatis. Hort. cliff.* 84. *Roy. lugdb.* 218. *Atriplex foetida. Bauh. pin.* 119. *Vulvaria. Dalech. hist.* 543. *Habitat in Europæ cultis oleraceis.* ☉
15. **CHENOPODIUM** foliis integerrimis ovatis, caule decumbente, cymis dichotomis aphyllis axillaribus. *Chenopodium foliis ovatis integerrimis, caule decumbente. Vir. cliff.* 21. *Fl. fvec.* 217, 223. *Roy. lugdb.* 218. *Dalib. parif.* 77. *Blitum foliis ovatis. Hort. cliff.* 28. *Blitum polyspermum. Bauh. pin.* 118. *Polysporon cassiani, bassii, anguillare. Lob. hist.* 128. *Habitat in Europæ cultis.* ☉
16. **CHENOPODIUM** foliis lineari-lanceolatis planis integerrimis. *Hort. cliff.* 86. *Hort. ups.* 55. *Roy. lugdb.* 220. *Linaria scoparia. Bauh. pin.* 212. *Oxyris. Dod. pempt.* 151. *Habitat in Græcia, Japonia, China.* ☉
17. **CHENOPODIUM** foliis subulatis semicylindricis. *Chenopodium foliis subulatis; superne planis, subtus convexis. Hort. cliff.* 86. *Fl. fvec.* 218, 224. *Roy. lugdb.* 220. *Kali minus album, semine splendente. Bauh. pin.* 289. *Morif. hist.* 2. p. 610. f. 5. t. 33. f. 3. *Habitat in Europæ maritimis.* *maritimum.*
18. **CHENOPODIUM** foliis lanceolatis subcarnosis, integerrimis, corymbis dichotomis aristatis axillaribus. *Chenopodium foliis lineari-lanceolatis, racemis dichotome ramosis setis terminatis. Ait. goett.* 1752. *β. Chenopodium virginicum, foliis linearibus obtusis subcanaliculatis, pedunculis axillaribus dichotomis muticis. Sp. pl.* 1. p. 222. *Habitat in Sibiria, Virginia.* ☉ *aristatum.*
- Linnei Spec. plant. pars I.* X *Cau-*

*Caulis spithameus, inequalis ramoso-patulus. Folia carnosâ, lanceolata, obtusa. Corymbi axillares, dichotomi, folio longiores, terminati spinis inermibus. Flores sessiles in divaricationibus.*

## BETA.

- maritima.* 1. BETA caulibus decumbentibus.  
Beta caulibus decumbentibus, foliis triangularibus petiolatis. *Mill. dict.*  
Beta sylvestris maritima. *Bauh. pin. 118. Raj. angl. 4. p. 127.*  
*Habitat in Angliâ, Belgii littoribus maris.*
- vulgaris.* 2. BETA caule erecto.  
Beta. *Hort. cliff. 83. Hort. upf. 56. Mat. med. 113. Roy. lugdb. 220.*
- rubra.* α Beta rubra vulgaris. *Bauh. pin. 118.*  
β. Beta rubra major. *Bauh. pin. 118.*  
γ. Beta rubra, radice rapæ. *Bauh. pin. 118.*  
δ. Beta lutea major. *Bauh. pin. 118.*  
ε. Beta pallide virens major. *Bauh. pin. 118.*
- Cicla.* ζ. Beta alba vel pallescens, quæ Cicla officinarum. *Bauh. pin. 118.*  
η. Beta communis viridis. *Bauh. pin. 118.*  
*Habitat - - - - - ♂, forte a maritima, in exoticis, prognata.*

## SALSOLA.

- kali.* 1. SALSOLA herbacea decumbens, foliis subulatis spinosis, calycibus marginatis axillaribus.  
Salsola foliis pungentibus. *Hort. cliff. 86. Fl. suec. 206, 225. Roy. lugdb. 220. Gron. virg. 28.*  
Kali spinosa affinis. *Bauh. pin. 289. Morif. hist. 2. f. 5. t. 33. f. 11.*  
*Tragum. Cam. epit. 779.*  
*Habitat in Europæ littoribus maris. ☉*  
*Planta decumbens. Folia subulata, crassa, spinosa, subtus & margine, uti ipse caulis, scabra s. hispida. Calyces cincti margine foliaceo.*
- Tragus.* 2. SALSOLA herbacea erecta, foliis subulatis spinosis lævibus, calycibus ovatis. *Amæn. acad. 4. p. 311.*  
Kali spinosum cochleatum. *Bauh. pin. 289.*

Kali

*Kali spinosum*, foliis longioribus & angustioribus.

*Tournef. inst.* 247.

*Drypis theophrasti. Tabern. hist.* 423.

*Habitat in Europa australi.* ☉

3. *SALSOLA* herbacea, foliis subulatis mucronatis, calycibus explanatis. rosacea.

*Salsola* foliis conico-subulatis mucronatis, calycibus corolliformibus obtusis coloratis crenatis. *Hort. ups.* 56.

*Kali humile*, alis purpureis florem rosaceum mentientibus. *Buxb. cent.* 1. p. 9. t. 14. f. 2.

*Habitat ad Asiæ septentrionalis salinas.* ☉

4. *SALSOLA* frutescens prostrata, foliis linearibus prostratis. inermibus. *Amxn. acad.* 2. p. 346. \*

*Salsola* lignosa, ramis filiformibus adicendentibus, floribus conglomerato-spicatis. *Læfl. it.* 131. \*

*Habitat in Asia boreali, Hispania.* ☿

5. *SALSOLA* frutescens; foliis ovatis acutis carnosiss. vermiculata.

*Salsola* fruticosa, floribus spicatis alternis solitariis.

*Læfl. it.* 129. \*

*Kali fruticosum*, ericæ folio. *Buxb. cent.* 1. p. 8. t. 14. f. 1?

*Kali geniculatum* aphyllanthes, glavis paleaceis floribus, hispanicum. *Barr. rar.* 501. t. 215.

*Habitat in Hispania.* ☿

6. *SALSOLA* herbacea patula, foliis inermibus. *Guett. Soda.*

*flamp.* 426. *Sauv. monsp.* 7.

*Kali majus*, cochleato semine. *Bauh. pin.* 287.

*Kali. Dod. pempt.* 81.

*Soda, Kali magnum*, sedi medii folio. *Lob. ic.* 394.

*Habitat in Europæ australis salis.* ☉

7. *SALSOLA* diffusa herbacea, foliis teretibus glabris, sativa. floribus conglomeratis. *Læfl. it.* 132. \*

*Kali hispanicum* supinum annum, sedi foliis brevioribus. *Juss. ass. paris.* 1715. p. 74.

*Kali minus alterum.* *Bauh. pin.* 283.

*Habitat in Hispaniæ australis maritimis.* ☉

8. *SALSOLA* diffusa herbacea, foliis teretibus obtusis hirsutis tomentosis.

*Chenopodium hirsutum.* *Sp. pl.* 1. p. 221.

*Kali minus villosum.* *Bauh. pin.* 289.

- Kali parvum hirsutum. *Bauh. hist. 3. p. 702.*  
*Habitat in Daniæ, Montpellii maritimis.* ☉
- Salsa. 9. SALSOLA patula herbacea, foliis semiteretibus obtusis, ramis reflexis.  
 Chenopodium foliis linearibus obtusis subtus convexis, caule ramofo: ramis deflexis. *Hort. upf. 55. Sp. pl. 1. p. 221.*  
 Chenopodium maritimum, foliis sedi teretibus. *Buxb. cent. 1. p. 21. t. 31. f. 1.*  
*Habitat ad Afracanam.* ☉
- altissima. 10. SALSOLA erecta ramosissima herbacea, foliis filiformibus acutiusculis.  
 Chenopodium altissimum. *Sp. pl. 221. Hort. upf. 55. n. 3.*  
 Chenopodium altissimum, foliis succulentis. *Buxb. cent. 1. p. 21. t. 31. f. 2.*  
 Kali gramineo folio. *Bauh. pin. 289. prodr. 123. Burf. XVI. 20.*
- β- Salsola foliis filiformibus inermibus mucronatis, caule ramosissimo. *Hort. upf. 56.*  
 Lerchea foliis filiformibus acutis. *Hall. gatt. 2. p. 22.*  
*Habitat ad Salinas Italiæ, Saxoniæ, Afracani.* ☉
- fruticosa. 11. SALSOLA erecta fruticosa, foliis filiformibus obtusiusculis.  
 Chenopodium foliis linearibus teretibus carnosis, caule fruticoso. *Hort. cliff. 86. Sp. pl. 1. p. 221. Roy. lugdb. 220. Guett. stamp. 2. p. 425.*  
 Chenopodium, sedi folio minimo, frutescens perennæ. *B. Duham. arb. 1. p. 163. t. 62.*  
 Lerchea foliis obtusis. *Hall. gatt. 21.*  
 Kali species Vermicularis marina arborefcens. *Bauh. hist. 3. p. 704.*  
 Anthyllis chamæphithydes frutescens. *Bauh. pin. 232.*  
 Sedum minus arborefcens. *Munt. hist. 469.*  
*Habitat in maritimis Galliæ, Hispaniæ, Persiæ.* ‡  
 Semina cochleæ forma præcedentes ad Salsolas revocat apud Michel. *florent. 128.*

## ANABASIS.

- tamariscifolia. 1. ANABASIS foliis subulatis, fructibus exsuccis.  
 Kali fruticosum hispanicum, tamariscifolio. *Tournef. inst. 247.*  
*Habitat in Hispania.* ‡

Fru-

*Frutex ramulis albis, glaberrimis. Folia subulata, triquetra, Tamarisci. Flores subspicati. solitarii, axillares. Calyx triphyllus, major. Petala quinque, ovata, calyce minora. Stamina quinque, corolla longiora. Germen conicum desinens in Stylum subulatum, Stigmatibus tribus, obtusis.*

2. ANABASIS aphylla: articulis emarginatis. *Amæn. aphylla. acad. 2. p. 347. Gron. orient. 73.*

*Kali bacciferum, falicorniae facie. Buxb. cent. 1. p. 11. t. 18.*

*Kali geniculatum alterum f. minus. Bauh. pin. 289.*

*Habitat ad maris Caspii litora & circa Tripolin. b*

3. ANABASIS foliis subclavatis. *Amæn. acad. 2. p. 347. foliosa.*

*Kali bacciferum, foliis clavatis. Buxb. cent. 1. p. 19.*

*t. 19. f. 1.*

*Habitat ad maris Caspii litora. ☉*

## CRESSA.

1. CRESSA. *Amæn. acad. 1. p. 395. Sauv. monsp. 92. erratica.*

*Anthyllis. Alp. exot. 157. t. 156. Raj. hist. 215. Magn. char. 212.*

*Quamoclit minima humifusa palustris, herniariae folio:*

*Tournes. cor. 4.*

*Chamaepithys incana, exiguo folio. Bauh. pin. 249.*

*Lysimachiae spicatae purpureae affinis, flosculis in cacumine caulis, pluribus quasi in nodos junctis. Pluk.*

*alm. 236. t. 43. f. 6.*

*Habitat in Cretae litoribus salis, Romae ad ostia.*

## TRIANTHEMA.

1. TRIANTHEMA. *Sauv. meth. 127. Kniph. orig. 4. Portulaca- præfat.*

*strum.*

*Portulaca curassavica procumbens, capparidis folio,*

*flore muscoso, capsula bifurca. Herm. par. 213. t.*

*213. Raj. suppl. 506.*

*Portulacæ affinis, folio subrotundo, flore pentapetalo*

*dilute purpureo. Sloan. jam. 89. hist. 1. p. 205. Raj.*

*suppl. 506.*

*Kali curassavicum procumbens, foliis subrotundis.*

*Pluk. alm. 202. t. 95. f. 4.*

*Habitat in Jamaica, Curassao.*

*Flores non vidi; an stamina 5-10? an Stylus 1?*

X3

GOM-

## GOMPHRENA.

- globosa. 1. GOMPHRENA caule erecto, foliis ovato-lanceolatis, capitulis solitariis, pedunculis diphyllis. *Hort. cliff.* 86. *Hort. upf.* 57. *Fl. zeyl.* 115. *Vir. cliff.* 22. *Roy. lugdb.* 418.  
*Amarantho* affinis indiæ orientalis, floribus conglomeratis, ocymastri folio. *Breyn. cent.* 109. t. 51. *Comm. hort.* 1. p. 85. t. 45.  
 Flos globosus. *Rumph. amb.* 5. p. 289. t. 100. f. 2.  
*Habitat in India.* ☉
- perennis. 2. GOMPHRENA foliis lanceolatis, capitulis diphyllis, flosculis perianthio proprio distinctis.  
*Amaranthoides* perenne, floribus stramineis radiatis. *Dill. elth.* 24. t. 20. f. 22.  
*Habitat in Bonaria.* 24
- hispida. 3. GOMPHRENA caule erecto, capitulis diphyllis, foliis crenatis.  
 Nin-angani. *Rheed. mal.* 9. p. 141. t. 72.  
*Habitat in Malabaria.*  
*Flores cærulescentes.*
- brasiliens. 4. GOMPHRENA caule erecto, foliis ovato oblongis, capitulis aphyllis globosis pedunculatis. *Aman. acad.* 4. p. 310.  
*Amarantho* affinis brasiliens, glomeratis parvisque floribus *Breyn. cent.* 1. 52.  
*Habitat in Brasilia.*
- ferrata. 5. GOMPHRENA caule erecto brachiato, capitulis solitariis terminalibus sessilibus, calycibus ferratis.  
*Habitat in America.*
- interrupta. 6. GOMPHRENA caule erecto, spica interrupta. *Roy. lugdb.* 419.  
*Habitat in America.*
- flava. 7. GOMPHRENA pedunculis oppositis bifidis tricapitatis, capitulo intermedio sessili.  
*Gomphrena* pedunculis ad alas geminatis tricapitatis. *Hort. cliff.* 27.  
*Habitat in Vera cruce.*

## BOSEA.

- yervamora. 1. BOSEA. *Hort. cliff.* 84. *Roy. lugdb.* 223.

*Tilia* forte arbor racemosa, folio longiore subtus albicante nervis purpureis insignito, flore pentapetalo purpureo. *Sloan. jam.* 135. *hist.* 2. p. 19. t. 158. f. 3. *Raj. dendr.* 88.

*Frutex peregrinus horto bosiano yervamora dictus.*

*Walt. hort.* 24. t. 10.

*Habitat in Canariis insulis.* §

## ULMUS.

1. **ULMUS** foliis duplicato-ferratis: basi inæqualibus. *campestris.*

*Ulmus fructu membranaceo. Hort. cliff.* 83. *Fl. svec.* 219, 226. *Mat. med.* 105. *Roy. lugdb.* 223. *Dalib. parif.* 82.

*Ulmus campestris & theophrasti. Bauh. pin.* 246.

*Ulmus. Dod. pempt.* 837.

*Habitat in Europa ad pagos, a Gades in Gesticiam.* §

2. **ULMUS** foliis æqualiter ferratis: basi inæqualibus. *americana.*

*Ulmus fructu membranaceo, foliis simplicissime ferratis. Gron. virg.* 145. \*

*Habitat in Virginia.* §

3. **ULMUS** foliis æqualiter ferratis: basi æqualibus. *pumila.*

*Ulmus humilis. Amm. ruth.* 260.

*Ulmus pumila, foliis parvis, glabra, cortice fungoso.*

*Pluk. alm.* 293.

*Habitat in Sibiria.* §

## N A M A.

1. **NAMA** caule erecto glabro, foliis linearibus, floribus racemosis. *Zeylanica.*

*Nama, Amæn. acad.* 1. p. 395. *Fl. zeyl.* 117. t. 2. \*

*Kniph. orig.* 4. *præf.*

*Alfine veronicæ affinis, gratiolæ foliis, maderaspatana. Pluk. alm.* 22. t. 130. f. 2.

*Habitat in India.* ☉

2. **NAMA** caule decumbente, foliis obovatis, floribus solitariis. *jamaicensis.*

*Nama reclinata villosa, foliis ovatis, petiolatis marginatis recurrentibus, floribus solitariis. Brown. jam.* 185. t. 18. f. 2.

*Habitat in Jamaica.*

*Corolla monopetala, cylindrica, quinquecrenata, longitudine calycis. Brown.*



## HYDROLEA.

spinosa.

## 1. HYDROLEA.

*Planta lacustris f. palustris spinosa. Less. ii. 310.*

*Habitat in America meridionali. 2.*

*Caules erecti, pedales: Ramis alternis. Folia lanceolata, sessilia, subundulata, viscosa: Spinae axillares, patentes. Flores terminales, subsessiles, pauci, conferti.*

## HEUCHERA.

americana.

1. HEUCHERA. *Hort. cliff. 82. Gron. virg. 29. Roy. lugdb. 437.*

*Cortusa americana, flore squalide purpureo. Herm. parad. 131. t. 131.*

*Sanicula f. Cortusa americana spicata, floribus squalide purpureis. Pluk. alm. 332. t. 58. f. 3.*

*Habitat in Virginia. 2.*

## SWERTIA.

perennis.

## 1. SWERTIA corollis quinquefidis, foliis radicalibus ovalibus.

*Swertia. Hort. cliff. 53. Roy. lugdb. 433.*

*Gentiana corollarum laciniis nectario gemino notatis. Monnier. obs. 154.*

*Gentiana palustris latifolia. Bauh. pin. 188.*

*Gentiana XII. punctato flore. Clus. hist. 1. p. 316.*

*Habitat in Helvetiæ, Baviaræ, Galliæ alpinis. 2.*

difformis.

## 2. SWERTIA corollis quinquefidis: terminali sexfida, pedunculis longissimis, foliis linearibus.

*Gentiana foliis linearibus acuminatis, pedunculis longissimis nudis unifloris oppositis. Gron. virg. 30.*

*Habitat in Virginia.*

rotata.

## 3. SWERTIA corollis quinquefidis, foliis lanceolato-linearibus.

*Habitat in Sibiria. D. Gmelin. ☉*

*Planta spithamea. Folia lanceolato-linearia; Rami alterni, pedunculis unifloris; Corollæ planæ.*

corniculata.

4. SWERTIA corollis quadrifidis quadricornibus. *Aman. acad. 2. p. 344. \**

*Habitat in Sibiria, Gmelin; Canada, Kalm. ☉*

## 5. SWER-

5. **SWERTIA** corollis quadrifidis ecornibus. *Anan. dichotoma.*  
*acad. 2. p. 345. \**  
*Habitat in Sibiria. D. Gmelin. ☉*

## GENTIANA.

\* *Corollis quinquefidis subcampaniformibus.*

1. **GENTIANA** corollis quinquefidis rotatis verticillatis, calycibus spathaceis. *Hall. helv. 479.*  
*Gentiana floribus lateralibus confertis pedunculatis,*  
*corollis rotatis. Hort. cliff. 80. Fl. succ. 201, 227.*  
*Mat. med. 110. Roy. lugdb. 432. Sauv. monsp. 126.*  
*Gentiana major lutea. Bauh. pin. 187. Fl. lapp. 96.*  
*Gentiana. Cam. epit. 415.*  
*Asterias. Rencal. spec. 64. t. 63.*  
*Habitat in Alpibus Norvegicis, Helveticis, Apenninis,*  
*Austriacis, Pyrenæis, Tridentinis. 2.*
2. **GENTIANA** corollis quinquefidis campaniformibus purpurea, verticillatis, calycibus subspathaceis.  
*Gentiana corollis campaniformibus verticillatis, foliis*  
*imis petiolatis ellipticis. Hall. helv. 479.*  
*Gentiana major purpurea. Bauh. pin. 187.*  
*Gentiana major, flore purpureo. Oed. dan. t. 50.*  
*Gentiana major alia. Cam. epit. 416.*  
*Coilantha. Rencal. spec. 65.*  
*Habitat in alpibus Helveticis, Pyrenæis, Norvegicis. 2.*  
*Planta, quam sub nomine Gentianæ majoris luteæ. flo-*  
*ribus punctis carentibus campanulæ forma C. B. 187.*  
*meminit, hujus species erat, lecta a Bursero.*
3. **GENTIANA** corollis quinquefidis campanulatis punctatis, verticillatis, calycibus quinqueidentatis.  
*Gentiana major, flore punctato. Bauh. pin. 187.*  
*Habitat in Helvetiæ, Sibiria, Austria, Stiria, Rhe-*  
*tia alpinis. 2.*  
*Hujus varietatem legit A. Gmelinus in Sibiria foliis*  
*lineari-lanceolatis, Floribus terminalibus ternis, pe-*  
*dunculis propriis, corollis flavis punctis sparsis purpu-*  
*rascentibus.*
4. **GENTIANA** corollis quinquefidis campanulatis oppositis sessilibus, foliis amplexicaulibus. *asclepiadea.*  
*Gentiana foliis ovato-lanceolatis floribus campanula-*  
*tis in alis sessilibus. Hall. helv. 478.*

*Gentiana floribus lateralibus solitariis sessilibus, corollis erectis. Hort. cliff. 80. Roy. lugdb. 432.*

*Gentiana asclepiadis folio. Bauh. pin. 187.*

*Dasystephana. Rencal. spec. 67. t. 68.*

*Habitat in alpinis Helvetiæ, Pannoniæ, Mauritaniz. 2.*

*Folia & caulis cruciatim opposita; Calyces longitudine Corollæ.*

Pneumonanthe.

5. **GENTIANA** corollis quinquefidis campanulatis oppositis pedunculatis, foliis linearibus.

*Gentiana floribus terminalibus raris, corollis erectis plicatis, foliis linearibus. Hort. cliff. 80. Fl. suec. 202, 228. Roy. lugdb. 432. Dalib. parif. 81.*

*Gentiana angustifolia autumnalis major. Bauh. pin. 188.*

*Gentiana palustris angustifolia. Bauh. pin. 188.*

*Cyana. Rencal. spec. 69. t. 63.*

*Habitat in Europæ pascuis humidiusculis. 2.*

saponaria.

6. **GENTIANA** corollis quinquefidis campanulatis ventricosis verticillatis, foliis trinerviis.

*Gentiana floribus ventricosis campanulatis erectis quinquefidis, foliis ovato-lanceolatis. Gron. virg. 29.*

*Gentiana major virginiana, floribus amplis ochroleucis. Pluk. alm. 166. t. 186. f. r.*

*Gentiana virginiana, saponariæ folio, flore cæruleo longiore. Moris. hist. 3. p. 484. f. 11. t. 5. f. 4. Catesb. car. 1. p. 70. t. 70.*

*Habitat in Virginia. 2.*

villosa.

7. **GENTIANA** corollis quinquefidis campanulatis ventricosis, foliis villosis.

*Gentiana floribus ventricosis campanulatis erectis quinquefidis, foliis oblongis acuminatis leviter villosis. Gron. virg. 145.*

*Habitat in Virginia. 2.*

acaulis.

8. **GENTIANA** corolla quinquefida campanulata caulem excedente.

*Gentiana corolla campanulata caulem longitudine excedente. Hort. cliff. 81. Roy. lugdb. 432. Hall. helv. 477. Sauv. monsp. 136.*

*Gentiana alpina latifolia, magno flore. Bauh. pin. 187. prodr. 97.*

- β. *Gentiana alpina angustifolia, magno flore. Bauh. pin. 187.*

Thy-

*Thylacitis. Reneal. spec. 70. t. 68.*

*Habitat in alpinis Helveticis, Austriacis, Pyrenaicis. 24*

9. GENTIANA corollis quinquefidis coronatis crenatis, pedunculo terminali longissimo dichotomo. *exaltata.*

*Centaureum minus maritimum, amplo flore cæruleo.*

*Plum. spec. 3. ic. 81. f. 1.*

*Habitat in America. ☉*

\* \* *Corollis quinquefidis infundibuliformibus.*

10. GENTIANA corolla quinquefida infundibuliformi *verna.*  
caulem excedente, foliis radicalibus confertis majoribus.

*Gentiana flore unico tubuloso, foliis ad terram congestis acutis. Hall. helv. 476.*

*Gentiana alpina verna major. Bauh. pin. 188.*

*Gentianella verna minor. Clus. hist. 315.*

*Ericoila. Reneal. spec. 75. t. 68.*

*Habitat in alpinis Helvetiæ, Austriæ, Rhetiæ, Pyrenæis. 24*

11. GENTIANA corolla quinquefida infundibuliformi *bavarica.*  
serrata, foliis obtusis.

*Gentiana foliis ovatis, caule unifloro, flore serrato. Hall. helv. 476.*

*Gentiana caule unifloro, flore hypocrateriformi quinquefido: laciniis subrotundis crenatis. Hort. cliff. 81.*

*Gentianella elegantissima bavarica. Cam. hort. 65. t. 15. f. 2.*

*Habitat in alpinis Helvetiæ, Baviaræ, Sveciæ.*

12. GENTIANA corollis quinquefidis infundibuliformibus *aurea*  
acuminatissimis: fauce imberbi mutica, ramis oppositis.

*Gentiana alpina pumila, flore aureo. Barr. rar. 3. t. 104. f. 1.*

*Habitat in alpinis Burdegalenfibus & Lapponiæ Norvegicæ. D. C. Solander. ☉*

*Caulis erectus, spithameus. Rami ad radicem plures, parvi, & laterales oppositi valde erecti. Folia Radicalia ovata, glabra, parva; Caulina similia, majora, sessilia, obtusiuscula. Flores terminales pauci in capitulum. Calyces angusti, subpedunculati, in lacinias 3 subulatas divisi. Corollæ infundibuliformes, tubo longitudine calycis; Limbo flavo: laciniis*

*niis 5, valde acuminatis, integerrimis, absque dentibus interjectis.*

*aiuala.*

13. **GENTIANA** corollis quinquefidis infundibuliformibus, ramis unifloris alternis.

*Gentiana* corollis infundibuliformibus quinqueidentatis, ramis alternis unifloris. *Fl. suec.* 204, 231.

*Gentiana* corolla infundibuliformi: denticulo laciniis interposito. *Fl. lapp.* 95. \*

*Gentiana* humillima, caule ramoso, tubo floris longissimo. *Hall. helv.* 475. t. 7. f. 5.

*Gentiana alpina æstiva, centaureæ minoris folio.* *Bauh. pin.* 188.

*Gentiana IX. Oed. dan.* t. 17.

*Habitat in Lapponiæ, Helvetiæ, Pyrenæis summis al-  
pibus.* ☉

*aquatica.*

14. **GENTIANA** corollis quinquefidis infundibuliformibus terminalibus sessilibus, foliis margine membranaceis.

*Gentiana* foliis margine membranaceis basi coadunatis. *Amman. acad.* 2. p. 343.

*Gentiana humilis aquatica verna.* *Amman. ruth.* 4. t. 1. f. 1.

*Habitat in Sibiria, Amman.* ☉

*utriculosa.*

15. **GENTIANA** corollis quinquefidis hypocrateriformibus, calycibus plicato-alatis. *Scop. carn.* 299.

*Gentiana* utriculis ventricosus. *Bauh. pin.* 188.

*Gentianella cærulea cordata.* *Column. ecphr.* 220. t. 221.

*Gentianella annua, azureo flore.* *Barr. rar.* 18. t. 48.

*Habitat in Alpibus Helvetiæ, Austriæ, Italiæ.* ☉

*Exacoides.*

16. **GENTIANA** corollis quinquefidis hypocrateriformibus, calycibus membranaceo-carinatis, caule dichotomo, foliis cordatis.

*Centaurium perfoliatum, florum calyce membranaceo ventricoso.* *Burm. afr.* 208. t. 74. f. 5.

*Centaurium luteum flore amplo, capsula quatuor alis membranaceis.* *Pluk. alm.* 208. t. 275. f. 4. *Seb. mus.*

1. t. 22. f. 1.

*Habitat ad Cap. b. spei.* ☉

*Centaurium.*

17. **GENTIANA** corollis quinquefidis infundibuliformibus, caule dichotomo.

Gen-

*Gentiana foliis lineari-lanceolatis, caule dichotomo, corollis infundibuliformibus quinquefidis. Hort. cliff. 81. Fl. fvec. 205, 232. Mat. med. 112. stylis simplicibus. Vir cliff. 21. Roy. lugdb. 7. Dalib. parif. 81.*

*Centaurium minus. Bauh. pin. 278.*

*Erithræa. Renealm. spec. 77. t. 76.*

- β. *Centaurium minus pumilum ramosissimum. Lt. al. 157. Rosen. scan. 10.*

*Centaurium minus palustre ramosissimum, flore purpureo. Vaill. parif. 32. t. 6. f. 1.*

*Habitat in Europæ apricis, præsertim maritimis. ☉*

*Centaurium palustre minimum, flore inaperto. Vaill. parif. 32. t. 6. f. 2. vel huic affinis vel *Sarothræ* species ut *D. Aymen*.*

18. GENTIANA corollis quinquefidis infundibuliformibus, floribus alternis sessilibus. *spicata.*

*Gentiana corollis infundibuliformibus quinquefidis laxè spicatis, foliis lanceolatis. Sauv. monsp. 132.*

*Centaurium minus spicatum album. Bauh. pin. 278. prodr. 130. t. 130. Bauh. hist. 3. p. 353.*

*Centaurium minus album. Tabern. ic. 780.*

- β. *Centaurium minus spicatum, flore rubello. Tournef. inst. 122.*

*Habitat in Euganeis & Monspelii, Italiæ præter humidis herbosis. ☉*

19. GENTIANA corollis quinquefidis infundibuliformibus, floribus verticillatis, caule simplicissimo. *verticillata.*

*Centaurium minus ad alas floridum. Plam. spec. 3. is. 81. f. 2.*

*Habitat in America.*

20. GENTIANA corollis quinquefidis infundibuliformibus, caule acutangulo, foliis ovatis amplexicaulibus. *quinquefolia.*

*Habitat in Pensylvania. Kalm.*

*Planta magnitudine & statura sequentis. Caulis indivisus, tetragonus angulis membranaceis. Folia ovata, sessilia, amplexicaulia, trinervia, acuta. Pedunculi oppositi, apice ferentes Flores quinque, subpedicellatos. Calyces brevissimi, angustii. Corollæ infundibuliformes, subcæruleæ: limbo parvo, faucē pervia.*

- aphylla.** 21. *GENTIANA* corollis quinquefidis hypocrateriformibus, caule aphyllis. *Jacq. amer.* 17. *hist.* t. 60. f. 2.  
*Habitat in America.*  
*Planta minima filiformis vix ramosa.*
- Amarella.** 22. *GENTIANA* corollis quinquefidis hypocrateriformibus fauce barbatis.  
*Gentiana* corollis hypocrateriformibus fauce barbatis.  
*Hort. cliff.* 81. *Fl. svec.* 203, 229. *Roy. lugdb.* 432.  
*Dalib. parif.* 81. *Mat. med.* 111.  
*Gentiana* floribus confertis faucibus membranula laciniata clausis, foliis ovato-acuminatis. *Guet. flam.* 303.  
*Gentiana pratensis*, flore lanuginoso. *Bauh. pin.* 188.  
*Gentiana autumnalis* ramosa. *Bauh. pin.* 188.  
*Opsantha*, *Reneal. spec.* 71.  
*Habitat in Europæ pratis.* ☉
- \* \* \* *Corollis non quinquefidis.*
- campestris.** 23. *GENTIANA* corollis quadrifidis fauce barbatis.  
*Gentiana* corolla hypocrateriformi tubo villis clauso, calycis foliis alternis majoribus. *Fl. lapp.* 94. *Fl. svec.* 203. *β.* 230.  
*Gentiana purpurea minima.* *Column. ecphr.* 1. p. 223. t. 221. *Barr. ic.* 97. f. 2.  
*Gentianella alpina* verna minor. *Bauh. pin.* 188.  
*Erythalla*, *Renealm. spec.* 72.  
*Habitat in Europæ pratis ficcis.* ☉  
*Præcedenti nimis affinis* ♂ fere varietas aut hybrida.
- ciliata.** 24. *GENTIANA* corollis quadrifidis margine ciliatis.  
*Gentiana* corollis hypocrateriformibus laciniis margine barbatis *Hort. cliff.* 81.  
*Gentiana angustifolia* autumnalis minor, floribus ad latera pilosis. *Bauh. pin.* 188.  
*Gentianella cærulea* oris pilosis. *Bauh. pin.* 108.  
*Gentianella cærulea* fimbriata angustifolia autumnalis.  
*Column. ecphr.* 1. p. 222. t. 221. f. 1.  
*Habitat in Helvetiæ, Italiæ, Canadæ montibus.*
- Cruciata.** 25. *GENTIANA* corollis quadrifidis imberbibus, floribus verticillatis sessilibus.  
*Gentiana* floribus confertis terminalibus, corollis quadrifidis imberbibus interjecto denticulo. *Hort. cliff.* 81. *Roy. lugdb.* 432. *Dalib. parif.* 80.

Gen-

*Gentiana cruciata. Bauh. pin. 188.*

*Gentiana minor. Cam. epit. 417.*

*Tretorrhiza. Renealm. spec. 74. t. 73.*

*Habitat in Pannoniæ, Apenninorum, Helvetiæ montosis apricis, ad vias, in sterilibus. 24.*

*Corollæ denticuli bifidi sunt.*

26. *GENTIANA* corollis quadrifidis, floribus acaulis, sessilibus, foliis ovatis.

*Gentianoides flore luteo. F. will. peruv. 3. p. 20. t. 14.*

*Habitat in Chili.*

27. *GENTIANA* corollis quadrifidis imberbibus, caule dichotomo filiformi.

*Gentiana* corollis infundibuliformibus quadrifidis, pedunculis ternis, foliis linearibus. *Sauv. monsp. 132.*

*Gentiana* caule dichotomo, foliis lineari-lanceolatis, floribus infundibuliformibus quadrifidis longissime pedunculatis. *Guett. stamp. 1. p. 305. Dalib. parif. 82.*

*Centaurium palustre luteum minimum. Raj. hist. 1092.*

*Vaill. parif. 32. t. 6. f. 3.*

*Centaurium pusillum luteum. C. B. 278. Magn. monsp. 232.*

*Habitat in Gallia. ☉*

28. *GENTIANA* corollis octifidis, foliis perfoliatis. perfoliata, *Læfl. hisp. 33.*

*Gentiana* caule dichotomo, foliis connatis, corollis octifidis. *Hort. cliff. 81. E 496. Roy. lugdb. 433.*

*Dalib. parif. 82. Sauv. monsp. 136. Læfl. it. 133. \**

*Centaurium luteum perfoliatum. Bauh. pin. 278.*

*Morif hist. 2. p. 565. t. 5. f. 26. f. 1, 2.*

*Centaurium parvum, flore luteo. Clus. hist. 2. p. 180.*

*Chlora. Reneal. spec. 80. t. 76.*

*Habitat in Gallia, Anglia, Hispania, oriente. ☉*

## PHYLLIS.

1. *PHYLLIS* stipulis dentatis. *Roy. lugdb. 92.*

nobla.

*Phyllis. Hort. cliff. 87. Hort. upf. 57.*

*Valerianella canariensis frutescens Simpla nobla dicta.*

*Dill. elth. 405. t. 299. f. 386.*

*Bupleuroides quæ Arbor umbellifera. Walth. hort. 11. t. 6.*

*Habitat in Canariis. ‡*

2. *PHYL-*



indica.

2. **PHYLLIS** stipulis integerrimis. *Roy. lugd.* 92. †  
*Habitat in India.* ‡  
*Flores hujus mihi ignoti sunt.*

## ERYNGIUM.

foetidum.

1. **ERYNGIUM** foliis radicalibus subensiformibus serratis; floralibus multifidis, caule dichotomo.  
*Eryngium foliis gladiatis utrinque laxe serratis, denticulis subulatis.* *Gron. virg.* 30.  
*Eryngium foetidum, foliis inferioribus angustis serratis; superioribus laciniatis & aculeatis.* *Brown. jam.* 185.  
*Eryngium americanum foetidum.* *Herm. lugdb.* 236. t. 237.  
*Eryngium, foliis angustis serratis, foetidum.* *Sloan. jam.* 127. *hist.* 1. p. 264. t. 156. f. 3, 4.  
*Habitat in Virginia, Jamaica, Mexico, Surinamo.* 24  
*Planta foetida, antihysterica, Serpentina adversa. Capitula cylindrica.*

aquaticum.

2. **ERYNGIUM** foliis gladiatis serrato-spinosis: floralibus indivisis, caule simplici.  
*Eryngium foliis gladiatis utrinque laxe serratis, denticulis subulatis.* *Hort. cliff.* 88. *Roy. lugdb.* 529.  
*Eryngium foliis gladiatis utrinque laxe serratis: summis tantum dentatis subulatis.* *Gron. virg.* 146.  
*Eryngium americanum, yuccæ folio, spinis ad oras molliusculis.* *Pluk. alm.* 13. t. 175. f. 4. *Raj. suppl.* 239. *Morif. hist.* 3. p. 167. f. 7. t. 37. f. 21.  
*Scorpii spina.* *Hern. mex.* 222.  
 3. *Eryngium lacustre virginianum, floribus, ex albido cæruleis, caule & foliis ranunculi flammæ minoris.* *Pluk. alm.* 137. t. 396. f. 3.  
*Habitat in Virginia.* 24  
*Facies & folia Brometiæ minoris, sed hæc ciliata spinis capillaribus flexilibus mitibus. Paleæ Florum majores sunt.*

planum.

3. **ERYNGIUM** foliis radicalibus ovalibus planis crenatis, capitulis pedunculatis. *Hort. cliff.* 87. *Hort. upf.* 37. *Roy. lugdb.* 92. *Dalib. parif.* 83.  
*Eryngium latifolium planum.* *Bauh. pin.* 87.  
*Eryngium pannonicum latifolium.* *Cus. hist.* 2. p. 158.  
*Habitat in Russia, Polonia, Austria, Helvetia.* 24  
 4 ERYN-

4. **ERYNGIUM** foliis radicalibus oblongis incis, cau- pusillum.  
le dichotomo, capitulis sessilibus. *Hort. cliff.* 87.  
*Roy. lugdb.* 93.  
*Eryngium planum minus.* *Bauh. pin.* 386.  
*Eryngium pusillum planum moutoni.* *Clus. hist.* 2. p.  
158.  
*Habitat in Hispania, Oriente.*
5. **ERYNGIUM** foliis radicalibus subrotundis plicatis maritimum.  
spinosis capitulis pedunculatis. *Hort. cliff.* 87. Fl.  
*succ.* 220, 233. *Roy. lugdb.* 93.  
*Eryngium maritimum.* *Bauh. pin.* 386.  
*Eryngium marinum.* *Clus. hist.* 2. p. 169. *Cam. epit.*  
448.  
*Habitat ad Europæ littora arenosa maritima.*
6. **ERYNGIUM** foliis radicalibus cordatis oblongis, alpinum.  
caulinis pinnatifidis, capitulis subcylindricis.  
*Eryngium alpinum cæruleum, capitulis diffaci.* *Bauh.*  
*pin.* 386.  
*Eryngium cæruleum genevense.* *Lob. ic.* 2 p. 23.  
*Habitat in alpinis Helvetiæ, Italiæ.* 24
7. **ERYNGIUM** foliis radicalibus cordatis, caulinis tri- uspida-  
palmatis auriculis retroflexis, paleis tricuspidadis. tum.  
*Amæn. acad.* 3. p. 405. \*  
*Eryngium foliis radicalibus ovatis crenatis petiolatis,*  
*capitulis pedunculatis.* *Gron. orient.* 76.  
*Eryngium syriacum ramosius, capitulis minoribus cæ-*  
*ruleis.* *Morif. hist.* 3. p. 166. f. 7. t. 37. f. 13.  
*Eryngium capitulis psyllii.* *Bocc. sic.* 80.  
*Habitat in Hispania, Sicilia, Oriente.* ♂
8. **ERYNGIUM** foliis amplexicaulibus pinnato-lacinia- campestre.  
tis. *Hort. cliff.* 87. *Mat. med.* 114. *Roy. lugdb.* 93.  
*Eryngium vulgare.* *Bauh. pin.* 386.  
*Eryngium campestre vulgare.* *Clus. hist.* 2. p. 157.  
*Habitat in Germaniæ, Galliæ, Hispaniæ, Italiæ incul-*  
*tis.* 24
9. **ERYNGIUM** foliis trifidis basi subpinnatis. amethysti-  
*Eryngium montanum amethystinum.* *Bauh. pin.* 386. num.  
*Morif. hist.* 3. p. 165. f. 7. t. 25. f. 2.  
*Eryngium totum cæruleum.* *Besl. cyst.* 0. 11. t. 8. f. 4.
- β. **Eryngium** minus trifidum hispanicum. *Barr. ic.* 36.  
*Bocc. mus. t.* 71.

*Habitat in Styriæ montibus. 2.*

*Folia linearia, semirifida, inferne subpinnata. Involucrum basi obvallatum setis subulatis interjectis.*

*Varietas β minor, involucri foliolis trifidis distincta.*

## HYDROCOTYLE.

valgaris.

1. **HYDROCOTYLE** foliis peltatis, umbellis quinquefloris.

*Hydrocotylè foliis peltatis orbiculatis undique emarginatis. Hort. cliff. 88. Fl. fvec. 221, 234. Roy. lugdb. 93. Hall. helv. 425. Dalib. parif. 83.*

*Ranunculus aquaticus, cotyledonis folio. Bauh. pin. 180.*

*Cotyledon aquatica. Lob. ic. 387.*

*Habitat in Europæ inundatis. 2.*

umbellata.

2. **HYDROCOTYLE** foliis peltatis, umbellis multifloris. *Brown. jam. 185.*

*Hydrocotyle repens, flore albo, cotyledonis folio. Gron. virg. 30.*

*Hydrocotyle maxima, folio umbilicato, floribus in umbellam nascentibus. Plum. spec. 7 ?*

*Cotyledon aquatica. Sloan. jam. 93. hist. 1. p. 212.*

*Erva de capitaon. Marcgr. bras. 27.*

*Acaricoba. Pif. bras. 90.*

*Habitat in America.*

*Planta simillima precedenti, hinc cum ea mixta; differt scapo foliis duplo longiore, flosculis non 5, sed pluribus quam viginti, in umbellam simplicem digestis.*

americana.

3. **HYDROCOTYLE** foliis reniformibus sublobatis crenatis. *Læfl. it. 281.*

*Habitat in America septentrionali & meridionali.*

*Plantæ facies primæ & magnitudo. Folia antice sine vix hiantæ ad centrum usque dissecta, margine in 9 lobos obsoletos divisa, quorum singuli tribus crenis minoribus notata. Umbella quinqueflora.*

asiatica.

4. **HYDROCOTYLE** foliis reniformibus æqualiter crenatis. *Hort. cliff. 88. Fl. zeyl. 118. Roy. lugdb. 94.*

*Valerianella zeylanica palustris repens, hederæ terrestris folio. Herm. par. 238. t. 238.*

*Ranunculo affinis umbelliferis accedens, chelidonii minoris folio, zeylanica minor. Pluk. alm. 314. t. 106. f. 5.*

*Pes equinus*. *Rumph. amb.* 5. p. 455. t. 169. f. 1.

*Codagam*. *Rheed. mal.* 10. p. 91. t. 46.

*Habitat in India.*

*Differt a præcedente marginis denticulis f. crenis equalibus, consistentia foliorum duplo crassiore & fere incana; petiolis pluribus ad genicula caulis, cum in præcedente folia solitaria.*

5. HYDROCOTYLE foliis linearibus, umbellis multifloris. chinensis.

*Habitat in China.*

*Magnitudo præcedentium; Caulis repens. Folia linearia, levia, obtusa, plana, sæpius bina ad articulos. Scapi longitudine folii. Umbella multiflora.*

## SANICULA.

1. SANICULA foliis radicalibus simplicibus, flosculis omnibus sessilibus. *Fl. succ.* 222, 235. *Mar. med.* 115. europæa.

*Sanicula flosculis omnibus sessilibus. Gron. virg.* 147.

*Sanicula. Hort. cliff.* 88. *Roy. lugdb.* 93.

*Sanicula officinarum. Bauh. pin.* 319.

*Diapensia. Cam. epit.* 763.

*Habitat in Europæ sylvis montosis.* 24

2. SANICULA foliis radicalibus compositis: foliolis ovatis. *Gron. virg.* 146. canadensis

*Sanicula canadensis, amplissimo laciniato folio. Tournef. inst.* 326.

*Habitat in Virginia.* 24

*Structura ita præcedentis, ut vix differentia, sed planta decuplo sæpe omnibus partibus major.*

3. SANICULA flosculis masculis pedunculatis, hermaphroditis sessilibus. *Gron. virg.* 31. marilandica

*Sanicula foliis septilobatis inæqualibus, flosculis masculis pedunculatis. Hort. upf.* 57.

*Sanicula marilandica, caule & ramulis dichotomis, echinis minimis in eodem communi pediculo ternis.*

*Raj. suppl.* 260.

*Habitat in Marilandia, Virginia.* 24

## ASTRANTIA.

1. ASTRANTIA. foliis quinquelobis: lobis trifidis. major. *Hall. helv.* 439.

Y 2

Astran-

*Astrantia. Hort. cliff. 88.*

*Helleborus niger, faniculæ folio. Bauh. pin. 186.*

*Veratrum nigrum. Dod. pempt. 387.*

*Habitat in alpinis Helvetiæ, Hetruriæ, Bohemiæ, Pyrenæis. 2.*

minor.

2. **ASTRANTIA** foliis digitatis serratis.

*Astrantia foliis digitatis septenis integris dentatis. Hall. helv. 440.*

*Helleborus, faniculæ folio, minor. Bauh. pin. 186. prodr. 97. Raj. hist. 475.*

*Helleborus minimus alpinus, astrantiæ flore. Bocc. sic. 10. t. 6.*

*Habitat in Alpibus Helvetiæ, Hetruriæ, Pyrenæis. 2.*

## BUPLEURUM.

\* *Herbacea.*

rotundifolium.

1. **BUPLEURUM** involucris universalibus nullis, foliis perfoliatis. *Hort. upf. 64. Mat. med. 136.*

*Bupleurum foliis ovatis perfoliatis. Hort. cliff. 104. Roy. lugdb. 109. Hall. helv. 437.*

*Perfoliata vulgatissima arvensis. Bauh. pin. 277.*

*Perfoliata. Dod. pempt. 104.*

β. *Perfoliata minor, ramis inflexis. Bauh. pin. 277. prodr. 130.*

*Perfoliata flore multiplici. Bauh. pin. 277.*

*Perfoliata crispa f. muscosa. Cam. hort. t. 37.*

*Habitat inter Europæ australis segetes. ☉*

stellatum.

2. **BUPLEURUM** involucellis coadunatis, universali triphylo.

*Bupleurum alpinum, angustis nervosis foliis, calyce universali leviter inciso. Hall. helv. 438. t. 10.*

*Perfoliata alpina angustifolia media. Bauh. pin. 277. prodr. 129. Burf. VI. 5.*

*Habitat in alpinis Helveticis.*

*Folia radicalia linearia; summa ovata, lanceolata, sessilia; involucris universalis foliola ovata, equalia: partialis 5, coalita ultra medium.*

petrosum.

3. **BUPLEURUM** involucellis coadunatis, universali pentaphyllo.

*Bupleurum foliis angustis & nervosis, umbella universali quinquefolia, peculiari sexfidastellata. Hall. helv. 438.*

*Perfoliata alpina, gramineo folio. Bauh. pin. 277.*

Se-

*Sedum petraeum*, bupleuri folio. *Pon. bald.* 247.

*Habitat in alpidibus Helvetiae*, Baldi. 24.

Folia linearia, sessilia. Involucri universalis foliola 5, equalia, latiora; partialis 5, coalita ad medium.

4. **BUPLEURUM** involucellis pentaphyllis orbiculatis, universali triphylo ovato, foliis amplexicaulibus cordato-lanceolatis. angulosum.

Perfoliata alpina angustifolia major, folio anguloso.

*Bauh. pin.* 277. *prodr.* 129. *Burf. VI.* 4.

5. **Bupleurum** alpina angustifolia minor. *Bauh. pin.* 277. *prodr.* 129. *Burf. VI.* 6.

*Habitat in Pyrenæis. β. in Valfellæ alpidibus.* 24.

Planta affinis maxime 2, uti patuit ex *Burseri* collectione, licet involucri partialis foliola in hac distincta. In varietate β. folia angustissima striata, in α. vero lanceolata, nec nervoso-striata.

3. **BUPLEURUM** involucellis pentaphyllis ovatis: longifolium, universali subpentaphyllo, foliis amplexicaulibus. *Bupleurum* foliis imis petiolatis, superioribus ovatis perfoliatis. *Hall. helv.* 437.

Perfoliata montana latifolia. *Bauh. pin.* 277.

Perfoliata montana. *Cam. hort.* 120. t. 38.

*Habitat* Goettingæ, inque monte Jura Helvetiæ.

Folia ovata, profunde amplexicaulia: radicalia petiolata; Involucr. univers. 3. vel pentaphyllum; partiale pentaphyllum, longitudine flosculorum.

6. **BUPLEURUM** involucellis pentaphyllis acutis, universali subpentaphyllo, foliis lanceolatis, caule flexuoso. falcatum.

*Bupleurum* foliis radicalibus pedunculatis: summo subrotundo, caulinis linearibus sessilibus. *Gnett. stamp.* 59.

*Bupleurum* foliis petiolatis falcato-ellipticis acutis. *Hall. helv.* 439.

*Bupleurum* folio subrotundo f. vulgatissimum. *Bauh. pin.* 278.

*Bupleurum.* *Riv. pent.* 44.

*Auricula leporis umbella lutea.* *Bauh. hist.* 3. p. 200. f. 1. *Isophyllum.* *Cord. hist.* 69.

*Habitat in Misinæ, Valfellæ sepibus.* 24.

Involucrum universale pentaphyllum, sepe diphyllum, inaequale; partiale pentaphyllum, mucronatum. Umbelle parvæ. Caulis altus flexuosus.

odontites. 7. **BUPLEURUM** involucellis pentaphyllis acutis, universalis triphylo, flosculo centrali altiore, ramis divaricatis.

*Bupleurum* foliis gramineis, calyce peculiari quinquefido aristato rigido. *Hall. helv.* 438.

Perfoliata minor angustifolia, bupleuri folio. *Bauh. pin.* 277.

Perfoliata minima, bupleuri folio. *Column. ecphr.* 1. p. 84. t. 247.

*Habitat in alpinis Vallesiae; in Italiae rupibus, vineis.* ☉

*Folia lineari-lanceolata. Caulis ramosissimus, angulatus, divaricatus ramis basi nodosis. Involucrum univ. subpentaphyllum: foliolis latioribus rigidis venosis acutis; parziale pentaphyllum simile; flosculi 9: intermediu petiolus duplo altior.*

femicompositum. 8. **BUPLEURUM** umbellis compositis simulque simplicibus. *Amæn. acad.* 3. p. 405. \*

*Habitat in Hispania.* ☉

*Affine B. Odontiti, vix tamen pro varietate assumendum.*

ranunculoides. 9. **BUPLEURUM** involucellis pentaphyllis lanceolatis longioribus: universalis triphylo, foliis caulinis lanceolatis.

*Bupleurum* foliis amplexicaulibus, inferioribus linearibus, summo cordato-oblongo. *Hort. cliff.* 104. *Hall. helv.* 427. *Roy. lugdb.* 110.

Perfoliata alpina angustifolia minima. *Bauh. pin.* 277. *prodr.* 130. *Burf. XVI.* 8.

Perfoliatum gramineum helveticum alpinum pumilum, facie ranunculi. *Lob. illustr.* 137. *Park. theatr.* 333. t. 338.

*Habitat in Helvetia, & Pyrenæis.* 24

*Folia Radicula linearia, acuta; Caulina lanceolata, sessilia.*

rigidum. 10. **BUPLEURUM** caule dichotomo subnudo, involucris minimis acutis.

*Bupleurum* foliis lanceolatis rigidis: radicalibus obtusis, caulinis acutis. *Roy. lugdb.* 110. *Sauv. monsp.* 76.

*Bupleurum* folio rigido. *Bauh. pin.* 278.

*Bupleurum* alterum latifolium. *Dod. pempt.* 633.

*Habitat Monspelii.* 24

*Folia Radicalia lato-lanceolata, nervosa, rigida. Caulis dichotomus, subnudus: foliolis minutissimis, linearibus, acutis. Umbellæ terminales, bi-f. trifida, totidem foliolis acerosis. Umbellula parvæ, sex-floræ.*

11. **BUPLEURUM** umbellis simplicibus alternis pen-  
taphyllis subtrifloris.

*Bupleurum foliis linearibus acutis sessilibus. Hort. cliff.*

104. *Roy. lugdb.* 110.

*Bupleurum tertium minimum. Column. ecphr.* 1. p. 85.  
t. 247. *Morif. hist.* 3. p. 300. f. 9. t. 14. f. 4.

*Bupleurum angustissimo folio. Bauh. pin.* 278.

*Habitat in Germania, Anglia, Gallia, Italia.* ☉

*Caulis ramosissimus; Ramis alternis. Folia linearia, acuta. Umbellulæ axillares, parvæ.*

12. **BUPLEURUM** caule erecto paniculato, foliis li-  
nearibus, involucris triphyllis, involucellis penta-  
phyllis.

*Bupleurum annuum angustifolium. Magn. monsp.* 42.  
*Allion. nic.* 192.

*Bupleurum minus angustifolium monspeliense. Raj.*  
*hist.* 474.

*Bupleurum angustifolium. Dod. pempt.* 474.

*Habitat in Gallia, Italia.* ☉

*Caulis sexpedalis, junceus, lævis: Ramis alternis, erectis. Folia graminea, bifaria, lævia, lætescentia. Involucrum lanceolatum, longitudine umbellæ. Involucella minora. Umbella radiis 4. umbellula 5. f. 6 flosculis luteis.*

\* \* *Frutescentia.*

13. **BUPLEURUM** frutescens, foliis amplexicaulibus villosum  
dentatis subtus villosis.

*Perfoliata foliis oblongis sinuosis subtus incanis. Burm.*  
*af.* 196. t. 71. f. 2.

*Habitat in Æthiopia.* ☿

14. **BUPLEURUM** frutescens, foliis ovatis inter-  
fruticosa gerrimis.

*Bupleurum, foliis obverse ovatis in petiolum attenuatis.*

*Hort. cliff.* 104. *Roy. lugdb.* 109. *Sauv. monsp.* 60.  
*Gron. orient.* 77.

*Seseli æthiopicum, salicis folio. Bauh. pin.* 161.



- Sese'i æthiopicum, frutex. *Dod. pempt.* 312.  
*Habitat in Galliæ australis & Orientis saxosis maritimis.* ‡
- Frutice-  
scens. 15. **BUPLEURUM** frutescens, foliis linearibus, involucro universali partialique. *Amæn. acad.* 4. p. 269. \*  
*Læfl. ii.* 134. \*  
*Bupleurum hispanicum arborescens, gramineo folio.*  
*Tournef. inst.* 310.  
*Bupleurum fruticans angustifolium hispanicum.* *Barr.*  
*rar.* 623. t. 1255.  
*Habitat in Hispaniæ collibus.* ‡
- difforme. 16. **BUPLEURUM** frutescens, foliis vernalibus decompositis planis incis, æstivalibus filiformibus angulatis trifidis.  
*Bupleurum foliis radicalibus pinnato-incis, caulinis fasciculatis tetragonis.* *Roy. lugdb.* 110.  
*Bupleurum frutescens, foliis ex uno puncto plurimis junceis tetragonis.* *Burm. afr.* 195. t. 71. f. 1?  
*Habitat in Æthiopia.* ‡  
*Folia hyemalia Petroselini, æstivalia Spartii. Involucrum foliolis subulatis, pluribus, basi connexis, marcescentibus; involucella foliolis quinis.*

## ECHINOPHORA.

- spinosa. 1. **ECHINOPHORA** foliolis subulato-spinosis integerrimis.  
*Echinophora foliis decompositis.* *Wach. ultr.* 200.  
*Caucalis caule lignoso, foliolis subulato-spinosis integerrimis.* *Roy. lugdb.* 96. *Sauv. monsp.* 258.  
*Crithmum maritimum spinosum.* *Bauh. pin.* 288.  
*Crithmum spinosum.* *Dod. pempt.* 705.  
*Habitat ad littora maris, præsertim mediterranei.* 24
- tenuifolia. 2. **ECHINOPHORA** foliolis incis inermibus.  
*Echinophora foliis supradecompositis.* *Wach. ultr.* 200.  
*Caucalis caule lignoso, foliis incis.* *Roy. lugdb.* 96.  
*Pastinaca sylvestris angustifolia, fructu echinato.*  
*Bauh. pin.* 151.  
*Pastinaca echinophora apula.* *Column. ecphr.* 1. p. 98.  
t. 101.  
*Habitat in Apuliæ maritimis apricis, salsis.* 24

## TORDYLIUM.

1. **TORDYLIUM** involucris umbella longioribus. *Hort. syriacum. cliff. 90. Hort. upf. 58. Roy. lugdb. 95. Sauv. meth. 230. Gron. orient. 78.*  
*Tordyllum syriacum humilius, semine granulato majore. Moris. hist. 3. p. 317. f. 9. t. 16. f. 7.*  
*Gingidium foliis pastinacæ latifoliæ. Bauh. pin. 151.*  
*Habitat in Syria. ☉*
2. **TORDYLIUM** involucris partialibus longitudine officinale. florum, foliis ovatis laciniatis. *Hort. cliff. 90. Hort. upf. 58. Roy. lugdb. 95. Mat. med. 145. Sauv. monsp. 230.*  
*Seseli creticum minus. Bauh. pin. 161.*  
*Seseli creticum. Dod. pempt. 314.*  
*Habitat in Narbona, Italia, Sicilia. ☉*
3. **TORDYLIUM** umbellulis remotis, foliis pinnatis: apulum. pinnis subrotundis laciniatis. *Hort. cliff. 90. Roy. lugdb. 94.*  
*Tordyllum apulum minimum. Column. ecphr. 1. p. 122. t. 124.*  
*Seseli creticum minimum. Bauh. pin. 161.*  
*Habitat in Italiæ, Apuliæ incultis.*  
*Umbellulæ singulæ flosculi marginales tres majores f. radiati.*
4. **TORDYLIUM** umbellis confertis radiatis, foliolis maximum. lanceolatis inciso-ferratis. *Hort. cliff. 90. Roy. lugdb. 94. Sauv. monsp. 230, 259.*  
*Tordyllum. Riv. pent. 1.*  
*Caucalis maxima, sphondylii aculeato femine. Bauh. pin. 152.*  
*Caucalis major. Clus. hist. 2. p. 101.*  
*Seseli creticum majus. Bauh. pin. 161.*  
*Habitat in Italiæ ruderalis sepibus.*
5. **TORDYLIUM** umbellis confertis nudiusculis; foliolis latifolium. liis pinnatis: foliolis lanceolatis inciso-ferratis. *Mill. dist. t. 8.*  
*Caucalis umbella universali trifida: partialibus pentaspermis, foliis pinnatis ferratis. Hort. cliff. 91. Roy. lugdb. 95. Dalib. parif. 84. Sauv. monsp. 236. Gron. orient. 80.*  
*Caucalis arvensis echinata latifolia. Bauh. pin. 152.*

*Lappula canaria latifolia. Bauh. hist. 3. p. 81.*

*Habitat in Gallia, Italia, Oriente inter segetes. ☉*

**Anthriscus.** 6. **TORDYLIUM** umbellis confertis, foliolis ovato-lanceolatis pinnatifidis. *Hort. cliff. 90. Fl. fvec. 224, 236. Roy. lugdb. 94.*

*Caucalis umbella conferta, foliolis ovato-lanceolatis pinnatifidis. Guett. stamp. 1. p. 81.*

*Caucalis semine aspero, flosculis rubentibus. Bauh. pin. 152. prodr. 80.*

*Caucalis minor, flore rubente. Moris. hist. 3. p. 308. f. 9. t. 14. f. 8.*

*Habitat in Europæ septentrionalis arvis ruderalis. ♂*

**nodosum.** 7. **TORDYLIUM** umbellis simplicibus sessilibus, seminibus exterioribus hispidis.

*Caucalis umbellis sessilibus simplicibus. Hort. upf. 58.*

*Hort. cliff. 91. Roy. lugdb. 95. Dalib. parif. 85.*

*Caucalis nodosa, echinato semine. Bauh. pin. 152. prodr. 80. Bauh. hist. 3. p. 83.*

*Habitat in Gallia, Italia ad vias. ☉*

## CAUCALIS.

**grandiflora.** 1. **CAUCALIS** involucris pentaphyllis: foliolo unico duplo majore. *Hort. cliff. 91. Roy. lugdb. 95. Sauv. monsp. 258.*

*Caucalis involucro universali pinnatifido, partialibus indivisis, umbella multifida. Fl. fvec. 225. Dalib. parif. 84.*

*Caucalis arvensis echinata, magno flore. Bauh. pin. 152.*

*Echinophora flore magno. Riv. pent. 25.*

*Echinophora pycnocarpus. Column. ecphr. 1. p. 91. t. 94.*

*Habitat in Europa australiori inter segetes.*

**dauceoides.** 2. **CAUCALIS** involucro universali longitudine umbellæ lanceolato.

*Caucalis dauci sylvestris folio, echinato magno fructu.*

*Magn. monsp. 292.*

*Habitat Montpellier. D. Sauvages.*

*Folia supradecomposita Dauci vulgaris. Involucrum universale pentaphyllum: foliolis lanceolatis, longitudine totius umbellæ, acuminatis. Involucra partialia lato-lanceolata, lateribus membranacea, acuta. Flo-*

*Flosculi copiosi albi : hermaphroditi 3 in radio ; flosculis disci , f. masculis parvis , æqualibus ; radii flosculis petalis maximis bipartitis albis. In præcedente involucrum umbella dimidio brevius 3 radii petala rotundiora.*

3. CAUCALIS involucro universali monophyllo, partialibus triphyllis. *Mähr. E. N. C. 1742. vol. 6. p. 401. †* mauritanica

*Caucalis mauritanica vulgari similis, sed semine majore. Walth. hort. 127.*

*Habitat in Mauritania.*

4. CAUCALIS umbellis trifidis umbellulis trispermis, involucris triphyllis. *Hort. cliff. 61. Hort. upf. 58. Roy. lugdb. 95. Sauv. monsp. 258. platycarpus*

*Caucalis monspelica echinata, magno fructu. Bauh. pin. 153.*

*Echinophora altera asperior platycarpus. Col. ecphr. 1. p. 95. t. 94.*

*Habitat in Italia, Monspelli. ☉*

5. CAUCALIS umbellis patentibus, foliolis partialibus supradecompositis laciniatis : lacinulis linearibus. *Roy. lugdb. 95. orientalis.*

*Caucalis orientalis altissima, folio ferulæ. Tournef. cor. 23.*

*Habitat in Oriente. ♂*

6. CAUCALIS feminum aculeis triglochidibus uncinatis : pillis verticillatis hispidis. *Guett. stamp. 80. leptophylla.*  
*Dalib. parif. 85.*

*Caucalis involucro universali monophyllo, laciniis foliorum perangustis. Hall. helv. 448.*

*Caucalis arvensis echinata, parvo flore & fructu. Bauh. pin. 152.*

*Echinophora tertia leptophyllon purpurea. Column. ecphr. 1. p. 96. t. 97.*

*Habitat in Gallia, Italia, Anglia. ♂*

## ARTEDIA.

1. ARTEDIA seminibus squamatis. *Hort. cliff. 89. squamata.*  
*Hort. upf. 58. Roy. lugdb. 96. Gron. orient. 81.*

*Thapsia orientalis, anethi folio, semine eleganter creato. Tournef. cor. 22. itin. 3. p. 298.*

Gin-

*Gingidium foeniculi folio. Bauh. pin. 151. Raj. hist. 448. 466.*

*Gingidium rauwolfii. Cam. hort. 67. t. 16.*

*Habitat in Libano. ☉*

### DAUCUS.

Carota.

1. DAUCUS seminibus hispidis, petiolis subtus nervosis. *Hort. cliff. 89. Hort. ups. 59. Fl. suec. 223, 257. Mat. med. 142. Roy. lugdb. 97. Gron. virg. 31. Dalib. parif. 85.*

*Daucus vulgaris. Clus. hist. 2. p. 198.*

*Pastinaca tenuifolia sylvestris diolcoridis. Bauh. pin. 151.*

- β. *Pastinaca tenuifolia sativa, radice lutea. Bauh. pin. 151.*

- γ. *Daucus sativus, radice atrorubente. Tournef. inst. 307.*

*Habitat in Europæ campis exaridis. ♂*

Mauritanicus.

2. DAUCUS seminibus hispidis, flosculo centrali sterili carnosio, receptaculo communi hemisphærico.

*Daucus hispanicus, umbella magna. Tournef. inst. 308.*

*Pastinaca tenuifolia sylvestris, umbella majore. Bauh. pin. 151?*

*Habitat in Hispania, Italia, Mauritania. ☉*

*Similis D. Carotæ, sed Umbella amplior; universalis non ad basin divisa, verum ipsa basi hemisphærico-folida. Umbellæ in centro flosculus sterilis, carnosus, atropurpureus, reliquis longe major. Foliorum petioli striati nervis albis. Caulis magis hirtus.*

Vismaga.

2. DAUCUS seminibus nudis. *Hort. cliff. 89. Roy. lugdb. 97. Sauv. monsp. 257. Gron. orient. 83.*

*Gingidium umbellæ oblonga. Bauh. pin. 151.*

*Gingidium alterum. Dod. pempt. 792.*

*Habitat in Europa australi, Mauritania, Libano.*

Gingidium.

4. DAUCUS radiis involucri planis: laciniis recurvis. *Roy. lugdb. 97.*

*Daucus montanus lucidus. Tournef. inst. 307.*

*Gingidium folio chærophylli. Bauh. pin. 151.*

*Gingidium. Matth. comm. 372. t. 373.*

*Pastinaca tenuifolia marina, foliis obscure virentibus & quasi lucidis. Magn. monsp. 199.*

*Pastinaca folio cœnanthes. Bocc. sic. 74.*

*Habitat Monspelii.*

5. DAUC-

## 5. DAUCUS seminibus aculeatis.

muricatus.

*Artedia muricata* seminibus aculeatis. *Hort. cliff.* 89.*Sp. pl.* 1. p. 242.*Caucalis major daucoides tingitana.* *Morif. hist.* 3. p. 308. f. 9. t. 14. f. 4. *Raj. hist.* 469. *Herm. par.* 111. t. III.*Echinophora tingitana.* *Riv. pent.* 27.β. *Caucalis pumila maritima.* *Bauh. pin.* 153.

maritimus.

*Lappula canaria* f. *Caucalis maritima.* *Bauh. hist.* 3. p. 81.*Habitat in Mauritania.* β. in littoribus maris Mediterranei. ♂. ☉*Simillimus D. Carotæ* Caule, Foliis, Involucris, Involucellis, Umbella per maturitatem contracta; Flosculi umbellularum 3. f. 5. fertiles, reliquis sterilibus; Flores non radiati. Semina cristis 4, longitudinalibus, valde dentatis, ante maturitatem rubris glochidibus.

## AMMI.

1. *Ammi foliis inferioribus pinnatis lanceolatis ferratis, superioribus multifidis linearibus.* *Hort. upf.* 59.*Ammi lacinii foliorum caulis lanceolatis.* *Hort. cliff.*89. *Roy. lugdb.* 96. *Sauv. monsp.* 232.*Ammi majus.* *Bauh. pin.* 159.*Ammi vulgare.* *Dod. pempt.* 415.*Habitat in Europæ australis, Orientis vineis, arvis.*

glaucofolium.

2. *AMMI foliorum omnium lacinulis lanceolatis.* *Guett.**flamp.* 2. p. 433. \*†*Ammi petræum glaucifolium perenne.* *Morif. hist.* 3. p. 295.*Daucus petræus glaucifolius.* *Bauh. hist.* 3. p. 58.*Habitat in Gallia.* 24

## BUNIMUM.

1. *BUNIMUM.* *Hort. cliff.* 91. *Roy. lugdb.* 107.*Bulbocastanum.**Bunium bulbo globoso.* *Sauv. monsp.* 256.*Bulbocastanum majus, folio apii.* *Bauh. pin.* 162.*Nucula terrestris.* *Lob. hist.* 429.*Habitat in Germaniæ, Angliæ, Galliæ saxosis.*

## CONIUM.

1. *CONIUM* seminibus striatis. *Hort. cliff.* 92. *Fl. suec. maculatum.*226, 238. *Roy. lugdb.* 107.

Cicu.

*Cicuta major.* *Bauh. pin.* 160.

*Cicutaria major vulgaris.* *Clus. hist.* 2. p. 200.

*Habitat in Europæ cultis, agris, ruderatis.* ♂

africanum. 2. *CONIUM* feminibus aculeatis. *Hort. cliff.* 92. \*

*Roy. lugdb.* 107.

*Caucalis africana, folio minore rutæ.* *Barh. lugdb.* 1.

p. 63. t. 63.

*Habitat in Africa.* ☉

*Planta glauca ut Sefeli, foliis Fumariæ. Umbella quadrisida: involucro lanceolato, brevi, margine membranaceo: Umbellulæ multifloræ: involucellis pentaphyllis, brevibus, acutis. Petala alba, equalia, intus carinata, non emarginata. Flosculi nonnulli masculi. Semina oblonga, muricata, magis gibba quam lata, stylis persistentibus.*

Royenl. 3. *CONIUM* feminibus radiato-spinosis. *Roy. lugdb.* 107.

*Habitat* - - - - -

## SELINUM.

sylvestre. 1. *SELINUM* radice fusiformi multiplici. *Hort. cliff.*

93. *Hort. upf.* 59. *Roy. lugdb.* 106.

*Apium sylvestre, lacteo succo turgens.* *Bauh. pin.* 153.

*Tyffeliolum plinii.* *Lob. hist.* 409. *Morif. hist.* 3. p. 319. f. 9. t. 17. f. 2.

*Habitat in Hircynia, Gallia.* 24

palustre. 2. *SELINUM* sublaetescens, radice unica. *Hall. helv.* 443.

*Selinum foliolis & lacinulis oblongo-linearibus.* *Guett. stamp.* 2. p. 70.

*Selinum palustre lactescens.* *Fl. lapp.* 110. *Fl. svec.* 227, 239. *Hort. cliff.* 92. *Roy. lugdb.* 116.

*Sefeli palustre lactescens.* *Bauh. pin.* 162. *prodr.* 85.

*Habitat in Europæ septentrionalis paludibus.* 24

Carvifolia. 3. *SELINUM* caule sulcato acutangulo, involucro universali caduco, pistillis fructus reflexis.

*Selinum foliolis pinnatim laciniatis.* *Gmel. sibir.* 1. p. 204. t. 48.

*Angelica tenuifolia.* *Riv. pent.* *Rupp. jen.* 277.

*Dill. giff.* 137. \*

*Carvifolia.* *Bauh. pin.* 158.

*Ha-*

*Habitat in Sibiriae campis a Jacco ad Obium, in Germania. 24.*

*Caules multi, erecti, tripedales, sulcati, angulis acutis. Folia supradecomposita: foliolis pinnatifidis, mucronatis. Pedunculus sub umbella fere crispatus. Involucrum universale deciduum. Petala alba, non subtus rubra. Semina medio 3 sulcis, Stylis longis reflexis.*

4. **SELINUM** umbellis confertis, involucrio universali reflexo, seminum costis quinque membranaceis. monnieri.

*Amœn. acad. 4. p. 269. \**

*Habitat in Gallia australi. ☉*

*Laserpitii structura in Seminibus; sed habitus ☿ reliqua Selini.*

### ATHAMANTA.

1. **ATHAMANTA** foliis bipinnatis planis, umbellis hemisphaericis, seminibus hirsutis. *Fl. suec. 229, 240.\** Libanotis.

*Libanotis pinnulis circa costam decussatis. Hall. herb. 451. opus. 250.*

*Libanotis minor, apii folio, minor. Bauh. pin. 157. prodr. 77.*

*Daucus montanus, pimpinellæ saxifragæ hircinæ folio. Pluk. alm. 129. t. 173. f. 1.*

*Habitat in Sveciæ, Germaniæ pratis siccis apricis. 24.*

2. **ATHAMANTA** foliis subbipinnatis: foliolis deorsum imbricatis, umbellis lenticularibus. condensata.

*Habitat in Sibiria. 24.*

*Radix perennis. Caulis simplex, pedalis, levis, inferne angulatus, superne sulcatus, præsertim pedunculo. Rami duo vel tres, floriferi, ex summis alis foliorum. Petioli canaliculati, basi membranacei, subtus angulati. Folia subbipinnata, foliolis alternim pinnatifidis lobo infimo majore, unde quasi cruciatæ evadunt: oppositiones ad basin ☿ inferior lacinia se extendit supra folium inferius; subtus nitida. Umbella densissima, utrinque convexa, antequam floret rufescens; flores tamen albi, exceptis antheris fuscis. Petala inflexa. Involucrum Universale sæpe deficit ☿ tantum rudimenta ostendit. Partiale longitudine florum. Foliola sunt imbricata, nec remota. Germina villosa, nec hirta. Antheræ ☿ receptacula stesculorum purpurea.*



Cervaria.

3. **ATHAMANTA** foliolis pinnatis decussatis inciso-angulatis, seminibus nudis.

*Athamanta* foliis pinnatis pinnatifidis: radicalibus decussatis. *Gmel. fibir.* 2. p. 186.

*Selinum* foliis radicalibus ovatis inæqualiter serratis.

*Hort. cliff.* 92. *Hort. upf.* 59.

*Daucus montanus*, apii folio, major. *Bauh. pin.* 150.

*Dauci* tertium genus. *Fuch. hist.* 233.

*Cervaria. Riv. pent.* 12.

- β. *Selinum* foliis simpliciter pinnatis inciso-angulatis.

*Hort. upf.* 60. \* *Sp. pl.* 1. p. 244.

*Habitat in* Helvetiæ, Austriæ, Alsatæ, Gallo-provinciæ, Sibiricæ montibus. 24

Oreofellum.

4. **ATHAMANTA** foliolis divaricatis. *Fl. svec.* 2. n. 241.

*Selinum* foliolis ovato-acutis acute serratis & incis. *Hort. cliff.* 92. *Roy. lugdb.* 106.

*Selinum* foliolis lacinulisque oblongo-linearibus. *Guet. flump.* 1. p. 70.

*Selinum* pinnis ad angulos obtusos pinnatis: pinnulis incis. non serratis. *Hall. helv.* 444. *Fl. svec.* n. 228.

*Apium montanum*, folio ampliore. *Bauh. pin.* 153.

*Apium montanum nigrum.* *Bauh. pin.* 153. *Segu. ver.* 2. p. 31.

*Oreofelinum. Clus. hist.* 2. p. 195. *Riv. pent.* 8.

*Habitat in* Germaniæ, Galliæ, Angliæ collibus apicis. 24

ficula.

5. **ATHAMANTA** foliis inferioribus nitidis, umbellis primordialibus subsessilibus, seminibus pilosis. *Hort. upf.* 60.

*Athamanta* foliolis multifidis planis, seminibus villosis. *Hort. cliff.* 93. *Roy. lugdb.* 97.

*Daucus secundus ficulus*, sophiæ folio. *Zan. hist.* 80.

*Habitat in* Sicilia. 24

*Folia inferiora nitida, superiora subvillosa. Caulis pubescens.*

cretensis.

6. **ATHAMANTA** foliolis linearibus planis hirsutis, petalis bipartitis, seminibus oblongis hirsutis. *Mat. med.* 143.

*Athamanta* foliis capillaribus, seminibus hirsutis. *Hort. cliff.* 93. *Roy. lugdb.* 97.

*Libanotis* foliis tenuissime pinnatis laciniis petiolatis. *Hall. helv.* 451. *opusc.* 241.

Dau-

- Daucus alpinus*, multifido longoque folio. *Bauh. pin.* 150.  
*Daucus creticus*, semine hirsuto. *Bauh. hist.* 3. p. 56.  
*Daucus foliis foeniculi tenuissimis*. *Bauh. pin.* 150.  
*Daucus creticus*. *Cam. epit.* 536. *Hall. opusc.* 242.  
*Habitat in Helvetia.* 21
7. *ATHAMANTA* foliis multipartitis: laciniis linearibus teretiusculis acuminatis. *Roy. lugdb.* 98.  
*Myrrhis annua*, semine striato villosa, incana. *Morif. umb.* 67.  
*Myrrhis tenuifolia annua cretica*, semine lanugine alba pubescente, *Morif. hist.* 3. p. 302.  
*Habitat in Creta.* ☉
8. *ATHAMANTA* seminibus membranaceo-striatis, foliis supradecompositis lævibus multifidis. chinensis.  
*Habitat - - - Chinensem dixit Barthram, qui semina misit ex Virginia.*
- Caulis angulatus, lævis, erectus, parum flexuosus. Folia Cherophylli, lævia. Umbella minus expansa, alba. Semina singula 5 alis longitudinalibus parvis; involucri duplex. Statura Selini monnieri.*
9. *ATHAMANTA* foliolis capillaribus, seminibus Meum. giabris striatis. *Hort. cliff.* 93. *Hort. ups.* 60. *Mat. med.* 144. *Roy. lugdb.* 97. *Sauv. monsp.* 256.  
*Meum foliis anethi.* *Bauh. pin.* 148.  
*Meum.* *Dod. pempt.* 305.  
*Habitat in Alpibus Italiæ, Hispaniæ, Helvetiæ.* 21

## PEUCEDANUM.

1. *PEUCEDANUM* foliis quinquies tripartitis filiformibus linearibus. officinale.  
*Peucedanum foliis quinquies tripartitis linearibus subulatis integerrimis.* *Hort. cliff.* 93. *Hort. ups.* 60. *Mat. med.* 126. *Roy. lugdb.* 98. *Sauv. monsp.* 182, 257.  
*Peucedanum.* *Bauh. hist.* 3. p. 36. *Raj. hist.* 416.
2. *Peucedanum Germanicum.* *Bauh. pin.* 149.  
3. *Peucedanum majus italicum.* *Bauh. pin.* 149.  
*Habitat in Europæ australioris pratis pingvibus.* 21
2. *PEUCEDANUM* foliolis linearibus ramosis. *Hort. alpestre.* *cliff.* 94. *Roy. lugdb.* 98.

*Ferula*, foliis libanotidis brevioribus, alpestris, umbellis amplissimis. *Boerh. lugdb.* 1. p. 65.

*Habitat* - - -

- Silvaus. 3. PEUCEDANUM** foliolis pinnatifidis: laciniis oppositis, involucro universali diphylo. *Hort. cliff.* 94. *Roy. lugdb.* 98. *Sauv. monsp.* 257. *Leif. hal.* 232. *Seseli pratense. Bauh. pin.* 162. *Raj. angl.* 3. p. 216. *Siler alterum pratense. Dod. pempt.* 310.

*Habitat in* Helvetiæ, Narbonæ, Germaniæ, Angliæ subhumidis.

*Caulis bipedalis* ♂ ultra. *Foliola lanceolata*, quasi carinata: extimis coadunatis. *Flores lutei*, extus albi.

- alsaticum. 4. PEUCEDANUM** foliolis pinnatifidis: lacinulis trifidis obtusiusculis.

*Selinum* foliolis pinnatim laciniatis: lacinulis trifidis obtusis. *Hort. cliff.* 92.

*Daucus alsaticus. Bauh. prodr.* 77. *Hall. opusc.* 318. \*

*Umbellifera alsatica magna*, umbella parva sublutea. *Bauh. hist.* 3. p. 106. *Raj. hist.* 414.

*Habitat in* Alsatiæ, Palatinatus dumetis humidiusculis.

- nodosum. 5. PEUCEDANUM** foliolis alternatim multifidis. *Hort. cliff.* 94. *Roy. lugdb.* 98.

*Silaum* (quod *Ligusticum creticum*, folio fœniculi, caule nodoso. *Tournef. ccr.* 23.) *Boerh. lugdb.* 1. p. 51.

*Habitat in* Creta.

## CRITHMUM.

- maritimum. 1. CRITHMUM** foliolis lanceolatis carnosis. *Hort. cliff. Hort. upf.* 61. *Roy. lugdb.* 98. *Sauv. monsp.* 258.

*Crithmum*, *Fœniculum maritimum minus. Bauh. pin.* 288.

*Fœniculum marinum* f. *Empetrum* f. *Calcifraga. Lob. ic.* 394.

*Habitat ad oceani Europæi littora.* 2.

- pyrenaicum. 2. CRITHMUM** foliolis lateralibus bis trifidis. *Hort. cliff.* 98. *Roy. lugdb.* 58.

*Apium pyrenaicum*, thapsiæ facie. *Tournef. inst.* 305.

*Habitat in* pyrenæis.

## HASSELQUISTIA.

1. HASSELQUISTIA. *Amæn. acad.* 4. p. 270. \* egyptiaca.  
*Pastinaca orientalis*, foliis eleganter incisus. *Buxb. cent.*  
 3. p. 16. t. 27 ?  
*Habitat in Arabia.*

## CACHRYS.

1. CACHRYS foliis bipinnatis, foliolis acutis multifidis, seminibus sulcatis lævibus. *Hort. cliff.* 94. *Roy. lugdb.* 99. *Sauv. monsp.* 105. Libanotis.  
 Cachrys femine fungoso fulcato plano minore, foliis peucedani angustis. *Morif. hist.* 3. p. 267. f. 9. t. 1. f. 3. 6.  
*Libanotis ferulæ folio*, semine anguloso. *Bauh. pin.* 158.  
*Habitat in Sicilia, Monspeli.* 2
2. CACHRYS foliis bipinnatis: foliolis linearibus acutis, seminibus sulcatis hispidis.  
 Cachrys femine fungoso fulcato aspero, foliis peucedani latiusculis. *Morif. hist.* 3. p. 267. f. 9. t. 1. f. 3.  
*Hippomarathrum creticum.* *Bauh. pin.* 147. *prodr.* 76.  
*Hippomarathrum ficulum.* *Bocc. sic.* 37.  
*Habitat in Sicilia, Hispania.*  
*Foliola linearia. Folia suprema opposita. Umbella universalis terminalis, sessilis. Involucris involucellisque ramosis. Reliquæ umbellæ pedunculatæ, Involucris involucellisque simplicibus. Semina crassissima, sulcata, hispida.*

## FERULA.

1. FERULA foliolis linearibus longissimis simplicibus. communis.  
*Hort. cliff.* 95. *Hort. upf.* 61. *Roy. lugdb.* 99. *Sauv. monsp.* 257.  
*Ferula femina plinii.* *Bauh. pin.* 148.  
*Ferula. Dod. pempt.* 321.  
*Habitat in Europa australi.* 2
2. FERULA foliis supradecompositis: foliolis lanceolato-linearibus planis. glauc.  
*Hort. cliff.* 95. *Roy. lugdb.* 99.  
*Ferula folio glauco*, semine lato oblongo. *Bauh. hist.* 3. p. 45.  
*Habitat in Sicilia, Italia.* 2
3. FERULA foliolis laciniatis: lacinulis tridentatis singulatis. singulatis.  
*Z 2* inæ.

- inæqualibus nitidis. *Hort. cliff.* 95. *Hort. upf.* 61.  
*Roy. lugdb.* 99.
- Ferula tingitana*, folio latissimo lucido. *Herm. par.* 165. t. 165.  
*Habitat in Hispania, Barbaria.* ♂
- Herulago.* 4. *FERULA* foliis pinnatifidis: pinnis linearibus planis trifidis. *Hort. cliff.* 95. *Roy. lugdb.* 99.  
*Ferula latiore folio.* *Morif. hist.* 3. p. 369. f. 9. t. 15. f. 1.  
*Ferulago latiore folio.* *Bauh. pin.* 148.  
*Habitat in Sicilia.*
- orientalis.* 5. *FERULA* foliorum pinnis basi nudis, foliolis setaceis. *Hort. cliff.* 95. *Roy. lugdb.* 100.  
*Ferula orientalis*, folio & facie cachryos. *Tournef. cor.* 22. *itin.* 3. p. 239. t. 239.  
*Habitat in Oriente.*
- meoides.* 6. *FERULA* foliorum pinnis utrinque appendiculatis, foliolis setaceis. *Hort. cliff.* 95. *Roy. lugdb.* 100.  
*Laserpitium orientale*, folio mei, flore luteo. *Tournef. cor.* 23.  
*Habitat in Oriente.*
- modiflora.* 7. *FERULA* foliolis appendiculatis, umbellis subsessilibus.  
*Ferula foliorum alis utrinque auctis: foliolis linearibus integerrimis, umbellis terminatricibus subsessilibus.* *Roy. lugdb.* 100. *Hort. upf.* 61.  
*Libanotis ferulæ folio & semine.* *Bauh. pin.* 158.  
*Panax asclepium, ferulæ facie.* *Lob. ic.* 783.  
*Habitat in Europa australi, Istria.* 24
- canadensis.* 8. *FERULA* lucida canadensis. *Gron. virg.* 147. *Hort. upf.* 61.  
*Habitat in Virginia.* 24
- Asia foetida.* 9. *FERULA* foliolis alternatim sinuatis obtusis. *Mat. med.* 118.  
*Asia foetida disjunensis umbellifera ligustico affinis.* *Kempf. amzn.* 535. t. 536.  
*Habitat in Persia.* 24

## LASERPITIUM.

- latifolium.* 1. *LASERPITIUM* foliolis cordatis inciso-serratis.  
*Hort. cliff.* 96. *Fl. svec.* 230, 242. *Mat. med.* 123.  
*Roy. lugdb.* 101. *Hall. helv.* 440.

- Libanotis latifolia altera s. vulgarior. Bauh. pin. 157.*  
*Libanotis latifolia major. Bauh. pin. 157.*  
*Seseli æthiopicum herba. Dod. pempt. 312.*  
*Habitat in Europæ nemoribus siccis. 24*
2. **LASERPITIUM** foliolis trilobis incis. trilobum.  
*Libanotis latifolia, aquilegiæ folio. Bauh. pin. 157.*  
*prodr. 83.*  
*Angelica foliolis tripartitis: lobis superne incis ob-*  
*tusis. Roy. lugdb. 104.*  
*Ligusticum rauvolfii, foliis aquilegiæ. Bauh. 3. p. 148.*  
*Pluk. phyt. 223. f. 7.*  
*Habitat in Gargano monte. 24*
3. **LASERPITIUM** foliolis cuneiformibus furcatis. gallicum.  
*Laserpitium foliolis ramulosis sessilibus. Roy. lugdb. 101.*  
*Laserpitium foliolis quinquelobis. Hort. cliff. 96.*  
*Laserpitium gallicum. Bauh. pin. 156. Raj. hist. 426.*  
*Laserpitium foliis angustioribus dilute virentibus con-*  
*jugatum dispositis. Raj. hist. 426.*  
*Laserpitium e regione massiliæ allatum, Bauh. hist. 3.*  
*p. 137.*  
*Habitat in Europa australi. 24*
4. **LASERPITIUM** foliolis lanceolatis integerrimis ses- angustifo-  
 silibus. *Hort. cliff. 96. Roy. lugdb. 102.* litum.  
*Laserpitium angustifolium majus, segmentis longiori-*  
*bus & indivisis. Moris. hist. 3. p. 321. f. 9. t. 19. f. 9.*  
*Laserpitium foliis longioribus dilute virentibus con-*  
*jugatum positis. Pluk. alm. 207. t. 198. f. 4.*  
*Habitat in Europa australi. 24*
5. **LASERPITIUM** foliolis lanceolatis integerrimis: prutenicum.  
 extimis coalitis. *Hort. cliff. 96. Bæhm. lipf. 208.*  
*Laserpitium foliis radicalibus patentissimis. Roy. lugdb.*  
*103.*  
*Laserpitium daucoides prutenicum, viscoso semine.*  
*Breyn. cent. 167. t. 84.*  
*Laserpitium minus. Rivin. pent. t. 23. Rupp. jen. 277.*  
*Hall. gatt. 177.*  
*Habitat in Borussia, Lipsiæ. 24*  
*Habitus Selni Caruifoliæ, sed Semina pubescentia.*
6. **LASERPITIUM** foliolis ovali lanceolatis integerri- Siler.  
 mis petiolatis. *Hort. cliff. 96. Mat. med. 124.*  
*Ligusticum quod Seseli officinarum. Bauh. pin. 162.*  
*Siler montanum. Moris. hist. 3. p. 276. f. 9. t. 3. f. 1.*  
*Habitat in Austria, Helvetia, Gallia. 24*

- peucedanoi- 7. **LASERPITUM** foliolis lineari-lanceolatis venoso-  
des. striatis distinctis. *Amén. acad.* 4. p. 310. \*  
Laserpitium peucedanoides, foliorum segmentis angu-  
stissimis. *Segu. veron.* 3. p. 227. t. 17.  
Laserpitium exoticum, lobis angustissimis integris.  
*Pluk. a'm.* 207. t. 96. f. 2.  
*Habitat in Baldi valle Vaccaria.* 24.
- ferluaceum. 8. **LASERPITUM** foliolis linearibus.  
Cachrys orientalis, ferulæ folio tenuiore, fructu alato  
plano. *Tournef. cor.* 23. it. 2. p. 121. t. 121.  
*Habitat in Oriente.*  
Folia quinquepinnata, valde supradecomposita, angu-  
stissima, linearia, levia, laxa. Semina alis exarata.
- Chironium. 9. **LASERPITUM** foliolis oblique cordatis, petiolis  
hirsutis.  
Panax Heracleum. *Morif. hist.* 3. p. 315. f. 9. t. 17. f. 11  
Panax pastinacæ folio. *Bauh. pin.* 156.  
Panaces peregrinum. *Dod. pempt.* 309.  
*Habitat in Montpelii.* 24  
Folia integra, cordata, lobo superiore brevior, cre-  
nata, obtusa. Petioli valde hirsuti. Involucrum  
parvum, tam universale, quam parziale; hinc di-  
versa a Pastinaca Costina.

### HERACLEUM.

- Sphondy- 1. **HERACLEUM** foliolis pinnatifidis. *Hort. cliff.* 103.  
lium. *Fl. suec.* 231, 243. *Roy. lugdb.* 113.  
Sphondylium vulgare hirsutum. *Bauh. pin.* 157.  
β. Sphondylium hirsutum, foliis angustioribus. *Bauh.*  
*pin.* 157. *prodr.* 83. *Riv. pent.* 5.  
*Habitat in Europæ nemorosis.* ♂
- sibiricum. 2. **HERACLEUM** foliis pinnatis: foliolis quinis: in-  
termediis sessilibus, corollulis uniformibus. *Hort.*  
*upf.* 65. \*  
Pastinaca foliis simpliciter pinnatis: foliolis pinnatif-  
dis. *Gmel. sibir.* 1. p. 218.  
*Habitat in Sibiria.* ♂  
*Flores hujus viridi-lutei, at prioris albi.*
- Panaces. 3. **HERACLEUM** foliis pinnatis; foliolis quinis: in-  
termediis sessilibus, floribus radiatis. *Hort. upf.* 65. \*  
Heracleum foliolis palmatis serratis. *Hort. cliff.* 103.  
*Roy. lugdb.* 113.

He-

*Heracleum foliis pinnatifidis. Gmel. fibr. 1: p. 213.*

*Panax sphondylii folio f. Heracleum. Bauh. pin. 157.*

*Habitat in Apenninis, Sibiria. ♂*

4. **HERACLEUM** foliis pinnatis utrinque scabris, flo- austriacum-  
ribus radiatis.

*Sphondylium alpinum parvum, Bauh. pin. 157. prodr.*  
*83. Burf. VIII: 21.*

*Habitat in Alpibus Austriæ.*

*Planta pedalis, Folia pinnata, foliolis 5, ovato-oblongis, sessilibus, serratis, utrinque scabris, parum incis-  
is, petiolis inferne membranaceis. Umbella alba,  
radiata. Differt a tertia parvitate, foliis non subtus  
tomentosis, nec extimis coadunatis; foliolis demum  
magis integris.*

5. **HERACLEUM** foliis simplicibus, floribus radiatis. alpinum.

*Sphondylium alpinum glabrum. Bauh. pin. 157. prodr.*  
*83. t. 83.*

*Sphondylium montanum glabrum, albo flore. Barr.*  
*ic. 55.*

*Habitat in Helvetiæ Alpibus, innotuit C. Bauhino*  
*1595.*

*Scapus nudus, Folia figura Rubi Chæmæmori f. Ficus.*

## LIGUSTICUM.

1. **LIGUSTICUM** foliis multiplicibus: foliolis super- Levisticum,  
ne incis. *Hort. cliff. 97. Hort. upf. 62. Mat. med.*

*119. Roy. lugdb. 104. Sauv. monsp. 261.*

*Ligusticum vulgare. Bauh. pin. 157.*

*Levisticum vulgare. Morif. hist. 3. p. 275. f. 9. t. 3.*  
*f. 1.*

*Habitat in Apenninis Liguriæ. 2.*

2. **LIGUSTICUM** foliis biternatis.

*Ligusticum foliis duplicato-ternatis. Hort. cliff. 97. Fl.*

*sec. 232, 244. Roy. lugdb. 104. Gron. virg. 31.*

*Ligusticum humilius scothicum. Pluk. alm. 217. t. 96.*  
*f. 2.*

3. **Apium maritimum. Fl. lapp. 107.**

*Seseli maritimum scothicum. Herm. par. 227. t. 227.*

*Habitat ad littora maris in Anglia, Svecia, Canada. 2.*

3. **LIGUSTICUM** foliis decompositis incis: radica- cornubiente  
libus ternatis lanceolatis integerrimis. *Amzn. acad.*

*4 p. 210 †*



- Smyrniolum tenuifolium nostras. Raj. suppl. 254. angl. 3. p. 209. t. 8.*  
*Saxifraga cornubiæ. Pet. herb. t. 26. f. 9.*  
*Habitat in Cornubia.*
- peloponnesiacum. 4. **LIGUSTICUM** foliis multiplicato pinnatis: foliolis pinnatis incis. *Hort. cliff. 97. Roy. lugdb. 104. Sauv. monsp. 250.*  
*Cicutaria latifolia foetida. Bauh. pin. 161.*  
*Seseli peloponnesiense. Cam. epit. 514.*  
*Habitat in Peloponnesiacis & Rhæticis montibus nemorosis. ♂*
- austriacum. 5. **LIGUSTICUM** foliis bipinnatis: foliolis confluentibus incis. integerrimis.  
*Seseli montanum, cicutæ folio, glabrum. Bauh. pin. 161.*  
*Seseli 1 montanum pannonicum. Clus. hist. 2. p. 193.*  
*Habitat in Austriæ alpinis.*
- peregrinum. 6. **LIGUSTICUM** umbellæ primariæ involucro subnullo, lateralibus basi membranaceis; radiis subramosis.  
*Habitat in Lusitania. ♂*  
*Caulis bipedalis rigidus angulatus levis. Folia supradecomposita: summa ternata: Foliolis tripartitis: laciniis cuneatis incis. glabris petroselinis, sed majoribus. Petioli lateribus membrana alba. Involucrum universale umbellæ triplo brevius: radiis linearibus, basi utrinque albo-membranaceis, vix ramosis. Umbella prima sœpissime nullum involucrum habet. Involucella linearis, longitudine umbellulæ. Petala flavescentia, inflexo-emarginata. Filamenta corollæ longiora, antheris albis. Germina lævia, vix striata.*

## ANGELICA.

- Archangeli-ca. 1. **ANGELICA** foliorum impari lobato. *Fl. lapp. 101. Fl. vec. 233, 245. Mat. med. 120. Hort. cliff. 97. Roy. lugdb. 103.*  
*Angelica sativa. Bauh. pin. 155.*  
*Angelica major. Dod. pempt. 318.*  
*Habitat in Alpibus Lapponiæ, ad rivulos. ♂*  
*Umbella universalis copiosa, radiis angulatis; Partiales globosæ Involucrum sæpe unifolium; Involucella polyphylla, linearis, reflexa. Petala ovata, acuta, viridia, caduca. Stamina alba.*

2. AN-

2. **ANGELICA** foliis æqualibus ovato-lanceolatis serratis. *Hort. cliff.* 97. *Fl. fvec.* 234, 246. *Mat. med.* 121. *Gron. virg.* 31. *Hall. helv.* 444. *Roy. lugdb.* 103. *Angelica sylvestris major. Bauh. pin.* 155. *Angelica sylvestris. Dod. pempt.* 318. *Fl. lapp.* 102. *Habitat in Europæ frigidioris subhumidis sylvaticis.* 24.
3. **ANGELICA** extimo foliorum pari coadunato: foliolo terminali petiolato. *Roy. lugdb.* 103. *atropurpurea.* *Angelica canadensis atropurpurea. Corn. canad.* 198. 1. 199. *Habitat in Canada.*
4. **ANGELICA** foliolis æqualibus ovatis inciso-serratis. lucida. *Hort. cliff.* 97. *Hort. upf.* 62. *Roy. lugdb.* 103. *Angelica lucida canadensis. Corn. canad.* 196. 1. 197. *Morif. hist.* 3. p. 281. f. 9. t. 3. f. 8. *Habitat in Canada.* ♂

## SIUM.

1. **SIUM** foliis pinnatis, umbellis terminalibus. *Hort. latifolium.* *cliff.* 98. *Fl. fvec.* 235, 247. *Roy. lugdb.* 104. *Hall. helv.* 435. *Sium latifolium. Bauh. pin.* 154. *Sium Riv. pent.* 78.
- β. *Sium* f. *Apium* palustre, foliis oblongis. *Bauh. pin.* 154. *Habitat in Europæ rivulis & ad ripas paludosas.* 24
2. **SIUM** foliis pinnatis, umbellis axillaribus sessilibus. nodiflorum. *Hort. cliff.* 98. *Fl. fvec.* 2. n. 248. *Roy. lugdb.* 105. *Sauv. monsp.* 231. *Sium aquaticum procumbens, ad alas floridum. Morif. hist.* 3. p. 283. f. 9. t. 5. f. 3. *Habitat in Europa ad ripas fluviorum. Involucrum universale in hac specie sæpe deest.*
3. **SIUM** foliis pinnatis, floralibus ternatis. *Hort. cliff.* *Sisarum.* 98. *Hort. upf.* 62. *Roy. lugdb.* 104. *Sisarum germanorum. Bauh. pin.* 155. *Sisarum. Dod. pempt.* 681. *Habitat in China?* 24
4. **SIUM** foliis serratis pinnatis: ramis ternatis. *Ninif.* *Vide Mat. med.* 117. †

362 — PENTANDRIA DIGYNIA.

- Sisyrinchium montanum* coræense, radice non tuberosa.  
*Kämpf. aman.* 817. t. 818.  
*Habitat in China.* 24
- rigidius. 5. *SIUM* foliis pinnatis: foliolis lanceolatis subintegerrimis.  
*Pimpinella* foliolis lanceolatis glabris acuminatis sæpius integerrimis, rarius serratura notatis. *Gron. vîrg.* 32.  
*Oenanthe maxima virginiana*, præoniæ feminae foliis.  
*Morif. hist.* 3. p. 288. f. 11. t. 7. f. 1.  
*Habitat in Virginia.* 24  
*Caulis rigidulus. Folia pinnata oppositionibus 5 f. 6 & impari. Foliola lanceolata, rigidula, versus apicem dentibus aliquot. Petioli canaliculati. Flores parvi.*
- Falcaria. 6. *SIUM* foliolis linearibus decurrentibus connatis. *Hort. cliff.* 98. *Roy. lugdb.* 105. *Sauv. monsp.* 232.  
*Eryngium arvense*, foliis ferratis. *Bauh. pin.* 386.  
*Ammi perenne repens.* *Morif. hist.* 3. p. 294. f. 9. t. 8. f. 1.  
*Habitat in Flandria, Helvetia, Bohemia, Alsatia, Gallia, Oriente.* 24
- græcum. 7. *SIUM* foliis omnibus bipinnatis. *Hort. cliff.* 98. \*. *Roy. lugdb.* 105.  
*Ligusticum græcum*, folio apii. *Tournef. cor.* 23.  
*Habitat in Græcia.*
- ficulum. 8. *SIUM* foliis radicalibus ternatis, caulinis bipinnatis.  
*Roy. lugdb.* 105.  
*Myrrhis* foliis pastinacæ læte virentibus. *Tournef. cor.* 22.  
*Daucus*, pastinacæ folio, ficulus. *Zan. hist.* 78. t. 30.  
*Habitat in Sicilia.*

S I S O N.

- Anomum.* 1. *SISON* foliis pinnatis, umbellis erectis. *Roy. lugdb.* 105. *Sauv. monsp.* 232.  
*Sison* foliis pinnatis. *Hort. cliff.* 98.  
*Sison* quod *Amomum* officinis nostris. *Bauh. pin.* 154.  
*Petrofelinum macedonicum fuchsi.* *Dod. pempt.* 697.  
*Habitat in Angliæ humectis lutosis.*
- segetum. 2. *SISON* foliis pinnatis, umbellis cernuis. *Roy. lugdb.* 105. *Hort. upf.* 63.

*Sium*

- Sium* terrestre, umbellis rarioribus. *Morif. hist.* 3. p. 283. f. 9. t. 5. f. 6.  
*Habitat in Angliæ agris inter segetes.* ♂
2. *SISON* foliis ternatis. *Hort. cliff.* 99. *Gron. virg.* 147. canadense.  
*Roy. lugdb.* 106. *Kalm. it.* 3. p. 259.  
*Myrrhis canadensis trilobata.* *Morif. hist.* 3. p. 301. f. 9. t. 11. f. 4.  
*Myrrhis canadensis.* *Riv. pent.* t. 53.  
*Habitat in America septentrionali.* 24  
*Folia Radicalia ternata, serrata: folio medio trifido; lateralibus bifidis. Caulina ternata, serrata, lanceolata, lucida. Umbella quadrifida, inæqualis, nuda. Umbellulæ subseptemfloræ, involucello minimo. Petala alba, equalia.*
4. *SISON* foliis tripinnatis: radicalibus linearibus; caulinis setaceis: stipularibus longioribus. *Amml.*  
*Sison* foliolis caulinis subcapillaribus. *Hort. ups.* 63. *Mat. med.* 140.  
*Fœniculum lusitanicum minimum acre.* *T. Shaw. afr.* 232. t. 232.  
*Ammi lacinulis foliorum caulis capillaribus.* *Hort. cliff.* 81. *Roy. lugdb.* 96.  
*Ammi parvum, foliis fœniculi.* *Bauh. pin.* 159.  
*Ammi. Cam. epit.* 522.  
*Habitat in Lusitania, Apulia, Ægypto.* ☉  
*Folia Radicalia linearia, supradecomposita, plana: Foliolis sessilibus, replicato-imbricatis; Caulina supradecomposito-pinnata, setacea: ex his quæ vaginam petiolarem terminant trichotoma, reliquis longiora.*
5. *SISON* repens, umbellis bifidis. *Fl. svec.* 2. n. 249. *R. inundatum.* *scan.* 397. \*  
*Sium* foliis submersis capillaribus, emerfis pinnatis. *Roy. lugdb.* 105. *Sauv. meth.* 232.  
*Sium minimum, foliis imis ferulaceis.* *Morif. hist.* 3. p. 223. f. 9. t. 9. f. 5.  
*Sium minimum umbellatum, folio varians.* *Fluk. phyt.* 61. f. 3.  
*Habitat in Europæ inundatis.*
6. *SISON* foliolis verticillatis capillaribus. *verticillatum.*  
*Carum foliolis setaceis verticillatis, radice napiformi.* *Sauv. monsp.* 168.

Bunium bulbis oblongis. *Sauv. monsp.* 256.

Oenanthe foliis pinnatis: foliolis lineariter laciniatis.

*Dalib. parif.* 90.

Daucus pratensis millefolii palustris folio. *Bauh. pin.* 150.

Daucus pratensis. *Dalech. hist.* 718.

Carvi foliis tenuissimis, asphodeli radice. *Tournef. inst.* 90.

*Habitat in Gallia, Pyrenæis.* 2

Involucrum huic universale & partiale. Foliola verticillatim in petiolo communi disposita.

## BUBON.

macedoni-  
cum.

1. BUBON foliolis rhombeo-ovatis crenatis, umbellis numerosissimis. *Hort. cliff.* 95. *Hort. upf.* 62. *Mat. med.* 133. *Roy. lugdb.* 100.

*Apium macedonicum. Bauh. pin.* 154.

*Petroselinum macedonicum. Lob. ic.* 702.

*Habitat in Macedonia, Mauritania.* ♂

Galbanum.

2. BUBON foliolis rhombeis dentatis glabris striatis, umbellis paucis. *Hort. cliff.* 96. *Mat. med.* 132. *Roy. lugdb.* 100.

*Anisum africanum frutescens, folio anisi, galbaniferum. Pluk. alm.* 31. t. 12. f. 2.

*Ferula africana galbanifera, foliis & facie ligustici.*

*Herm. par.* 163. t. 163.

*Habitat in Æthiopia.* ♂

gummifera.

3. BUBON foliolis glabris: inferioribus rhombeis serratis; superioribus pinnatifidis tridentatis. *Roy. lugdb.* 100.

*Ferula africana galbanifera, folio myrrhidis. Comm. hort.* 2. p. 115. t. 58.

*Habitat in Æthiopia.* ♂

rigidus.

4. BUBON foliolis linearibus. *Hort. cliff.* 95. *Roy. lugdb.* 100.

*Ferula durior s. rigidis & brevissimis foliis. Bocc. mus.* 2. p. 84. t. 76. *Barr. rar.* 61. t. 77.

*Habitat in Sicilia.* 2

## CUMINUM.

1. CUMINUM. *Mat. med.* 139.

Cuminum.

Cuminum semine longiore. *Bauh. pin.* 146.Cuminum lativum. *Cam. epit.* 518.*Habitat in Aegypto, Aethiopia.* ☉

## OENANTHE.

## 1. OENANTHE stolonifera, foliis caulinis pinnatis fistulosa. liformibus fistulosis.

Oenanthe foliis caulinis fistulosis teretibus. *Hort.**cliff.* 99. *Fl. fvec.* 235, 250. *Roy. Lugdb.* 108. *Hall.**helv.* 432.Oenanthe aquatica. *Bauh. pin.* 162. *it. scan.* 243.Oenanthe. *Riv. pent.* 66.*Habitat in Europæ fossis paludibus.* 242. OENANTHE foliis omnibus multifidis obtusis sub-æqualibus. *Hort. cliff.* 99. *Fl. fvec.* 237, 251. *Roy. lugdb.* 107. crocata.Oenanthe chærophylli foliis. *Bauh. pin.* 162.Oenanthe cicutæ facie, succo viroso crocante. *Lob.**adv.* 326. *Afl. angl.* 1747. n. 480. p. 235. t. 3.*Habitat in Europæ paludibus.* 24*Umbelle univers. radii angulati. Involucrum univ. polyphyllum magnum.*3. OENANTHE umbellularum pedunculis marginalibus longioribus ramosis masculis. *Hort. upf.* 63. prolifera.

Oenanthe flosculis disci sessilibus, radii pedunculo ramo-

so elevato. *Vir. cliff.* 24. *Roy. lugdb.* 108.

Oenanthe flosculis radiantibus umbellarum proliferis.

*Hort. cliff.* 99.Oenanthe prolifera apula. *Bauh. pin.* 163. *Morif. hist.*

3. p. 289. f. 9. t. 7. f. 5.

*Habitat in Sicilia, Apulia.* 24*Umbelle universalis radii laterales, 3 f. 4 divaricati, horizontales, flosculis masculis. Disci flosculi subsessiles, conferti in hemisphaerium, hermaphroditi.*4. OENANTHE fructibus globosis. *Hort. cliff.* 99. globulosa.*Roy. lugdb.* 108.Oenanthe lusitanica, semine crassiore globoso. *Tour-**nef. infl.* 313.*Habitat in Lusitania.*

## 5. OE-

pimpinell-  
loides.

5. **OENANTHE** foliolis radicalibus cuneatis fissis: caulinis integris linearibus longissimis canaliculatis. *Hort. cliff.* 99. *Roy. lugdb.* 103. *Sauv. monsp.* 259.  
*Oenanthe apii* folio. *Bauh. pin.* 162.  
*Oenanthe pastinacæ* sylvestris folio, semine atriplicis. *Bauh. pin.* 162.  
*Oenanthe aquatica pimpinellæ saxifragæ* divifura. *Pluk. alm.* 268. t. 49. f. 4. mala.  
*Habitat* Montpelii & in Europa australi.  
*Caulis* semipedalis, angulatus. *Folia* Radicalia omnino *Petroselinii*, sed crassiora, bipinnata, lata: foliolis cuneiformibus, incis, divisis. *Caulina* minus composita, vix bipinnata: foliolis linearibus, canaliculatis, longissimis. *Umbella* inæqualis. *Involucrum* universale partialiaque subulato-setacea, *Corollæ* albæ, parum radiatæ.

### PHELLANDRIUM.

- aquaticum. 1. **PHELLANDRIUM** foliorum ramificationibus divaricatis.  
*Phellandrium.* *Hort. cliff.* 100. *Fl. fvec.* 238, 252. *Roy. lugdb.* 108. *Dod. pempt.* 591.  
*Cicutaria palustris tenuifolia.* *Bauh. pin.* 161.  
 β. *Millefolium aquaticum umbellatum*, cibriandri folio. *Bauh. pin.* 216.  
*Millefolium aquaticum.* *Matth. diofc.* 2. p. 484.  
*Habitat* in Europæ fossis. ♂  
*Varietas* β. in fluviis amplioribus profundioribus vide *Raj. angl.* 3. p. 216.  
 Mutellina. 2. **PHELLANDRIUM** caule subnudo, foliis bipinnatis. Sefeli caule vix ramoso, umbella purpurea, pinnulis acute multifidis. *Hall. helv.* 431.  
*Meum alpinum*, umbella purpurascente. *Bauh. pin.* 148. *Hall. opusc.* 288.  
*Mutellina.* *Bauh. hist.* 3. p. 66. *Cam. epit.* 8.  
*Habitat* in Helvetia, Carniola. 2.  
*Folia* *Chærophylli.* *Involucella* longitudine umbellarum.

### CICUTA.

- virosa. 1. **CICUTA** umbellis oppositifoliis, petiolis marginatis obtusis.

Ci-

*Cicuta*. *Hort. cliff.* 100. *Mat. med.* 129. *Fl. fvec.* 239;  
253. *Roy. lugdb.* 109.

*Cicuta aquatica*. *Fl. lapp.* 103.

*Sium erucæ folio*. *Bauh. pin.* 154.

*Habitat in paludibus Europæ sterilibus.* 2

2. *CICUTA* ramis bulbiferis.

bulbifera.

*Ammi foliorum lacinulis capillaribus, caule angulato.*

*Gron. virg.* 31.

*Umbellifera aquatica*, foliis in minutissima & plane  
capillaria segmenta divis. *Raj. suppl.* 260.

*Habitat in Virginia, Canada.*

3. *CICUTA* foliorum ferraturis mucronatis, petiolis maculata.  
membranaceis apice bilobis.

*Egopodium foliolis lanceolatis acuminatis serratis.*

*Gron. virg.* 32.

*Angelica caribæarum elatior, olusatrici folio, flore albo, seminibus luteis striatis, cumini odore & sapore.*

*Pluk. alm.* 31. t. 76. f. 1.

*Angelica virginiana*, foliis acutioribus, semine striato  
minore, cumini sapore & odore. *Morif. hist.* 3. p. 281.

*Habitat in Virginie aquosis.*

ÆTHUSA.

1. *ÆTHUSA*. *Hort. cliff.* 100. *Fl. fvec.* 240, 254. *Cynapium.*

*Roy. lugdb.* 109. *Hall. helv.* 433.

*Cicuta minor, petroselinio similis.* *Bauh. pin.* 160.

*Cicutaria fatua.* *Lob. ic.* 280.

*Habitat inter Europæ olera.* ☉

CORIANDRUM.

1. *CORIANDRUM* fructibus globosis. *Hort. cliff.* sativum.

100. *Hort. ups.* 63. *Mat. med.* 135. *Roy. lugdb.* 109.

*Sauv. monsp.* 260.

*Coriandrum majus.* *Bauh. pin.* 158.

*Coriandrum.* *Cam. epit.* 523.

*Habitat in Italiæ agris.* ☉

2. *CORIANDRUM* fructibus didymis. *Hort. cliff.* 100. testiculatum

*Roy. lugdb.* 109. *Sauv. monsp.* 260.

*Coriandrum minus testiculatum.* *Bauh. pin.* 158. *Pluk.*

*alm.* 120. t. 169. f. 2.

*Coriandrum minus odorum.* *Bauh. hist.* 3. p. 91.

β. Co-



β. *Coriandrum sylvestre foetidissimum. Bauh. pin. 158.*

*Segu. veron. 2. p. 28.*

*Habitat in Europæ australis agris. ☉*

*Odor C. sativo virosior. Caulis angulatus. Umbella sepius simplex, rarius composita. Involucrum Universalis unifolium; Partialia nulla. Petala non radiata. Antheræ purpurascens. Germina inferne didyma.*

## SCANDIX.

*odorata.*

1. **SCANDIX** seminibus sulcatis angulatis. *Hort. cliff. 101. Hort. upf. 64. Roy. lugdb. 111. umbellarum disco masculino.*

*Myrrhis major, Cicutaria odorata. Bauh. pin. 160.*

*Myrrhis. Dod. pempt. 701.*

*Habitat in alpinis Alvarniæ, inque confinibus agri Veronensis & Vicentini.*

*Pecten. ♀*

2. **SCANDIX** seminibus lævibus rostro longissimo. *Hort. cliff. 101. Hort. upf. 64. Roy. lugdb. 111.*

*Scandix, semine rostrato. vulgaris. Bauh. pin. 152.*

*Pecten veneris. Cam. epit. 302.*

*Habitat inter Germaniæ & Europæ australioris segetes. ☉*

*Cerefolium.*

3. **SCANDIX** seminibus nitidis ovato-subulatis, umbellis sessilibus lateralibus. *Hort. cliff. 110. Fl. fvec. 242, 255. Mat. med. 137. Gron. virg. 32. Roy. Lugdb. 111.*

*Chærophyllo seminibus lævibus, umbellis ad nodos sessilibus. Bæhm. lipf. 492.*

*Chærophyllo sativum. Bauh. pin. 152.*

*Chærefolium. Dod. pempt. 700.*

*Habitat in agris & arvis Europæ australioris. ☉*

*Anthriscus.*

4. **SCANDIX** seminibus ovatis hispidis, corollis uniformibus, caule lævi.

*Scandix seminibus hispidis. Hort. cliff. 101. Fl. fvec. 241. Roy. lugdb. 111. 242, 256.*

*Myrrhis sylvestris, seminibus asperis. Bauh. pin. 160.*

*Myrrhis sylvestris æquicolorum. Column. ecphr. 1. p. 110. t. 112.*

*Habitat in Europæ aggeribus terrenis. ☉*

5. **SCAN-**

3. SCANDIX seminibus subulatis hispidis, floribus ra-  
diatis, caulibus lævibus.  
Scandix seminibus rostratis patulis pedunculo bre-  
vioribus. *Sama monsp.* 260.  
Scandix cretica minor. *Bauh. pin.* 152.  
Scandix, semine rostrato, Italica. *Bauh. prodr.* 78.  
Anisomarathrum, *Col. ecphr.* 1. p. 89. t. 90.  
*Habitat in G. Narbonensi, Italia, Creta.* ☉
6. SCANDIX seminibus subcylindricis hispidis, caule nodosa.  
hispido, geniculis tumidis.  
Chærophylum articulis tumidis, umbella universali  
trifida. *Hort. cliff.* 102. *Roy. lugdb.* 112.  
Cerefolium annuum nodosum, semine aspero majore.  
*Morif. hist.* 3. p. 303. f. 9. t. 10. f. 4.  
Myrrhis nodosa annua, semine aspero. *Morif. blaf.*  
288.  
*Habitat in Sicilia.* ☉  
Caulis scaber aculeis mollibus, reflexis: articulis super-  
ne incrassatis. Umbella radiis 2 f. 3. Umbellulæ  
radiis 5 f. 6. Fructus longi, cylindrici, utrinque  
obtusi, scabri.
7. SCANDIX seminibus pedunculo villosa breviori-  
bus. *Roy. lugdb.* 111.  
Scandix orientalis, flore maximo. *Tournef. cor.* 23.  
*Habitat in Oriente.*
8. SCANDIX seminibus nitidis ovato-subulatis, foliis  
decompositis. *Gzon. virg.* 147.  
Cerefolium virginianum procumbens, fumarie foliis.  
*Morif. hist.* 3. p. 303. f. 9. t. 11. f. ult.  
*Habitat in Virginia.*

# CHÆROPHYLLUM.

1. CHÆROPHYLLUM caule striato geniculis tumi-  
diusculis. *Fl. suec.* 2. n. 257.  
Chærophylum flosculis omnibus fertilibus, caule  
æquali. *Sp. pl.* 1. p. 258.  
Chærophylum seminibus lævibus nitidis, petiolis  
rameis æqualibus. *Hort. cliff.* 101. *Fl. suec.* 243,  
257. *Roy. lugdb.* 112. *Mat. med.* 143.  
Chærophylum sylvestre perenne, cicutæ folio.  
*Tournef. inst.* 114. *Fl. lapp.* 104.  
Myrrhis sylvestris, seminibus lævibus. *Bauh. pin.*  
160.  
*Linnaei Spec. plant. pars I.*      Aa      Cisu-

*Cicutaria vulgaris*. *Dod. pempt.* 701.

*Habitat in Europæ pomariis* ♂ *cultis.* 24

*Caulis subfulcatus. Pedunculus communis teres. Petala integra, plana, exteriora majora.*

bulbosum. 2. **CHÆROPHYLLUM** caule lævi: geniculis tumidis. *Leyf. hal.* 258.

*Chærophylllum* foliis supradecompositis: caulibus articulis lævibus superne incrassatis. *Hort. ups.* 64.

*Chærophylllum* radice turbinata carnosa. *Hort. cliff.* 102. *Roy. lugdb.* 112.

*Cicutaria bulbosa.* *Bauh. pin.* 161.

*Myrrhis annua*, semine striato lævi, tuberosa nodosa coniophyllon. *Pluk. alm.* 249. t. 206. f. 2.

*Myrrhis foetens.* *Riv. pent.* 49.

*Habitat in Germania, Hungaria, Helvetia, Norvegia.* ♂

*Radix turbinata f. obovata. Caulis sexpedalis, lævis, rufo-fusco-maculatus: articulis glabris apice tumetibus, at infimis hispidis. Folia incisa, supra glabra, petiolis ♂ carina pilis albis vagis subtus adpersa. Involucella foliolis 5 f. 7., subulatis, inæqualibus, basi fere coadunatis. Petala alba, obcordata, inæqualia. Flosculi disci nonnulli abortientes.*

temulum. 3. **CHÆROPHYLLUM** caule scabro: geniculis tumidis.

*Chærophylllum* caule maculato: geniculis tumidis.

*Hort. cliff.* 102. *Fl. svec.* 244, 258. *Roy. lugdb.* 112.

*Gort. gelr.* 59.

*Chærophylllum sylvestre.* *Bauh. pin.* 152.

*Myrrhis annua vulgaris*, caule fusco. *Morif. hist.* 3. p. 302. f. 9. t. 10. f. 7.

*Habitat ad Europæ arvosa, vias* ♂ *sepes.* ♂

*Umbellæ virgines dormientes cernuæ. Umbellulae in medio abortientes.*

aureum. 4. **CHÆROPHYLLUM** caule æquali, foliolis incisis, seminibus coloratis striatis.

*Myrrhis perennis alba minor*, foliis hirsutis, semine aureo. *Morif. umb.* 282. *Rupp. jen.* 3. p. 282. t. 5. \*

*Myrrhis radice nigra perenni*, foliis hirsutis, seminibus flavis oblique striatis. *Hall. gert.* 184. \*

*Habitat in Germania.* 24

5. **CHÆ-**

5. CHÆROPHYLLUM caule æquali, foliolis incisib. hirsutum. acutis, seminibus subulatis.

Chærophyllum foliolis dissectis, petiolis ramiferis universalibus utrinque membrana auctis. *Hort. cliff.* 101.

*Roy. lugdb.* 111. *Sauv. monsp.* 262.

Cerefolium latifolium hirsutum album & rubrum.

*Morij. hist.* 3. p. 304. f. 9. t. 10. f. 6.

Cicutaria latifolia hirsuta. *Bauh. hist.* 3. p. 182.

Cicutaria palustris latifolia alba. *Bauh. pin.* 161.

Cicutaria palustris latifolia rubra. *Bauh. pin.* 161.

Myrrhis palustris. *Riv. pent.* 50.

Myrrhis seminibus strictis longissimis. *Hall. helv.* 453.

Myrrhis broccenbergentis. *Bauh. pin.* 160. *Hall. apusc.* 132.

Seseli montanum, cicuta folio, subhirsutum. *Bauh. pin.* 161. *prodr.* 85.

*Habitat in Alpibus Helvetiæ.* 24

6. CHÆROPHYLLUM caule æquali, foliolis cordatis serratis integris.

Chærophyllum foliolis lanceolato-ovatis serratis integris. *Hort. cliff.* 102. *Roy. lugdb.* 112.

Cerefolium, rugoso angelicæ folio, aromaticum. *Borc. mus.* 2. p. 29. t. 19.

Angelica podagrariæ folio. *Riv. pent.* - - -

Angelica sylvestris hirsuta inodora. *Bauh. pin.* 156. *prodr.* 82.

*Habitat in Lusatia, Misnia.* 24

Statura & Folia Ægopodii Podagrariæ. Flores plerique masculi; pauci hermaphroditi.

7. CHÆROPHYLLUM fruticosum.

arborescens.

Cicuta arbor virginiana. *Pluk. mant.* 49.

Cicutaria arbor virginiana. *Raj. suppl.* 257.

*Habitat in Virginia.* 6

Gaulis frutescens. Folia Chærophylli sylvestris, ampla, supradecomposita, pinnis pinnatifidis, levissimis, incisoferratis. Umbellæ tenues, albæ; involucrio partiali. Flosculi omnes fertiles.

## IMPERATORIA.

1. IMPERATORIA. *Hort. cliff.* 103. *Hort. apsc.* 65. *Ostruthiana.* *Mat. med.* 122.

Imperatoria. *Bauh. pin.* 156.

A a 2

Ma-

*Magistrantia. Cam. epit. 592.*

*Habitat ad radices Alpium Helvetiæ, Austriæ, Alverniæ. 24*

*Involucrum uno alterove foliolo tenuissimo. Folia floralia opposita: stipula petiolaris membranacea, ventricosa, una intra alteram.*

## SESELI.

*Pimpinelloldea. 1. SESELI caule declinato, umbellis nubilibus nutantibus.*

*Habitat in Europa australiore. 24*

*Radix perenns. Folia pinnata, pinnis multifidis, segmentis alternis: hæc etiam alternatim secta, segmentis linearibus, planis extrorsum versis. Caulis pedalis, teres, ante florescentiam decumbens, sub florescentia ferme erectus foliolo uno alterove: petiolo membranaceo, spatheiformi, oblongo; e quo pedunculus simplex. Umbellæ ante florescentiam nutantes ut in Pimpinella saxifraga: involucri universali nullo; partiali lineari angusto, longitudine fere umbellulæ. Semina ovalia stris 3. elevatis.*

*montanum. 2. SESELI petiolis ramiferis membranaceis oblongis integris, foliis caulinis angustissimis. Hort. cliff. 102. Roy. lugdb. 112. Dalib. parif. 92. Sauv. monsp. 256. Meum latifolium adulterinum. Bauh. pin. 148.*

*Saxifraga montana minor Italica, foliis in breviores partes divis. Morif. hist. 3. p. 272.*

*Habitat in Galliæ, Italiæ collibus. 24*

*Carvisfolia Vaill. parif. t. 5. f. 2. Caulis pedalis, teres, levis. Folia Radicalia breviora, dancina, bipinnata, ad costam sæpe cruciata, foliolis subtripartitis. Caulina vagina petiolari integra, pinnata: foliolis ternatis: pinnis tripartitis, linearibus acutiusculis, remotiusculis, longioribus, canaliculatis. Umbellæ pedicellis brevibus. Semina pubescentia. Huic vel sequenti proxima.*

*glaucum. 3. SESELI petiolis ramiferis membranaceis oblongis integris: foliolis singularibus binatisque canaliculatis lævibus petiolo longioribus. Guett. stamp. 64. Fœniculum sylvestre, glauco folio. Tournef. inst. 311. Daucus, glauco folio, similis fœniculo tortuoso. Bauh. hist. 3. p. 16.*

*Saxifraga montana minor glauca & rigidior. Morif. hist. 3. p. 273.*

*Habitat in Gallia, 24.*

*Flores ante explicationem purpurascens, explicati albi antheris purpurascens.*

4. SESELI petiolis rameis membranaceis ventricosis ~~anatum.~~ emarginatis. *Hort. cliff. 103. Roy. lugdb. 112. Sauv. monsp. 256.*

*Fœniculum sylvestre annuum, tragofelini odore, umbella alba. Vaill. parif. 54. t. 9. f. 4.*

*Libanotis tenuifolia germanica. Bauh. pin. 158. Burf.*

*Habitat in Pannonia, Gallia, Germania. ☉*

*Caulis strictus, spithameus & ultra, striatus. Folia bipinnata, laciniata. Flores albicanti-subviolacei.*

5. SESELI petiolis membrana destitutis. *Roy. lugdb. Ammoides. 112.*

*Fœniculum lusitanicum minimum acre. Tournef. inst. 312.*

*Saxifraga annua acris, millefolii terrestris folio tenuissimo. Morif. hist. 3. p. 272.*

*Ammoides. Bauh. pin. 159. Morif. hist. 3. p. 295.*

*Habitat in Lusitania, Italia. ☉*

*Planta parva, ramosissima. Folia fere Millefolii. Umbellæ multæ radiis 4 f. 5, divaricatis in ambitu & 2 f. 3. parvis in disco; absque involucri. Involucella radiis 2 f. 3, parvis. Petala alba, retusa. Antheræ subviolaceæ. Semina striata.*

6. SESELI petiolis ramiferis membranaceis oblongis pumilum. integris: foliolis ternatis linearibus subcarnosis petioli longitudine. *Gerard.*

*Daucus montanus, multifido folio, selini semine.*

*Bauh. pin. 150.*

*Fœniculum minimum patulum. Clus. hist. 2. p. 200.*

*Habitat in Austria, Gallo-provincia.*

*Planta parva, tota umbellis paniculata. Folia pinnata: foliolis ternatis.*

7. SESELI caule alto rigido, foliolis linearibus fasciculatis. *tortuosum.*

*Oenanthe striata rigida. Hort. cliff. 99. Sauv. monsp. 257.*

*Seseli massiliense, fœniculi folio. Bauh. pin. 101.*

*Fœniculum tortuosum. Bauh. hist. 3. p. 16.*

*Habitat in Europa australi 24*

*Caulis rigidus, quasi lignosus, foliis multoties latior. Foliola partialia, non duo tantum opposita, sed sæpius quatuor: interioribus minoribus, unde fasciculata.*

**Turbith. 8. SESPLI** involucrio universali monophyllo, seminibus stratis viliosis styliatis. *Amzn. acad. 4. p. 319. \**

*Seseli tertium. Bark lugdb. 1. p. 50.*

*Thapsi fœniculi folio. Bauh. pin. 148.*

*Habitat in Europa australi. 24*

**Hippomarathrum. 9. Si SELI** involucri cellis coniato-monophyllis.

*Laurus montanus multifido brevique folio. Bauh. pin. 150.*

*Hippomarathrum. Riv. pent. 66. Rupp. jen. 3. p. 280. Kram. aust. 82. \**

*Habitat in Austria.*

*Folia bipinnata: primis trifidis, linearibus. Caulis junceus, glaucus, ut tota planta. Suprema folia tantum vagina constant. Involucrum universale nullum; involucella monophylla, pelvisformia, margine crenata s. foliolis in unum connatis, qua nota ab omnibus umbellatis differt. Germina sessilia. Corollæ albae, sæpe subtus purpurascentes.*

**pyrenæum.**

**SESPLI** foliis duplicato-pinnatis; foliolis incisim acutis, involucri setaceis umbellula longioribus.

*Carvi alpinum. Bauh. pin. 158. prodr. 84. Burs. VIII. 36.*

*Habitat in Pyrenæis.*

*Caulis pedalis, teres, striatus. Folia radicalia duplicato-pinnata, laxa, parum divaricata: Foliolis pinnatis, longitudinaliter incisim, acuminatis, cuneiformibus; Folium in caule sæpe unicum, ex cujus ala exit ramus; petioli radicales simplices; caulinus latomembraneus. Umbella universalis nuda, oblonga, pedunculis circiter octo; partialis conferta, brevissima, uniformis: involucrio polyphylo, setaceo, longiore umbellulis.*

**saxifragum.**

**II. SESPLI** caule filiformi divaricato, foliis duplicato-ternatis linearibus, umbellis subsexfidis.

*Pimpinella saxifraga tenuifolia. Bauh. pin. 160. prodr. 84. Burs. VIII. 5.*

**Pim-**

*Pimpinella tenuifolia*. Riv. pent.

*Habitat ad Lacum Genevensem, inque Germania.*

*Planta pedalis. Caulis filiformis, ramosus & subramosus, divaricatus, teres, tenuis. Folia parva, duplicato-ternata, linearia, raro lobata. Umbellæ nuda, pedunculis sex sæpius; umbellulæ involucrate flosculis sex ad octo: involuella setacea.*

12. SESELI caule elongato geniculis callosis, foliis duplicato-pinnatis: pinnis lanceolato-linearibus distantibus. *elatum.*

*Daucus montanus, folio fœniculi longiore. Magn. monsp. 294.*

*Apium montanum, folio tenuiore. Bauh. pin. 153.*

*Habitat in Austria, Gallia.*

*Caulis (ex Horto) humanæ altitudinis, levis, geniculis callosis albidis. Foliola glauca, linearilanceolata; ad radicem majora; in caule pauca, laxiora. Umbellæ respectu plantæ paucioribus radiis & parvæ. Flores albi.*

## THAPSIA.

1. THAPSIA foliolis dentatis villosis basi [coadunatis. villosa.

*Hort. cliff. 105. Roy. lugdb. 113.*

*Thapsia latifolia villosa. Bauh. pin. 148.*

*Thapsia. 1. Clus. hist. 2. p. 192.*

*Habitat in Hispania, Lusitania, Galloprovincia, Cætio Montpelienfi. 2.*

*Petiolis & Folia villosa sunt*

2. THAPSIA foliolis multifidis basi angustatis. *Hort. ferida.*

*cliff. 105. Roy. lugdb. 113.*

*Thapsia carotæ folio. Bauh. pin. 148.*

*Thapsia carotæ effigie. Bauh. hist. 3. p. 187, Moris.*

*hist. 3. p. 319. f. 9. t. 18. f. 7.*

*Habitat in Hispania. 2.*

3. THAPSIA foliis digitatis: foliolis bipinnatis setaceo-multifidis. *Hort. cliff. 106. Roy. lugdb. 113. Asclepium.*

*Thapsia foliis pinnatifidis linearibus ad pedunculum communem radiatis. Roy. lugdb. 114.*

*Thapsia tenuifolia, petiolis radiatis. Moris. hist. 3. p. 319. f. 9. t. 18. f. 9.*



*Thapsia Turbit garganicum, semine latissimo. Bauh. hist. 2. p. 50.*

*Panax Asclepium, semine folioso. Bauh. pin. 158.*

*Panax Asclepium apulum. Column. ceph. 1. p. 87. t. 86.*

*Habitat in Apulia, Oriente. 21*

*Folia digitata, foliolis bipinnatis capillaceis.*

trifoliata.

4. *THAPSIA foliis ternatis ovatis.*

*Sium folio infimo cordato, caulinis ternatis, omnibus crenatis. Gron. virg. 31.*

*Habitat in Virginia.*

*Semina singula gaudent alis 5, longitudinalibus, membranaceis, & involucri nulla sunt uti in precedente, cum qua genere conjungo, quamvis facie diversam.*

## PASTINACA.

fativa.

1. *PASTINACA foliis simpliciter pinnatis. Hort. cliff. 105. Hort. upf. 66. Fl. svec. 2. n. 259. Mat. med. 130. Roy. lugdb. 114. Gort. gelr. 60.*

*Pastinaca sylvestris latifolia. Bauh. pin. 155.*

β. *Pastinaca fativa latifolia. Bauh. pin. 155.*

*Habitat in Europæ australioris ruderatis & pascuis.*

Opopanax.

2. *PASTINACA foliis decompositis pinnatis. Hort. cliff. 105. Mat. med. 131. Roy. lugdb. 114.*

*Panax Costinum. Bauh. pin. 156.*

*Panax Heracleum. Moris. hist. 3. p. 315. f. 9. t. 17. f. 2.*

*Habitat in Italia, Sicilia,*

## SMYRNIUM.

perfoliatum

1. *SMYRNIUM foliis caulinis simplicibus amplexicaulibus. Hort. cliff. 104. Hort. upf. 66. Roy. lugdb. 114.*

*Smyrnum peregrinum rotundo s. oblongo folio.*

*Bauh. pin. 154. prodr. 82.*

*Smyrnum amani montis. Dod. pempt. 698.*

*Habitat in Italia, Creta. ♂*

ægyptiacum

2. *SMYRNIUM foliis floralibus bints simplicibus cordatis integerrimis. Amæn. acad. 4 p. 270. \**

*Habitat in Ægypto.*

Olusatrum.

3. *SMYRNIUM foliis caulinis ternatis petiolatis serratis. Hort. cliff. 105. Roy. lugdb. 114. Sauv. monsp. 262.*

Hip-

*Hippofelinum theophrasti* f. *Smyrnum theophrasti*.

*Bauh. pin.* 154.

*Hippofelinum. Dod. pempt.* 698.

*Habitat in* Scotia, Wallia, Gallia, Hispania, Belgio. ♂

*Folia Radicalia triternata: caulina ternata: suprema opposita ternata. Foliorum vaginæ lacerato-ciliatæ. Involucella brevissima. Flores disci masculi; radiales hermaphroditi.*

4. **SMYRNIUM** foliis pinnatis serratis: posticis ternatis, flosculis omnibus fertilibus.

*Ægopodium foliis caulinis summis novenis. Hort. cliff.* 107. *Roy. lugdb.* 115.

*Angelica acadienfis, flore luteo. Dod. ast.* 55. *Raj. hist.* 1368.

*Angelica humilior & minor, flore luteo. Moris. hist.* 3. p. 281.

β. *Smyrnum foliis caulinis decompositis acuminatis. Gron. virg.* 148.

*Smyrnum aureum, lobis ternis quinisque, marianum. Pluk. mant.* 173.

*Habitat in America boreali.*

5. **SMYRNIUM** foliis caulinis duplicato-ternatis integerrimis. integerri-  
mum,

*Smyrnum foliis caulinis ternatis petiolatis: foliolis oblongo-ovatis integerrimis. Gron. virg.* 148.

*Habitat in* Virginia. 24

*Caulis sesquipedalis, lævis, parum ramosus. Folia fere Aquilegiæ, triternata, oblongo-ovata, glabra, integermina, glauco-viridia. Flosculi disci umbellularum masculi.*

## ANETHUM.

1. **ANETHUM** fructibus compressis. *Hort. cliff.* 106. *graveolens.*

*Hort. upf.* 66. *Mat. med.* 146. *Roy. lugdb.* 116.

*Anethum hortense. Bauh. pin.* 147.

*Habitat inter* Lusitaniam & Hispaniam *segetes.* ☉

2. **ANETHUM** fructibus ovatis. *Hort. cliff.* 106. *Hort. Foeniculum. upf.* 66. *Mat. med.* 147. *Roy. lugdb.* 116.

*Foeniculum dulce. Bauh. pin.* 147.

β. *Foeniculum vulgare germanicum. Bauh. pin.* 147.

- γ. *Fœniculum vulgare italicum*, femine oblongo gustu-  
acuto. *Bauh. pin.* 147.  
δ. *Fœniculum sylvestre.* *Bauh. pin.* 147.  
*Habitat in Narbonæ, Aremoriæ, Maderæ rupibus*  
*cretaceis.* ♂

## CARUM.

- Carvi.* 1. **CARUM.** *Dod. pempt.* 299. *Fl. lapp.* 105. *Fl. svec.*  
245, 260. *Mat. med.* 138. *Hort. cliff.* 106. *Roy.*  
*lugdb.* 116. *Hall. helv.* 428.  
*Carum pratense*, *Carvi officinarum.* *Bauh. pin.* 158.  
*Habitat in Europæ borealis pratis.* ♂

## PIMPINELLA.

- saxifraga.* 1. **PIMPINELLA** foliis pinnatis: foliolis radicalibus  
subrotundis; summis linearibus.  
*Pimpinella foliolis subrotundis.* *Fl. svec.* 246, 261.  
*Mat. med.* 125.  
*Pimpinella.* *Hort. cliff.* 106. *Roy. lugdb.* 115.  
*Pimpinella saxifraga minor.* *Bauh. pin.* 160.  
β. *Pimpinella saxifraga major altera.* *Bauh. pin.* 159.  
*major.* γ. *Pimpinella saxifraga major*, umbella candida. *Bauh.*  
*pin.* 159.  
δ. *Pimpinella saxifraga major*, umbella rubente. *Bauh.*  
*pin.* 159.  
*hircina.* ε. *Pimpinella saxifraga tenuifolia.* *Bauh. pin.* 159.  
*Habitat in Europæ pascuis siccis.* 2.  
*Has inter α — ε limites non reperi; β enim sata in*  
*solo latiori produxit γ.*  
*glauc.* 2. **PIMPINELLA** foliis supradecompositis, caule an-  
gulato ramosissimo.  
*Tragofelinum foliis tenuissime divisis; pinnulis ulti-*  
*mis tripartitis.* *Hall. helv.* 430. \*  
*Seseli petiolis ramiferis membranaceis oblongis inte-*  
*gris: foliolis biqatis ternatisque.* *Guett. stamp.* 1.  
*p.* 34.  
*Habitat in Helvetia, Gallia, Italia,*  
*Caret involucellis, quo a Seseli differt.*  
*peregrina.* 3. **PIMPINELLA** foliis radicalibus pinnatis crenatis,  
summis cuneiformibus incisis.  
*Anisum foliis radicalibus pinnatis.* *Hort. cliff.* 107.  
*Hort. upf.* 67. *Roy. lugdb.* 115. *Sauv. meth.* 231.  
Api-

*Apium peregrinum*, foliis subrotundis. *Bauh. pin.* 153.  
*Daucus tertius diofcoridis*. *Column. ecphr.* 1. p. 108.  
 1. 109.

*Habitat in Italiæ pascuis sterilibus.* 24

4. **PIMPINELLA** folis radicalibus trifidis incis, Anisum.  
*Anisum* foliis radicalibus simplicibus. *Hort. cliff.* 107.

*Hort. upf.* 67. *Mat. med.* 141. *Roy. lugdb.* 116.

*Anisum herbariis*. *Bauh. pin.* 159.

*Anisum vulgare*. *Clus. hist.* 2. p. 202.

β. *Cuminum* semine rotundiore & minore. *Bauh. pin.*  
 146. *Raj. extr.* 63.

*Habitat in Ægypto.* ☉

## A P I U M.

1. **APIUM** foliolis caulinis linearibus, involucellis m- Petroseli.  
 nutis. *Hort. cliff.* 108. *Hort. upf.* 67. *Mat. med.* 127. num.  
*Roy. lugdb.* 116.

*Apium hortense*, *Petroselinum vulgo*. *Bauh. pin.* 153.

*Apium hortense*. *Dod. pempt.* 694.

β. *Apium* vel *Petroselinum crispum*. *Bauh. pin.* 153.

*Habitat in Sardinia juxta scaturigines.* ♂

2. **APIUM** foliolis caulinis cuneiformibus, umbellis graveolens.  
 sessilibus. *Hort. cliff.* 107. *Fl. svec.* 248, 252. *Mat.*  
*med.* 128. *Roy. lugdb.* 115.

*Apium palustre* f. *Apium officinarum*. *Bauh. pin.* 154.

β. *Apium dulce*, *Celleri italorum*. *Tournef. inf.* 305.

*Habitat in Europæ humectis, præsertim maritimis.* ♂

## Æ G O P O D I U M.

1. **ÆGOPODIUM** foliis caulinis summis ternatis. *Hort.* Podagraria.  
*cliff.* 107. *Fl. succ.* 247, 263. *Roy. lugdb.* 115.

*Angelica sylvestris minor*. f. *erratica*. *Bauh. pin.* 155.

*Herba gerardi*. *Dod. pempt.* 320.

*Habitat in Europa ad sepes inque pomariis* 24

*Folia summa opposita sunt.*

## TRIGYNIA.

### R H U S.

1. **RHUS** foliis pinnatis obtusiuscule serratis ovalibus Coriaria  
 subtus villosis. *Mat. med.* 150.

*Rhus* 3

*Rhus foliis pinnatis ferratis. Hort. cliff. 110. Hort. upf. 68. Roy. lugdb. 243. Sav. monsp. 227. Gron. orient. 91.*

*Rhus folio ulmi. Bauh. pin. 414.*

*Rhus coriaria. Dod. pempt. 779.*

*Habitat in Europa australi, Syria, Palæstina. §.*

typhinum. 2. RHUS foliis pinnatis lanceolatis argute ferratis: subtus tomentosis. *Amæn. acad. 4. p. 311. \**

*Rhus virginianum. Bauh. pin. 517. Dill. elth. 253*

*Habitat in Virginia. §*

javanicum. 3. RHUS foliis pinnatis ovatis acuminatis ferratis, subtus tomentosis.

*Habitat in China. Osbeck.*

glabrum. 4. RHUS foliis pinnatis ferratis lanceolatis utrinque nudis. *Kalm. it. 2. p. 211.*

*Rhus foliis pinnatis ferratis. Gron. virg. 148. Cold. noveb. 63.*

*Rhus virginicum, panicula sparsa, ramis patulis glabris. Dill. elth. 323. t. 243. f. 314. Catesb. car. 3. p. 4. t. 4.*

*Rhus angustifolium. Bauh. pin. 414.*

*Sumach angustifolium. Bauh. prodr. 158. Burf. XXII. 77.*

*Habitat in America septentrionalis altioribus ad agros*  
*§ Sepes. §*

Vernix. 5. RHUS foliis pinnatis integerrimis, petiolo integro æquali. *Mat. med. 151. Kalm. it. 2. p. 211.*

*Rhus foliis pinnatis integerrimis. Hort. cliff. 110. Hort. upf. 68. Gron. virg. 148. Roy. lugdb. 244. Cold. noveb. 64.*

*Toxicodendron foliis alatis, fructu rhomboide. Dill. elth. 390. t. 292. f. 377.*

*Arbor americana, alatis foliis, succo lacteo venenata. Pluk. alm. 45. t. 145. f. 1.*

*Arbor venenata. Kalm. it. 2. p. 211.*

*Arbor Poyson-wood-Tree. Act. angl. n. 367. p. 145.*

*Site f. Sitz-diu. Kämpf. amæn. 791. t. 792.*

*Habitat in America septentrionali, Japonia. §*

Copallinum 6. RHUS foliis pinnatis integerrimis, petiolo membranaceo articulado. *Roy. lugdb. 24. Mat. med. 152.*

Rhus

*Rhus elatior*, foliis impari-pinnatis, petiolis membranceis articulatis. *Gron. virg.* 149.

*Rhus virginianum*, lentisci foliis. *Raj. hist.* 1799.

*Rhus oblongiorum* similis americana, gummi candidum fundens non ferrata, foliorum rachi medio alata. *Pluk. alm.* 318. t. 55. f. 1.

*Habitat in America septentrionali.* ♀

7. RHUS foliis pinnatis quinatis integerrimis subrotundis glabris. *Metopium.* *Amaz. acad.* 5. p. 395. \*

*Metopium* foliis subrotundis pinnato-quinatis, racemis alaribus. *Brown. jam.* 177. t. 13. f. 3.

*Terebinthus maxima*, pinnis paucioribus majoribus atque rotundioribus, fructu racemoso sparso. *Sloan. jam.* 167. *hist.* 2. p. 90. t. 199. f. 3. *Raj. dendr.* 51.

*Borbonia* fructu corallino, flore pentapetalo. *Phen. ic.* 61. *Habitat in America.* ♀

8. RHUS foliis ternatis: foliolis petiolatis ovatis nudis integerrimis, caule radicante. *Hort. cliff.* 110. *Gron. virg.* 33. *Roy. lugdb.* 244. *Kalm. it.* 2. p. 296, 214. *Toxicodendron amplexicaule*, foliis minoribus glabris. *Dill. elih.* 390. radicans.

β. *Toxicodendron rectum*, foliis minoribus glabris. *Dill. elih.* 389. t. 291. f. 375.

*Habitat in Virginia, Canada.* ♀

9. RHUS foliis ternatis: foliolis petiolatis angulatis pubescentibus, caule radicante. *Toxicodendrum.*

*Rhus* foliis ternatis: foliolis petiolatis ovatis acutis pubescentibus; nunc integris, nunc sinuatis. *Gron. virg.* 149.

*Toxicodendron triphyllum*, folio sinuato pubescente. *Tournef. inst.* 611.

*Vitis canadensis.* *Munt. phytogr.* t. 60.

*Edera trifolia canadensis.* *Cornut. canad.* 96. t. 97. *Barr. ic.* 228.

*Arbor trifolia venenata virginiana*, folio hirsuto. *Raj. hist.* 1799.

*Habitat in Virginia, Canada.* ♀

Utraque hæc & antecedens dioica, utraque in tenellis plantis caule radicat.

10. RHUS foliis ternatis: foliolis petiolatis ovatis remotissime serratis subtus tomentosis. *Amaz. acad.* 5. p. 395. \* Cominis.

*Cominia arborea*, foliis undulatis pinnato-ternatis, floribus minimis, racemis terminalibus. *Brown. jam.* 205.

*Baccifera indica trifolia*, fructu rotundo monopyreno. *Sloan. jam.* 170. *hist.* 2. p. 100. t. 208. f. 1. *Raj. hist.* 1593.

*Malago Maram. Rheed. mal.* 5. p. 49. t. 25.

*Habitat in Indiis.* §

tomentosum 11. *RHUS* foliis ternatis: foliolis subpetiolatis rhombeis angulatis subtus tomentosis.

*Rhus* foliis ternatis: foliolis ovatis utrinque acutis dentatis: lateralibus petiolatis. *Hort. cliff.* 111.

*Rhus* foliis ternatis: foliolis petiolatis ovatis acutis dentatis. *Vir. cliff.* 25. *Roy. lugdb.* 244.

*Rhus africanum trifoliatum majus*, foliis obtusis & incisifis hirsutis pubescentibus. *Pluk. alm.* 319. t. 219. f. 7.

*Vitex trifolia minor indica serrata.* *Comm. hort.* 1. p. 279. t. 92.

*Habitat ad Cap b. Spei.* §

Cobbe. 12. *RHUS* foliis ternatis: foliolis ovatis acuminatis serratis, pedunculis tomentosis.

*Rhois trifoliata frutex salwaccensis*, floribus juliformibus. *Pet. mus.* 678. *Raj. dendr.* 58.

*Kobbe. Fl. Zeyl.* 441.

*Habitat in Zeylona.* §

angustifolium. 13. *RHUS* foliis ternatis: foliolis petiolatis linearilanceolatis integerrimis subtus tomentosis. *Hort. cliff.* 111. *Roy. lugdb.* 244.

*Rhus africanum trifoliatum majus*, foliis subtus argenteis acutis & margine incisifis. *Pluk. alm.* 319. t. 219. f. 6.

*Rhus fruticosum*, foliis trifidis linearibus acuminatis. *Burm. afric.* 251. t. 91. f. 1.

*Habitat in Æthiopia.* §

lucidum. 14. *RHUS* foliis ternatis: foliolis sessilibus cuneiformibus lævibus. *Vir. cliff.* 25. *Hort. cliff.* 111. *Hort. upf.* 68. *Roy. lugdb.* 243.

*Rhus africanum trifoliatum minus glabrum*, splendet folio subrotundo integro. *Pluk. alm.* 319. t. 219. f. 9.

*Rhus*

*Rhus arboreum trifoliatum latifolium.* *Burm. afr.* 252.  
p. 91. f. 2.

*Vitex trifolia minor indica rotundifolia.* *Comm. hort.*  
1. p. 181. f. 93.

*Habitat ad Cap. b. Spei.* ♀

15. RHUS foliis simplicibus obovatis.

*Cotinus.*

*Cotinus foliis obverse ovatis.* *Hort. cliff.* 111. *Roy.*  
*lugdb.* 243. *Gron. orient.* 92.

*Cotinus coriaria.* *Dod. pempt.* 780.

*Cocconilea* f. *Coccygia.* *Bauh. pin.* 415.

*Habitat in Lombardia Italiae* ♂ *ad radices Apenninorum, in Carniola.* ♀

*Bractea paniculae filiformes, pilosae, longiores pedicellis. Baccae glabrae, referentes dimidium fructus Bursae pastoris.*

## VIBURNUM.

1. VIBURNUM foliis integerrimis ovatis: ramificationibus venarum subtus villosis-glandulosis. *Tinus.*

*Viburnum foliis ovatis integerrimis.* *Hort. upf.* 69.  
*Sauv. monsp.* 136.

*Tinus.* *Hort. cliff.* 109.

*Tinus* 1. 2. 3. *Clus. hist.* 1. p. 49.

*Laurus Tinus* f. *sylvestris*: trium generum. *Bauh.*  
*hist.* 3. p. 418.

*Laurus sylvestris, corni feminae foliis subhirsutis.* *Bauh.*  
*pin.* 461.

β. *Laurus sylvestris, foliis venosis.* *Bauh. pin.* 461.

γ. *Laurus sylvestris, folio minore.* *Bauh. pin.* 461.

*Habitat in Lusitania, Hispania, Italia.* ♀

2. VIBURNUM foliis integerrimis lanceolato-ovatis. *nudum.*

*Tinus foliis ovatis in petiolos terminatis integerrimis.*

*Gron. virg.* 33.

*Habitat in Virginia.* ♀

*Gymæ in hac nude, in præcedenti involucrate.*

3. VIBURNUM foliis subrotundis crenato-serratis glabris. *prunifolium*  
*Gron. virg.* 33.

*Viburnum canadense glabrum.* *Vaill. afl.* 1722. p. 200.

*Mespilus prunifolia virginiana non spinosa, fructu nigricante.* *Pluk. alm.* 249. t. 46. f. 2.

*Habitat in Virginia, Canada.* ♀



- Cominia arborea*, foliis undulatis pinnato-ternatis, floribus minimis, racemis terminalibus. *Brown. jam.* 205.
- Baccifera indica trifolia*, fructu rotundo monoppyreno. *Sloan. jam.* 170. *hist.* 2. p. 100. t. 208. f. 1. *Raj. hist.* 1593.
- Malago Maram. *Rheed. mal.* 5. p. 49. t. 25.
- Habitat in Indiis.* §
- tomentosum 11. *RHUS* foliis ternatis: foliolis subpetiolatis rhombeis angulatis subtus tomentosis.
- Rhus* foliis ternatis: foliolis ovatis utrinque acutis dentatis: lateralibus petiolatis. *Hort. cliff.* 111.
- Rhus* foliis ternatis: foliolis petiolatis ovatis acutis dentatis. *Vir. cliff.* 25. *Roy. lugdb.* 244.
- Rhus africanum trifoliatum majus*, foliis obtusis & incisifis hirsutis pubescentibus. *Pluk. alm.* 319. t. 219. f. 7.
- Vitex trifolia minor indica serrata.* *Comm. hort.* 1. p. 279. t. 92.
- Habitat ad Cap b. Spei.* §
- Cobbe. 12. *RHUS* foliis ternatis: foliolis ovatis acuminatis serratis, pedunculis tomentosis.
- Rhois trifoliata frutex salwaccensis*, floribus juliformibus. *Pet. mus.* 678. *Raj. dendr.* 58.
- Kobbe. *Fl. zeyl.* 441.
- Habitat in Zeylona.* §
- angustifolium. 13. *RHUS* foliis ternatis: foliolis petiolatis linearilanceolatis integerrimis subtus tomentosis. *Hort. cliff.* 111. *Roy. lugdb.* 244.
- Rhus africanum trifoliatum majus*, foliis subtus argenteis acutis & margine incisifis. *Pluk. alm.* 319. t. 219. f. 6.
- Rhus fruticosum*, foliis trifidis linearibus acuminatis. *Burm. afric.* 251. t. 91. f. 1.
- Habitat in Æthiopia.* §
- lucidum. 14. *RHUS* foliis ternatis: foliolis sessilibus cuneiformibus lævibus. *Vir. cliff.* 25. *Hort. cliff.* 111. *Hort. upf.* 68. *Roy. lugdb.* 243.
- Rhus africanum trifoliatum minus glabrum*, splendente folio subrotundo integro. *Pluk. alm.* 319. t. 219. f. 9.

Rhus

*Rhus arboreum trifoliatum latifolium.* *Byrm. afr.* 252.  
p. 91. f. 2.

*Vitex trifolia minor indica rotundifolia.* *Comm. hort.*  
1. p. 181. f. 93.

*Habitat ad Cap. b. Spei.* ☿

15. RHUS foliis simplicibus obovatis.

*Cotinus.*

*Cotinus foliis obverse ovatis.* *Hort. cliff.* 111. *Roy.*  
*lugdb.* 243. *Gron. orient.* 92.

*Cotinus coriaria.* *Dod. pempt.* 780.

*Cocconillea* f. *Coccygia.* *Bauh. pin.* 415.

*Habitat in Lombardia Italiæ & ad radices Apenninorum, in Carniola.* ☿

*Bractæ panicula filiformes, pilosæ, longiores pedicellis. Baccæ glabræ, referentes dimidium fructus Burse pastoris.*

## VIBURNUM.

1. VIBURNUM foliis integerrimis ovatis: ramificationibus venarum subtus villosoglandulosis. *Tinus.*

*Viburnum foliis ovatis integerrimis.* *Hort. ups.* 69.  
*Sauv. monsp.* 136.

*Tinus.* *Hort. cliff.* 109.

*Tinus* 1. 2. 3. *Clus. hist.* 1. p. 49.

*Laurus Tinus* f. *sylvestris*: trium generum. *Bauh. hist.* 3. p. 418.

*Laurus sylvestris, corni feminae foliis subhirsutis.* *Bauh. pin.* 461.

β. *Laurus sylvestris, foliis venosis.* *Bauh. pin.* 461.

γ. *Laurus sylvestris, folio minore.* *Bauh. pin.* 461.

*Habitat in Lusitania, Hispania, Italia.* ☿

2. VIBURNUM foliis integerrimis lanceolato-ovatis. *nudum.*

*Tinus foliis ovatis in petiolos terminatis integerrimis.*  
*Gron. virg.* 33.

*Habitat in Virginia.* ☿

*Cymæ in hac nudæ, in præcedenti involucratæ.*

3. VIBURNUM foliis subrotundis crenato-ferratis glabris. *prunifolium*  
*bris.* *Gron. virg.* 33.

*Viburnum canadense glabrum.* *Vaill. act.* 1722. p. 200.

*Mespilus prunifolia virginiana non spinosa, fructu nigricante.* *Pluk. alm.* 249. t. 46. f. 2.

*Habitat in Virginia, Canada.* ☿

- dentatum. 4. **VIBURNUM** foliis ovatis dentato-serratis plicatis.  
*Habitat in Virginia.* §
- Lantana. 5. **VIBURNUM** foliis cordatis serratis venosis subtus tomentosis. *Vir. cliff.* 25. *Hort. ups.* 68. *Roy. lugdb.* 242. *Saur. monsp.* 136.  
*Viburnum.* *Hort. cliff.* 107.  
*Viburnum vulgo.* *Bauh. pin.* 249.  
*Lantana.* *Dod. pempt.* 701.  
*Habitat in Europæ australioris sepibus argillofis.* §
- acerifolium. 6. **VIBURNUM** foliis lobatis, petiolis lævibus.  
*Opulus.* *Gron. rirg.* 149.  
*Habitat in Virginia.* §
- Opulus. 7. **VIBURNUM** foliis lobatis, petiolis glandulosis.  
*Opulus.* *Hort. cliff.* 109. *Fl. svec.* 249, 264. *Roy. lugdb.* 243.  
*Sambucus aquatica, flore simplici.* *Bauh. pin.* 450.  
*Sambucus palustris.* *Dod. pempt.* 846.
- rostrum. 8. **Sambucus** aquatica, flore globofo pleno. *Bauh. pin.* 456.  
*Habitat in Europæ pratis humidiusculis.* §
- Lentago. 8. **VIBURNUM** foliis serrulatis ovatis acuminatis glabris, petiolis marginatis.  
*Habitat in Canada.* *Kalm.* §  
*Folia crassa Tini, glabra, denticulis minimis serrulata; petiolis longitudinaliter marginatis. Gemma terminalis subulata, basi ventricosa.*
- Cassinoides. 9. **VIBURNUM** foliis ovatis crenatis glabris, petiolis eglandulatis carinatis.  
*Viburnum phillyreæ folio.* *Duham. arb.* 2. p. 350.  
*Cassine foliis ovato-lanceolatis serratis oppositis deciduis, floribus corymbosis.* *Mill. dist.* 1. 83. f. 1.  
*Cassine vere perquam, similis arbuscula, phillyreæ foliis antagonistis.* *Pluk. mant.* 40. *Angl. hortul.* 16. t. 20.  
*Habitat in America septentrionali.* §  
*Folia infima obovata; proxima ovata; superiora lanceolata.*

## CASSINE.

- Peragua. 1. **CASSINE** foliis oblongis serratis.  
*Cassine.* *Hort. cliff.* 72. *Mat. med.* 152. *Roy. lugdb.* 242.

*Celastrus* foliis subrotundis dentatis, flore ac fructu racemoso. *Burm. afr.* 239. t. 85.

*Phillyrea capensis*, folio celastris. *Dill. elth.* 315. t. 239. f. 305.

*Frutex æthiopicus*, alaterni foliis. *Seb. thes.* 1. p. 46. t. 29. f. 5.

*Habitat in Æthiopia.* b

2. **CASSINE** foliis subovatis integerrimis.

Maurocenia

*Maurocenia Hort. cliff.* 108.

*Frangula sempervirens*, folio rigido subrotundo. *Dill. elth.* 146. t. 121. f. 147.

*Cerasus capensis*, fructu rubro, folio fere obtuso. *Pet. gaz.* 1. 57. f. 4.

*Cerasus africana*, foliis plerumque in summo sinuatis, fructu rubro. *Pluk. alm.* 49. t. 158. f. 2.

*Habitat in Æthiopia.* b

## SAMBUCUS.

1. **SAMBUCUS** cymis trifidis, stipulis foliaceis, caule herbaceo.

*Sambucus* caule herbaceo simplici. *Vir. cliff.* 25. *Fl. fvec.* 251, 265. *Mat. med.* 149. *Hort. cliff.* 110. *Roy. lugdb.* 245.

*Sambucus humilis* f. *Ebulus*. *Bauh. pin.* 456.

*Ebulus. Fuchf. hist.* 65.

- β. *Sambucus humilis* f. *Ebulus* folio laciniato. *Bauh. pin.* 456.

*Habitat in Europa.* 2.

2. **SAMBUCUS** cymis quinquepartitis, foliis sub-bipinnatis, caule frutescente.

*Habitat in Canada. D. Kalm.* 2.

*Media inter S. Ebulum & nigram, habitus S. nigrae, at folia pinnata: foliolis inferioribus ternatis, omnibus magis ovatis. Fruticat, sed plerumque perit supra terram hyeme. Cymæ structura uti in nigra. Stipulae filiformes, truncatae, ut in S. nigra; nec in foliola expansa, ut in Ebulo.*

3. **SAMBUCUS** cymis quinquepartitis, caule arboreo. nigra.

*Sambucus* caule arboreo ramoso, floribus umbellatis.

*Roy. lugdb.* 243. *Fl. fvec.* 250, 265. *Mat. med.* 148.

*Sambucus* caule perenni ramoso. *Hort. cliff.* 109.

*Sambucus* fructu in umbella nigro. *Bauh. pin.* 456.

*Sambucus*. *Dod. pempt.* 845.

3. *Sambucus* fructu in umbella viridi. *Bauh. pin.* 456.

laciniata. γ. *Sambucus* laciniato folio. *Bauh. pin.* 456. *Dod. pempt.* 845.

*Habitat in Germania, Japonia.* β

racemosa. 4. *SAMBUCUS* racemis compositis ovatis, caule arboreo.

*Sambucus* caule arboreo ramofo, floribus racemosis.

*Roy. lugdb.* 243. *Sauv. monsp.* 222.

*Sambucus* racemosa rubra. *Bauh. pin.* 456.

*Sambucus* racemosa, acinis rubris. *Bauh. hist.* 1. p. 549.

*Habitat in Europæ australis montibus.* β

## SPATHELIA.

simplex. 1. *SPATHELIA.*

*Spathe* caudice simplici, fronde pinnata comosa, racemo simplicissimo laxo terminali. *Brown. jam.* 187.

*Aceri* aut *Paliuro* affinis arbor caudice non ramofo, foliis forbi, floribus racemosis purpureis, fructu tribus membranulis alato. *Sloan. jam.* 138. *hist.* 2. p. 28. t. 171. *Raj. dendr.* 94.

*Habitat in Jamaica.* β

## STAPHYLEA.

pinnata. 1. *STAPHYLEA* foliis pinnatis. *Hort. cliff.* 112. *Hort. upf.* 69. *Roy. lugdb.* 436.

*Staphylodendron.* *Dalech. hist.* 102.

*Pistacia* sylvestris. *Bauh. pin.* 401.

*Habitat in Europæ australioris succulentis.*

*Styli duo, Fructus bilocularis.*

trifolia. 2. *STAPHYLEA* foliis ternatis. *Hort. cliff.* 112. *Roy. lugdb.* 437. *Cold. noveb.* 62.

*Staphylodendron* triphyllum, vasculo tripartito. *Gron. vrg.* 34.

*Staphylodendron virginianum* trifoliatum. *Herm. lugdb.* 230.

*Pistachia* virginica sylvestris trifolia. *Morif. blas.* 295.

*Habitat in Virginia.* β

*Hec trigyna est.*

## TAMARIX.

gallia. 1. *TAMARIX* floribus pentandris. *Hort. cliff.* 111. *Roy. lugdb.* 436. *Sauv. monsp.* 45.

Ta-

*Tamarix pedunculis nudis, floribus pentandris. Hort. ups. 69 Mat. med. 154.*

*Tamarix altera, folio tenuiore, f. gallica. Bauh. pin. 485.*

*Tamariscus narbonensis. Lob. ic. 218.*

*Habitat in Gallia, Hispania, Italia. b*

2. **TAMARIX** floribus decandris. *Hort. cliff. 111. Roy. germanica. lugdb. 436.*

*Tamarix fruticosa, folio crassiore, f. germanica. Bauh. pin. 485.*

*Tamariscus germanica. Lob. ic. 218.*

*Habitat in Germaniæ locis inundatis. b*

## TURNERA.

1. **TURNERA** floribus sessilibus petiolaribus, foliis ba- ulmifolia, si biglandulosis.

*Turnera foliis serratis, petiolis floriferis. Vir. cliff. 20. Hort. cliff. 112. t. 10. Roy. lugdb. 434. Sp. pl. 1. p. 965.*

*Turnera frutescens ulmifolia. Plum. gen. 15. Mart. cent. 49. t. 49.*

*Cistus urticæ folio, flore luteo, vasculis trigonis. Sloan. jam. 86. hist. 1. p. 202. t. 127. f. 4, 5. Raf. dendr. 492.*

- β. *Turnera frutescens, folio longiore & mucronato.*

*Mart. cent. 49. t. 49.*

*Habitat in Jamaica, ♂ in America calidiore. ♂*

2. **TURNERA** floribus sessilibus petiolaribus, foliis Pumilea. eglandulosis. *Amæn. acad. 5. p. 395. \**

*Pumilea minima hirsuta, foliis angustis profunde serratis. Brown. jam. 188.*

*Chamæ Cistus luteus, foliis parvis serratis. Pet. gaz. 59. t. 38. f. 9.*

*Chamæ Cistus urticæ folio, flore luteo. Sloan. jam. 87. hist. 1. p. 202. t. 127. f. 6.*

*Habitat in Jamaica. ☉*

3. **TURNERA** floribus pedunculatis axillaribus, foliis cistoides, apice serratis.

*Pumilea subhirsuta simplex, foliis linearibus crenatis.*

*Brown. jam. 189.*

*Helianthemoides. Berh. lugdb. 2. p. 269.*

*Helianthemum betonicae* foliis, caule hirsuto. *Plum. spec. 7. icon. 150. f. 1.*

*Chamaecistus* caule hirsuto, folio oblongo angusto sinuato, flore luteo pediculo insidente. *Sloan. jam. 87. hist. 1. p. 202. t. 127. f. 7.*

*Habitat in America meridionali, Jamaica, Surinami.* ☉

*Radix fibrosa.* Caulis *digitalis*, *simplicissimus*, *erectus*, *pilosus*. Folia *subpetiolata*, *lanceolata*, *versus apicem paucis crenis notata*, *subtus tomentosa*. Pedunculi *axillares*, *solitarii*, *uniflori*, *geniculo notati*, *longitudine floris*. Calyx *tomentosus*.

## TELEPHIUM.

Imperati.

1. **TELEPHIUM** foliis alternis. *Hort. ups. 70.*

*Telephium* foliis oblongo-ovatis, racemis secundis terminalibus. *Vir. cliff. 20. Hort. cliff. 73. Roy. lugdb. 434.*

*Telephium repens*, folio non deciduo. *Bauh. pin. 287.*

*Telephium legitimum.* *Clus. hist. 2. p. 67.*

*Cistus* folio majoranæ. *Bauh. pin. 405.*

*Habitat in Gallo-provincia.* 2.

oppositifolium.

2. **TELEPHIUM** foliis oppositis. †

*Telephium myofotidis* foliis amplioribus conjugatis. *Shaw. afric. 572. f. 572.*

*Habitat in Barbaria.*

## CORRIGIOLA.

littoralis.

1. **CORRIGIOLA.** *Hort. ups. 70.*

*Corrigiola* caule fasciculis florum terminato. *Hort. cliff. 112. Roy. lugdb. 420. Hall. helv. 391.*

*Polygonum littoreum* minus, flosculis spadiceo-albicantibus. *Bauh. pin. 281. prodr. 131. Moris. hist. 2. p. 593. f. 5. t. 29. f. 1.*

*Habitat in Galliae, Germaniae littoribus arenosis.* ☉

## PHARNACEUM.

Cerviana.

1. **PHARNACEUM** pedunculis subumbellatis lateralibus æquantibus folia linearia.

*Pharnaceum glabrum*, pedicellis folia æquantibus.

*Phil. bot. 258. \* Sp. pl. 1. p. 272.*

*Cerviana. Minuart. monogr. 1.*

Tri-

*Trichlis pentastemon*, floribus umbellatis. *Hall. gatt.* 26.

*Alfine pumila*, gallii facie, foliis glaucis. *Amm. ruth.* 84.

*Alfine Saxifraga* ind. orient. foliis circa caulem radiatis, flore herbido & muscoso. *Pluk. mant.* 9. t. 332. f. 11.

*Spergula purpurea minima*. *Buxb. cent.* 3. p. 33. t. 62. f. 2.

*Habitat* Rostockii, in Russia, Hispania, Asia. ☉

2. PHARNACEUM pedunculis communibus longissimis, foliis linearibus. *Hort. cliff.* 492. \* *Roy. lugdb.* 221.

*Habitat* in Africa:

*Figura forte habetur* in *Pluk. phyt.* 304. f. 4.

3. PHARNACEUM pedunculis unifloris lateralibus, foliis linearibus.

*Mollugo* foliis septenis linearibus. *Hort. cliff.* 28.

*Alfine procumbens*, gallii facie, africana. *Herm. lugdb.* 19. t. 21.

*Rubia stellaris* f. *Asperula* minor humisparfa, flosculis albis, æthiopica. *Pluk. mant.* 163. t. 331. f. 4.

*Habitat* in Æthiopia.

4. PHARNACEUM racemis bipartitis terminalibus, foliis obcordatis. *Amæn. acad.* 6. afr. 6. \*

*Habitat* ad Cap. b. spel.

## ALSINE.

1. ALSINE petalis bipartitis, foliis ovato-cordatis. *Fl. media. lapp.* 186. *Fl. suec.* 369, 267. *Hort. cliff.* 173 *Gron. virg.* 161. *Roy. lugdb.* 449.

*Alfine media*. *Bauh. pin.* 250.

*Alfine minor*. *Dod. pempt.* 29.

*Habitat* in Europæ cultis. ☉

2. ALSINE foliis setaceis, calycibus aristatis, petalis mucronatis integris brevibus.

*Arenaria* foliis setaceis mucronatis, corollis brevissimis, calycibus inferne mucronatis. *Sp. pl.* 1. p. 424.

*Alfine* foliis fasciculatis tenuissimis durisque, petalis integris. *Hall. helv.* 389. t. 7. f. 2. *Segu. ver.* 3. p. 173.



*Habitat in Helvetia.*

*Stamina huic 5. Segu. l. c.*

- segetalis. 3. **ALSINE** petalis integris, foliis filiformibus, Spergula foliis filiformibus unum latus spectantibus, stipulis membranaceis vaginantibus, pedunculis umbellatis. *Guett. stamp. 299. Dalib. parif. 133.*  
 Alsine segetalis, gramineis foliis unum latus spectantibus. *Vall. parif. 8. t. 3. f. 3.*

*Habitat Parisiis. ☉*

*Vigiliæ hujus ab hora 3 in 9. Seguiet.*

## DRYPIS.

- spinosa. 1. **DRYPIS.** *Ger. emac. 1312.*

*Drypis italica aculeata, floribus albis umbellatis compactis. Mich. gen. 24. t. 23.*

*Drypis theophrasti f. anguillara. Dalech. hist. 1480. Løb. ic. 789. Tabern. ic. 144. Bauh. hist. 3. p. 388.*

*Spina alba foliis vidua, Bauh. pin. 388.*

*Carduus foliis tenuissime spinosis ad instar juniperi. Moris. hist. 3. p. 161. f. 7. t. 32. f. 8.*

*Habitat in Mauritania, Italia. ♂*

*Caules primi procumbentes, tetragoni, brachiati, superne dichotomi. Folia subulata, subtrigona, mucronata; dichotomiæ lanceolata, utrinque tridentata. Corollæ coronatæ Silenes, purpurascens. Capsulæ circumscissæ. D. Mygind.*

## BASELLA.

- rubra. 2. **BASELLA** foliis planis, pedunculis simplicibus,

*Basella. Rheed. mal. 7. p. 45. t. 24. Fl. zeyl. 119.*

*Cuscuta foliis subcordatis. Hort. cliff. 39.*

*Gandola rubra. Rumph. amb. 5. p. 417.*

*Habitat in India. ♂. ☉*

- alba. 3. **BASELLA** foliis ovatis undatis, pedunculis simplicibus folio longioribus.

*Basella flore albo & caulibus viridibus. Thron. carol. 11.*

*Gandola alba. Rumph. amb. 5. p. 417.*

*Mirabili peruvianæ affinis tinctoria, betæ folio, scandens. Pluk. alm. 252. t. 63. f. 1.*

*Habitat in China, Amboina. ♂*

*Caulis herbaceus, sed perennans.*

3. **BASELLA** foliis subcordatis, pedunculis confertis lucida.  
ramosis.

*Habitat in India.* ☉

*Affines nimium sunt hæ tres Species, sed constantes se servant in hortis.*

### SAROTHRA.

1. **SAROTHRA.** *Amæn. acad.* 3. p. 11. *Kalm. it.* 2. *Gentianol-*  
*des.* p. 253.

*Gentiana* caule ramisque ramosissimis, foliis subulatis minimis. *Gron. virg.* 29.

*Centaurium* minus spicatum, angustissimo folio, f. scoparium marilandicum novum. *Pluk. mant.* 43. t. 324. f. 4.

*Habitat in Virginie, Pensylvaniæ apricis glareosis.*

*Gentiana* inaperta an hujus generis?

## TETRAGYNIA.

### PARNASSIA.

1. **PARNASSIA.** *Fl. lapp.* 108. *Fl. svec.* 252. 268. *Hort. palustris.*  
*cliff.* 113. *Roy. tugdb.* 420. *Hall. helv.* 316. *Dalib.*  
*parif.* 96.

*Gramen* parnassii albo simplici flore. *Bauh. pin.* 309.

*Hepatica* alba. *Cord. hist.* 53.

*Pyrola* rotundifolia palustris, flore unico ampliore.

*Morif. hist.* 3. p. 505. f. 12. t. 10. f. 3.

*Habitat in Europæ uliginosis.* 24

### EVOLVULUS.

1. **EVOLVULUS** foliis subrotundis, caule repente, nummulari-  
floribus subsessilibus. rius.

*Convolvulus* nummularius. *Sp. pl.* 157.

*Convolvulus* minor repens, nummulariæ folio, flore cæruleo. *Sloan. jam.* 58. *hist.* 1. p. 157. t. 99. f. 9.

*Raj. suppl.* 382.

*Convolvulus* herbaceus repens, foliis subrotundis, floribus singularibus axillaribus. *Brown. jam.* 153.

*Habitat in Jamaicæ, Barbados pratis.*

2. **EVOLVULUS** foliis cordatis obtusis mucronatis gangeticus  
petiolatis villosis, caule diffuso, pedunculis unifloris.

*Amæn. acad.* 4. p. 306. n. 121.

*Habitat in India.*

- alfinoides. 3. *EVOLVULUS* foliis obcordatis obtusis petiolatis pilosis, caule diffuso, pedunculis trifloris. *Fl. zeyl.* 76.  
*Convolvulus* foliis ovatis obtusis, caule filiformi. *Hort. cliff.* 68.  
*Anagallis hirsuta* minor, foliis alternis, floribus solitariis. *Burm. zeyl.* 11. t. 6. f. 1. & t. 9. f. 1.  
*Vistnu-clandi.* *Rheed. mal.* 11. p. 131. t. 64.  
*Habitat in* Malabaria, Zeylona, Bismagaria, Bahama.
- unifolius 4. *EVOLVULUS* foliis lanceolatis villosis sessilibus, caule erecto, pedunculis trifloris longis. *Syst. nat.* 10. p. 923. *Amæn. acad.* 4. p. 306. n. 122. \*  
*Convolvulus herbaceus erectus*, foliis linearibus, pedunculis longis tenuissimis bibracteatis alaribus. *Brown. jam.* 152. t. 10. f. 2.  
*Habitat in* Jamaica. ☉
- tridentatus. 5. *EVOLVULUS* foliis lineari-cuneiformibus tricuspidatis: basi dilatata dentatis. *Sp. pl.* 157.  
*Convolvulus minor procumbens*, acetosella foliis tricuspidatis. *Pluk. mant.* 117. t. 167. f. 5.  
*Sendera-Clandi.* *Rheed. mal.* 11. p. 133. t. 65.  
 β. *Convolvulus indicus barbatus minor*, foliorum apicibus lunulatis. *Pluk. alm.* 117. t. 276. f. 6.  
*Habitat in* India.

## PENTAGYNIA.

## ARALIA.

- arbores. 1. *ARALIA* caule arboreo, foliis simplicibus, umbellæ universalis radiis uniglandulosis. *Amæn. acad.* 5. p. 369. \*  
*Aralia arborea*, foliis nitidis oblongo-ovatis, umbella laxa radiis singulis glandula notatis. *Brown. jam.* 189.  
*Hedera arbor*, folio hastato. *Plum. spec.* 18. ic. 148.  
*Habitat in* Jamaica. †
- spinosa. 2. *ARALIA* arborescens, caule foliolisque aculeatis. *Vir. cliff.* 26. *Gron. virg.* 34.  
*Aralia caule aculeato.* *Hort. cliff.* 113. *Roy. lugd.* 92.  
*Angelica arborescens spinosa* f. *Arbor indica fraxini* folio, cortice spinoso. *Comm. hort.* 1. p. 89. t. 47. *Raj. hist.* 1798.

Christophoriana arbor aculeata virginienſis. *Pluk. alm.* 98. t. 20.

*Habitat in Virginia.* ‡

3. *ARALIA* caule petiolisque aculeatis, foliolis inermibus villoſis. chinenſis.

Frutex aquoſus maſ. *Rumph. amb.* 4. p. 105. t. 44.?

*Habitat in China.* Osbeck. †

Caulis ſimplex, aculeis ſpineſſimus, nudus, in ſummitate tantum folioſus. Petioli compoſiti, tripartiti, aculeati, villoſi. Folia partialia pinnata & inſuper foliola 2 ad ramificationes petioli. Foliola ſ. pinnae ovatae, ſerratae, ſubtus praefertim villoſae, duplo majores, quam in praecedente. Panicula ramoſa umbellulis numeroſis. Differt itaque a praecedenti, quod nulli aculei foliolis per diſcum ſecundum venas adſperſi: quod folia non glabra ſed duplo majora. Habitus etiam diverſus.

4. *ARALIA* caule folioſo herbaceo laevi. *Hort. upſ.* 70. racemoſa. Cold. noveb. 67.

*Aralia* racemis umbelliferis lateralibus. *Vir. cliff.* 26.

*Aralia* ex alis florifera. *Hort. cliff.* 113. *Roy. lugdb.* 92.

*Panaces* carpimon ſ. racemoſa canadenſis. *Corn. canad.*

74. t. 75. *Barr. ic.* 705.

Christophoriana canadenſis racemoſa & ramoſa. *Moriſ. hiſt.* 1. p. 9. f. 1. t. 2. f. 9.

*Habitat in Canada.* 24

5. *ARALIA* caule nudo, foliis binis ternatis. *Hort. cliff.* nudicaulis. 113. *Gron. virg.* 34. *Roy. lugdb.* 94.

*Aralia* caule nudo, radice repente. Cold. noveb. 66.

Christophoriana, virginiana zarzae radicibus ſurculoſis & fungoſis. *Pluk. alm.* 98. t. 238. f. 5.

*Habitat in Virginia, ſimillima (forte eadem) in Iava.* 24

Caulis breviffimus & fere nullus. Folia duo, decompoſita petiolis longis: Folium ternatum: Foliolis pinnatis: pinnae quinque, ſerratae. Scapus inter duo folia ortus, aphyllus, trifidus, ſ. umbellulas tres ſerens.

## BARRERA.

1. *BARRERA.*

*Erica* capitata ſ. nodiflora, corioſis foliis rectis, aethiopica. *Pluk. mant.* 69. t. 346. f. 7.

capenſis.

*Habitat in Æthiopia. †*

*Frutex; Rami per spatia determinata conferti, erecti, rufescentes adspersi pilis albis. Folia alterna, approximata, ovato-lanceolata, integerrima, pilis albis velut ciliata; Capitula florum terminalia; Simillima est habitu Diosmæ ciliatæ.*

## STATICE.

*Armeria.* 1. **STATICE** scapo simplici capitato, foliis linearibus. *Hort. cliff. 115. Fl. suec. 253, 269. Gron. virg. 150. Roy. lugdb. 194.*

*Limonium aphyllocaulon gramineum globosum. Moris. hist. 3. p. 601. f. 15. t. 1. f. 29.*

*Caryophyllus montanus major, flore globoso. Bauh. pin. 211.*

β. *Cariophyllus montanus minor. Bauh. pin. 211.*

*Habitat in Europæ & Americæ septentrionalis campis. 24*

*Limonium.* 2. **STATICE** scapo paniculato tereti, foliis lævibus. *Statice caule nudo ramofo. Hort. cliff. 115. Fl. suec. 254, 270. Gron. virg. 150. Roy. lugdb. 192.*

*Limonium maritimum majus. Bauh. pin. 192.*

β. *Limonium maritimum minus, oleæ folio. Bauh. pin. 192.*

γ. *Limonium maritimum minimum. Bauh. pin. 192. prodr. 99. Bocc. sic. 25. t. 19.*

*Habitat in Europæ & Virginie maritimis.*

*cordata.* 2. **STATICE** scapo paniculato, foliis spathulatis retusis. *Sauv. monsp. 15. Allion. niceens. 162.*

*Limonium maritimum minus, foliis cordatis. Bauh. pin. 192. prodr. 99.*

*Limonium minimum cordatum f. folio retuso. Barr. ic. 805.*

*Habitat ad mari; Mediterranei littora. 24*

*echioides.* 4. **STATICE** scapo paniculato tereti, foliis tuberculatis. *Statice foliis obverse-ovatis verrucosis, caule ramofo nudo. Sauv. monsp. 15.*

*Limonium minus annuum, bullatis foliis. Magn. monsp. 157. t. 156.*

*Habitat Monspelii. ☉*

*reticulata.* 5. **STATICE** scapo paniculato prostrato, ramis sterilibus retroflexis nudis, foliis cuneiformibus.

Sta-

**Statice** foliis obverse ovatis, ramis nudis humifusis, intricatis. *Sauv. monsp.* 15.

**Limonium** minus, virgulis retiformiter se invicem implicatis. *Pluk. alm.* 221. t. 42. f. 4.

**Limonium** reticulatum. *Raj. hist.* 396. *Bocc. sic.* 82. t. 44.  
*Habitat in Melita.*

6. **STATICE** scapo dichotomo ancipiti, foliis lanceo- spectosa.  
latis mucronatis, floribus aggregatis.

**Limonium** foliis lanceolatis mucronatis radicalibus, caule ancipiti dichotomo, stipulis simplicibus. *Hort. ups.* 71. *Gmel. sibir.* 2. p. 221. t. 91. f. 1.

**Limonium** floribus elegantissimis. *Raj. hist.* 397.

**Limonium** elatius, plantaginis foliis procumbentibus in aculeum terminatis, floribus albis in capitula congestis. *Amm. ruth.* 129.

*Habitat in Tataria.* ♂

*Calyses etiam in hac indivisi sunt.*

7. **STATICE** scapo dichotomo, foliis lanceolatis mucronatis, floribus alternis distantibus. tatarica.  
*Gmel. sibir.* 2. p. 223. t. 92.

**Limonium** orientale, plantaginis folio, floribus umbellatis. *Tournef. cor.* 25. *Boerh. lugdb.* 1. p. 76. t. 76.

**Limonium** elatius, plantaginis foliis procumbentibus in aculeum terminatis, floribus albis spicatis. *Amm. ruth.* 130.

*Habitat in Tataria,* ♂

8. **STATICE** scapo paniculato, foliis subulatis mucronatis. Echinus.

**Limonium** foliis caulinis subulatis pungentibus. *Roy. lugdb.* 192.

**Limonium** orientale frutescens, caryophylli folio in aculeum rigidissimum abeunte. *Tournef. cor.* 25.

**Limonium** cespitosum, foliis aculeatis. *Buxb. cent.* 2. p. 18. t. 10.

β. **Limonium** græcum, juniperi folio. *Tournef. cor.* 25. **Echinus** f. *Tragacantha* altera. *Alp. exot.* 57. t. 56.

*Habitat in Græciæ & Mediæ desertis.*

Folia imbricata, subulata, supra plana, margine scabra, apice mucronato-spinosa; inferiora reflexa.

9. **STATICE** scapo ramofo flexuoso, corymbis terminalibus, foliis nervosis. flexuosa.

Statice foliis ovalibus muticis nervosis, caule nudo alternatim ramoso flexuoso, corymbis terminantibus. *Gmel. sibir. 2. p. 217. t. 89. f. 1.*

*Habitat in Sibiria.*

fusfruticosa. 10. **STATICE** caule fruticoso: superne nudo ramoso, capitulis sessilibus, foliis lanceolatis vaginantibus.

Statice foliis e lanceolato-linearibus: caulinis basi vaginantibus, capitulis alternis sessilibus. *Gmel. sibir. 2. p. 216. t. 88. f. 2. 3.*

*Habitat in Sibiria. D. Gmelin. 5*

monopetala. 11. **STATICE** caule fruticoso folioso, floribus solitariis, foliis lanceolatis vaginantibus.

Statice foliis caulinis lanceolato-linearibus. *Hort. cliff. 116. Roy. lugdb. 192.*

Limonium lignosum. *Bocc. sic. 34, 35. t. 16, 17.*

*Habitat in Sicilia.*

*Gallas sepe fert ex Phalena. Reaum. inf. 3. t. 39. f. 1-4.*

Limonium hispanicum, multifido folio. *Tournef. inst. 242. etiam monopetalum est.*

ferulacea. 12. **STATICE** caule fruticoso ramoso, ramulis imbricatis paleis apice pilo terminatis.

Limonium ferulaceo folio. *Grisl. luss. Morif. hist. 3. f. 15. t. 1. f. 23. Pluk. alm. 221. t. 28. f. 4.*

*Habitat in Barbaria, Lusitania, Hispania. Alströmer.*

Caules teretes, sublignosi, nudi paleis alternis, paniculati. Ramuli ramosissimi, fasciculati, filiformes, imbricati paleis minutissimis, apice setula terminatis. Flores subimbricati, ascendentes, secundi, lutei.

aurea. 13. **STATICE** caule fruticoso folioso ramoso, foliis subulatis.

Statice caulibus fruticosis teretiusculis, foliis subulatis caulinis, floribus ex corymboso fasciculatis. *Gmel. sibir. 2. p. 218.*

Limonium maritimum humile, ephedrae facie. calycibus florum aureis. *Amm. ruth. 132. t. 18. f. 2.*

*Habitat in Dauriae campis montosis.*

sinuata. 14. **STATICE** caule herbaceo, foliis radicalibus alternatim pinnato-sinuatis: caulinis ternis triquetris subulatis decurrentibus. *Hort. upf. 71. Gron. orient. 96.*

Sta-

- Statice* foliis caulinis decurrentibus. *Hort. cliff.* 116.  
*Limonium peregrinum*, foliis asplenii, *Bauh. pin.* 192.  
*Limonii species*, *Rauv. itin.* 313. t. 314.  
*Limonium rauvolfianum*. *Clus. cur.* 33. t. 33.  
 β. *Limonium africanum*, caule elato, foliis integris hirsutis, petalo pallide flavo, calyce amœne purpureo. *Mart. cent.* 84. t. 48.  
 γ. *Limonium caulibus alatis*, asplenii foliis minus asperis, calycibus acutioribus flavescentibus. *Shaw. afric.* 363.  
*Habitat in Sicilia, Palæstina, Africa.* ♀  
 Differt a *Caule ramisque inferne teretibus*, nec alatis.  
 β. *Foliis minus sinuatis.* γ. *Caulium alis membranaceis integris* & *bracteis obtusis glabris*, nec acutis hispidis.

# LINUM.

\* *foliis alternis.*

1. LINUM calycibus capsulisque mucronatis, petalis crenatis, foliis lanceolatis alternis, caule subfoliario. usitatissimum.  
*Linum ramis foliisque alternis lineari-lanceolatis, radice annua.* *Hort. cliff.* 114. *Roy. lugdb.* 433. *Philos. bot.* 261. \*  
*Linum foliis alternis lanceolatis integerrimis, calycibus acuminatis angulatis, capsulis mucronatis.* *Hort. upf.* 71. *Mat. med.* 156.  
*Linum arvense.* *Bauh. pin.* 214. *Raj. hist.* 1073.  
 β. *Linum sativum.* *Bauh. pin.* 214.  
 γ. *Linum sativum humilius, flore majore.* *Barh. lugdb.* 1. p. 284.  
 ζ. *Linum sativum latifolium africanum, fructu majore.* *Tournef. inst.* 339.  
*Habitat hodie inter segetes Europæ australis.* ☉  
*Calycis margines interiores subciliati.*  
 2. LINUM calycibus capsulisque obtusiusculis, foliis alternis lanceolatis integerrimis. *Mill. dict. t.* 166. f. 2.  
*Linum foliis alternis lanceolatis integerrimis, calycibus apice obtusis, capsulis muticis.* *Hort. upf.* 72.  
*Linum perenne, ramis foliisque alternis lineari-lanceolatis.* *Hort. cliff.* 14. *Roy. lugdb.* 434. *Sauv. monsp.* 53.  
*Linum perenne majus cæruleum, capitulo majore.* *Meris. hist.* 2. p. 573.



- Linum sylvestre cæruleum perenne erectius, flore & capitulo majore. Raj. angl. 3. p. 362.*  
*Habitat in Sibiria & Cantabrigiæ. 24*
- viscosum. 3. **LINUM** foliis lanceolatis pilosis quinquenerviis. †  
*Linum sylvestre latifolium, caule viscoso, flore rubro. Bauh. pin. 214. Raj. hist. 1074.*  
*Habitat in montibus Bononiensibus, Sumano, Augustæ vindelicorum, Ingolstadii. 24*
- hirsutum. 4. **LINUM** calycibus hirsutis acuminatis, sessilibus alternis, caule corymbofo.  
*Linum sylvestre latifolium hirsutum cæruleum. Bauh. pin. 339. Moris. hist. 2. p. 573. f. 5. t. 26. f. 5. Raj. hist. 1073.*  
*Linum sylvestre latifolium. Clus. hist. 1. p. 317.*  
*Habitat in Austriæ, Tataria editis graminosis.*  
*Simillimum L. nodifloro, sed Caulis teres; Folia ciliata f. hirsuta, latiora: caulina alterna; ramea opposita. Flores sessiles, alterni. Calyces foliacei, longi. Folia ramea f. floralia & Calyces ciliati setis brevissimis, glandulosis, præsertim versus apicem.*
- narbonense. 5. **LINUM** calycibus acuminatis, foliis lanceolatis sparsis strictis scabris acuminatis, caule tereti basi ramofo.  
*Linum sylvestre cæruleum, folio acuto. Bauh. pin. 214. prodr. 107. Raj. hist. 1073.*  
*Linum sylvestre angustifolium, cæruleo amplo flore. Magn. monsp. 161. Sauv. monf. 54.*  
*Habitat in Galloprovincia, Monipellii, unde Burserus attulit C. Bauhino. 24*  
*Calyces magni, lanceolati, latere membranacei, apice exquisitissime acuminati & quasi mucrone subulato instructi.*
- tenuifolium 6. **LINUM** calycibus acuminatis, foliis sparsis lineari-setaceis retrorsum scabris.  
*Linum perenne, foliis angustioribus, calyce aristato. Hall. helv. 371.*  
*Linum sylvestre angustifolium, flore magno. Bauh. pin. 214.*  
*Linum sylvestre V. angustifolium. Clus. hist. 1. p. 318.*  
 β. **LINUM** sylvestre angustifolium, flore magno intenso cæruleo. *Tournef. infl. 340.*

7. *Linum sylvestre angustifolium*, flore magno violaceo.  
*Tournef. inst.* 340.
8. *Linum sylvestre angustifolium*, flore magno lineis  
purpureis distincto. *Tournef. inst.* 340.
9. *Linum sylvestre angustifolium*, flore magno cæruleo  
& velut cinereo. *Tournef. inst.* 340.  
*Linum sylvestris* V. *angustifolii* aliud genus. *Clus. hist.* 1.  
p. 318.
2. *Linum sylvestre angustifolium*, floribus dilute purpu-  
rascentibus vel carneis. *Bauh. pin.* 214.  
*Linum sylvestre* VI. *angustifolium*. *Clus. hist.* 1. p.  
318.  
*Linum oxyphyllum multicaule*. *Bosc. mus.* 2. p. 169.  
t. 125.
7. *Linaria*, capillaceo folio, altera. *Bauh. pin.* 213.
9. *Linum angustifolium album*, ramusculis per terram  
sparsis. *Tournef. inst.* 340.
1. *Linum sylvestre*, angustis & densioribus foliis, flore  
minore. *Bauh. pin.* 214.  
*Habitat in Galliæ, Helvetiæ, aridis herbosis.* 24  
*Calyces ovati, valde acuminati, uti Folia aculeis mi-  
nutissimis serrati.*
7. **LINUM** calycibus rotundatis obtusis, foliis lineari-  
bus acutis. austriacum.  
*Linum sylvestre angustifolium*, foliis rarioribus. *Bauh.*  
*pin.* 214.  
*Linum sylvestre* 11. *Tabern. ic.* 822.  
*Habitat in Austria inferiore.*
8. **LINUM** calycibus acutis alternis, capsulis muticis, virginia-  
num.  
panicula filiformi, foliis alternis lanceolatis; radi-  
calibus ovatis.  
*Linum* ramis foliisque alternis lanceolatis sessilibus  
nervo longitudinali instructis. *Gron. virg.* 35.  
*Habitat in Virginia, Pennsylvania,*  
*Caulis filiformis, pedalis, paniculatus. Folia radicalia*  
*ovalia, integerrima, lævia; caulina alterna, lanceo-*  
*lata. Flores alterni: pedunculis brevissimis. Caly-*  
*ces acuti. Capsulæ globosæ, muticæ; Corollæ luteæ.*
9. **LINUM** calycibus subferrato-scabris lanceolatis flavum.  
subsessilibus, panicula ramis dichotomis.  
*Linum sylvestre latifolium luteum.* *Bauh. pin.* 214.

- Linum latifolium luteum. Bauh. hist. 3. p. 454.*  
*Linum sylvestre III. latifolium. Clus. hist. 1. p. 317.*  
*Habitat in Austria.*
- strictum. 10. **LINUM** calycibus subulatis, foliis lanceolatis strictis mucronatis: margine scabris.  
*Linum foliis asperis umbellatum luteum. Magn. monsp. 164. Ray. hist. 1076 Sauv. monsp. 53.*  
*Lithospermum, linariæ folio, monspeliensium. Bauh. pin. 259.*  
*Passerina lobelii. Bauh. hist. 3. p. 454.*  
*Habitat in Montpelii, Hispaniæ, Siciliæ squalidis. ☉*  
*Planta corymbosa, stricta. Folia margine scabra; Calycēs subulati, fructu duplo longiores.*
- sesfruticosum. 11. **LINUM** foliis linearibus acutis scabris, caulibus suffruticosis.  
*Linum angustifolium fruticosum valentinum; foliis rigidis & aculeatis. Tournef. inst. 340.*  
*Habitat in Regno Valentino. ‡*
- arborescens. 12. **LINUM** foliis cuneiformibus, caulibus arborescentibus.  
*Linum arborescens. Alp. exot. 19. t. 13.*  
*Habitat in Creta, Italia. 24*
- campanulatum. 13. **LINUM** foliorum basi utrinque puncto glanduloso.  
*Linum foliis imis spatulatis, floribus luteis. Sauv. monsp. 76.*  
*Linum sylvestre luteum, foliis subrotundis. Bauh. pin. 212. Tournef. inst. 340. Magn. char. 261.*  
*Rapunculus nemorosus angustifolius, parvo flore. Bauh. pin. 93.*  
*Campanula lutea linifolia montis lupi, flore volubilis. Lob. ic. 414.*  
*Campanula linifolia lutea. Bauh. hist. 2. p. 817.*  
*Habitat in Galloprovinciæ montibus, Montpelii in monte lupi. 24*  
*Caules simplices, digitales, angulati, terminati sæpius*  
*Floribus tribus, magnis, pedunculatis. Calyces lanceolati, acuminati.*
- maritimum. 14. **LINUM** calycibus ovatis acutis muticis, foliis lanceolatis: inferioribus oppositis.  
*Linum caule simplici, ramis foliisque inferioribus oppositis | linearis lanceolatis. Hort. cliff. 114. \* Roy. lugdb. 434. Sauv. monsp. 147.*

*Linum foliis ovatis, floribus racemosis. Guett. stamp.*  
2. p. 499.

*Linum maritimum luteum. Bauh. pin. 214.*

*Linum luteum narbonense. Bauh. hist. 3. p. 454. Raj.*  
*hist. 1074.*

*Linum sylvestre. Dod. pempt. 534.*

*Habitat in Austria ad Thermas badenses, Monspeli*  
*& in Oriente, ut in Italia ad stagna marina.*

*Planta tota glaberrima. Calycis foliola ovata, acutiuscula.*

14. **LINUM** calycibus acutis, foliis lineari-lanceolatis gallicum.  
alternis, paniculae pedunculis bifloris.

*Linum foliis lineari-lanceolatis, pedunculis bifidis,*  
*floribus luteis. Sauv. monip. 53.*

*Linum sylvestre minus, flore luteo. Bauh. pin. 214.*

*Habitat Monspeli.*

*Pistilla interdum tria, saepius quinque observata a D.*  
*Sauvages.*

\* \* *Foliis oppositis.*

15. **LINUM** foliis oppositis lineari-lanceolatis, floribus africanum  
terminalibus pedunculatis. *Roy. lugdb. 434.*

*Linum africanum luteum, foliis conjugatis. Boerh.*  
*lugdb. 1. p. 284.*

*Habitat in Africa.*

17. **LINUM** foliis floriferis oppositis lanceolatis, flori- mod. flor. um.  
bus alternis sessilibus, calycibus longitudine folio-  
rum. *Hort. cliff. 114.*

*Linum luteum ad singula genicula floridum. Bauh.*  
*pin. 214. Morif. hist. 2. p. 574. f. 5. t. 26. f. 11. Raj.*  
*hist. 1025.*

*Linum luteum sylvestre latifolium. Column. ecphr. 2.*  
*p. 79. t. 80.*

*Habitat in Italiae pratis argillosis.*

*Caulis angulatus, laevis, bifidus f. trifidus. Folia a-*  
*trinque ad basin glandula ovata fusca: caulina, alterna*  
*lanceolata, laevia; ramea opposita, angustiora. Flo-*  
*res alterni, sessiles. Calyx linearis, foliaceus, lon-*  
*gitudine foliorum. Corolla flavescentes. Capsula glo-*  
*bosa, acuminata.*

18. **LINUM** foliis oppositis ovato-lanceolatis, caule di- catharticum  
chotomo, corollis acutis. *Hort. cliff. 372. Fl. suec.*  
*255, 271. Mat. med. 157. Roy. lugdb. 424.*

*Linum pratense*, flosculis exiguis. *Bauh. pin.* 214.  
*Spergula bifolia*, lini capitulis. *Loef. pruss.* 261. t. 86.  
*Habitat in Europæ septentrionalis pascuis succulen-*  
*tis.* ☉

Radiola.

19. *LINUM* foliis oppositis, caule dichotomo, floribus  
 tetrandris tetragynis. *Hort. cliff.* 114. *Fl. suec.* 256,  
 272. *Roy. lugdb.* 434.

*Chamæ Linum vulgare.* *Vaill. parif.* 33. t. 4. f. 6.

*Linocarpum*, serpilli folio, multicaule & multiflorum.  
*Mich. gen.* 23. t. 21.

*Radiola vulgaris serpillifolia.* *Raj. angl.* 3. p. 161. t. 15.  
 f. 3.

*Polygonum minimum* f. *Millegrana minima.* *Bauh.*  
*pin.* 282.

*Habitat in Europæ sabulo inundato.* ☉

quadrifolli-  
um.

20. *LINUM* foliis quaternis.

*Linum africanum*, rubiæ foliis quaternis stellatis. *Raj.*  
*suppl.* 525.

*Habitat in Æthiopia.*

verticilla-  
rum.

21. *LINUM* foliis verticillatis.

*Linum sylvestre*, flore cæcio, foliis verticillatis. *Borr.*  
*mus.* 2. p. 49. t. 42. *Raj. suppl.* 524. *Barr. rar.* 685.  
 t. 1226.

*Habitat in Italia.* ☉

*Calyces* ☿ *folia superiora villosa.*

## ALDROVANDA.

vesiculosa.

1. *ALDROVANDA.* *Mont. aël. bonon.* 2. p. 3. p. 404.  
 t. 12.

*Lenticula palustris indica*, foliis subrotundis binis ca-  
 pillamentis ad imum barbatis. *Pluk. a'm.* 211. t. 41.  
 f. 6.

*Habitat in Italiæ* ☿ *Indiæ paludosis.* *D. Monti.*

## DROSERA.

rotundifolia

1. *DROSERA* scapis radicatis, foliis orbiculatis. *Fl.*  
*lapp.* 109. *Fl. suec.* 257, 273. *Mat. med.* 158. *Fl. zeyl.*  
 120. *Gron. virg.* 35. *Roy. lugdb.* 120.

*Ros folis folio rotunda.* *Bauh. pin.* 357.

*Salsiflora* f. *Ros folis.* *Thal. herc.* t. 9. f. 1.

*Habitat in Europæ, Asiæ, Americæ paludibus.*

2. *DRO-*

2. DROSEBA scapis radicatis, foliis oblongis. *Fl. lapp.* longifolia  
110. *Fl. svec.* 258, 274. *Roy. lugdb.* 417.  
Ros folis folio oblongo. *Bauh. pin.* 357.  
Salsifiora f. Sponfa folis f. Ros folis f. Rorella. *Thal.*  
*herc.* 116. t. 9. f. 2.  
*Habitat in Europa ubique cum præcedente; an itaque*  
*satis diversa species?*
3. DROSEBA scapis radicatis, foliis subulatis subtus lufitanica,  
convexis, floribus decandris  
Ros folis lufitanicus, foliis alphodeli minoris. *Morif.*  
*hist.* 3. p. 620. f. 15. t. 4. f. 4. *Pluk. alm.* 323. t. 117.  
f. 2. *Raj. suppl.* 551.  
*Habitat in Lufitania.*  
Generis *Drosera* esse Speciem negat *Rajus*. Stamina  
huic constantia observavit *Alstræmer*; adeoque  
Oxalls diceretur, si flores essent umbellati & fo-  
liorum proprietas non detineret cum Capsula unilo-  
culari.
4. DROSEBA scapis radicatis, foliis lanceolatis. capensis.  
Drosera foliis ad radicem longissimis, floribus spica-  
catis. *Burm. afr.* 209. t. 75. f. 1.  
Ros folis africanus, foliis prælongis, caule nudo al-  
tissimo. *Raj. suppl.* 515.  
Ros folis africanus, folio lato & longo. *Herm. afr.* 19.  
*Habitat in Æthiopia.*
5. DROSEBA caule simplici folioso, foliis lanceolatis. cistiflora.  
*Amæn. acad.* 6. afr. 7. \*  
Drosera foliis ad caulem oblongis alternis, flore am-  
plo purpureo. *Burm. afr.* 210. t. 75. f. 2.  
Ros folis folio angusto, flore amplo. *Breyn. prodr.* 3.  
t. 22. f. 2.  
Ros folis africanus, cisti flore albo, caule folioso.  
*Raj. suppl.* 515.  
*Habitat ad Cap. b. spei.*
6. DROSEBA caule ramoso folioso, foliis linearibus. indica.  
*Fl. zeyl.* 121.  
Ros folis ramosus, caule folioso. *Burm. zeyl.* 207. t.  
94. f. 1.  
Araca-puda. *Rheed. mal.* 10. p. - - t. 20.  
*Habitat in India.*

## CRASSULA.

- coccinea. 1. **CRASSULA** foliis planis cartilagineo-ciliatis basi connato-vaginantibus. *Vir. cliff.* 26. *Hort. cliff.* 116. *Roy. lugdb.* 454.  
*Cotyledon africana* frutescens, flore umbellato coccineo. *Comm. rar.* 24. t. 24. *Bratl. succ.* 5. p. 7. t. 50.  
*Cotyledon africana* frutescens, flore carneo amplo. *Breyn. prodr.* 3. p. 30. t. 20. f. 1.  
*Habitat in Æthiopia.* §
- perfoliata. 2. **CRASSULA** foliis lanceolato-subulatis sessilibus connatis plano-canaliculatis: subtus convexis. *Hort. cliff.* 116. *Roy. lugdb.* 455.  
*Crassula altissima* perfoliata. *Dill. elth.* 114. t. 96. f. 113. *Mill. ic.* 108.  
*Aloë africana* caulescens perfoliata glauca & non spinosa. *Comm. præl.* 74. t. 23.  
*Habitat in Æthiopia.* §
- subulata. 3. **CRASSULA** caule herbaceo, foliis subulatis teretibus patentibus. †  
*Sedum africanum* umbellatum album. *Herm. lugdb.* 550. t. 552.  
*Habitat in Æthiopia.* ☉
- tetragona. 4. **CRASSULA** foliis subulatis obsolete tetragonis. *Hort. cliff.* 116. \* *Roy. lugdb.* 455.  
*Cotyledonoides africanum.* *Bratl. succ.* 5. p. 18. t. 11. f. 41.  
*Habitat in Æthiopia.* §  
*Caulis levis, tripedalis. Folia opposita, patentia, subincurvata. Pedunculus terminalis, nudus, teres: Cyma trifida, ramosissima, fastigiata. Corollæ albæ: Antheris purpureis.*
- centauroides. 5. **CRASSULA** caule herbaceo dichotomo, foliis cordatis sessilibus, pedunculis unifloris. *Amæn. acad.* 6. afr. 8. \*  
*Sedoides africana* annua centauroides. *Herm. parad.* 169.  
*Habitat in Æthiopia.* ☉
- dichotoma. 6. **CRASSULA** caule herbaceo dichotomo, foliis ovato-lanceolatis, pedunculis unifloris. *Amæn. acad.* 6. afr.

*Sedum africanum* annuum, centaurii minoris folio, flore aureo. *Herm. lugdb.* 550. t. 553. \*

*Habitat in Æthiopia.* ☉

7. *CRASSULA* caule herbaceo erecto, ramis dichotomis foliis oppositis obovatis strigosis, pedunculis unifloris. *Amæn. acad.* 6. afr. 9. \*

*Habitat in Æthiopia.*

8. *CRASSULA* caule herbaceo prostrato, foliis oppositis ovatis gibbis imbricatis, floribus sessilibus foliariis. *Amæn. acad.* 6. afr. 10. \*

*Ficoides africana* annua minima muscosa. *Herm. par.* 170.

*Habitat in Æthiopia.* ☉

9. *CRASSULA* foliis oppositis obtuse ovatis, integerri- cultrata, mis: hinc angustioribus. *Hort. cliff.* 496. *Roy. lugdb.* 455.

*Crassula anacampserotis* folio. *Dill. elth.* 115. t. 97. f. 114.

*Habitat in Æthiopia.* †

10. *CRASSULA* foliis oppositis ovalibus planiusculis distinctis ciliatis, corymbis terminalibus. *Hort. cliff.* 496. *Roy. lugdb.* 455.

*Crassula caulescens*, foliis sempervivi cruciatis. *Dill. elth.* 116. t. 98. f. 116.

*Habitat in Æthiopia.* 24

11. *CRASSULA* foliis oppositis patentibus connatis scabra, scabris ciliatis, caule retrorsum scabro.

*Crassula mesembryanthemi* facie, foliis longioribus asperis. *Dill. elth.* 117. t. 99. f. 117.

*Cotyledon africana* frutescens, foliis asperis angustis acuminatis, flore virescente. *Mart. cent.* 24. t. 24.

*Habitat in Africa?*

12. *CRASSULA* foliis serrato-dentatis planis alternis, alternifolia, caule simplicissimo, floribus pendulis. *Hort. cliff.* 497. \* *Roy. lugdb.* 455.

*Crassula* foliis oblongis acutis dentatis, flore ex alis solitario flavo. *Burm. afr.* 58. t. 24. f. 1.

*Cotyledon*, flore luteo, media. *Herm. lugdb.* 191.

*Habitat in Æthiopia.*

13. *CRASSULA* foliis subulatis radicalibus, caule nudicaulis do. *Hort. cliff.* 116. *Roy. lugdb.* 455.



- Crassula eespitosa longifolia*, *Dill. elth.* 116. t. 98. f. 115.  
*Habitat in Æthiopia.* 24
- punctata, 14. *CRASSULA* foliis oppositis ovatis punctatis ciliatis: inferioribus oblongis.  
*Crassula* caule flaccido, foliis connatis cordatis succulentis, floribus confertis. *Mill. dict.*  
*Telephium frutescens*, floribus spicatis minimis, folio triangulari crasso. *Raj. suppl.* 118?  
*Habitat in Æthiopia.* *Burmans.*  
*Caulis teres, lavis, simplex. Folia opposita, disticha, oblonga, sessilia, carnosae, excavato-punctata, subtus convexa, tenerrime ciliata: Floralia ovata. Corymbi axillares, brevissimi, fastigiati. Corollæ campanulatae, albæ, limbo reflexo. Antheræ purpureæ.*
- orbicularis, 15. *CRASSULA* caule flaccido prolifero determinato folioso. foliis patentissimis imbricatis. *Hort. cliff.* 496. *Roy. lugdb.* 455.  
*Crassula orbicularis repens*, foliis sempervivi. *Dill. elth.* 119. t. 100. f. 118.  
*Habitat in Æthiopia.* 24
- pellucida, 16. *CRASSULA* caule flaccido repente, foliis oppositis, *Crassula*, portulacæ facie, repens. *Dill. elth.* 119. t. 100. f. 119.  
*Habitat in Æthiopia.*
- Portulacaria 17. *CRASSULA*? foliis obovatis oppositis, caule arboreo.  
*Anacamperos* caule arboreo, foliis cuneiformibus oppositis *Hort. cliff.* 207. *Hort. ups.* 147.  
*Crassula portulacæ facie arborescens.* *Dill. elth.* 120. t. 101. f. 120.  
*Habitat in Æthiopia.* 5  
*Fructificatio quantum novi, nondum detecta est; maxime seduceret habitus, si Crassula non esset. Nomino hanc, ut excitem alios in flores ejus inquirere, quibus datur Cap. b. spei adeundi occasio.*

## SIBBALDIA.

- procumbens. 1. *SIBBALDIA* procumbens, foliis tridentatis. *Fl. lapp.* 111. *Fl. suec.* 260, 275. *Hall. herb.* 342. *Roy. lugdb.* 276.

## PENTANDRIA POLYGYNIA.

407

*Fragariæ sylvestri affinis planta, flore luteo. Sibb. scot.*

2. p. 25. t. 6. f. 1. *Oed. dan. t. 32.*

*Pentaphylloides fruticosum minimum procumbens, flore luteo. Pluk. alm. 284. t. 212. f. 3.*

*Habitat in Alpibus Lapponiæ, Helvetiæ, Scotiæ. 24*

2. **SIBBALDIA** erecta, foliolis linearibus multifidis. erecta.

*Pentaphylloides foliis tenuissime laciniatis, flosculis carneis. Amm. ruth. 112. t. 15.*

*Habitat in Sibiria. D. Gmelin.*

## P O L Y G Y N I A

### MYOSURUS.

1. **MYOSURUS.**

*minimus.*

*Myosurus foliis integerrimis. Hort. cliff. 117. Fl. svec. 261, 276. Roy. lugdb. 492.*

*Holosteo affinis Cauda muris. Bauh. pin. 190.*

*Cauda muris. Dod. pempt.*

*Habitat in Europæ collibus apricis aridis. ☉*



## Classis VI.

HEXANDRIA  
MONOGYNIA.

## BROMELIA.

Ananas.

1. BROMELIA foliis ciliato-spinosis mucronatis, spica comosa.

Bromelia foliis spinosis, fructibus connatis caulem cingentibus. *Hort. cliff.* 127. *Hort. upf.* 73. *Roy. lugdb.* 24.Carduus brasiliensis, foliis aloës. *Bauh. pin.* 384.Ananas acostæ. *Comm. hort.* 1. p. 109. t. 57.Ananas aculeatus, fructu ovato carne albida. *T. Trew. chret.* t. 2.Anassa. *Rumph. amb.* 5. p. 227. t. 81.Kapa-Tliskka. *Rheed. mal.* 11. p. 1. t. 1, 2.

- β. Ananas aculeatus, fructu pyramidato: carne aurea.
- Tournef. inst.*
- 653.

- γ. Ananas aculeatus, fructu conico: carne aurea.
- Pluk. spec.*
- 20.

- δ. Ananas lucide virens, folio vix ferrato.
- Dill. clth.*
25. t. 21. f. 22.

Ananas non aculeatus Pitta dictus. *Plum. spec.* 20.

Habitat in nova Hispania, Surinamo. 24

Pinguin.

2. BROMELIA foliis ciliato-spinosis mucronatis, racemo terminali.

Bromelia foliis aculeatis, racemo laxo terminali. *Hort. cliff.* 129. *Roy. lugdb.* 25. *Brown. jam.* 193. n. 4.Ananas americana sylvestris altera minor. *Pluk. mant.* 29. t. 258. f. 4.Pinguin. *Dill. clth.* 320. t. 240. f. 311.

Habitat in Jamaica, Barbados. 24

karatas.

3. BROMELIA acaulis, floribus aggregatis sessilibus subradicalibus.
- Jacq. amer.*
- 18.

- Bromelia foliis ciliato-spinosis mucronatis, panicula diffusa.
- Sp. pl.*
1. p. 285.

Bromelia foliis spinosis oblique recurvis, fructibus dispersis racemosis. *Hort. upf.* 73.Karatas foliis altissimis angustissimis & aculeatis. *Plum. gen.* 10?

Ca-

Caraguata-acanga. *Pis. bras.* 190. t. 191.  
 Mexocotl f. Manguei. *Hern. mex.* 272. *Morif. hist.*  
 2. f. 4. t. 22. f. 7.

*Habitat in America meridionali.* 24

*Fructus inter folia aculeata ad ipsam radicem locata, ut  
 difficillime ob spinas recurvatas eruatur.*

4. BROMELIA foliis ferrato-spinosis obtusis, spicis al- lingulatis.  
 ternis.

*Bromelia ramosa & racemosa, foliis arundinaceis fer-  
 ratis. Plum. gen.* 26. *ic.* 64. *f.* 1.

*Habitat in America meridionali.* 24

5. BROMELIA foliis radicalibus dentato-spinosis, cau- nudicaulis.  
 linis integerrimis.

*Bromelia pyramidata, aculeis nigris. Plum. gen.* 46.  
*ic.* 62.

*Habitat in America meridionali.* 24

## TILLANDSIA.

1. TILLANDSIA culmo paniculato. utriculatis.

*Tillandsia foliis basi conniventibus in utriculum. Hort.  
 cliff.* 129.

*Viscum caryophylloides maximum, flore tripetalo pal-  
 lide luteo, semine filamentoso. Sloan. jam.* 76. *hist.*

1. p. 88. *Raj. suppl.* 405.

*Visci modo arboribus indicis adnascens. Baub. pin.*  
 423.

*Caraguata multiplici spica, flore albo. Plum. gen.* 10.

*Habitat in America meridionalis arboribus.* 24

4. TILLANDSIA foliis superne ferrato-spinosis, spica ferrata,  
 comosa,

*Tillandsia parasitica maxima, foliis amplioribus obtu-  
 sis ciliato-subspinosis, racemo pyramidato. Brown.  
 jam.* 195.

*Caraguata clavata & spicata, foliis ferratis. Plum.  
 gen.* 10. *ic.* 75. *f.* 1.

*Habitat in America meridionali.*

9. TILLANDSIA foliis lanceolato-lingulatis integerrimis: basi ventricosus. lingulatis.  
*Jacq. hist.* 62.

*Tillandsia media parasitica, foliis oblongis obtusis, flo-  
 ribus comosis terminalibus. Brown. jam.* 194.

*Caraguata latifolia & clavata. Plum. gen.* 10. *ic.* 74.

*Viscum caryophylloides maximum*, capitulis in summitate conglomeratis. *Sloan. jam. 77. hist. 1. p. 189. t. 120. Raj. suppl. 405.*

*Habitat in Americæ meridionalis arboribus vetustis.*

tenuifolia.

4. **TILLANDSIA** foliis filiformibus integerrimis, spica simplici laxa. *Roy. lugdb. 25. Jacq. hist. 1. 63.*

*Tillandsia parasitica parva*, foliis tenuissimis erectis, spica breviori simplici disticha. *Brown. jam. 194.*

*Renealmia spica multiplici angustifolia*, flore cæruleo. *Plum. gen. 37. ic. 238. f. 2.*

*Viscum caryophylloides minus*, foliorum imis viridibus apicibus subrubicundis, flore tripetalo purpureo, semine filamentoso. *Sloan. jam. 77. hist. 1. p. 190. t. 122. f. 1.*

*Habitat in Americæ meridionalis arboribus.*

paniculata.

5. **TILLANDSIA** foliis radicalibus brevissimis, culmo subnudo, ramis subdivisis adscendentibus.

*Tillandsia parasitica major*, foliis attenuatis basi ventricosus, racemo laxo spatioso assurgenti. *Brown. jam. 194.*

*Renealmia ramosissima*, foliis variegatis & circinatis. *Plum. gen. 27. ic. 237.*

*Habitat in America meridionali.*

polystachia.

6. **TILLANDSIA** culmo spicis imbricatis lateralibus.

*Viscum caryophylloides angustifolium*, floribus cæruleis. *Catesb. car. 2. p. 89. t. 89?*

*Renealmia spica multiplici*, flore albo. *Plum. gen. 37.*

- β. *Renealmia*, spica multiplici, alia angustifolia. *Plum. gen. 37.*

*Habitat in America calidiore.*

monostachia

7. **TILLANDSIA** foliis linearibus canaliculatis reclinatissimis, culmo simplici imbricato, spica simplici.

*Renealmia non ramosa squamata*, & floribus niveis.

*Plum. gen. 37. ic. 238. f. 1.*

*Habitat in America meridionali.*

recurvata.

8. **TILLANDSIA** foliis subulatis scabris reclinatis, culmis unifloris, glumis bifloris. *Roy. lugdb. 25.*

*Tillandsia parasitica parva pruinosa*, scapo tenui bifloro. *Brown. jam. 194.*

*Viscum caryophylloides minus*, foliis pruinæ instar can-

candicantibus, flore tripetalo purpureo, semine filamentoso. *Sloan. jam. 77. hist. 1. p. 190. t. 121. f. 1. Raj. suppl. 406.*

*Habitat in Jamaica arboribus.*

♂ **TILLANDSIA** filiformis ramosa intorta scabra. *usneoides.*  
*Hort. cliff. 129. Gron. virg. 36. Barr. æquin. 99. Roy. lugdb. 25.*

**Renealmia** parasitica, caule filiformi ramoso geniculato longissimo, foliis subulatis. *Brown. jam. 192.*

**Viscum** caryophylloides tenuissimum e ramis arborum musci in modum dependens, foliis pruinæ instar candicantibus, flore tripetalo, semine filamentoso. *Sloan. jam. 77. hist. 1. p. 191. t. 122. f. 2. 3. Raj. suppl. 406.*

**Cuscuta**, ramis arborum innascens, caroliniana, filamentis lanugine tectis. *Pluk. alm. 126. t. 26. f. 5. Camanbaya. Marcgr. bras. 46. Pet. gaz. t. 62. f. 12. Habitat in Virginie, Jamaica, Brasiliæ arboribus.*

**Renealmiæ Plumieri** solo fructu, eoque minus diverso, differunt ab ejusdem Caraguatis, consultius visum hæc duo genera combinare, quam difficile & tamen ambigue distinguere.

## BURMANNIA.

1. **BURMANNIA** spica gemina. *Burm. zeyl. 50. t. 20. f. disticha.*  
*1. Fl. zeyl. 128.*

**Burmattia** spica duplici. *Hort. cliff. 128. Roy. lugdb. 125.*

**Planta** zeylanica aquatica, lato & brevi gramineo folio. *Raj. suppl. 559.*

*Habitat in Zeylonæ paludosis.*

2. **BURMANNIA** flore gemino.

**Burmattia** flore duplici. *Hort. cliff. 128.*

**Burmattia** scapo bifloro. *Gron. virg. 36.*

*Habitat in Virginie paludosis.*

biflora.

## TRADESCANTIA.

1. **TRADESCANTIA** erecta lævis, floribus congestis. *virginiana.*

**Tradescantia.** *Hort. cliff. 127. Hort. upf. 73. Gron. virg. 36. Roy. lugdb. 37.*

**Ephe-**

*Ephemerum phalangoides tripetalum non repens virginianum gramineum. Moris. hist. 3. p. 605. f. 15. t. 2. f. 4.*

*Allium f. Moly virginianum. Bauh. pin. 506.*

*Habitat in Virginia, 24*

malabarica. 2. *TRADESCANTIA erecta laevis, pedunculis solitariis longissimis.*

*Tali-pulli. Rheed. mal. 9. p. 123. t. 63.*

*Habitat in Malabarica.*

geniculata. 3. *TRADESCANTIA hirsuta procumbens. Jacq. amer. 18. hist. t. 64.*

*Ranunculus f. Damasonium repens, parnassiae foliis villosis. Plum. ic. 116. f. 2.*

*Habitat in America meridionali.*

### PONTERDERIA.

ovata. 1. *PONTERDERIA foliis ovatis, floribus capitatis. Narukila. Rheed. mal. 11. p. 67. t. 34. Raj. suppl. 573. Habitat in Malabaricae aquis. 24*

cordata. 2. *PONTERDERIA foliis cordatis, floribus spicatis. Pontederia floribus spicatis. Hort. cliff. 133. Gron. virg. 37. Roy. lugdb. 37. Cold. noveb. 69. Gladiolus lacustris virginianus caeruleus, sagittae folio. Pet. gaz. 1. t. 1. f. 12.*

*Sagittae similis planta palustris virginiana, spica florum caerulea. Moris. hist. 3. p. 618. f. 15. t. 4. f. 8.*

*Plantagini aquaticae quodammodo accedens, floribus caeruleis hyacinthi spicatis. Plum. mant. 152. t. 349 f. ult. Habitat in Virginiae aquis. 24*

hastata. 3. *PONTERDERIA foliis hastatis, floribus umbellatis. Pontederia floribus umbellatis. Fl. zeyl. 129.*

*Sagittae quodammodo similis planta maderaspatana, floribus medio caule quasi ex utriculo prodeuntibus. Pluk. alm. 326. t. 220. f. 8. Moris. hist. 3. p. 618. f. 15. t. 4. f. 7.*

*Carimgola. Rheed. mal. 11. p. 91. t. 44.*

*Habitat in India. 24*

### HÆMANTHUS.

coccineus. 1. *HÆMANTHUS foliis linguiformibus planis laevibus. Roy. lugdb. 42.*

Hæ-

*Hæmanthus* foliis obtusis basi truncatis. *Comm. hort.*

2. p. 127. t. 64.

*Narcissus indicus puniceus. Ferr. cult.* 137.

*Narcissus indicus serpentarius. Hern. mex.* 885. t. 899.

*Habitat ad Cap. b. spei.* 24

2. *HÆMANTHUS* foliis linguiformibus ciliatis. ciliaris.

*Lilium africanum sphaericum*, floribus obsolete puniceis minoribus. *Hern. lugdb.* 375.

*Bulbus oblongus aethiopicus*, foliis guttatis & cilli instar pilosis. *Breyn. cent.* t. 39.

*Habitat ad Cap. b. spei. Burmannus.* 24

*Bulbus magnus. Folia præcedentis, sed ciliata ( ut in Amaryll. guttata, cui folia angustiora E. fere narcissina ) : Cilia in hac fusca, in Amaryllide pallida. Flores nobis non protulit; ex herbario Oldenlandi, in quo flores Hæmanthi eidem paginæ cum foliis adglutinati, hujus generis judicio.*

3. *HÆMANTHUS* foliis lanceolato-ovatis undulatis puniceis. erectis. *Hort. cliff.* 127. *Hort. upf.* 38. *Roy. lugdb.* 42.

*Hæmanthus colchici foliis*, perianthio herbaceo. *Dill. elth.* 167. t. 140. f. 2. *Trew. chret.* 44.

*Hyacintho affinis africana*, caule maculato. *Seb. mus.* t. p. 20. t. 12. f. 1, 2, 3.

*Satyrium e guinea. Swert. flor.* 1. p. 62. f. 3. *Morif. hst.* 3. p. 491. f. 12. t. 12. f. 11. *Rudb. elyf.* 2. p. 210. f. 3.

*Habitat in Guinea.* 24

4. *HÆMANTHUS* foliis linearibus carinatis. *Mill* carinatus. *dict.* t.

*Habitat ad Cap. b. spei.* 24

## GALANTHUS.

1. *GALANTHUS. Hort. cliff.* 134. *Hort. upf.* 73. *Roy. nivalis. lugdb.* 35.

*Leucojum hulbosum trifolium minus. Bauh. pin.* 56.

*Erangelia. Reneal. spec.* 97. t. 96.

*Habitat ad radices Montium Veronæ, Tridenti, Carniolæ, Viennæ.* 24

1. LEU.



## LEUCOJUM.

- vernum. 1. **LEUCOJUM** spatha uniflora, stylo clavato.  
*Leucojum. Hort. cliff. 135. Hort. upf. 74. Roy. lugdb. 35.*  
*Leucojum bulbosum vulgare. Bauh. pin. 55. Rudb.*  
*elys. 2. p. 95. f. 1.*  
*Leucojum bulbosum. Clus. hist. 1. p. 169.*  
*Habitat in Germaniæ, Helvetiæ, Italiæ umbrosis pratis, ad rivulos.*
- æstivum. 2. **LEUCOJUM** spatha multiflora, stylo clavato.  
*Leucojum bulbosum majus 1. multiflorum. Bauh. pin. 55.*  
*Leucojum bulbosum ferotinum majus. 1. Clus. hist. 1.*  
*p. 170.*  
*Polyanthemum. Renealm. spec. 99. t. 100.*  
*Habitat in Pannonia, Hetruria, Monipellii. 24*  
*Folia hujus lata ut in L. verno, nec filiformia ut in L. autumnali.*
- autumnale. 3. **LEUCOJUM** spatha multiflora, stylo filiformi.  
*Lafl. it. 136.*  
*Leucojum bulbosum autumnale. Bauh. pin. 56.*  
*Leucojum bulbosum autumnale tenuifolium. Clus. hist.*  
*1. p. 170.*  
*Tricophyllum. Reneal. spec. 101. t. 100.*  
*Habitat in Lusitania. 24*

## NARCISSUS.

- Poëticus. 1. **NARCISSUS** spatha uniflora, nectario rotato brevissimo scarloso crenulato. *Hort. upf. 74.*  
*Narcissus foliis ensiformibus, floris nectario rotato brevissimo. Hort. cliff. 134. Roy. lugdb. 35. Sauv. monf. 17.*  
*Narcissus albus, circulo purpureo. Bauh. pin. 48.*  
*Narcissus medio purpureus. Dod. pempt. 223.*
- β. *Narcissus medio purpureus multiplex. Bauh. pin. 54.*  
*Habitat in G. Narbonensi, Italia. 24*
- Pseudo-Narcissus. 2. **NARCISSUS** spatha uniflora, nectario campanulato, erecto crispo æquante petala ovata,  
*Narcissus foliis ensiformibus, florum nectario longitudine petalorum. Hort. cliff. 134. Hort. upf. 74. Sauv. monsp. 17.*  
*Narcissus sylvestris pallidus, calyce luteo. Bauh. pin. 52.*  
*Narcissus luteus sylvestris. Dod. pempt. 227. f. 12.*  
 β. Nar-

- β. *Narcissus sylvestris multiplex*, calyce carens. *Bauh.*  
*pin.* 54.  
 γ. *Narcissus luteus sylvestris*, duplici f. triplici tubo au-  
 reo. *Bauh. pin.* 54.  
*Habitat in Gallia, Angliæ, Hispaniæ, Italiæ nemo-*  
*ribus.* 24.  
 3. *NARCISSUS* spatha uniflora, nectario campanulato bicolor.  
 margine patulo crispo æquante petala.  
*Narcissus albus*, calyce flavo, alter. *Bauh. pin.* 52.  
*Narcissus major totus luteus*, calyce prælongo. *Bauh.*  
*pin.* 52. *Rud. elys.* 2. p. 71. f. 9.  
*Habitat in Europa australi.* 24.  
*Similis N. Pseudo-Narcisso, sed Petala alba; Necta-*  
*rium saturate luteum, majus: limbo patulo, undulato,*  
*crenato.*  
 4. *NARCISSUS* spatha uniflora, nectario obconico e- minor.  
 recto crispo sexfidō æquante petala lanceolata.  
*Narcissus parvus totus luteus.* *Bauh. pin.* 53. *Rudb.*  
*elys.* 2. p. 72. f. 11.  
*Narcissus sylvestris pallidus minimus.* *Barr. ic.* 976.  
*Pseudo-Narcissus minor hispanicus latifolius.* *Clus. hist.*  
*1. p.* 165.  
*Bulbocodium minus.* *Bauh. hist.* 2. p. 597.  
*Habitat in Hispania.* 24.  
*Affinis maxime N. Pseudo-Narcisso, sed triplo omni-*  
*bus partibus minor. Scapus vix striatus. Spatha vi-*  
*rens. Flos magis nutans. Petala basi inter se distin-*  
*cta, lanceolata, recta, nec obliqua aut ovata. Necta-*  
*rii margo sexfidus, undulatus, crispus.*  
 5. *NARCISSUS* spatha uniflora, nectario cylindrico moschatus.  
 truncato subrepando æquante petala oblonga.  
*Narcissus totus albus, nutante flore, longa tuba.* *Barr.*  
*ic.* 945, 946, 954, 953, 921, 922.  
*Narcissus albus*, calyce flavo, moscari odore. *Bauh.*  
*pin.* 52. *Rudb. elys.* 2. p. 69. f. 6. & 73. f. 15, 16.  
*Narcissus flavus*, tubo rotundo. *Bauh. pin.* 52. *Rudb.*  
*elys.* 2. p. 68. f. 3. 4.  
*Habitat in Hispania.* 24.  
*Nectarium cylindricum, ore obsolete repando, non denta-*  
*to aut crispo.*  
 6. *NARCISSUS* spatha multiflora nectario campanula- calachinus.  
 to subcrenato æquante petala, foliis planis.

Nar.

*Narcissus angustifolius flavus*, magno caule. *Bauh. pin.* 51. *Rudb. elys.* 2. p. 60. f. 5.

*Narcissus IX. angustifolius* 1. *Clus. hist.* 1. p. 158.

*Habitat in Europa australi, Oriente.* 24

*Simillimus N. Tazettæ, sed Petala paulo majora & acutiora. Nectarium longitudine petalorum.*

Tazetta.

7. **NARCISSUS** spatha multiflora, nectario campanulato truncato brevior petalis, foliis planis.

*Narcissus foliis ensiformibus, florum nectario campanulato erecto petalis longe brevior.* *Hort. cliff.* 134.

*Narcissus medio luteus, copioso flore: odore gravi.* *Bauh. pin.* 50. *Rudb. elys.* 2. p. 57. f. 11.

*Narcissus luteus polyanthos lusitanicus.* *Bauh. pin.* 50.

*Narcissus latifolius, flore prorsus albo* 1, 2. *Clus. hist.* 1. p. 155.

*Habitat in G. Narbonensis, Lusitanæ, Hispaniæ maritimis subhumidis.* 24

*Nectarium triplo brevius est petalis.*

odorus.

8. **NARCISSUS** spatha submultiflora, nectario campanulato sexfido lævi dimidio brevior petalis. *Auzn. acad.* 4. p. 311.

*Narcissus polyanthos, flore minore stellato toto luteo.*

*Rudb. elys.* 2. p. 60. f. 7.

*Habitat in Europa australi.* 24

*Flos luteus, triplo major flore N. Tazettæ, varians spatha uniflora & multiflora. Nectarium non fimbriatum, sed ore diviso lobis sex obtusis.*

triandrus.

9. **NARCISSUS** spatha subuniflora nectario campanulato crenulato, dimidio brevior petalis, staminibus ternis.

*Narcissus juncifolius, albo flore reflexo.* *Clus. app. alt.*

*Narcissus albus, oblongo calyce.* *Bauh. pin.* 53. *Rudb. elys.* 2. p. 72. f. 12.

*Habitat in Pyrenæis.* 24

*Magnitudo N. poetici, sed folia dimidio angustiora, canaliculata. Spatha (mihi) uniflora. Corolla tota nivea. Petala ovato-oblonga. Nectarium campanulatum, corolla dimidio brevius, margine recto inæqualiter crenulato. Stamina tria (mihi ut Clusio), raro sex, quæ tamen in quibusdam vidi individuis. Antheræ luteæ, nectario breviores.*

10. NAR-

10. NARCISSUS spatha submultiflora, nectario campanulato subtrifido integerrimo, dimidio brevior petalis.
- Narcissus angustifolius pallidus, calyce flavo. *Bauh. pin. 51. Rudb. elys. 2. p. 61. f. 3.*  
*Habitat in Europa australi. 2.*  
*Similis N. Jonquillæ, sed Nectarium cylindricum, longitudine plus, quam dimidiæ petalorum, non crispum, sed obsolete trilobum.*
11. NARCISSUS spatha multiflora, nectario hemisphærico crenato, brevior petalis, foliis semiteretibus.
- Narcissus foliis subulatis, florum nectario brevissimo. *Hort. cliff. 134. Roy. lugdb. 35. Sauv. monsp. 19.*  
 Narcissus juncifolius luteus minor. *Bauh. pin. 51.*  
 Narcissus juncifolius, oblongo calyce luteus major. *Bauh. pin. 51.*  
 Narcissus juncifolius minor & 2. *Clus. hist. 1. p. 159.*  
*Habitat inter Hispalim & Gades, inter Guadaloupam & Toletum, inque Oriente locis uliginosis. 2.*
12. NARCISSUS spatha uniflora, nectario brevissimo serotinus. sexpartito. *Læst. it. 19.*
- Narcissus albus autumnalis minimus. *Bauh. pin. 51. Rudb. elys. 2. p. 64. f. 8.*  
 Narcissus serotinus. *Clus. hist. 1. p. 162.*  
 Narcissus autumnalis minor. *Clus. hisp. 251. t. 252.*  
*Habitat in Hispania, Italia, Barbaria 2.*
13. NARCISSUS spatha uniflora, nectario turbinato petalis majore, genitalibus declinatis Bulbocodium.
- Narcissus foliis subulatis, nectario maximo patulo, genitalibus declinatis. *Hort. cliff. 134. Roy. lugdb. 35.*  
 Narcissus montanus alter, flore fimbriato. *Bauh. pin. 53.*  
 Narcissus montanus juncifolius, calyce aureo. *Bauh. pin. 53. Rudb. elys. 2. p. 75. f. 5. 7.*  
 Pseudo-Narcissus juncifolius 2, flavo flore. *Clus. hist. 1. p. 166.*  
*Habitat inter Ulyssiponem & Hispalim. 2.*

PANCRATIUM.

1. PANCRATIUM spatha uniflora, petalis reflexis, zeylanicum. *Fl. zeyl. 126.*  
*Linnaei Spec. plant. pars I. Dd Nar-*

- Narcissus zeylanicus*, flore albo hexagono odorato.  
*Herm. lugdb.* 691. t. 693. *Comm. hort.* 1. p. 75. t. 38. *Rudb. elys.* 2. p. 179. f. 7.  
*Lilium indicum.* *Rumph. amb.* 6. p. 161. t. 70. f. 2.  
*Catulli-pola.* *Kheed. mal.* 11. p. -t. 40.  
*Habitat in India.* 21
- mexicanum. 2. **PANCRATIUM** spatha biflora. *Hort. cliff.* 133.  
*Roy. lugdb.* 34.  
*Pancratium mexicanum*, flore gemello candido. *Dill. elth.* 299. t. 222. f. 289.  
*Habitat in Mexico.* 21
- caribæum. 3. **PANCRATIUM** spatha multiflora, foliis lanceolatis. *Hort. cliff.* 133. *Brown. jam.* 194. n. 1.  
*Narcissus americanus*, flore multiplici albo hexagono odorato. *Comm. hort.* 2. p. 173. t. 87.  
*Narcissus totus albus latifolius polyanthos major odoratus.* *Sloan. jam.* 115. *hist.* 1. p. 244. *Mart. cent.* 27. t. 27.  
*Habitat in Jamaica, Caribæis.* 21
- maritimum. 4. **PANCRATIUM** spatha multiflora, petalis planis, foliis lingulatis. *Mill. dict.* t. 197.  
*Narcissus maritimus.* *Bauh. pin.* 54.  
*Lilio-Narcissus albus maritimus minor.* *Morif. hist.* 2. p. 365. f. 4. t. 10. f. 28. \*  
*Hemerocallis valentina.* *Clus. hist.* 1. p. 167.  
*Habitat in Hispaniæ maritimis circa Valentiam & infra Monspelium.* 21
- carolinianum. 5. **PANCRATIUM** spatha multiflora, foliis linearibus staminibus nectarii longitudine.  
*Lilio-Narcissus polyanthos*, flore albo. *Catesb. car.* 3. p. 5. t. 5.  
*Habitat in Jamaica, Carolina.* 21
- illyricum. 6. **PANCRATIUM** spatha multiflora, foliis ensiformibus, staminibus nectario longioribus. *Roy. lugdb.* 34.  
*Pancratium* foliis ensiformibus, spatha multiflora, floribus magnis candidis fragrantibus. *Trew. chret.* t. 27.  
*Narcissus illyricus liliaceus.* *Bauh. pin.* 55. *Seb. thes.* 1. p. 17. t. 8. f. 1.

*Lilionarcissus hemerocallidis* facie. *Best. syst. vern.* 3.  
t. 16. f. 1.

*Habitat in Illyria?* 2.

7. **PANCRATIUM** spatha multiflora, foliis ovatis amboinense, nervosis petiolatis.

*Narcissus amboinensis*, folio latissimo subrotundo.

*Comm. hort.* 1. p. 77. t. 39. *Rudb. clyf.* 2. p. 238.  
f. 17.

- β. **Pancrati** foliis ovatis acuminatis petiolatis, spatha multiflora, floribus minoribus candidis fragrantibus.  
*Trev. chret.* t. 28.

*Cepa sylvestris.* *Rumph. amb.* 6. p. 160. t. 70. f. 1.

*Habitat in Amboina.*

## CRINUM.

1. **CRINUM** foliis ovato lanceolatis acuminatis sessilibus planis. latifolium.

*Sjovanna-pola-tali.* *Rheed. mal.* 11. p. 77. t. 39. *Rudb. clyf.* 2. p. 91. f. 12.

*Habitat in Asiæ arenosis.* 2.

Folia 3. f. 4. digito transversos lata, a basi sensim attenuata. Antheræ filiformes, lineares, incumbentes, longitudine dimidii filamenti f. corollæ.

2. **CRINUM** foliis carinatis. *Fl. zeyl.* 127. *Mill. dict.* asiaticum.  
t. 110. *Osborn. it.* 94.

*Lilium zeylanicum bulbiferum & umbelliferum.* *Herm. lugdb.* 682. t. 683. *Rudb. clyf.* 2. p. 182. f. 10.

*Radix toxicaria.* *Rumph. amb.* 11. p. 155. t. 69.

*Habitat in Malabaria, Zeylona, America.* 2.

*Bulbi intra capsulas generantur.*

3. **CRINUM** corollarum apicibus introrsum unguiculatis. americanum  
*Crinum.* *Hort. cliff.* 127. *Hort. upf.* 76. *Roy. lugdb.* 37.

*Lilio-Alphodelus americanus sempervirens maximus polyanthus albus.* *Comm. rar.* 14. t. 14. *Dill. elik.* 194. t. 160. f. 195.

- β. *Lilio-Alphodelus americanus sempervirens minor albus.* *Comm. rar.* 15. t. 15.

*Habitat in America.* 2.

4. **CRINUM** foliis sub lanceolatis planis, corollis obtusis. africanum.

*Polyanthes floribus umbellatis. Vir. cliff. 29. Hort. cliff. 126. Roy. lugdb. 27. Mill. dict. t. 210.*

*Hyacinthus africanus tuberosus, flore cæruleo umbellato. Breyn. prodr. 1. p. 39. icon. 23. t. 10. Comm. hort. 2. p. 133. t. 67. Seb. thes. 1. p. 29. t. 19. f. 4.*

*Hyacintho affinis, tuberosa radice, africana, umbella cærulea inodora. Pluk. alm. 187. t. 195. f. 1.*

*Habitat in Æthiopia. 2.*

## AMARYLLIS.

expensis. 1. **AMARYLLIS** spatha uniflora remotissima, corolla æquali, staminibus pistilloque rectis. *Amen. acad. 6. afr. 11. \**

*Sisyrinchium indicum. Cornut. canad. 168. Mori. hist. 2. p. 421. f. 4. t. 23. f. 9. Rudb. elys. 2. p. 236. f. 17.*

*Habitat ad Cap. b. Spei. 2.*

lutea. 2. **AMARYLLIS** spatha uniflora, corolla æquali, staminibus declinatis. *Hort. cliff. 135. Roy. lugdb. 36.*

*Narcissus autumnalis major. Clus. hist. 1. p. 164.*

*Colchicum luteum r. majus. Bauh. pin. 69.*

*Habitat in Hispania, Italia, Thracia. 2.*

Aramastro. 3. **AMARYLLIS** spatha uniflora, corolla æquali, pistillo declinato. *Hort. cliff. 135. Gron. virg. 36. Roy. lugdb. 36.*

*Lilio-Narcissus virginienis. Catesb. car. 3. p. 12. t. 12.*

*Lilio-Narcissus vernus angustifolius, flore purpurascente. Barr. ic. 994.*

*Lilio-Narcissus liliiflorus carolinianus, flore albo singulari cum rubedine diluto. Pluk. alm. 220. t. 43. f. 3.*

*Lilio-Narcissus indicus pumilus monanthos albus. Moris. hist. 2. p. 266. f. 4. t. 24. f. 4.*

*Habitat in Virginia. 2.*

formosissima. 4. **AMARYLLIS** spatha uniflora, corolla inæquali, petalis tribus genitalibusque declinatis. *Hort. cliff. 135.*

*Hort. upf. 75. Aët. stockh. 1742. p. 93. t. 6. Roy. lugdb. 36.*

*Lilio-Narcissus jacobæus, flore sanguineo nutante. Dill. elth. 195. t. 162. f. 196.*

*Narcissus jacobæus major. Rudb. elys. 2. p. 89. f. 10.*

*Habitat in America meridionali, innotuit europæis 1593. 2.*

*Squamæ totidem quot filamenta, e receptaculo ortæ, filamentorum basi æstæ adfident; an sic in reliquis?*

5. AMARYLLIS spatha multiflora, corollis campanu- reginz.  
 latis æqualibus undulatis, genitalibus declinatis.  
*Mill. dict. t. 24.*  
*Lilium americanum, puniceo flore, Belladonna di-*  
*ctum. Herm. par. 194. t. 194.*  
*Habitat in Caribæis.*  
*Differt ab A. Belladonna marginibus petalorum undulatis,*  
*nec ad unguis recurvatis.*
6. AMARYLLIS spatha multiflora, corollis campanu- Belladonna  
 latis æqualibus ungue reflexis, genitalibus declina-  
 tis. *Hort. diff. 135. Roy. lugdb. 36. Mill. dict. t. 23.*  
*Lilio-Narcissus polyanthos, flore incarnato: fundo*  
*ex luteo albescente. Sloan. jam. 115. hist. 1. p. 244.*  
*Seb. thef. 1. p. 25. t. 17. f. 1.*  
*Lilium rubrum. Merian. furin. 22. t. 22.*  
*Habitat in Caribæis, Barbados, Surinama. 24.*
7. AMARYLLIS spatha multiflora, corollis revolutis, farniensis.  
 genitalibus erectis. *Hort. upf. 75.*  
*Amaryllis spatha multiflora, corollis æqualibus pa-*  
*tentissimis revolutis, genitalibus longissimis. Hort.*  
*cliff. 131. Roy. lugdb. 36.*  
*Narcissus japonicus, rutilo flore. Corn. canad. 157. t.*  
*158. Rudb. clyf. 2. p. 23. f. 14. Ehret. select. t. 9. f. 3.*  
*Kämpf. amæn. 872. Seb. mus. 1. t. 17. f. 3.*  
*Lilium farniense. Dugl. monogr. t. 1. 2.*  
*Habitat in Japonia, nunc in Sarnia insula Angliæ. 24.*
8. AMARYLLIS spatha multiflora, corollis campanu- zeylanica.  
 latis æqualibus, genitalibus declinatis, scapo tere-  
 ti ancipiti. *Roy. lugdb. 36.*  
*Lilio-Narcissus zeylanicus latifolius, flore niveo ex-*  
*terne linea purpurea striato. Comm. hort. 1. p. 73. t.*  
*37. Rudb. clyf. 2. p. 181. f. 9.*  
*Tulipa javana. Rumph. amb. 5. p. 30. t. 105.*
- β. Lilio-Narcissus africanus, scillæ foliis, flore niveo  
 linea purpurea striato. *Ehret. pict. 5. f. 2. Trew.*  
*ehret. t. 13.*  
*Habitat in Zeylona. 24.*
9. AMARYLLIS spatha multiflora, corollis campanu- longifolia.  
 latis æqualibus, genitalibus declinatis, scapo com-  
 presso longitudine umbellæ. *Roy. lugdb. 36. Ehret.*  
*pict. t. 13.*



*Lilium africanum* humile longissimis foliis, polyanthos saturato colore purpurascens. *Herm. parad.* 195. t. 195.

*Habitat in Aethiopia.* 24.

orientalis. 10. AMARYLLIS *spatha multiflora*, corollis inaequalibus, foliis lingviformibus. *Buttn. canon.* 215.

*Amaryllis spatha multiflora*, foliis ovato-oblongis obtusis. *Roy. lugdb.* 37.

*Lilio Narcissus indicus maximus sphaericus*, floribus plurimis rubris liliaceis. *Morif. hist.* 2. p. 568. f. 4. t. 10. f. 35.

*Narcissus indicus orientalis.* *Swert. flor.* 31. f. 1.

*Brunsvigia Heisl.*

*Habitat in India.* 24.

ciliaris. 11. AMARYLLIS *spatha multiflora*, foliis ciliatis.

*Habitat in Aethiopia.*

*Folia plana, linearia, pollicis latitudine, bipalmaria longitudine, adeoque figura Amaryllidis, non vero Hemanthi, sed margine undique ciliata, ciliis albis nec ferrugineis; adeoque longe recedit ab Hemantho ciliari; neutra tamen nobis floruit.*

## BULBOCODIUM.

vernum. 1. BULBOCODIUM foliis lanceolatis. *Roy. lugdb.* 41.

*Bulbocodium.* *Hort. cliff.* 133.

*Colchicum vernum hispanicum.* *Bauh. pin.* 69. *Rudb. elyf.* 2 p. 128. f. 2.

*Colchicum vernum.* *Cus. hist.* 2. app. 203.

*Habitat in Hispania.* 24.

## IAPHYLLANTHES.

monspeliensis. 1. APHYLLANTHES. *Hort. cliff.* 493. *Sauv. monsp.* 8.

*Aphyllanthes monspeliensium.* *Lob. adv.* 190. *Bauh. hist.* 3. p. 336.

*Caryophyllus caeruleus monspeliensium.* *Bauh. pin.* 209. *Morif. hist.* 2. p. 562. f. 5. t. 25. f. 12.

*Habitat Monspeli prope Castelneuf locis montosis saxosis sterilibus.*

*Radix repens. Culmi nudi, simplices, cincti basi vaginis uti Juncus. Gluma bivalvis, biflora; Gluma propria etiam bivalvis. Juncus esset si Corolla careret,*

ALLIUM.

\* *Foliis caulinis planis.*

1. **ALLIUM** caule planifolio umbellifero, umbella globosa, staminibus tricuspидatis, petalis carina scabris. *Ampeloprasum.*  
*Allium* staminibus alterne trifidis, foliis gramineis, floribus sphaerice congestis, radice laterali solida.  
*Hall. al. 5. opusc. 344.*  
*Porrum* radice laterali solida, foliis linearibus, caule erecto, umbella subglobosa. *Roy. lugdb. 40.*  
*Allium* sphaerico capite, folio latiore f. *Scorodoprasum* alterum. *Bauh. pin. 74. Rudb. clyf. 2. p. 151.*  
*Scorodoprasum* latifolium spontaneum italicum, floribus dilute purpureis odoratis. *Mich. gen. 25. t. 24. f. 5.*  
*Habitat in Oriente, inque insula Holms Angliae. 24.*  
*Stamina corolla paulo longiora.*
2. **ALLIUM** caule planifolio umbellifero, staminibus tricuspидatis, radice tunicata. *Porrum.*  
*Hort. upf. 77. Mart. med. 165.*  
*Allium* staminibus alterne trifidis, foliis gramineis, floribus sphaerice congestis, radice tunicata cauli circumnata. *Hall. all. 7.*  
*Porrum* radice ambiente tunicata oblonga solitaria.  
*Hort. cliff. 136. Roy. lugdb. 40.*  
*Porrum* commune capitatum. *Bauh. pin. 72.*  
*Porrum* sativum latifolium. *Bauh. pin. 72.*  
*Habitat - - -*  
*Affinis nimium praecedenti ♂ forte sola varietas. Petala carina scabra.*
3. **ALLIUM** caule planifolio umbellifero, umbella globosa, staminibus tricuspидatis corolla duplo longioribus. *lineare.*  
*Allium* staminibus alterne trifidis, radice tunicis reticulatis obducta. *Hall. all. 9.*  
*Porrum* caule tereti folioso, foliis linearibus planis, umbella globosa, staminibus corolla longioribus.  
*Gmel. sibir. 1. p. 56. t. 13. ♂ 14. f. 1.*  
*Habitat in Siberia.*
4. **ALLIUM** caule planifolio umbellifero staminibus rotundum tricuspидatis, umbella subglobosa, floribus lateralibus nutantibus.

*Allium* staminibus alterne trifidis, spica sphaerica, foliis gramineis, radice multiplici. *Hall. opusc.* 350. n. 8.

*Allium* montanum, capite rotundo. *Bauh. pin.* 75.

*Scorodoprasum* latifolium spontaneum italicum, floribus dilute purpureis odoratis. *Mich. gen.* 25. t. 24. f. 1.

*Habitat in Europa australiori* 24.

*Viæerialis.* 5. *ALLIUM* caule planifolio umbellifero, umbella rotundata, staminibus lanceolatis corolla longioribus, foliis ellipticis. *Mar. med.* 163.

*Allium* foliis caulinis lanceolatis, floribus umbellatis. *Roy. lugdb.* 39.

*Allium* radice oblonga reticulo obducta. *Hall. all.* 20.

*Allium* montanum latifolium maculatum. *Bauh. pin.* 74.

*Allium* anguinum. *Cam. epit.* 329.

*Habitat in Alpibus Helvetiæ, Italiæ.*

*Huic Folia in toto genere latissima colchici.*

*subhirsutum* 6. *ALLIUM* caule planifolio umbellifero, foliis inferioribus hirsutis, staminibus subulatis. *Osbeck. it.* 49.

*Allium* foliis radicalibus subhirsutis, caulinis glabris, floribus umbellatis. *Hall. all.* 18.

*Allium* foliis lanceolato-linearibus, florum umbella fastigiata. *Roy. lugdb.* 39.

*Moly* angustifolium umbellatum. *Bauh. pin.* 75.

*Moly* dioscoridis. *Clus. hist.* 1. p. 192.

*Habitat in Africa, Italia, Hispania, Oriente.*

*magicum.* 7. *ALLIUM* caule planifolio umbellifero, ramulo bulbifero staminibus simplicibus.

*Allium* caule tereti, propagine ex ala. *Hort. cliff.* 137.

*Roy. lugdb.* 39. *Sauv. monsp.* 18.

*Allium* foliis caulinis lanceolatis, floribus umbellatis, ex ala bulbiferum. *Hall. all.* 19.

*Moly* latifolium triflorum. *Bauh. pin.* 75.

*Moly* indicum, flore purpureo. *Svert. Fl.* 61.

8. *Caucasum.* *Lob. ic.* 162.

*Habitat* - -

*Caulis teres, nudus, pedalis, infirmus. Folia radicalia canaliculata, ampla, obtusa. Capitulum bulbiferum. Sauv.*

*obliquum.* 9. *ALLIUM* caule planifolio umbellifero, staminibus filiformibus flore triplo longioribus, foliis obliquis.

*Allium radice tunicata, foliis planis linearibus caulinis, capitulo umbellato. Gmel. Sibir. 1. p. 49. t. 9.*

*Habitat in Sibiria.*

*Caulis bipedalis, teres. Folia alterna, plano-canaliculata, obliqua: Vaginis teretibus. Umbella globosa; Spatha globosa, purpurea, acuminata. Petala ovata, erecta, concava, flava carina viridi. Filamenta alba, corolla duplo longiora. Antheræ luteæ. Germen subglobosum.*

9. **ALLIUM** caule subplanifolio umbellifero, staminibus subulatis longioribus, umbella globosa, foliis linearibus subconvexis. *ramosum.*

*Allium scapo nudo tereti farcto, foliis semicylindricis, staminibus corolla longioribus. Gmel. Sib. 1. p. 52. t. 9. f. 1.*

*Habitat in Sibiria. G. Demidoff.*

*Caulis ferme bipedalis, teres. Folia alterna, planiuscula, canaliculata, subtus convexa, margine obtusa. Umbella globosa. Petala ovata, purpurascens carina saturatiore. Filamenta purpurea, corolla parum longiora. Antheræ violaceæ. Germen subrotundum: maculis 3, viridibus. E medio caule ex ala folio opposita. Ramus nudus, longitudine caulis.*

10. **ALLIUM** caule planifolio bulbifero, bulbo composito, staminibus tricuspidatis. *fativum.* *Hort. ups. 76. Mat. med. 164.*

*Allium radice bulbo multipartito, capitulo bulbifero, foliis linearibus. Hort. cliff. 137. Roy. lugdb. 39.*

*Allium staminibus alterne trifidis, foliis gramineis, capite bulbifero, radicibus in unum bulbum congruentibus. Hall. all. 1.*

*Allium sativum. Bauh. pin. 73.*

*Allium. Cam. epit. 328.*

*Habitat in Sicilia.*

11. **ALLIUM** caule planifolio bulbifero, foliis crenulatis: vaginis ancipitibus, staminibus tricuspidatis. *Scoroditis, prasum.* *Hort. ups. 77.*

*Allium capitulo bulboso erecto, foliis planis subcrenatis: vaginis ancipitibus. Fl. suec. 266, 278. \* It. el. 60.*

*Allium montanum bicornae latifolium, flore dilute purpurascens. Bauh. pin. 74. Rudb. cl. 2. p. 155. \**

β. *Allium* staminibus alterne trifidis, capite bulbifero, scapo ante maturitatem contorto. *Hall. all.* 2.

*Allium* sativum alterum f. *Allioprasum* caulibus summo circumvoluto. *Bauh. pin.* 73.

*Allii* genus *Ophioscorodon* dictum quibusdam. *Bauh. hist.* 2. p. 559.

*Porrum* radice laterali cordata solida, pedunculo revoluta, capite bulbifero. *Hort. elys.* 136.

*Habitat in Oekandia, Dania, Pannonia.*

areolarium. 12. *ALLIUM* caule planifolio bulbifero, vaginis teretibus, spatha mutica, staminibus tricuspidatis. *It. scan.* 227. *Fl. fvec.* 2. n. 279. \*

*Allium* staminibus alterne trifidis, capite bulbifero, foliis gramineis leviter ferratis. *Hall. all.* 3.

*Allium* montanum bicornae latifolium, flore dilute purpurascens. *Bauh. pin.* 74.

*Porrum* planifolium, staminibus alterne trifidis, umbella bulbifera, *Rupp. hall.* 154. t. 2. f. 2.

*Habitat in Thuringia; in Falstria Scaniae.*

*Caulis ante anthesin cernuus.*

carinatum. 13. *ALLIUM* caule planifolio bulbifero, staminibus subulatis. *Fl. fvec.* 2. n. 281.

*Allium* umbella bulbifera, vagina bicornae, foliis carinatis. *Hall. all.* 27. t. 1. f. 2.

*Ampeloprasum* proliferum. *Lob. ic.* 156. \*

β. *Allium* montanum bicornae angustifolium, flore dilute purpurascens. *Bauh. pin.* 74.

*Allium* f. *Moly* montanum 2. *Clus. hist.* 1. p. 193.

*Habitat in Germania.*

\* *Foliis caulinis teretibus.*

sphaerocephalon. 14. *ALLIUM* caule teretifolio umbellifero, foliis semiteretibus, staminibus tricuspidatis corolla longioribus.

*Allium* staminibus alterne trifidis, foliis fistulosis, spica sphaerica non bulbifera suaverubens. *Hall. opusc.* 353. n. 10.

*Allium* montanum, capite rotundo. *Bauh. pin.* 75.

*Rudb. elys.* 1. p. 157. f. 7.

*Allium* f. *Moly* montanum, purpureo flore. *Clus. hist.* 1. p. 195.

*Allii* genus forte *Scorodoprasum* alterum. *Bauh. hist.* 2. p. 564.

Scorodoprasum montanum juncifolium, capite rotundo dilute janthino, floribus paucis. *Mich. gen.* 25. 1. 24. f. 2.

*Habitat in Italia, Sibiria, Helvetia.*

15. **ALLIUM** caule subteretifolio umbellifero, umbella globosa, staminibus simplicibus corolla longioribus, spatha subulata. *purviflorum.*

*Habitat in Europa australi.* 21.

*Facies A. sphaerocephali.* Bulbus oblongus, tunicatus. Caulis teres, striatus, foliosus. Folia longitudine fere caulis, subulata. Umbella globosa: Floribus minimis. Petalis exterioribus ovatis, tristioribus; interioribus duplo longioribus, lanceolatis, purpureiscentibus. Stamina simplicia, corolla longiora, uti etiam Stylus. Spatha subulata, corolla longior, diphylla. Adeoque simillimum *A. sphaerocephalo*, sed stamina simplicia.

16. **ALLIUM** caule subteretifolio umbellifero, pedunculis exterioribus brevioribus, staminibus tricuspidatis. *Hort. ups.* 78. *descendens.*

Allium staminibus alterne trifidis, foliis fistulosis, capite suave rubente sphaerico non bulbifero, radice laterali. *Hall. all.* 8.

Allium staminibus alterne trifidis, foliis fistulosis, capite sphaerico non bulbifero atropurpureo. *Hall. all.* p. 11. t. 2. f. 1.

Allium f. *Moly latifolium*, capite sphaerico, flore purpureo. *Rudb. elys.* 2. p. 160. f. 20.

*Moly atropurpureum.* *Rudb. elys.* 2. p. 165. f. 13.

*Habitat in Helvetia.*

17. **ALLIUM** caule teretifolio umbellifero, umbella fastigiata subsexflora, petalis acutis, staminibus simplicibus, foliis setaceis. *moschatum.*

Allium foliis teretibus caulinis, floribus umbellatis. *Hall. all.* 17.

*Moly moschatum*, capillaceo folio. *Bauh. pin.* 76. *prodr.* 28. t. 28. *Rudb. elys.* 2. p. 166. f. 14. \* *Sauv. monsp.* 10.

*Habitat in Galloprovinciæ, Narbonæ, Hispaniæ aprici elevatis.*

18. **ALLIUM** caule subteretifolio umbellifero, floribus pallens. *pen-*

pendulis truncatis, staminibus simplicibus corollam æquantibus.

Allium umbella non bulbifera lutea, vagina bicorni, foliis teretibus. *Hall. all.* 21.

Allium montanum bicorni, flore pallido odore. *Bauh. pin.* 75.

Gethioides sylvestre. *Colum. ecphr.* 2. p. 6. t. 7. f. 2.

Habitat in Italia, Hispania, Montpelii, Pannonia. 24.

Caulis bipedalis, teres, lavis. Folia alterna, semiteretia, fistulosa, heptagona, erecta. Spatha subulata, nervosa. Corolla campanulata, truncata, alba. Petalis obovatis, obtusissimis, erectis, concavis, carina viridi. Stamina simplicia, longitudine petalorum. Flores mox a florescentia pendunt. Stylus vix ullus, post anthesin excresecens, brevissimus.

Flavum.

19. ALLIUM caule teretifolio umbellifero, floribus pendulis, petalis ovatis, staminibus corolla longioribus.

Allium foliis teretibus, vagina bicorni, umbella lutea pendula. *Hall. opusc.* 24.

Allium juncifolium bicorni luteum. *Rudb. elyf.* 2. p. 157. f. 6.

Habitat Montpelii. 24.

Simillimum *A. pallescens*; Sed differt Petalis ovatis flavissimis, nec obovatis albidis. Staminibus 3 pistillo corolla duplo longioribus, nec brevioribus.

paniculatum.

20. ALLIUM caule subteretifolio umbellifero, pedunculis capillaribus effusis, staminibus simplicibus, spatha longissima.

Allium foliis teretibus, vagina bicorni, umbella pendula suave purpurea. *Hall. opusc.* 386. n. 25.

Allium montanum bicorni, floribus triquetris purpureis pedunculis longis & reflexis. *Segu. ver.* 2. p. 70.

Habitat in Sibiria, Austria, Italia, Oriente. 24.

Folia subulata, canaliculata. Flores pedunculis capillaribus, longissimis, laxis. Petala purpurea, obovata, longitudine staminum. Spatha apice subulata, longissimo.

vinale.

21. ALLIUM caule teretifolio bulbifero, staminibus tricuspideatis.

Allium staminibus alterne trifidis, foliis fistulosis, capite bulbifero, radice multiplici. *Hall. all.* 4.

Allium sylvestre campestre purpurascens. *Bauh. pin.* 74.

Al-

*Allium campestre juncifolium capitatum purpurascens majus.* Bauh. pin. 74.

*Porrum sylvestre vinearum.* Bauh. pin. 72.

*Porrum sylvestre, gemino capite.* Bauh. pin. 72.

*Habitat in Germania.*

*Capitulum arctum bulbis acuminatis recurvis, sæpe in longiora acumina crinitis. Corollæ parvæ, violacæ. Stamina setis duabus, longis, extra florem eminentibus.*

22. **ALLIUM** caule teretifolio bulbifero, foliis scabris semiteretibus subtus sulcatis, staminibus simplicibus. oleraceum.

*Allium umbella bulbifera, vagina bicorni, foliis teretibus.* Hall. all. 23. f. 4.

*Cepa capitulo bulboso prolifero, foliis semiteretibus novemangularibus scabris fistulosis.* Fl. suec. 265, 280. \*

*Porrum radice solida, caule erecto, capitulo bulbifero.* Roy. lugdb. 40. Gort. geldr. 69.

*Allium montanum bicorni, flore exalbido.* Bauh. pin. 75.

*Habitat in Suecia, Germania.*

\* \* *Foliis radicalibus, scapo nudo.*

23. **ALLIUM** scapo nudo ancipiti, foliis linearibus planis, staminibus tricuspidatis. antenn. Hort. upf. 80.

*Allium staminibus alterne trifidis, foliis latissimis, capite sphaerico, radice laterali solida.* Hall. all. 6.

*Porrum scapo nudo ancipiti, antequam floruerit nutante, foliis ensiformibus: hinc paullo convexioribus.* Gmel. fibir. 1. p. 55. t. 18.

*Habitat in Sibiria.*

*Variat ætate & solo præ reliquis, ut vix dignoscatur.*

24. **ALLIUM** scapo nudo tereti, foliis subulatis, umbella globosa, staminibus tricuspidatis. Afcalonium,

*Cepa foliis subulatis, radicibus oblongis conglobatis.*

*Hort. cliff. 36. Hort. upf. 79.*

*Cepa afcalonica.* Moris. hist. 2. p. 382. f. 4. t. 14. f. 3.

*Cepa sterilis.* Bauh. pin. 72.

*Habitat in Palæstina.* Haeselquist. 24.

*Statura fere A. Schænoprasi. Scapus subnudus, teres, spithameus & ultra. Spatha membranacea. Umbella globosa, floribus A. Schænoprasi minoribus. Petala erecta, ovato-lanceolata, cærulea, carina atro-*



*trocerulea.* Stamina longitudine petalorum: alternis latissimis apice trifidis. Antheræ flavæ. Stylus stigmate obtuso.

senescens. 25. ALLIUM scapo nudo ancipiti, foliis linearibus subtus convexis lævibus, umbella subrotunda, staminibus subulatis. *Hort. upf.* 79.

Allium caule ancipiti teretiusculo, foliis ensiformibus: hinc paulo convexioribus. *Gmel. sibir.* 1. p. 53. t. 11. f. 2.

Allium foliis linearibus, florum umbella subglobosa. *Roy. lugdb.* 39.

Allium radice bulbosa, foliis gramineis, umbella erecta non bulbifera, vaginis brevissimis. *Hall. helv.* 297. all. 16. n. 2.

Allium montanum, foliis narcissi, majus. *Bauh. pin.* 75.

Habitat in Sibiria, Sicilia. 24.

Petala erecta, a se invicem dehiscencia s. distantia, canaliculata, inflexa. Folia obliqua & Stamina corolla paullo longiora huic, non vero sequenti.

nigrum. 26. ALLIUM scapo nudo tereti, foliis linearibus, umbella erecta, petalis erectis, spatha mucronata bifida.

Allium montanum bicorni latifolium, flore magno dilute purpurascente. *Rudb. elyf.* 2. p. 160. f. 21. *Burf.*

Habitat Algeriæ inque Galloprovincia.

Simile *A. senescenti*, sed duplo majus. Folia *Narcissina*. Flores quadruplo majores. Petala ovato-oblonga. Stamina lanceolata, corolla breviora. Spatha ovata, acuminata, vix ad basin in duas partes æquales divisa.

angulosum. 27. ALLIUM scapo nudo ancipiti, foliis linearibus canaliculatis subtus subangulatis, umbella fastigiata. *Hort. upf.* 79. \*

Cepa scapo nudo subangulato fæcto, foliis linearibus subtus angulosis, staminibus corolla brevioribus. *Gmel. sibir.* 1. p. 58. t. 14. f. 2.

Allium radice senescente lignosa transversa, umbella erecta. *Hall. all.* 16.

Allium petræum umbelliferum. *Bauh. hist.* 2. p. 564. t. 564.

*Allium montanum*, foliis narcissi, minus. *Bauh. pin.* 75.

*Habitat in* Sibiria, Germaniæ humidiusculis. 2.

Folia interdum subtus vix angulata, at Scapus superne parum angulato-striatus; Petala interiora apice patentia, omnia lateribus conniventia.

28. *ALLIUM* scapo nudo tereti, foliis linearibus, ca-  
pitulo bulbifero. *Kalm. it.* 3. p. 79.

*Habitat in* Canada. 2.

*Radix perennis*. Folia linearia, plana, lævia, recta, spithamea. Scapus subnudus, teres, lævis, vix foliis longior. Spatha ovata, mucronata, acutiusculi. Capitulum ex bulbis. Flosculi pauci, pedicellati, albi: Petalis ovalibus. Stamina simplicia, longitudine corollæ. Antheræ rufescentes.

29. *ALLIUM* scapo nudo semicylindrico, foliis lanceolatis petiolatis, umbella fastigiata.

*Allium* foliis lanceolatis, scapo nudo semicylindraceo; bulbo setis obvallato. *Fl. suec.* 163, 277. *L. gotl.* 169. \*

*Allium* foliis radicalibus petiolatis, floribus umbellatis.

*Roy. lugdb.* 39. *Hall. all.* 21. *Gmel. sib.* 1. p. 49.

*Allium* sylvestre latifolium. *Bauh. pin.* 74.

*Allium* ursinum. *Fuchs. hist.* 739. *Cam. epit.* 330.

*Habitat in* Europæ septentrionalioris nemorosis, roridis. 2.

30. *ALLIUM* scapo nudo, foliis triquetris, staminibus simplicibus.

*Allium* caule triangulo. *Tournef. inst.* 385. *Roy. lugdb.*

39.

*Allium* pratense, folio gramineo, flore prorsus albo, radice oblonga. *Rudb. elys.* 2. p. 159. f. 16.

*Moly* parvum, caule triangulo. *Bauh. pin.* 75.

*Moly* caule & foliis triangularibus. *Park. par.* 142. t.

143. f. 4

*Habitat in* Hispania. 2.

Folia triquetra latere interiore canaliculato.

31. *ALLIUM* scapo nudo inferne ventricoso longiore  
foliis teretibus. *Hort. upf.* 77. *Mat. med.* 166.

*Allium* staminibus alterne trifidis, caule ad terram ventricoso. *Hall. all.* 10.

*Cepa scapo nudo ventricoso foliis longiore, radice depressa. Hort. cliff. 137. Roy. lugdb. 40.*

*Cepa vulgaris. Bauh. pin. 71.*

*Habitat - - -*

Moly.

32. **ALLIUM** scapo nudo subcylindrico, foliis lanceolatis sessilibus, umbella fastigiata. *Hort. upf. 76.*

*Allium foliis radicalibus sessilibus lanceolatis, floribus umbellatis. Roy. lugdb. 38. Hall. all. 29.*

*Allium foliis lanceolatis, umbella laxa. Hort. cliff. 27.*

*Moly latifolium luteum, odore allii. Bauh. pin. 75.*

*Moly flavo flore. Swert. flor. 1. t. 60. f. 2.*

*Moly montanum latifolium, flavo flore. Clus. app. alt.*

*Habitat in Hungaria, Baldo, Monspelii, Pyrenæis. 24.*

*Petala patentia, tria exteriora magis patentia, interiora erecta.*

roseum.

33. **ALLIUM** scapo nudo umbellifero, pedicellis brevibus, petalis ovalibus, staminibus brevissimis, foliis linearibus.

*Allium roseum. Syst. nat. 296. n. 9.*

*Allium foliis ensiformibus, radice bulbillis stipata, floribus amplis umbellatis. Sauv. monsp. 42.*

*Allium sylvestre s. Moly minus, roseo amplo flore.*

*Magn. monsp. 11. t. 10. Rudb. elys. 2. p. 166. f. 17.*

*Habitat Monspelii, Pedemonti in vineis. 24.*

*Folia linearia, plana, longitudine vix Scapi. Scapus teres, nudus. Spatha obtusa. Umbellæ radii pedunculares floribus breviores. Petala obovato-oblonga, obtusa, atropurpurea, magna. Stamina simplicia, alba, brevissima. Bulbi ad fructum nulli. Flores majores, quam in reliquis.*

fistulosum.

34. **ALLIUM** scapo nudo folia adæquante, foliis teretibus ventricosis. *Hort. upf. 78.*

*Allium staminibus simplicibus, caule medio ventricoso. Hall. all. 13.*

*Cepa scapo longitudine foliorum, foliis ventricosis, radice oblonga. Hort. cliff. 136. Roy. lugdb. 40.*

*Cepa oblonga. Bauh. pin. 71. Dod. pempt. 687.*

*Habitat in - - -*

Schœnopræsum.

35. **ALLIUM** scapo nudo folia adæquante, foliis teretibus subulato-filiformibus. *Hort. upf. 78.*

*Allium staminibus simplicibus, foliis radicalibus teretibus scapi longitudine. Hall. all. 14.*

- Cepa scapis foliisque subulatis teretibus fistulosis æqualibus, spathis globosis. *Fl. succ.* 264, 282. *It. xl.* 54, 64.
- Cepa scapo longitudine figuraque foliorum, foliis subulatis filiformibus, spathis globosis. *Hort. cliff.* 136.
- Porrum sectivum juncifolium. *Bauh. pin.* 72.
- Schœnoprasum. *Dod. pempt.* 689.
- β. Cepa scapis foliisque teretibus, capitulis pyramidalis. *Gmel. sibir.* 1. p. 59. t. 15. f. 1.
- Cepa palustris altissima. *Buxb. cent.* 4. p. 27. t. 45.
- Habitat in alpibus Sibiria, Oelandia, locis rupestribus.*
36. ALLIUM scapo nudo tereti inani, foliis subulatis tenuissimum filiformibus, capitulis laxis paucifloris. *Gmel. sibir.* 1. p. 61. t. 15. f. 2, 3.
- Habitat in Sibiria.*
37. ALLIUM scapo nudo subnullo, capsulis cernuis, foliis planis ciliatis. *Chamæ Moly.*
- Moly humile, folio gramineo. *Bauh. pin.* 75. *Rudb. chys.* 2. p. 166. f. 15.
- Chamæ Moly, an Moly dioscoridis. *Column. ecphr.* 325. t. 326.
- Habitat in Italia.*

## LILIUM.

1. LILIUM foliis sparsis, corollis campanulatis: intus candidum. glabris. *Hort. cliff.* 120. *Hort. ups.* 80. *Mat. med.* 173. *Roy. lugdb.* 30. *Gron. orient.* 105.
- Lilium album, flore erecto, vulgare. *Bauh. pin.* 76.
- Lilium candidum. *Dod. pempt.* 197.
- β. Lilium album, floribus dependentibus, s. peregrinum. *Bauh. pin.* 76.
- γ. Lilium caule plano compresso. *Gron. orient.* 106.
- Lilii albi pulchri & ignoti species. *Bauh. pin.* 76.
- Habitat in Palestina, Syria, Gades.* 24
2. LILIUM foliis sparsis, corollis campanulatis erectis: intus scabris. *Hort. cliff.* 120. *Hort. ups.* 80. *Roy. lugdb.* 31. *Gmel. sibir.* 1 p. 41. *Gron. orient.* 104.
- Lilium purpureo-croceum majus. *Bauh. pin.* 76.
- β. Lilium purpureo-croceum, flore pleno.
- γ. Lilium purpureo-croceum minus. *Bauh. pin.* 77.
- δ. Lilium phœniceum. *Bauh. pin.* 77.
- Linnaei Spec. plant. pars I.* E e

ε. *Lilium bulbiferum latifolium majus. Bauh. pin. 77.*

ζ. *Lilium bulbiferum angustifolium. Bauh. pin. 77.*

η. *Lilium bulbiferum minus. Bauh. pin. 77.*

θ. *Lilium bulbiferum incanum. Bauh. pin. 77.*

*Habitat in Italia, Austria, Sibiria. 24*

pomponum 3. **LILIUM** foliis sparsis subulatis, floribus reflexis, corollis revolutis. *Hort. cliff. 120. Hort. upf. 81. Roy. lugdb. 31. Mill. dict. 1. 165. f. 2.*

*Lilium radice tunicata, foliis sparsis, floribus reflexis, corollis revolutis. Gmel. fibir. 1. p. 42.*

*Lilium rubrum angustifolium. Bauh. pin. 78.*

*Lilium rubrum præcox. Clus. hist. 1. p. 133.*

β. *Lilium miniatum odoratum angustifolium. Bauh. pin. 79. Rudb. elyf. 2. p. 176. f. 3.*

γ. *Lilium brevi & gramineo folio. Bauh. pin. 79.*

*Habitat in Pyrenæis, Sibiria. 24*

*Folia linearia, acuta, triquetra, canaliculata, sparsa. Nectarii rima valde dentata.*

chalconi- 4. **LILIUM** foliis sparsis lanceolatis, floribus reflexis, corollis revolutis. *Hort. cliff. 120. Hort. upf. 81. Roy. lugdb. 31.*

cam,

*Lilium byzantinum miniatum. Bauh. pin. 78.*

*Lilium rubrum f. miniatum byzantinum. Clus. hist. 1. p. 131.*

β. *Lilium byzantinum miniatum polyanthos. Bauh. pin. 78.*

γ. *Lilium purpureo-sanguineum, flore reflexo. Bauh. pin. 78.*

*Hemerocallis chalcedonica. Lob. ic. 169.*

*Habitat in Persia, Platina Carniola. 24*

*Racemus ante florentiam vix incurvatus ut in 3. 5.*

♂ *Caulis usque ad apicem foliis vestitus confertus.*

superbum. 5. **LILIUM** foliis sparsis lanceolatis, floribus ramoso-pyramidalis reflexis, corollis revolutis. *Mill. dict. n. 8.*

*Lilium, foliis sparsis, multiflorum, floribus reflexis fundo aureo limbo aurantio punctis nigricantibus, pedunculis singulis uno foliolo. Trew. chret. 2. t. 11.*

*Lilium f. Martagon canadense, flore luteo punctato.*

*Catesb. car. 2. p. 56. t. 56.*

*Habitat in America septentrionali. 24*

6. **LILIUM** foliis verticillatis, floribus reflexis, corollis revolutis. *Hort. cliff.* 120. *Hort. upf.* 81. *Gmel. fibr.* 1. p. 44. *Roy. lugdb.* 31.  
*Lilium*, floribus reflexis, montanum. *Bauh. pin.* 77.  
*Lilium* sylvestre. *Dod. pempt.* 201.
- β. *Lilium*, floribus reflexis, alterum hirsutum. *Bauh. pin.* 78.  
*Habitat in Hungaria, Helvetia, Sibiria, Lipsiæ.* 24
7. **LILIUM** foliis verticillatis, floribus reflexis, corollis campanulatis. *canadense.*  
*Lilium* angustifolium, flore flavo maculis nigris distincto. *Tournef. inst.* 371. *Barr. rar.* 778. t. 125.  
*Lilium* Martagon canadense maculatum. *Morif. hist.* 2. p. 408. f. 4. t. 20. f. 9. *Rudb. thyf.* 2. p. 177. f. 5.  
*Lilium* f. Martagon canadense, floribus magnis flavis non reflexis. *Catesb. car.* 3. p. 11. t. 11.  
*Habitat in Canada.* 24
8. **LILIUM** foliis verticillatis, floribus erectis, corolla philadelphica campanulata, petalis unguiculatis. *cum.*
- β. *Lilium* foliis verticillatis brevibus, corollis campanulatis, unguibus petalorum angustioribus, floribus erectis. *Mill. dict.* t. 165. f. 1.  
*Habitat in Canada.* 24  
*Simillimum est L. camtschatcensi, sed ejus petala ovata sessilia; hujus unguibus oblongis distinctis; separandum itaque secundum vivi autoptam Millerum.*
9. **LILIUM** foliis verticillatis, flore erecto, corolla camtschatcensis campanulata, petalis sessilibus. *Amzn. acad.* 2. p. 348. *cense.*  
*Habitat in Canada, Camtschatca.*

## FRITILLARIA.

1. **FRITILLARIA** racemo comoso inferne nudo, foliis integerrimis. *fo. imperialis.*  
*Hort. upf.* 82.  
*Petilium* foliis caulinis. *Hort. cliff.* 119. *Roy. lugdb.* 30.  
*Lilium* f. Corona imperialis: genus. *Bauh. pin.* 79.  
*Tufai. Clus. hist.* 1. p. 127, 128.  
*Habitat in Persia? Constantinopoli venit in Europam circa 1570.* 24
2. **FRITILLARIA** racemo comoso inferne nudo, foliis crenatis. *fo. regia.*

- Corona regalis, lilli folio crenato. *Dill. clth.* 110. t. 93.  
f. 109.  
*Habitat ad Cap. b. Spei.*
- perfica, 3. FRITILLARIA racemo nudiusculo, foliis obliquis.  
*Hort. upf.* 82.  
Fritillaria racemo nudo terminali. *Hort. cliff.* 119.  
Fritillaria radice rotunda. *Roy. lugdb.* 30.  
Lilium perficum. *Bauh. pin.* 79. *Rudb. elyf.* 2. p. 183.  
f. 1.  
Lilium fusianum. *Clus. hist.* 1. p. 130.  
*Habitat in Persia ? e Susis venit in Europam* 1573. 24  
*Folia infima sunt bina, f. terna, f. quaterna verticillatim opposita.*
- pyrenaica, 4. FRITILLARIA caule multifloro, foliis infimis oppositis. *Hort. cliff.* 81.  
Fritillaria flore minore. *Bauh. pin.* 64.  
Fritillaria pyrenæa. *Clus. hist.* 2. p. 256. *posth.* 10.  
*Habitat in Pyrenæis.* 24
- Meleagris, 5. FRITILLARIA caule subunifloro, foliis omnibus alternis. *Hort. upf.* 81. *Fl. fvec.* 2. n. 283.  
Fritillaria e foliorum alis florens. *Hort. cliff.* 119.  
Fritillaria radice depressa. *Roy lugdb.* 30.  
Fritillaria præcox purpurea variegata. *Bauh. pin.* 64.  
Meleagris. *Renal. spec.* 147. t. 146.  
β. Fritillaria alba variegata. *Bauh. pin.* 64.  
γ. Fritillaria alba præcox. *Bauh. pin.* 64.  
δ. Fritillaria serotina atropurpurea. *Bauh. pin.* 64.  
*Habitat in Gallia, Italia, Austria, Upsalia.* 24

## UVULARIA.

- amplexifolia, 1. UVULARIA foliis amplexicaulibus.  
Uvularia foliis cordato-oblongis. *Roy. lugdb.* 29.  
Smilax perfoliata ramosa, flore albo. *Barr. rar.* 58. t. 720.  
Polygonatum latifolium ramosum. *Bauh. pin.* 303.  
*Bauh. hist.* 3. p. 531. *Morif. hist.* 3. p. 537. f. 13. t. 4.  
f. 11. *Raj. hist.* 665.  
Polygonatum latifolium 4 ramosum. *Clus. hist.* 1. p. 276.  
Laurus alexandrina. *Cam. epit.* 936.  
*Habitat in Bohemiæ, Silesiæ, Saxoniæ, Delphinatus montibus.* 24

2. *UVULARIA* foliis perfoliatis. *Amæn. Acad.* 2. p. perfoliata.

337.

*Uvularia* caule perfoliato. *Gron. virg.* 37. *Cold. noveb.* 74.

*Uvularia* folio integerrimo. *Roy. lugdb.* 29.

*Uvularia. Hort. cliff.* 121.

*Polygonum* ramosum, flore luteo, majus. *Corn. canad.*

38. *Morif. hist.* 3. p. 538. f. 13. t. 4. f. 12.

*Polygonatum* latifolium perfoliatum brasiliæ. *Bauh. pin.* 303. *prodr.* 139. *Burf. XVII.* 61.

*Habitat* in Virginia, Canada. *Plantæ Burferi ex Toupinambault Brasiliæ a pharmacopæo allatæ, & C. B. communicatæ, videntur mihi omnes ex Canada fuisse.*

3. *UVULARIA* foliis sessilibus.

sessilifolia.

*Uvularia* foliis sessilibus, flore unico. *Cold. noveb.* 73.

*Habitat* in Canada. *Kalm.*

*Habitus præcedentis*; *Caulis bifidus, uniflorus. Folia lanceolato-ovata, alterna, at ubi Flos ex altero ramo exit, folia bina adsunt. Pedunculus nudus,*

## GLORIOSA.

1. *GLORIOSA. Hort. cliff.* 121. *Fl. zeyl.* 122.\* *Roy. superba. lugdb.* 29.

*Methonica* malabarorum. *Herm. lugdb.* 688. t. 689.

*Pluk. alm.* 249. t. 116. f. 3.

*Lilium* zeylanicum superbum. *Comm. hort.* 1. p. 69. t.

35. *Rudb. elyf.* 2. p. 178. f. 7.

*Mendon. Rheed. mal.* 7. t. 107. f. 57.

*Habitat* in Malabar. 24

## ERYTHRONIUM.

1. *ERYTHRONIUM. Hort. cliff.* 119. *Gmel. sib.* 1. *Dens canis* p. 39. t. 7. *Roy. lugdb.* 30.

*Dens canis* latiore rotundioreque folio. *Bauh. pin.* 87.

*Dens caninus. Dod. pempt* 203.

β. *Dens canis* angustiore longioreque folio. *Bauh. pin.* 87.

γ. *Erythronium* foliis ovato-oblongis glabris nigro-maculatis. *Gron. virg.* 151. *flore flavo.*

*Habitat* in Liguria, Allobrogibus, *Augustæ Taurinorum, Sibiria, Virginia.* 24



## TULIPA.

- sylvestris.* 1. TULIPA flore subnutante, foliis lanceolatis. *Fl. suec. 2 n. 284. \**  
*Tulipa minor lutea italica. Bauh. pin. 63.*  
*Tulipa minor lutea gallica. Bauh. pin. 63. Sauv. monsp. 306.*  
*Habitat Monspeli, inque Apenninis, Londini. 2.*  
*Distinguitur a sequente: Caule altiore, Folii angustioribus, Floris colore luteo subtus virescente, petalis acutioribus, Scapo versus florem inflexo & ante florescentiam nutante. Sed præprimis staminum filamentis basi valde attenuatis, & supra basin hirsutis. Petala interiora margine hirsuta sunt, & flos odorus.*
- gesneriana.* 2. TULIPA flore erecto, foliis ovato-lanceolatis.  
*Tulipa scapo triphylo, foliis ovato-lanceolatis. Hort. upf. 82.*  
*Tulipa. Hort. cliff. 118. Roy. lugdb. 31. Fl. suec. 262.*  
*Tulipa (genus fere totum). Bauh. pin. 57.*  
*Tulipa turcarum. Cord. hist. 213.*  
*Habitat in Cappadocia, unde in Europam 1559. Gesnero autore. 2.*
- breyniana.* 3. TULIPA caule multifloro-polyphylo, foliis linearibus.  
*Sisyrinchium, ex phœniceo suaverubente flore, æthiopicum. Breyn. cent. t. 36. Rudb. elys. 2. p. 13. f. 11.*  
*Habitat in Æthiopia. 2.*  
*Folia caulina 6. f. 7, alterna, lineari-lanceolata: summa sensim breviora, Flores ad apicem, 3. f. 4. minores quam plantæ Breynii cultæ. Petala basi angustata. Stylus nullus.*

## ALBUCA.

- major.* 1. ALBUCA foliis lanceolatis.  
*Ornithogalum canadense petalis alternis patentibus: interioribus erectis. Sp. pl. 1. p. 308.*  
*Ornithogalum luteo-virens indicum. Corn. canad. 160. Rudb. elys. 2. p. 140. f. 7. Raj. hist. 1154. Moris. hist. 2. p. 423. f. 4. t. 24. f. 7.*  
*Habitat ad Cap. b. Spei. 2.*
- minor.* 2. ALBUCA foliis subulatis.

*Ornithogalum africanum*, flore viridi altero alteri innato. *Herm. parad.* 209. t. 209.  
*Habitat ad Cap. b. Spei.* 24

## HYPOXIS.

1. **HYPOXIS** pilosa, capsulis ovatis. erecta.  
*Ornithogalum hirsutum* scapo angulato, pedunculis umbellatis villosis. *Sp. pl.* 1. p. 306.  
*Ornithogalum* scapo bifloro. *Roy. lugdb.* 31.  
*Ornithogalum vernum* luteum, foliis angustis hirsutis.  
*Gron. virg.* 37.  
*Ornithogalum virginianum* luteum. *Pet. gaz.* 1. t. 1. f. 3.  
*Ornithogalum herbaceum* luteum parvum virginianum, foliis gramineis hirsutis. *Pluk. alm.* 272. t. 350. f. 12.  
*Habitat in Virginia, Canada.* 24
2. **HYPOXIS** pilosa, capsulis clavatis. *Aman. acad.* 5. de cumbens.  
*p.* 396. \*  
*Ornithogalum*, foliis gramineis, floribus geminatis, pedunculis longissimis alaribus. *Brown. jam.* 195.  
*Anthericum* sessile, foliis linearibus planis, caule decumbente. *Mill. dist.* t. 39. f. 2.  
*Crocus* foliis & radice scorzonera. *Plum. ic.* 108. f. 1.  
*Habitat in America meridionali.* 24
3. **HYPOXIS** tubis florum longissimis, † fascicularis.  
*Allium sylvestre* f. *Moly* minus, amplo flore. *Russel. alep.* 34. t. 2.  
*Habitat ad Aleppo.* 24
4. **HYPOXIS** pilosa acaulis, fructificationibus subradicalibus. † sessilis.  
*Ornithogali virginici* facie herba tuberosa carolinensis.  
*Dill. elth.* 298. t. 220. f. 287.  
*Habitat in Carolina.* 24

## ORNITHOGALUM.

\* *flaminibus omnibus subulatis.*

1. **ORNITHOGALUM** scapo anguloso diphylo, pedunculis umbellatis simplicibus. *Fl. suec.* 270, 285. luteum.  
*Ornithogalum* scapo diphylo, pedunculis simplicibus terminalibus, filamentis omnibus subulatis. *Hort. cliff.* 124. *Roy. lugdb.* 31.

- Ornithogalum luteum.* *Bauh. pin.* 71.  
*Pyrrochiton.* *Reeal. spec.* 91. t. 90.  
*Habitat in Europæ cultis, macellis.* 24
- minimum. 2. *ORNITHOGALUM* scapo angulato diphylo, pedunculis umbellatis ramosis. *Fl. suec.* 271. 286. \*  
*Ornithogalum luteum minus.* *Bauh. pin.* 71.  
*Hypoxis.* *Reeal. spec.* 92.  
*Habitat in Europæ cultis oleraceis.*
- pyrenaicum 3. *ORNITHOGALUM* racemo longissimo, filamentis lanceolatis, pedunculis floriferis patentibus æqualibus: fructiferis scapo approximatis.  
*Ornithogalum* racemo longissimo, filamentis dilatato-linearibus, capsulis erectis. *It. scan.* 220.  
*Ornithogalum spica longissima*, filamentis triangularibus. *Hall. herb.* 294. *Gron. orient.* 110.  
*Ornithogalum angustifolium majus*, floribus ex albo virefcentibus. *Bauh. pin.* 70. *Rudb. chys.* 2. p. 134. f. 3.  
*Ornithogalum majus.* *Clus. hist.* 1. p. 187.  
*Ornithogalum pyrenaicum.* *Clus. cur.* 21.  
*Stachyoides.* *Reeal. spec.* 93. t. 90.  
*Habitat in Alpibus Helveticis, Genevensibus, Pyrenaicis, Carniolæ* 24
- narbonense. 4. *ORNITHOGALUM* racemo oblongo, filamentis lanceolatis membranaceis, pedunculis floribusque patentibus. *Amæn. acad.* 4. p. 312. \*  
*Ornithogalum majus spicatum*, flore albo. *Bauh. pin.* 70.  
*Ornithogalum narbonense.* *Dod. pempt.* 222.  
*Habitat in Galliæ australis & Italiæ agris.* 24
- comosum. 5. *ORNITHOGALUM* racemo brevissimo, bracteis lanceolatis longitudine florum, petalis obtusis, filamentis subulatis. *Amæn. acad.* 4. p. 312. \*  
*Ornithogalum spicatum* f. comosum, flore lacteo. *Bauh. pin.* 70. *Rudb. chys.* 2. p. 135. f. 1.  
*Ornithogali lactei species major.* *Best. cyfl.*  
*Habitat* - - - - 24
- latifolium. 6. *ORNITHOGALUM* racemo longissimo, foliis lanceolato-ensiformibus.  
*Ornithogalum latifolium & maximum.* *Bauh. pin.* 70.  
 Ornith-

*Ornithogalum* vel *Lilium alexandrinum*, floribus albis innumerabilibus. *Swert. floril.* 58.

*Habitat in Arabia, Ægypto.*

*Folia pedalia plus quam duos pollices lata.* 24

7. *ORNITHOGALUM* racemo conico, floribus numerosis adscendentibus. *Roy. lugdb.* 32. pyramidale.

*Ornithogalum angustifolium spicatum maximum.* *Bauh. pin.* 70. *Rudb. elys.* 2. p. 134. f. 4.

*Ornithogalum lacteum maximum.* *Besl. cyst. vern.* 5. t. 14. f. 2.

*Habitat in collibus Lusitaniæ.* 24

\* \* *Staminibus alternis emarginatis.*

8. *ORNITHOGALUM* floribus corymbosis, pedunculis scapo humilioribus, filamentis subemarginatis. *Roy. lugdb.* 32. arabicum.

*Ornithogalum umbellatum maximum.* *Bauh. pin.* 69. *Rudb. elys.* 2. p. 130. f. 1.

*Ornithogalum arabicum.* *Clus. hist.* 11. p. 186. *Besl. cyst. vern.* 5. t. 12. f. 1.

*Melanomphale.* *Rencal. spec.* 89. t. 90.

*Habitat Alexandria Ægypti, ad Cap. b. spei.* 24

*Stamina connata.* *Hall. gert.* 81.

9. *ORNITHOGALUM* floribus corymbosis, pedunculis scapo altioribus, filamentis emarginatis. *Hort. cliff.* 124. *Hort. upf.* 84. *Roy. lugdb.* 32. *Gron. orient.* 111. umbellatum.

*Ornithogalum umbellatum medium angustifolium.* *Bauh. pin.* 70.

*Eliocarmos.* *Rencal. spec.* 88. t. 87.

*Habitat in Germania, Gallia, Oriente.* 24

10. *ORNITHOGALUM* floribus secundis pendulis, nectario stamineo campaniformi.

*Ornithogalum floribus secundis pendulis, filamentis latis emarginatis.* *Hort. cliff.* 124. *Hort. upf.* 84.

*Roy. lugdb.* 32.

*Ornithogalum exoticum, magno flore minori innato.* *Bauh. pin.* 70. *Rudb. elys.* 1. p. 137. f. 12.

*Ornithogalum neapolitanum.* *Clus. app.* 2. p. 9. t. 9.

*Habitat in Italia Neapoli innotuit circa 1570.* 24

11. *ORNITHOGALUM* foliis cordato-ovatis. *Roy. capense.* *lugdb.* 31. *It. scan.* 73. \*

*Ornithogalum africanum*, plantaginis roseæ folio, radice tuberosa. *Comm. hort.* 2. p. 175. t. 88.

*Ornithogalo affinis* radice tuberosa, cyclaminis folio, flore pallide cæruleo. *Breyn. cent.* t. 41. *Rudb. elyf.* 1. p. 138. f. 14.

*Habitat ad Cap. b. Spei.* 24.

*Stamina connexa in cylindrum.*

## SCILLA.

*maritima.* 1. *SCILLA nudiflora*, bracteis refractis.

*Scilla* radice tunicata. *Hort. cliff.* 123. *Hort. upf.* 89.

*Mat. med.* 162. *Roy. lugdb.* 32.

*Scilla vulgaris*, radice rubra. *Bauh. pin.* 73.

*Scilla hispanica.* *Clus. hist.* 1. p. 171.

β. *Scilla radice alba.* *Bauh. pin.* 73. *Seb. musf.* 1. t. 44. f. 4. 5.

*Habitat ad Hispaniæ, Siciliæ, Styriæ litora arenosa.* 24.

*Folia lanceolata, stricta. Scapus ante folia, longissimus, multiflorus. Flores albi: Bracteæ lineari-lanceolatae, refractæ genu in medio, sursum flexæ, subtus calcaratæ.*

*Lilio-Hyacinthus.* 2. *SCILLA radice squamata.* *Hort. cliff.* 123. *Roy. lugdb.* 32.

*Hyacinthus stellaris*, folio & radice lillii. *Bauh. pin.* 46.

*Habitat in Biscatia, Aquitania, Hispania, Pyrenæis.* 24.

*italica.* 3. *SCILLA corymbo conferto hemisphærico.*

*Scilla* radice solida, floribus corymbosis confertissimis.

*Hort. cliff.* 123. *Roy. lugdb.* 32.

*Hyacinthus stellaris spicatus cinereus.* *Bauh. pin.* 46.

*Hyacinthus stellaris italicus.* *Besl. cyst. vern.* 42. f. 1.

*Hyacinthus stellatus cinerei coloris.* *Clus. hist.* 1. p. 184.

*Habitat - - -* 24.

*peruviana.* 4. *SCILLA corymbo conferto conico.*

*Hyacinthus indicus bulbosus stellatus.* *Bauh. pin.* 47.

*Rudb. elyf.* 1. p. 37. f. 1.

*Hyacinthus stellatus peruvianus.* *Clus. hist.* 1. p. 182.

*Eriophorus peruvianus.* *Clus. hist.* 1. p. 173.

*Habitat in Lusitania.* 24.

*Radix tunicata, lanuginosa. Sauv. Scapus teres. Corollæ senescentes recurvatæ, petalorum marginibus recurvatis,*

5. *SCIL-*

5. **SCILLA** floribus lateralibus alternis subnatis. *amœna.*  
*Hort. cliff. 123. Roy. lugdb. 33.*  
*Hyacinthus stellaris cœruleus amœnus. Bauh. pin. 46.*  
*Rudb. elyf. 2. p. 34. t. 7.*  
*Hyacinthus stellaris bizantinus, Bess. cyst. vern. 43. f. 2.*  
*Habitat forte Bizantii, unde venit in Europam 1590. 24*
6. **SCILLA** radice solida, floribus erectiusculis paucifloris. *bifolia.*  
*Hort. cliff. 123. Roy. lugdb. 33. Sauv. monsp. 18.*  
*Hyacinthus stellaris bifolius germanicus. Bauh. pin. 45.*  
*Hyacinthus cœruleus mas minor. Fuch. hist. 837.*
- β. *Hyacinthus stellatus, albo flore. Clus. hist. 1. p. 184.*  
*Habitat in Gallia, Germania. 24*  
*Flores sæpe 4. altitudine æquales.*
7. **SCILLA** foliis filiformibus linearibus, floribus corymbosis, pedunculis nudis adscendentibus longitudine floris. *autumnalis.*  
*Scilla radice solida, foliis setaceis, floribus fastigiatis, pedunculis arcuatis ex ala tuberculi mammillaris.*  
*Guett. stamp. 131. Dalib. parif. 102.*  
*Scilla radice solida, scapo multifloro, floribus capitatis. Sauv. monsp. 19.*  
*Hyacinthus stellaris autumnalis minor. Bauh. pin. 47.*  
*Hyacinthus autumnalis minor. Clus. 1. p. 185.*  
*Habitat in Hispania, Gallia, Veronæ, solo glareoso. 24*
8. **SCILLA** folio teretiusculo, latere subspicato. *unifolia.*  
*Bulbus monophyllus, flore albo. Bauh. hist. 2. p. 622.*  
*Habitat in Lusitania.*

## CYANELLA.

1. **CYANELLA.** *Capensis.*  
*Habitat ad Cap. b. spei. Royen. Barmannus. 24.*  
*Statura Scillæ. Flores irregulares.*

## ASPHODELUS.

1. **ASPHODELUS** caule folioso, foliis triquetris striatis. *luteus.*  
*Hort. cliff. 127. Hort. ups. 83. Roy. lugdb. 33.*  
*Asphodelus luteus flore & radice. Bauh. pin. 28.*  
*Asphodelus femina, Cam. epit. 372.*  
*Habitat in Sicilia. ♂*

Corollæ sexpartita : petalis 4 in crucem dispositis : duobus intra superiora tria petala positis. Stamina declinata , quorum tria breviora.

- astulosus. 2. ASPHODELUS caule nudo , foliis strictis subulatis striatis subastulosis. *Hort. cliff.* 83. \*  
 Asphodelus caule nudo , foliis strictis. *Roy. lugdb.* 34.  
 Asphodelus foliis fistulosis. *Bauh. pin.* 29.  
 Asphodelus minor. *Clus. hist.* 1. p. 197.  
*Habitat in Gallo - Provincia , Hispania , Creta.* 24

- ramosus. 3. ASPHODELUS caule nudo , foliis ensiformibus carinatis lævibus. *Mat. med.* 172.  
 Asphodelus caule nudo , foliis laxis. *Hort. cliff.* 127.  
*Roy. lugdb.* 24. *Sauv. monsp.* 20.  
 Asphodelus albus ramosus mas. *Bauh. pin.* 28.  
 Asphodelus albus non ramosus. *Bauh. pin.* 28.  
 Asphodelus. 1. 2. *Clus. hist.* 1. p. 197.  
*Habitat in Narbona , Lusitania , Hispania , Italia , Austria.* 24

## ANTHERICUM.

\* *Phalangium foliis canaliculatis , filamentis glabris.*

- serotinum. 1. ANTHERICUM foliis planiusculis , scapo unifloro-  
 Bulbocodium foliis subulato-linearibus. *Roy. lugdb.* 41.  
*Sp. pl.* 1. p. 294.  
 Bulbocodium alpinum juncifolium , flore unico intus albo extus squalide rubente. *Raj. angl.* 3. p. 374.  
*t.* 17. f. 1.  
 Pseudo-Narcissus gramineo folio. *Bauh. pin.* 51. *prodr.* 27. *Rudb. elys.* 2. p. 64. f. 9. *Burf. III.* 41.  
*Habitat in alpidibus Angliæ , Helvetiæ , Taureri rasti-  
 diensis , Wallæsiæ.* 24  
*Scapus bracteis f. foliis 3 f. 4 , lanceolatis , alternis ,  
 minutis , vaginantibus absque Spatha. Flos inferus.*  
 græcum. 2. ANTHERICUM foliis planis , scapo simplici , flo-  
 ribus corymbosis.  
 Bulbocodium græcum , myosotidis flore. *Tournef.*  
*cor.* 50.  
*Habitat in Oriente. Burmannus.* 24  
*Bulbus. Folia linearia , glabra , longitudine fere scapi.*  
*Scapus simplex Foliis 3 f. 4 , brevioribus. Corym-  
 bus terminalis , subquinqueflorus : pedunculo interme-  
 dio*

*dio unifloro ; lateralibus bifloris. Corolla alba A. ramosi. Stamina corolla breviora & Stylus staminibus brevior.*

3. **ANTHERICUM** foliis planis , scapo ramoso , corollis revolutis.

*Asphodelus foliis compressis asperis , caule patulo.*

*Tournef. inst. 343.*

*Habitat - - - - - 24*

4. **ANTHERICUM** foliis planis , scapo ramoso , corollis planis , pistillo recto.

*Anthericum foliis planis , corollis planis deciduis. It.*

*gottl. 178. \* Fl. succ. 267 , 289.*

*Anthericum caulibus ramosis , foliis planis. Guett.*

*stamp. p. 129.*

*Phalangium , parvo flore , ramosum. Bauh. pin. 29.*

*Phalangium majus. Cam. epit. 580.*

*Habitat in Europæ australioris rupibus calcareis. 24*

5. **ANTHERICUM** foliis planis , scapo simplicissimo , corollis planis , pistillo declinato.

*Lillago. Hort. upf. 83. It.*

*scan. 104. Fl. succ. 2. n. 290.*

*Anthericum caulibus non ramosis , foliis planis. Guett.*

*stamp. 1. p. 128.*

*Phalangium , parvo flore , non ramosum. Bauh. pin.*

*29. Moris. hist. 2. p. 333. f. 4. t. 1. f. 10.*

*Phalangium non ramosum. Lob. ic. 48.*

*Habitat in Helvetia , Germania , Gallia. 24*

6. **ANTHERICUM** foliis planis , scapo simplicissimo , corollis campanulatis , staminibus declinatis.

*Hemerocallis Liliastrum scapo simplici , corollis hexa-*

*petalis campanulatis. Hort. cliff. 128. Sp. pl. 1. p.*

*324. Hall. helv. 290.*

*Phalangium magno flore. Bauh. pin. 29.*

*Phalangium allobrogicum majus. Clus. cur. app. alt.*

*Phalangium. Dakch. hist. 852.*

*Habitat in Alpibus Helveticis , Allobrogicis. 24*

*Generis hujus naturalis Species est , sed quis character*

*uniet ?*

*\* \* Bulbine foliis carnosiss , Filamentis barbatis.*

7. **ANTHERICUM** foliis carnosiss teretibus , caule frutescens fruticoso.

An-



- Anthericum caulescens*, foliis pulposis teretibus subulatis supine convexo-planis. *Wach. ultr.* 305.  
*Bulbine caulescens*. *Hort. cliff.* 122. *Roy. lugdb.* 122.  
*Phalangium capense caulescens*, foliis cepitiis succosis.  
*Dill. elih.* 310. t. 231. f. 298.  
*Habitat ad Cap. b. Spei.* b
- alooides. 8. **ANTHERICUM** foliis carnosis subulatis planiusculis. *Hort. upf.* 83.  
*Anthericum foliis pulposis lanceolato-subulatis supine excavatis*. *Wach. ultr.* 304.  
*Bulbine acaulis*. *Hort. cliff.* 123. *Roy. lugdb.* 33.  
*Phalangium capense sessile*, foliis aloeformibus pulposis. *Dill. elih.* 312. t. 232. f. 300.  
*Habitat ad Cap. b. Spei.* 24
- Asphodeloides. 9. **ANTHERICUM** foliis carnosis subulatis semiteretibus strictis. *Hort. upf.* 83.  
*Anthericum acaule*, foliis pulposis teretibus subulatis supine convexo-planis. *Wach. ult.* 305.  
*Habitat in Æthiopia.* 24  
*Folia vix striata, sed stricta, margine aspera, dorso versus alterum latus gibbiora.*
- annuum. 10. **ANTHERICUM** foliis carnosis subulatis teretibus, scapo subracemoso. *Hort. upf.* 83. \*  
*Anthericum acaule*, foliis setaceis caule angustioribus. *Wach. ultr.* 305.  
*Habitat in Æthiopia.* ☉
- hispidum. 11. **ANTHERICUM** foliis carnosis compressis hispidis. *Asphodelus caule seminudo ramoso*, foliis planis lanceolatis. *Mill. dist.* t. 56?  
*Habitat ad Cap. b. Spei.* 24  
*Radix fibrosa. Folia subcarnosa, compressa, striata, semipedalia, recurvata, hispida undique pilis patulis. Scapus simplex, hispidus. Racemus bracteis lanceolatis, longitudine pedicellorum hispidis, uti Perdicelli. Flores albi carina viridi. Antheræ luteæ.*  
 \* \* \* *Nartheicum foliis ensiformibus.* .
- offisragum. 12. **ANTHERICUM** foliis ensiformibus, filamentis lanatis. *Fl. fvec.* 268, 287. *Gmel. fibir.* 1. p. 73. t. 18. f. 2.

*Anthericum* scapo folioso laxè spicato, filamentis vil-  
lofis. *Fl. lapp.* 136. *Roy. lugdb.* 45.

*Narthecium.* *Mähr. E. N. C.* 1742. p. 389. t. 5. f.

1. *Wach. ultr.* 303. *Gort. gedr.* 69.

*Pseudo-Asphodelus palustris anglicus.* *Bauh. pin.* 29.

*Asphodelus luteus palustris.* *Dod. pempt.* 208.

*Habitat in Europæ borealis uliginosis.* 24

13. *ANTHERICUM* foliis ensiformibus, perianthiis calyculatum  
trilobis, filamentis glabris, pistillis trigynis. *Fl. suec.*

269. 288. *L. gotl.* 194, 221. *Gmel. sib.* 1. p. 13. t. 18. f. 2.

*Anthericum* filamentis lævibus, perianthio trifido.

*Hort. cliff.* 140. *Gron. virg.* 39.

*Anthericum* scapo nudo capitato, filamentis glabris.

*Fl. lapp.* 139. t. 10. f. 3.

*Phalangium alpinum palustre, iridis folio.* *T. Segu.*

*ver.* 2. p. 61. t. 14.

*Pseudo-Asphodelus alpinus.* *Bauh. pin.* 29.

*Habitat in alpidibus Helvetiæ, Lapponiæ, Sibiriciæ.* 24

Genus nullam in tota classe *Liliacea* difficilius deter-  
minatur: 12 ossifragum & 13 calyculatum habitu

toto ita. uniuntur, ut demta fructificatione vix di-  
stinguantur, at Species 13 gerit Stamina glabra,

*Pericarpium* subglobosum, Stigmata tria sine stylis  
& *Perianthium* trigalve. Species 12, annectitur

speciebus 7, 8, 9, 10, 11. *Staminibus* hirsutis, at  
differt Germine pyramidato, *Seminibus* paleaceis,

*Corolla* persistente. Species 1 — 6. differunt a spe-  
ciebus 7 — 10. *Staminibus* glabris, habitu *Aspho-*

*deli*, adeoque vel sub uno eodemque genere traden-  
da, vel etiam sub quatuor distinctis: α. 1. — 6.

β. 7 — 11. γ. 8; δ. 12, 13. proponenda; hoc dum  
fieret, difficillime character specierum α obtineretur.

## LEONTICE.

1. *LEONTICE* foliis pinnatis: petiolo communi fim- Chrysogo-  
num.  
plici. *Hort. cliff.* 122. *Gron. orient.* 113.

*Leontopetalum* foliis costæ simplici innascentibus. *Tour-  
nef. cor.* 49.

*Leontopetalum* affinis, foliis quernis. *Bauh. pin.* 324.

*Morif. hist.* 2. p. 285. f. 4 t. 5. f. 3.

*Chrysogonum* dioscoridis. *Rauv. itin.* 119. *Raj. hist.*  
1326.

*Habitat inter Græciæ segetes.* 24

2. *LE-*

- Leontopetalum.** 2. **LEONTICE** foliis decompositis: petiolo communi trifido. *Hort. cliff.* 122. *Gron. orient.* 114.  
**Leontopetalon** foliis costæ ramosæ innascentibus. *Tournef. cor.* 49.  
**Leontopetalon.** *Bauh. pin.* 324. *Cam. epit.* 565. *Morif. hist.* 2. p. 285. f. 3. t. 15 f. 6. *Raj. hist.* 1326.  
*Habitat in Apulia; Hetruria, Creta.* 2
- Thallactroides.** 3. **LEONTICE** folio caulino triternato (27), florali biternato (9).  
**Leontice** folius supradecompositis. *Gron. virg.* 151. \*  
*Cold. noveb.* 75.  
*Habitat in Virginia.* 2  
*Modus crescendi Epim-dii.* Caulis nudus, terminatus  
Foliis tribus, patentibus, petiolatis: Petiolus singulus trifidus, adnatis novem foliolis, quorum lateralia duo, sessilia, biloba; intermedium petiolatum, trilobum; infra divisuram primam caulis exserit e latere folium Petiolos tres: singulis simplicibus triphyllis, infra quos Racemus simpl. x cum duobus lateralibus.
- Leontopetaloides.** 4. **LEONTICE** foliis simplicibus tripartito-multifidis, corollis monopetalis calyculatis †  
**Leontopetaloides.** *Amm. act.* 4. p. 211. t. 113.  
*Habitat in India.*  
*Habitus hujus generis, at flos videtur satis diversus. an sufficienter pro distincto genere, determinent auctores.*

## ASPARAGUS.

- officinalis.** 1. **ASPARAGUS** caule herbaceo erecto, foliis setaceis, stipulis duabus interioribus, una exteriori. *Fl. suec.* 272, 291. *Nat. med.* 169. *Hort. cliff.* 121. *Roy. lugdb.* 28.
- maritimus.** α. **Asparagus maritimus**, crassiore folio. *Bauh. pin.* 490.  
β. **Asparagus sylvestris**, tenuissimo folio. *Bauh. pin.* 490.
- altiss.** γ. **Asparagus sativa.** *Bauh. pin.* 489.  
**Asparagus.** *Cam. epit.* 259.  
*Habitat in Europæ arenosis.* 2  
Caules herbacei, erecti, paniculati. Stipulæ solitariae exteriores & binæ interiores minores, inter quas Folia terna, raro 4 f. 5, linearia, sed in summis ramis, ubi nulli flores; at in infimis ramulus. Pedunculi

*gemi, laxi, uniflori, penduli, geniculo articulat.*  
*Corolla campanulata, petalis interioribus longioribus.*

2. **ASPARAGUS** caule inermi, ramis declinatis, foliis declinatus, setaceis. *Roy. lugdb.* 29.

*Habitat in Africa.*

3. **ASPARAGUS** aculeis solitariis, foliis ensiformibus falcatis. *Fl. zeyl.* 123. \*

*Asparagus foliis falcatis ex uno puncto numerosis.*

*Burm. zeyl.* 36. t. 13. f. 2.

*Habitat in Zeylona.* ♀

4. **ASPARAGUS** aculeis solitariis, ramis reflexis retro- retrofractus, fractisque, foliis fasciculatis.

*Asparagus frutescens, foliis fasciculatis setaceis terminalibus, ramis reflexis retroflexis.* *Hort. cliff.* 121.

\* *Roy. lugdb.* 28.

*Asparagus africanus tenuifolius, viminalibus virgis, foliis loricis adinstar ex uno puncto numerosis stellatim positis.* *Pluk. amalth.* 40. t. 375. f. 3.

*Habitat in Africa.* ♀

5. **ASPARAGUS** aculeis solitariis, caule erecto, ramis asaticis, filiformibus, foliis fasciculatis.

*Asparagus aculeis alternis, ramis solitariis filiformibus, foliis setaceis fasciculatis lateralibus.* *Hort. cliff.*

122. \* *Roy. lugdb.* 28.

*Habitat in Asia.*

6. **ASPARAGUS** spinis retroflexis, ramis flexuosis, foliis fasciculatis angulatis muticis deciduis.

*Asparagus aculeis solitariis, ramulis flexuosis binis oppositis spinosis foliosis, foliis fasciculatis.* *Roy. lugdb.* 29.

*Asparagus aculeatus, spinis horridus.* *Bauh. pin.* 490.

*Corduba tertia.* *Clus. hist.* 2. p. 178.

*Habitat in Lusitania, Hispania.* ♀

7. **ASPARAGUS** caule inermi fruticoso, foliis aciformibus rigidulis perennantibus mucronatis æqualibus.

*Asparagus foliis aciformibus pungentibus, caule fruticoso inermi.* *Sauv. monsp.* 45.

*Asparagus foliis acutis.* *Bauh. pin.* 490.

*Asparagus sylvestris.* *Cam. epit.* 260.

*Habitat in Lusitania, Hispania, Oriente.* ♀

- aphyllus. 8. ASPARAGUS aphyllus. spinis subulatis striatis inæqualibus divergentibus *Hort. cliff. 122. Roy. lugdb. 29.* Asparagus aculeatus alter tribus aut quatuor spinis ad eundem exortum. *Bauh. pin. 490.* Corduba altera. *Clus. hist. 2. p. 178.*
- capensis. 9. ASPARAGUS spinis lateralibus terminalibusque, ramis aggregatis, foliis fasciculatis. Asparagus tolius quinis setaceis, spinis terminalibus lateralibusque, ramulis ternis quaternisque. *Hort. cliff. 121.* Asparagus aculeatus, triplici spina surrectus. *Pluk. alm. 54. t. 15. f. 4.* Habitat ad Caput b. Spei.  $\dagger$
- sarmentosus. 10. ASPARAGUS foliis solitariis lineari-lanceolatis, caule flexuoso, aculeis recurvis. *Fl. zeyl. 124. \** Asparagus aculeatus zeylanicus maximus sarmentosus. *Herm. lugdb. 62. t. 650. 63. Kaj. hist. 1877.* Schadaveli Kelangu. *Kheed. mal. 10. p. 19.* Habitat in Zeylona.  $\dagger$ . Affinis maxime *Medeolæ asparagoidi.*
- verticillatus. 11. ASPARAGUS foliis verticillatis  $\dagger$  Asparagus orientalis, foliis gallii. *Tournef. cor. 21.* *Buxb. cent. 5. app. 47. t. 37.* Habitat in Oriente, circa Derbentum  $\mathfrak{E}$  alibi.
- graminifolius. 12. ASPARAGUS foliis linearibus striatis, scapo racemoso. Habitat in Asia. Statura *Antherici phalangii.* Folia graminea, valde striata, spithamæa. Scapus vix foliis longior. Flores ad singulos dentes racemi 4. f. 5, pedicellis flore ipso brevioribus, quasi e gluma obtusa produntibus.
- terminalis. 13. ASPARAGUS inermis, foliis alternis lanceolatis petiolatis, racemo terminali composito. Terminalis. *Rumph. amb. 4. p. 79. t. 34.* Habitat in India.  $\dagger$  Folia fere Cannæ. Racemus terminalis ex aliquot racemis ramosis. Pedicelli alterni, solitarii, flore breviores, cincti ad basin quasi gluma obtusa. *Albar duri.*

14. ASPARAGUS? caudice simplici, foliis ensiformi- Draco,  
bus mucronatis subcarneosis imbricatis patentibus.  
Cordylina foliis inermibus integerrimis flaccidis. Roy.  
lugdb. 22.  
Palma foliis longissimis pendulis, absque pedunculis e  
caudice glabro enatis. Boerh. lugdb. 2. p. 169.  
Draco arbor. Bauh. pin. 503. Clus. hist. 1. p. 1.  
Habitat in India orientali. h  
Fructificationem, quam etiamnum Ulyssipone ♂ in Hi-  
spania producit, non obtinui, Flores Asparagi similes  
retulit Læffing.

CONVALLARIA.

\* Lil. conv. T. corollis campanulatis.

1. CONVALLARIA scapo nudo. Fl. lapp. 113. Fl. suec. majalis.  
273, 292. Mat. med. 167. Hort. cliff. 124. Roy.  
lugdb. 26 Gmel. sibir. 1. p. 34.  
Lilium convallium album. Bauh. pin. 304.  
Lilium convallium alpinum. Bauh. pin. 304.  
Lilium convallium latifolium. Bauh. pin. 304.  
Habitat in Europa septentrionali. 24

\* \* Polygonata T. corollis infundibuliformibus.

2. CONVALLARIA foliis verticillatis. Fl. lapp. 114. verticillata,  
Fl. suec. 275, 293. Hort. cliff. 125. Roy lugdb. 26.  
Polygonatum angustifolium non ramosum. Bauh. pin. 303.  
Polygonatum alterum. Dod. pempt. 345.  
3. Polygonatum angustifolium ramosum. Bauh. pin. 304.  
Habitat in Europæ septentrionalis saltibus, præcipue  
in 24

3. CONVALLARIA foliis alternis amplexicaulibus, Polygonata-  
caule ancipiti, pedunculis axillaribus subunifloris. tum.  
Mat. med. 168. Gmel. sibir. 1. p. 34. Phil. bot. 218.  
Convallaria foliis alternis, floribus axillaribus. Fl. suec.  
274, 294. Hort. cliff. 124. Gron. virg. 37. Roy. lugdb. 26.  
Convallaria foliis alternis, pedunculis pendulis uni-  
floris. Sauv. monsp. 42.  
Polygonatum floribus ex singularibus pedunculis. Bauh.  
pin. 3. p. 530.  
Polygonatum latifolium, flore majore odore. Bauh.  
pin. 303.

*Habitat in Europæ septentrionalis præcipitiis, rupibus.* 24.

multiflora.

4. CONVALLARIA foliis alternis amplexicaulibus, caule tereti, pedunculis axillaribus multifloris. *Phil. bot.* 218. *Fl. suec.* 2. p. 295.

*Convallaria foliis alternis, pedunculis pendulis multifloris. Sauv. monsp.* 42.

*Polygonatum latifolium maximum. Bauh. pin.* 303.

*Polygonatum latifolium. 1. Clus. hist.* 1. p. 275.

*Habitat in Europæ septentrionalis præcipitiis, rupibus.* 24.

\* \* \* *Smilaces T. corollis rotatis.*

racemosa.

5. CONVALLARIA foliis sessilibus, racemo terminali composito.

*Convallaria racemo composito. Roy. lugdb.* 26.

*Convallaria foliis alternis, racemo terminali. Hort. cliff.* 125. *Gron. virg.* 38.

*Polygonatum racemosum. Corn. canad.* 36. t. 37.

*Polygonatum ramosum & racemosum spicatum. Moris. hist.* 3. p. 537. f. 13. t. 4. f. 9.

*Polygonatum racemosum americanum, ellebori albi foliis amplissimis. Pluk. alm.* 301. t. 311. f. 2.

*Habitat in Virginia, Canada.* 24.

stellata.

6. CONVALLARIA foliis amplexicaulibus plurimis.

*Polygonatum virginianum erectum spicatum, flore stellato. Moris. hist.* 3. p. 536. f. 13. t. 4. f. 7.

*Polygonatum canadense spicatum fertile. Corn. canad.* 33. t. 33.

*Habitat in Canada.* 24.

trifolia.

7. CONVALLARIA foliis amplexicaulibus ternis, racemo terminali simplici.

*Convallaria floribus racemosis, foliis ovatis oblongis caulinis. Gmel. sibir.* 1. p. 36. t. 4.

*Phalangium veratri foliis. Amm. ruth.* 137.

*Habitat in Sibiria sibir.* 24.

bifolia.

8. CONVALLARIA foliis cordatis. *Fl. lapp.* 113. *Fl.*

*suec.* 276, 296. *Hort. cliff.* 125. *Roy. lugdb.* 26.

*Gmel. sibir.* 1. p. 35.

*Lilium convallium minus. Bauh. pin.* 304.

*Gramen parnassi. Cam. epit.* 744.

*Habitat in Europæ borealis pratis depressis, asperis.* 24.

POLIANTHES.

1. POLIANTHES.

*Polianthes floribus alternis. Hort. cliff. 127. Hort. upf. 76. Fl. zeyl. 225.*

*Hyacinthus indicus tuberosus, flore narciss. Bauh. pin. 47. Rudb. elys. 2. p. 39. f. 4.*

β. *Hyacinthus indicus tuberosus flore hyacinthi orientalis. Bauh. pin. 47. Rudb. elys. 2. p. 38. f. 2.*

*Hyacinthus indicus, tuberosa radice. Clus. hist. 1. p. 176.*

*Amica nocturna. Rumph. amb. 5. p. 285. t. 98.*

*Habitat in Iava, Zeylona. 24*

tuberosa.

HYACINTHUS.

1. HYACINTHUS corollis campanulatis sexpartitis apice revolutis. *Hort. cliff. 125. Roy. lugdb. 27. Sauv. monsp. 17.* non scriptus.

*Hyacinthus, oblongo flore, cæruleus major. Bauh. pin. 43.*

*Habitat in Angliæ, Galliæ, Hispaniæ, Italiæ nemonibus. 24*

*Bractæe junctæ sunt floribus longiores, & binæ. Petala apice revoluta. Stigma guttula madidum.*

2. HYACINTHUS corollis campanulatis sexpartitis, racemo cernuo.

*Hyacinthus hispanicus. Clus. hist. 1. p. 177.*

*Hyacinthus, oblongo flore suaviter rubente, minor. Bauh. pin. 44?*

*Habitat in Hispania. 24*

*Convenit cum H. non scripto habitu & facie, quamvis minor; differt vero Foliis magis linearibus minus lanceolatis, erectioribus; Racemo magis nutante; Corollis incarnatis nec cæruleis, teretioribus, cum petalorum lateribus ad basin minus plano-pateantibus, nec dorso linea elevata notatis, minusque revolutis (quamvis reflexa) quam in illo, Pistillum demum staminibus brevius est. Bractæe, binæ saturatius incarnatæ.*

3. HYACINTHUS corollarum exterioribus petalis subdistinctis, interioribus coadunatis. ferotinus.

*Hyacinthus obsoleto flore. Bauh. pin. 44.*

*Hyacinthus, obsoleto coloris, hispanicus ferotinus.*

*Clus. hist. 1. p. 177, 178.*

*Habitat in Hispania, Mauritania. 24*



- Racemus secundus. Flores tristi colore, nec tamen noctu ambrosiaci. Corolla sexfida: laciniis 3 alternis exterioribus fere ad basin usque separatis recurvisque.*
- viridis.** 4. **HYACINTHUS** corollarum exterioribus laciniis subulatis longissimis.  
*Habitat ad Cap. b. spei; vivam aluit D. Burmannus. Statura H. amethystini, sed Corollae virides: Laciniis tribus exterioribus alternis duplo longioribus toto flore, subulatis in acumen angustissimum.*
- amethystinus.** 5. **HYACINTHUS** corollis campanulatis semisextidis basi cylindricis. *Hort. upf. 85.*  
*Hyacinthus, oblongo caeruleo flore, minor. Bauh. pin. 44. Rudb. elyf. 2. p. 27. f. 8.*  
*Hyacinthus minor hispanicus angustifolius. Bauh. hist. 2. p. 587. Clus. cur. app. alt.*  
*Habitat in Hispania ? 24*
- orientalis.** 6. **HYACINTHUS** corollis infundibuliformibus semisextidis basi ventricosis. *Hort. upf. 85. Hort. cliff. 125. Roy. lugdb. 27 Gron. orient. 115.*  
*Hyacinthus orientalis, (spec. 1.—15) & plenus (1—3) Bauh. pin. 44.*  
*Hyacinthus orientalis major & minor. Dod. pempt. 216.*  
*Habitat in Asia, Africa. 24*
- Muscari.** 7. **HYACINTHUS** corollis ovatis omnibus aequalibus. *Hort. cliff. 126. Hort. upf. 85. Roy. lugdb. 28.*  
*Hyacinthus racemosus moschatus. Bauh. pin. 43.*  
*Muscari obsoletiore flore. Clus. hist. 1. p. 178.*  
*Habitat in Asia, ultra Bosphorum, inde in Europam ante 1554. 24*
- monstrosus.** 8. **HYACINTHUS** corollis subovatis: *Vir. cliff. 28. Roy. lugdb. 28.*  
 β- *Hyacinthus floribus paniculatis monstrosis. Hort. cliff. 126.*  
*Hyacinthus panicula caerulea. Bauh. pin. 42.*  
*Hyacinthus sannesius, panicula comosa. Col. ecphr. 2. p. 10. t. 12.*  
*Habitat - - - primum inventa in agro Papiensi, & juxta Boran Galliae. 24*  
*Videtur sequentis sola varietas ?*

9. **HYACINTHUS** corollis angulato - cylindricis: sum- racemosus.  
 mis sterilibus longius pedicelatis.  
*Hyacinthus corollis globosis: summis pedunculatis, foliis ensiformibus. Sauv. monsp. 17.*  
*Hyacinthus comosus, major purpureus. Bauh. pin. 42.*  
*Hyacinthus. Cam. epit. 798.*  
*Habitat in Galliæ & Europæ australis agris. 5*
10. **HYACINTHUS** corollis globosis uniformibus, botryoides.  
 foliis canaliculato-cylindricis strictis.  
*Hyacinthus corollis globosis. Hort. cliff. 126. Hort. wyl. 85. Roy. lugdb. 28. \**  
*Hyacinthus corollis globosis, foliis in cylindrum convolutis. Sauv. monsp. 19.*  
*Hyacinthus racemosus cæruleus major. Bauh. pin. 42.*  
*Hyacinthus botryoides purpureus 111. Clus. hist. 1. p. 181.*  
*Hyacinthus botryoides cæruleus amœnus. Lob. ic. 108.*  
*Habitat in Italia.*  
*Differt Floribus inodoris & foliis erectis a sequenti, cui Flores odori & Folia patenti decumbentia carinata.*
11. **HYACINTHUS** corollis ovatis: summis sessilibus, racemosus.  
 foliis laxis. Sauv. monsp. 17.  
*Hyacinthus racemosus cæruleus minor juncifolius Bauh. pin. 43.*  
*Hyacinthus botryoides 1. Clus. hist. 1. p. 181.*  
*Habitat in Europa australi. 4*
12. **HYACINTHUS** corollis sexpartitis: petalis tribus Orchioides.  
 exterioribus brevioribus.  
*Hyacinthus orchioides africanus. major. bifolius maculatus, flore sulphureo obsoleto majore Breyn. prodr. 3 p. 24 t. 11. f. 1, 2, 3.*  
*Orchis angustifolia maculata. Buxb. cent. 3 p. 10. t. 16.*  
*Orchis hyacinthoides, foliis caule & floribus maculatis, Buxb. cent. 3. p. 12. t. 20.*  
*Habitat in Æthiopia. 4*  
*Folia duo, paniculata, ovato-oblonga. Racemus Floribus copiosissimis, violaceis. Corollæ vix ad basin sexpartitæ: Petala exteriora alterna ovata, breviora; interiora, obovata, longiora.*
13. **HYACINTHUS** corollis lanatis, caule ramoso. lanatus.  
*Roy, lugdb. 27.*  
*Habitat - - -*

## A L E T R I S.

- farinosa. 1. **ALETRIS** acaulis, foliis lanceolatis membranaceis, floribus alternis.  
*Aletris. Amæn. acad. 3. p. 11. \**  
*Hyacinthus caule nudo, foliis linguiformibus acuminatis dentatis. Gron. virg. 38.*  
*Hyacinthus floridanus spicatus. Phuk. amalt. 119. t. 437. f. 2.*  
*Habitat in America septentrionali. 21.*
- capensis. 2. **ALETRIS** acaulis foliis lanceolatis undulatis, spica ovata, floribus nutantibus.  
*Habitat ad Cap. b. spei. 21.*
- Hyacinthodes. 3. **ALETRIS** acaulis, foliis lanceolatis carnosius, floribus geminatis.  
*Aloë hyacinthoides floribus sessilibus horizontalibus infundibuliformibus æqualibus limbo revolutis. Sp. pl. 321.*  
*Aloë foliis lanceolatis planis rectis radicalibus. Hort. cliff. 132. Hort. upf. 85. Fl. zeyl. 130.*  
*Katu Kapel. Rheed. mal. 11. p. 83. t. 42 ?*
- Zeylanica. a. *Aloë zeylanica pumila, foliis variegatis. Comm. hort. 2. p. 41. t. 21. Pluk. alm. 19. t. 256. f. 5.*  
*Aloë foliis exterioribus lanceolatis planis erectis radicalibus, interioribus longissimis subulatis arcuatis. Roy. lugdb. 23.*
- guineensis. β. *Aloë guineensis, radice geniculata, foliis e viridi & atro undulatis variegatis. Comm. hort. 2. p. 39. t. 21. præl. 84. t. 33.*  
*Habitat α in Zeylona, β in Guinea. 21.*  
*Flores non vidi; figura corollæ Hyacinthi; etiam poris germinis n. clariferis tribus. antiqua tamen hujus generis videtur; varietas α foliis quibusdam longissimis subulatis compressis, ab altera β foliis lanceolatis omnibus differt, nec tamen specie videtur aliena.*
- fragrans. 4. **ALETRIS** caulescens, foliis lanceolatis laxis.  
*Aloë africana arborecens, floribus albicantibus fragrantissimis. Comm. hort. 2. p. 7. t. 4. 31. p. 93. t. 49.*  
*Habitat in Africa. 5*

## Y U C C A.

- gloriosa. 1. **YUCCA** foliis integerrimis. *Vir. cliff. 29.*  
*Yucca foliis margine integerrimis. Hort. cliff. 130.*  
*Hort. upf. 88. Roy. lugdb. 29.*

Yucca

*Yucca foliis aloës. Bauh. pin. 91.*  
*Yucca indica, foliis aloës. Barr. rar. 70. t. 1194.*  
*Cordylne foliis pungentibus integerrimis. Roy. lugdb. 22.*  
*Habitat in Canada, Peru. §*

3. YUCCA foliis crenulatis strictis.

aliofolia.

*Yucca foliis crenulatis. Vir. cliff. 29.*  
*Yucca foliorum margine crenulato. Hort. cliff. 130. a.*  
*Yucca arborefcens, foliis rigidioribus rectis ferratis.*  
*Dill. clh. 435. t. 323. f. 416.*  
*Aloë, yuccæ foliis, caulefcens. Pluk. alm. 19. t. 256.*  
*f. 4*  
*Aloë americana, yuccæ folio, arborefcens. Comm.*  
*præl. 64. t. 14.*  
*Habitat in Jamaica, Vera Cruce. §*

3. YUCCA foliis crenatis nutantibus.

draconis.

*Yucca foliorum margine crenulato. Hort. cliff. 130. β.*  
*Hort. upf. 88.*  
*Yucca draconis folio ferrato. Dill. clh. 437. t. 324 f.*  
*417. Comm. præl. 42. 67. t. 16.*  
*Draconi arbori affinis americana. Bauh. pin. 506.*  
*Cordylne foliis pungentibus crenatis. Roy. lugdb. 22.*  
*Habitat in America. §*

4. YUCCA foliis serrato-filamentosis.

filamentosa.

*Yucca foliis lanceolatis acuminatis integerrimis margi-*  
*ne filamentosis. Gron. virg. 152. Trew. ehret. t. 37.*  
*Yucca foliis filamentosis. Morif. hift. 2. p. 419.*  
*Yucca virginiana, foliis per marginem apprime filatis.*  
*Pluk. alm. 396.*  
*Habitat in Virginia. §*

A L O E.

1. ALOE floribus pedunculis cernuis corymbosis sub-perfoliata.  
 cylindricis.

*Aloë foliis caulinis dentatis amplexicaulibus vaginan-*  
*tibus. Hort. cliff. 132. Hort. upf. 86. Roy. lugdb. 23.*

a. *Aloë africana caulefcens, foliis magis glaucis caulem*  
*amplectentibus & in mucronem obtufiorem definen-*  
*tibus. Comm. præl. 68. t. 17. rar. 44. t. 44.*

• β. *Aloë africana caulefcens, foliis minus glaucis, dor-*  
*si parte fuprema spinofa. Comm. præl. 69. t. 18.*

γ. *Aloë africana caulefcens, foliis glaucis latioribus &*  
*undique spinofis. Comm. præl. 70. t. 19.*

- δ. Aloë africana caulescens, foliis glaucis brevissimis, foliorum summitate interna & externa nonnihil spinosa. *Comm. præl.* 73. t. 22, rar. 45. t. 45.
- ε. Aloë africana caulescens, foliis glaucis: dorso integro spinoso. *Comm. præl.* 71. t. 20; *hort.* 2. t. 13. *Hall. gart.* 71. \*
- ζ. Aloë africana caulescens, foliis glaucis brevioribus: foliorum parte interna & externa nonnihil spinosa. *Comm. præl.* 71. t. 21.
- Aloë africana caulescens, foliis glaucis caulem amplectentibus. *Comm. hort.* 2. p. 27. t. 14.
- θ. Aloë africana caulescens, foliis spinosis maculis ab utraque parte albicantibus notatis. *Comm. hort.* 2. p. 9. t. 5.
- ι. Aloë africana caulescens perfoliata glauca & non spinosa. *Comm. præl.* 74. t. 23.
- κ. Aloë africana, foliis glaucis: margine & dorsi parte superiore spinosis, flore rubro. *Comm. præl.* 75. t. 24. *hort.* 2. p. 23. t. 12.
- λ. Aloë africana maculata spinosa major. *Dill. elth.* 17. t. 14 f. 15.
- μ. Aloë africana maculata spinosa minor. *Dill. elth.* 18. t. 14 f. 16.
- ν. Aloë africana mitriformis spinosa. *Dill. elth.* 21. t. 17. f. 19.
- ξ. Aloë succotrina angustifolia spinosa, flore purpureo. *Comm. hort.* 1. p. 91. t. 48.
- Aloë americana, ananæ floribus suave rubentibus. *Pluk. alm.* 19. t. 240. f. 4.
- humilis.* c. Aloë foliis erectis subulatis radicatis undique inerme spinosis. *Hort. cliff.* 131. *Roy. lugdb.* 24.
- Aloë africana humilis, spinis inermibus & verrucosis obsita. *Comm. præl.* 77. t. 26. rar. 46. t. 46.
- Habitat in Æthiopia. h*
- Plantam o ad hanc speciem pertinere docuere flores, quamvis habitu videatur aliena.*
- vera.* π. Aloë foliis spinosis confertis dentatis vaginantibus planis maculatis. *Hort. cliff.* 130. *Hort. upf.* 86. *Mat. med.* 160. *Roy. lugdb.* 22. *Brown. jam.* 197.
- Aloë vulgaris. *Bauh. pin.* 386.
- Kadanaku f. Catevala. *Rheed. mal.* 11. p. 7. t. 3.
- Habitat in Indiis, Africa, at π. in Italiæ, Siciliæ, Melitæ testis, muris. h*

A. veræ inflorescentia & Flores omnino A. perfoliatæ ejusque gemina varietas, quamvis Corolla fere minus divisa mihi apparuit.

2. ALOE floribus pedunculatis cernuis racemosis pri- variegata.  
maticis: ore patulo æquali.

Aloë foliis 'canaliculatis trifariam Imbricatis radicatis erectis: angulis cartilagineis ternis. *Hort. cliff.* 132. *Hort. upf.* 87. *Roy. lugdb.* 24.

Aloë africana humilis, foliis ex albo viridi variegatis. *Comm. præl.* 79. t. 28. rar. 27. t. 27. *Till. pisf.* 7. t. 7. *Habitat in Æthiopiæ argillofis.* 24

3. ALOE floribus pedunculatis pendulis ovata-cylindri- disticha.  
cis curvis.

Aloë foliis linguiformibus patulis distichis. *Hort. cliff.* 132. *Hort. upf.* 86. *Roy. lugdb.* 23.

Aloe africana, flore rubro, folio maculis albicantibus ab utraque parte notato. *Comm. hort.* 2. p. 15. t. 8.

4. Aloë africana flore rubro, folio triangulari & verrucis albicantibus ab utraque parte notato. *Comm. hort.* 2. p. 17. t. 9.

Aloë africana sessilis, foliis carinatis verrucosis. *Dill. elth.* 22. t. 18. f. 20. *Mill. dict.* t. 19.

5. Aloë africana, foliis planis conjugatis carinatis verrucosis, caule & flore corallii colore. *Mart. cent.* 11. t. 11.

6. Aloë africana arborefcens montana non spinosa, folio longissimo plicatili, flore rubro. *Comm. hort.* 2. p. 5. t. 3.

*Habitat in Africæ rupibus.* 7.

7. ALOE floribus sessilibus ovatis crenatis: segmentis spiralis.  
interioribus conniventibus.

Aloë foliis ovatis acuminatis caulinis quinquefariam imbricatis. *Hort. cliff.* 132. *Hort. upf.* 87. *Roy. lugdb.* 23.

Aloë africana erecta rotunda, folio parvo & in acumen acutissimum exeunte. *Dill. elth.* 16. t. 13. f. 14. *Comm. præl.* 83. t. 32.

*Habitat in Africæ campestribus.* 7

8. ALOE floribus sessilibus triquetris bilabiatis: labio in- retusa.  
feriore revoluto.

Aloë foliis rhomboidalibus ventricosus quinquefariam pos-

positis radicatis triquetris plano exstantibus. *Hort. cliff.* 132. *Hort. upf.* 87. *Roy. lugdb.* 24.

*Aloë africana*, brevissimo crassissimoque folio, flore viridi. *Comm. hort.* 2. p. 11. t. 6. *Till. pif.* 6. t. 5.

*Habitat in Africæ argillofis.* 24.

viscosa.

6. *ALOE* floribus sessilibus infundibuliformibus bilabiatis: lacinis quinque revolutis: summa erecta.

*Aloë foliis caniculatis trifariam imbricatis: caulinis apice reflexo-patulis.* *Hort. cliff.* 132. *Hort. upf.* 87. *Roy. lugdb.* 23.

*Aloë africana erecta triangularis & triangulari folio viscosa.* *Comm. præl.* 82. t. 31: *Till. pif.* 6. t. 5. *Dill. elth.* 15. t. 13. f. 13.

*Habitat in Æthiopiz campestribus.* 24

pumila.

7. *ALOE* floribus sessilibus bilabiatis: labio superiore erectiore, inferiore recurvato.

*Margaritifera.* a. *Aloë foliis ovato-subulatis acuminatis tuberculis cartilagineis undique adspersis.* *Hort. cliff.* 131. *Roy. lugdb.* 24.

*Aloë africana*, folio in summitate triangulari, margaritifera, flore subviridi. *Comm. hort.* 2. p. 19. t. 10.

*Aloë africana margaritifera*, folio undique verrucis numerosissimis. *Bradl. succ.* 3. p. 1. t. 21.

β. *Aloë africana margaritifera minor.* *Comm. hort.* 2. p. 21. t. 11. *Dill. elth.* 19. t. 16. f. 17.

γ. *Aloë africana margaritifera minima.* *Comm. præl.* 43. *Dill. elth.* 40. t. 16. f. 18.

*Arachnoides.* δ. *Aloë africana humilis arachnoidea.* *Comm. præl.* 78. t. 27.

ε. *Aloë foliis ovato-lanceolatis carnosissimis apice triquetris: angulis inerme dentatis.* *Hort. cliff.* 131. *Hort. upf.* 86. *Roy. lugdb.* 24.

*Aloë africana minima atroviridis*, spinis herbaceis numerosis ornata. *Boerh. lugdb.* 2. p. 131. t. 131. *Bradl. succ.* 3. p. 12. t. 30.

*Habitat in Æthiopiz campestribus.* 24

Flores in hoc genere specierum certissimi indices continent *Margaritifera* & *Arachnoidea*.

Uvaria.

8. *ALOE* floribus sessilibus reflexis imbricatis prismaticis.

*Aloë foliis linearibus radicalibus membranaceis.* *Hort. cliff.* 133. *Roy. lugdb.* 23.

*Aloë*

*Aloë africana*, folio triangulari longissimo & angustissimo, floribus luteis foetidis. *Comm. hort.* 2. p. 29. t. 15. *Seb. thes.* 1. p. 29. t. 19. f. 3.

*Habitat ad Cap. b. Spei.* 24

## AGAVE

1. AGAVE foliis dentato-spinosis, scapo ramofo. *Amæn. americana.* *acad.* 3. p. 22.

*Agave foliis spinoso-dentatis mucronatisque.* *Hort. upf.* 81.

*Aloë foliis lanceolatis dentatis spina terminatis radicalibus.* *Hort. cliff.* 130. *Roy. lugdb.* 22.

*Aloë folio in oblongum mucronem abeunte.* *Bauh. pin.* 286.

*Habitat in America calidiore. Cortusus plantam primum in Europa habuit 1561. ( Cam. hort. 11. ) hodie ab ea sepes in Lusitania.* ¶

2. AGAVE foliis dentatis, staminibus corollam æquan- vivipar-  
tibus.

*Aloë americana polygon.* *Comm. præl.* 65. t. 15.

*Aloë americana.* *Rumph. amb.* 5. p. 273. t. 94.

*Habitat in America.*

*Confer. Aloë americana sobolifera.* *Herm. lugdb.* 16. t. 17.

3. AGAVE foliis dentato-spinosis, scapo simplicissimo. *virginica.* *Amoen. acad.* 3. p. 22.

*Aloë foliis lanceolatis spina cartilaginea terminatis, floribus alternis sessilibus.* *Amæn. acad.* 3. p. 22.

*Habitat in Virginia.* 24

- 4 AGAVE foliis integerrimis. *Amæn. acad.* 3. p. 22. *fatida.*

*Aloë foliis integerrimis patentiusculis aculeo terminatis, radice caulescente.* *Hort. cliff.* 132.

*Aloë americana, viridi rigidissimo & foetido folio, Piet dicta indigenis.* *Comm. hort.* 2. p. 35. t. 18.

*Aloë americana, radice tuberosa, minor.* *Pluk. alm.* 19. t. 258. f. 2.

*Habitat in Curassao.* ¶

## ALSTROEMERIA.

1. ALSTROEMERIA caule erecto. *Amæn. acad.* 6. p. pelcaina. He-



*Hemerocallis* floribus purpurascens maculatis. *Fewill. peruv. 2. p. 711. t. 5.*  
*Habitat in Peru & Lima. 24.*  
*Folia resupinata. Flores pulcherrime variegati.*

- Ligta.** 2. *ALSTROEMERIA* caule adscendente †  
*Hemerocallis* floribus purpurascens striatis. *Fewill. peruv. 2. p. 710. t. 4.*  
*Habitat in Lima.*
- Salifolia.** 3. *ALSTROEMERIA* caule volubili †  
*Hemerocallis* scandens, floribus purpureis. *Fewill. peruv. 2. p. 713. t. 6.*  
*Habitat in Lima.*

## HEMEROCALLIS.

- flava.** 1. *HEMEROCALLIS* corollis flavis.  
*Hemerocallis* scapo ramoso, corollis monopetalis.  
*Hort. upf. 88.*  
*Hemerocallis* radice tuberosa, corollis monopetalis.  
*Hort. cliff. 128. Roy. lugdb. 26.*  
*Hemerocallis* radice tuberosa, corollis monopetalis luteis. *Gmel. sibir. 1. p. 37.*  
*Lillio-Asphodelus* luteo flore. *Cus. hist. 1. p. 137.*  
*Lilium luteum, asphodeli radice. Bauh. pin. 80.*  
*Habitat in Sibiriae, Hungariae campis uliginosis. 24*
- fulva.** 2. *HEMEROCALLIS* corollis fulvis.  
*Lillio-Asphodelus* puniceus. *Cus. hist. 1. p. 137.*  
*Lilium rubrum, asphodeli radice. Bauh. pin. 80.*  
*Habitat in China. 24.*  
*Specie diversa hac esse videtur, nec tamen differentiationem reperi; hac major, illa minor; hac floribus fulvis, illa flavis; hac Maturationis, illa Grossificationis mense floret.*

## ACORUS.

- Calamus.** 1. *ACORUS.* *Roy. lugdb. 6. Fl. suec. 277, 297. Mat. med. 16.*
- vulgaris.** a. *Acorus* verus s. *Calamus aromaticus officinarum.*  
*Bauh. pin. 34. Hort. cliff. 137.*  
*Typha aromatica, clava rugosa. Moris. hist. 3 p. 246.*  
*f. 8. t. 13. f. 4.*

β. *Aco-*

- β. *Acorus asiaticus*, radice tenuiore. *Herm. lugdb. 9. verus.*  
*Fl. zeyl. 132.*  
*Acorum. Rumph. amb. 5. p. 178. t. 72. f. 1.*  
*Waembu. Rheed. mal. 11. p. 99. t. 60.*  
*Habitat in Europæ, in Indiæ fossis paludosis. 2.*

## ORONTIUM.

1. *ORONTIUM. Amæn. acad. 3. p. 17. t. 1. f. 3. \* aquaticum.*  
*Kaln. it. 3. p. 54.*  
*Arum folio enervi ovato. Gron. virg. 113.*  
*Arum fluitans, pene nudo. Banist. virg. 1924. Catefb.*  
*car. 1. p. 82. t. 82.*  
*Habitat in Virginiæ, Canadæ paludibus, scaturigini-*  
*bus. 2.*

## CALAMUS.

1. *CALAMUS. Rotang.*  
*Arundo zeylanica spinosissima major, fructibus rotun-*  
*dis scabris acidis. Burm. zeyl. 36. Fl. zeyl. 468.*  
*Arundo nucifera Rotang dicta, fructu spadicei colo-*  
*ris striis purpureis venuste tessulato. Pluk. alm. 53.*  
*Arundo Rotang dicta. Pif. mant. 188.*  
*Fructus cannæ de Benghala. Bauh. pin. 405.*  
*Tsiere Tsiurel. Rheed. mal. 12. p. 121. t. 64, 65.*  
 α. *Palmjuncus Calapparius. Rumph. amb. 5. p. 88. t. 51.*  
 β. *Palmjuncus niger. Rumph. amb. 5. p. 121. t. 52.*  
 γ. *Palmjuncus albus. Rumph. amb. 5. p. 107. t. 53.*  
 δ. *Palmjuncus verus. Rumph. amb. 5. p. 105. t. 54.*  
 ε. *Palmjuncus viminalis. Rumph. amb. 5. p. 108. t. 55.*  
 ζ. *Palmjuncus equestris. Rumph. amb. 5. p. 110. t. 56. &*  
*t. 57. f. 1.*  
 η. *Palmjuncus zalacca. Rumph. amb. 5. p. 115. t. 57. f. 2.*  
 θ. *Palmjuncus Draco. Rumph. amb. 5. p. 114. t. 58. f. 1.*  
*Habitat in Indiæ sylvis juxta fluentia. 5.*

## JUNCUS.

\* *Culmis nudis.*

1. *JUNCUS culmo subnudo tereti mucronato, panicu-*  
*la terminali, involucrio diphylo spinoso.*  
*Juncus culmo pungente, panicula ex folii solitarii a-*  
*xilla. Sauv. monsp. 9.*  
*Juncus panicula universalis culmo brevior in fine co-*  
*arcta, secundariis paululum laxis, capsulis triquetre-*  
*subovatis nitentibus. Guett. stamp. 2. p. 412.*

Jun-

- Juncus acutus*, capitulis forghi. *Bauh. hist.* 2. p. 520.  
*Morif. hist.* 3. p. 232. f. 8. t. 10. f. 15.
- β. *Juncus* panicula universali culmo longiore, secundariis laxis longe pedunculatis: media præcipue. *Guett. stamp.* 2. p. 412.  
*Juncus acutus maritimus anglicus.* *Raj. hist.* 1303.  
*Morif. ibid. f.* 14.  
*Habitat in Angliæ, Galliæ, Italiæ maritimis paludosis.* 2
- conglomeratus. 2. *JUNCUS* culmo nudo stricto, capitulo laterali. *Roy. lugdb.* 44. *Fl. fvec.* 278, 298.  
*Juncus lævis*, panicula non sparsa. *Bauh. pin.* 12. *theatr.* 183.  
*Juncus lævis*, panicula conglomerata. *Scheuch. gram.* 343.  
*Juncus.* *Cam. epit.* 780.  
*Habitat in Europæ borealis uliginosis.* 2
- effusus. 3. *JUNCUS* culmo nudo stricto, panicula laterali. *Roy. lugdb.* 44. *Fl. fvec.* 279, 299.  
 a. *Juncus* culmo nudo acuminato ad basin squamato, floribus sessilibus. *Fl. lapp.* 116. *Hall. herb.* 254.  
*Juncoides alpinum*, flosculis junci glomeratis atrofuscis. *Scheuch. hist.* 323.  
 β. *Juncus* culmo nudo acuminato ad basin squamato, floribus pedunculatis. *Fl. lapp.* 117. *Gron. virg.* 38.  
*Juncus lævis*, panicula sparsa, major. *Bauh. pin.* 12. *theatr.* 182. *Scheuch. gram.* 341.  
*Juncus lævis.* *Dod. pempt.* 606.  
*Habitat in Europæ uliginosis, a in alpibus.* 2
- inflexus. 4. *JUNCUS* culmo nudo: apice membranaceo incurvo panicula laterali.  
*Juncus* culmo nudo, paniculam arcuatim tegente. *Sauv. monsp.* 9.  
*Juncus*, acumine reflexo, major. *Bauh. pin.* 12. *prodr.* 184. *Morif. hist.* 3. p. 233. f. 8. t. 10. f. 25. *Barr. ic.* 204. *Hall. herb.* 252.  
 β. *Juncus*, acumine reflexo, alter. *Bauh. pin.* 12. *prodr.* 22. *Scheuch. gram.* 345.  
 γ. *Juncus* panicula universali laterali culmo longiore, secundariis sparsis. *Guett. stamp.* 412.  
*Juncus acutus* panicula sparsa. *Bauh. pin.* 11.  
*Habitat in Europæ australi.* 2

5. **JUNCUS** culmo nudo filiformi nutante, panicula filiformis. laterali. *Roy. lugdb.* 44. *Gron. virg.* 152. *Fl. suec.* 280, 300.  
*Juncus parvus*, calamo supra paniculam longius producto. *Raj. angl.* 3. p. 432. *Fl. lapp.* 118.  
*Juncus lævis*, panicula spatiosa, minor. *Bauh. pin.* 12. *theatr.* 183. *Scheuch. gram.* 347. t. 7. f. 1.  
*Habitat in Europæ uliginojo-paludosis turfosis.* 24
6. **JUNCUS** culmo nudo, foliis floribusque tribus trifidus. minalibus *Fl. lapp.* 119. *Fl. suec.* 281, 301. *Roy. lugdb.* 43. *Hall. helv.* 253.  
*Juncus*, acumine reflexo, minor & trifidus. *Bauh. pin.* 12. *prodr.* 22. t. 22. *theatr.* 186  
*Juncoides alpinum trifidum.* *Scheuch. gram.* 325.  
*Habitat in Alpibus Lapponicis, Helveticis, Pyrenæis.* 24
7. **JUNCUS** culmo nudo, foliis setaceis, capitulis glomeratis aphyllis. *Fl. suec.* 282, 302. squarrosus.  
*Juncus foliis setaceis, culmo nudo, capitulis glomeratis terminalibus.* *Roy. lugdb.* 44.  
*Juncus montanus palustris.* *Raj. hist.* 1303. *Fl. lapp.* 121.  
*Gramen junceum, foliis & spica junci.* *Bauh. pin.* 5.  
*Gramen junceum, semine acuminato.* *Læf. pruss.* 15. t. 29.  
*Habitat in Europæ borealis cespitosis.* 24
- \* *Culmis foliosis.*
8. **JUNCUS** foliis nodoso-articulatis, petalis obtusis. articulatus. *Fl. suec.* 285, 303.  
*Juncus foliis articulosis, panicula inæquali, capitulis sessilibus, capsulis subulatis triquetris.* *Roy. lugdb.* 43.  
*Juncus foliis articulosis, floribus umbellatis.* *Tournef. inst.* 247. *Fl. lapp.* 120.  
*Gramen junceum, folio articulato, aquaticum.* *Bauh. pin.* 5. *prodr.* 12. *theatr.* 76. *Scheuch. gram.* 331.  
 β. *Gramen junceum, folio articulato, cum utriculis.* *Bauh. prodr.* 12. *Scheuch. gram.* 333.  
 γ. *Juncus alpinus, folio articulato.* *Scheuch. gram.* 333.  
*Gramen junceum, folio articulato, sylvaticum.* *Bauh. pin.* 5. *theatr.* 75. *Mori: hist.* 3. p. 227. f. 8. t. 9. f. 1. *Scheuch. gram.* 334.
- Linnaei Spec. plant. pars I.* Gg      δ. Jun-

8. Junci foliosi, foliis per ficitatem nodosis. *Mich. gen.*  
38. n. 1. - 13.

*Habitat in Europæ aquosis. 2*

nodosus. 9. JUNCUS foliis nodoso-articulatis, petalis mucronatis.

Juncus foliosus minor, pericarpis ovatis, americanum.  
*Gron. virg.* 15.

Gramen junceum elatius, pericarpis ovatis. *Pluk. alm.*  
179. t. 92. f. 9.

Gramen junceum virginianum, calyculis paleaceis, hincorne. *Morij. hist.* 3. p. 228. f. 8. t. 9. f. 15.

*Habitat in America septentrionali. 2*

Simillimus J. articulato, sed Flores magis capitati: capitulis rarioribus, majoribus & panicule rami simplices, nec supradecompositi; imprimis distinguitur petalis in acumen subulatum protrahitis.

bulbosus. 10. JUNCUS foliis linearibus canaliculatis, capsulis obtusis. *Fl. fvec.* 284, 304. *Mees. frif.* 160.

Juncus foliis linearibus canaliculato concavis, capsulis triangularibus. *Guett. stamp.* 134.

Juncus foliis angulatis, culmo subnudo, panicula terminali prolifera, capsulis ovatis. *Roy. lugdh.* 43.

Juncus parvus cum pericarpis rotundis. *Bauh. hist.* 2. p. 522.

Juncus repens apocarpus minor. *Barr. ic.* 114.

Gramen junceum, foliis & spica junci. *Bauh. pin.* 5. theatr. 78. *Scheuch, gram.* 322.

Gramen junceum, junci sparsa panicula. *Morij. hist.* 3. p. 227. f. 8. t. 9. f. 11.

*Habitat in Europæ pascuis sterilibus & ad vias. 2*

Radix repens, densa. Culmi filiformes, subcompressi.

Folia linearia, levia, subtus convexa & striata, supra canaliculata. Folium florale longitudine corymbi. Corymbus communis & partiales inaequales. Capsulae ovatae, fuscae, nitidae.

bulbosus. 11. JUNCUS culmo dichotomo, foliis angulatis, floribus solitariis sessilibus. *Roy. lugdh.* 43 *Fl. fvec.* 283. 305 *Hall. helv.* 254. *Gert. Ligr.* 52.

Juncus palustris humilior, panicula laxissima. *Rupp. jen.* 130. *Fl. Lapp.* 122.

- Gramen nemorosum, calyculis paleaceis, erectum.  
*Scheuch. gram.* 327.
- Gramen nemorosum, calyculis paleaceis. *Bauh. pin.*  
*7. theatr.* 100.
- Holosteum. *Lob. ic.* 18.
- β. Gramen nemorosum, calyculis paleaceis, repens  
*Scheuch. gram.* 329.
- γ. Gramen nemorosum, calyculis paleaceis, species  
 altera procerior. *Scheuch. gram.* 328.
- ζ. Gramen bufonium erectum angustifolium majus. *Barr.*  
*ic.* 264.
- ε. Gramen holosteum alpinum minimum. *Bauh. pin.* 7.  
*prodr.* 64. *theatr.* 101.
- Gramen juncoides minimum anglo-britannicum. *Raj.*  
*angl.* 3. p. 434.
- Habitat in Europæ inundatis.* ☉
- Pinheaded cobweb mushroom. *Pet. gaz.* 51. f. 7. *hujus*  
*infans, paucis lecta.*
12. JUNCUS foliis setaceis depressiusculis, pedunculis *stygius.*  
 geminis terminalibus, glumis solitariis subbifloris.  
*Habitat in Sveciæ paludibus, cæspitosis sylvaticis pro-*  
*fundis.* 24.
- Culmi *spithamei, simplices.* Folia lineari-subulata,  
 altitudine culmi, erecta, depressa, levia, tria: 2. ad  
 radicem. β. in medio culmi. γ. culmum instar spathæ  
 terminans. Pedunculi duo, terminales: Gluma biflo-  
 ra flosculis duobus, sessilibus, gluma longioribus. Pe-  
 tala exteriora angustiora, acutiora, concava, extus  
 purpurascens; Interiora pallida. Stamina filiformia.  
 Antheræ flavæ. Germen majusculum. Stylus albus.  
 Stigmata tria, reflexa.
13. JUNCUS folio subulato, gluma biflora terminali. biglumis.  
*Amæn. acad.* 2. p. 266. t. 3. f. 3. \*. *Fl. svec.* 2. n.  
 307.
- Habitat in Alpibus Lapponicis.* *D. Montin.* 24.
- Planta faciem gerit Schæni ferruginei.*
14. JUNCUS foliis planis, gluma triflora terminali. triglumis.  
*Fl. Svec.* 286; 306.
- Juncus gluma triflora culmum terminans. *Fl. lapp.* 115.  
 t. 10. f. 5.

- Juncus exiguus montanus mucrone carens.* *Bauh. pin.* 12. *prodr.* 22. *theatr.* 183. *Rudb. elys.* 1. p. 103. f. 8. *Habitat frequens in Alpibus Lapponicis, Tauro Rastadiensi.* 24
- Pilosus.** 15. *JUNCUS* foliis planis pilosis, corymbo ramoso. *Fl. Suec.* 287. 308.
- Juncus foliis planis, panicula diffusa, pedunculis proliferis & flosculis intermediis sessilibus.* *Roy. lugdb.* 43.
- Gramen nemorosum hirsutum latifolium majus.* *Bauh. pin.* 7. *Morif. hist.* 3. p. 225. f. 8. t. 9. f. 1.
- β. *Gramen hirsutum angustifolium perenne, lini utriculis.* *Larr. ic.* 740.
- γ. *Juncus foliis planis, culmo paniculato, floribus sparsis.* *Fl. lapp.* 124.
- Gramen hirsutum nemorosum angustifolium alpinum, paniculis obscure rufescentibus.* *Scheuch. gram.* 312. t. 6.
- δ. *Gramen nemorosum hirsutum latifolium majus.* *Scheuch. gram.* 317.
- Gramen hirsutum latifolium maximum, juncea panicula.* *Morif. hist.* 3. p. 225. f. 8. t. 9. f. 2.
- ε. *Gramen hirsutum angustifolium majus.* *Bauh. pin.* 7. *prodr.* 16. *theatr.* 105. *Morif. hist.* 3. p. 225. f. 8. t. 9. f. 3. *Scheuch. gram.* 318.
- ζ. *Gramen hirsutum latifolium minus.* *Bauh. theatr.* 102. *Habitat in Europæ sylvis, at varietates β. δ. ε. ζ. in Europæ australioribus; γ. in Alpibus.* 24
- niveus.** 16. *JUNCUS* foliis planis subpilosis, corymbis folio brevioribus, floribus fasciculatis.
- Juncus foliis planis angustioribus, panicula umbellata foliis brevioribus, floribus longissimis albis.* *Hall. helv.* 237.
- Juncoides montanum nemorosum, flore niveo.* *Scheuch. gram.* 320.
- Gramen hirsutum angustifolium minus, paniculis albis.* *Bauh. pin.* 7. *theatr.* 106. *Morif. hist.* 3. f. 8. t. 9. f. 39.
- Habitat in Alpibus Bohemicis, Helveticis, Rhæticis, Monspelii.* 24
- campestris.** 17. *JUNCUS* foliis planis subpilosis, spicis sessilibus pedunculatisque. *Fl. Suec.* 288, 309.

- Juncus foliis planis*, panicula rara, spicis sessilibus pendunculatisque. *Hort. cliff.* 137. *Roy. lugdb.* 42.  
*Gron. virg.* 38.
- Juncus villosus*, capitulis psyllii. *Tournef. inst.* 246.  
*Fl. lapp.* 126.
- Juncoides villosus*, capitulis psyllii. *Scheuch. gram.* 310.
- Gramen hirsutum*, capitulis psyllii. *Bauh. pin.* 7. *theatr.* 103.
- β. *Juncus foliis planis*, culmo paniculato, spicis ovatis.  
*Fl. lapp.* 127. t. 10. f. 2.
- γ. *Juncoides hirsutum*, capitulo glomerato. *Scheuch. gram.* 312.
- Gramen hirsutum*, capitulo globofo. *Bauh. pin.* 7. *theatr.* 104.
- δ. *Juncus montanum nemorosum villosum latifolium elatius*, capitulis florum candidis & splendentibus.  
*Mich. gen.* 42.
- ε. *Juncoides latifolium alpinum glabrum*, panicula sublutea splendente. *Scheuch. gram.* 314.
- ζ. *Gramen hirsutum elatius*, panicula juncea compacta.  
*Raj. angl.* 3. p. 416.
- η. *Juncoides bohemicum*, panicula minore nigricante, scapo super eandem erecto & longius producto,  
*Mich. gen.* 42.
- Habitat in Europæ pascuis siccioribus.* 21.
18. **JUNCUS** foliis planis, spica racemosa nutante. spicatus.  
*Fl. lapp.* 125. t. 10. f. 4. *Fl. succ.* 289, 310.
- Juncus alpinus latifolius*, panicula racemosa nigricante pendula. *Till. pis.* 91.
- Habitat in Lapponiæ Alpibus.* 24

ACHRAS.

1. **ACHRAS** floribus solitariis, foliis cuneiformi-lanceolatis. *mammosa.*  
*Achras. Læfl. it.* 186?
- Sapota*, fructu ovato majore. *Plum. gen.* 43.
- Sideroxylon inerme*, calycibus decaphyllis. *Jacq. amer.* 15.
- Malus Persica maxima*, foliis magnis integris longis, fructu maximo oblongo scabro: ossiculo partim rugoso partim glabro. *Sloan. jam.* 2. p. 124. t. 218.
- Arbor americana pomifera*, frondosis ramulis, foliis



amplis longioribus obtusis duris & venosis margine æquali. *Pluk. alm.* 39. t. 268. f. 2.

Fructus oblongus utrinque acuminatus s. conicus lævis splendens spadiceus. *Kaj. hst.* 1800.

β. *Achras* fructu elliptico scabro majori, floribus solitariis alaribus. *Brown. jam* 200. t. 19. f. 3?

*Habitat in America meridionali.* ‡

*Hujus Flores mihi non visi.*

Sapota. 2. *ACHRAS* floribus solitariis, foliis ovato-oblongis.

*Achras* brachiatus diffusus, fructu subrotundo cicatricula mucrole brevi. *Brown. jam* 200.

*Anona* foliis laurinis glabris viridi-fuscis, fructu minore. *Sloan. jam.* 206. *hist.* 2. p. 171. t. 230. *Kaj. dendr.* 78. *Catesb. car.* 2. p. 87. t. 87.

β. *Achras* caudice altissimo, fructu minori, semine mucronato. *Brown. jam.* 201.

Sapota fructu turbinato minori. *Plum. gen.* 43.

*Anona* maxima, foliis laurinis glabris viridi-fuscis, fructu minimo. *Sloan. jam.* 206. *hist.* 2. p. 172. t. 169. f. 2. *Kaj. dendr.* 79.

*Habitat in America meridionali.* ‡

salicifolia. 3. *ACHRAS* floribus confertis, foliis lanceolato-ovatis.

*Achras* foliis oblongis nitidis utrinque productis, floribus confertis, fasciculis infra frondes sparsis. *Brown. jam.* 201. t. 17. f. 4.

*Salicis* folio lato splendente arbor, floribus parvis palide luteis peptapetalis e ramulorum lateribus confertim exeuntibus. *Sloan. jam.* 170. *hist.* 2. p. 98. t. 206. f. 2.

*Habitat in America meridionali.* ‡

## RICHARDIA.

scabra. 1. *RICHARDIA.* *Houft. mss.*

*Habitat in Vera Cruce.* 24

*Planta ex stellatis XLIV.* Caulis altus, brachiatus, subarticulatus, obsolete tetragonus, hispidus: Filis rigidiusculis, sparsis, reflexis. Folia lanceolato-ovata, alternatim nervosa, integerrima, subpetiolata, scabra. Flores in capitula terminalia, passim etiam in verticillos congesti; capitula terminalia, radiata. Foliis 4 vel pluribus, scilicet alternis minoribus. Corollulæ ut in *Asperulis*, tenues, longæ, sed sex-fide

*fide* ♂ fructus ex 3 seminibus; sic differt a Sperma-  
toce numero & ceteris.

## BURSER A.

### 1. BURSER A. Jacq. hist. t. 65.

gummifera.

Terebinthus. *Brownii*. Jacq. amer. 18.

Terebinthus foliis cordato-ovatis pinnatis, cortice læ-  
vi rufescente, floribus masculis spicatis. *Brown. jam.*  
345.

Terebinthus major, betulæ cortice, fructu triangulari.

*Sloan. jam.* 2. p. 89. t. 199. *Catesb. car.* 1. p. 30. t. 30.

Pistacia foliis pinnatis deciduis, foliolis ovatis. *Sp. pl.*

1. p. 1026.

Simaruba. *Mat. med.* 535.

*Habitat in America calidiorē. ♀*

## PRINOS.

### 1. PRINOS foliis longitudinaliter serratis.

verucillatus

Prinos. *Gron. virg.* 39. \*

Alcanna major latifolia dentata. *Munt. phyt.* 213. t. 51.

Aquifolium foliis deciduis. *Duham. arb.* 1. p. 62. t. 23.

*Habitat in Virginie paludibus. ♀*

### 2. PRINOS foliis apice serratis.

glaber.

Cassine foliis lanceolatis alternis sempervirentibus, flo-  
ribus axillaribus. *Mill. dict.* t. 83. f. 2.

Cassine vera floridanorum arbuscula baccifera, alater-  
ni ferme facie, foliis alternatim sitis, tetrapylene.

*Catesb. car.* 2. p. 57. t. 57.

*Habitat in Canada. Kalm. ♀*

*Arbor. Foliis alternis, petiolatis, oblongis, consisten-  
tibus, glabris, versus apicem serraturis utrinque sæpius  
biniis, acutis, facie ferme Myricæ. Pedunculi axil-  
lares, parvi, sæpius triflori. Flores non vidi.*

## BERBERIS.

### 1. BERBERIS pedunculis racemosis. *Mat. med.* 290. vulgaris.

*Gron. orient.* 120.

Berberis spinis triplicibus. *Hort. cliff.* 122. *Fl. suec.*

290, 311. *Roy. lugdb.* 438. *Hall. helv.* 424.

Berberis dumetorum. *Bauh. pin.* 454.

Berberis vulgaris. f. *Crespinus. Cam. epit.* 86.

*Habitat in Europæ sylvis, Oriente, Libano. ♀*

Folia prima obovata, serrato-ciliata, inarticulata: stipulae utrinque dente capillari terminatae: Caulina alterna: horum infimum subpinnatifidum dentibus spinosis; reliqua mutata in spinas tripartitas. Folia secundaria bina, oblonga, serrata, inter folia infima & spinas intra se occultant minora folia. Ergo Foliis, anni praesentis, mutatis in spinas, insequentis anticipantia prodibit. An simile in ulla alia?

cretica. 2. BERBERIS pedunculis unifloris.

Berberis cretica, buxi folio. *Tournef. cor.* 42.

Berberis alpina cretica. *Baub. pin.* 454.

Lycium creticum. *Alp. exot.* 21. t. 20:

Lycium e Candia. *Pon. ital.* 137.

Habitat in Creta.  $\S$

## LORANTHUS.

Scurrula. 1. LORANTHUS pedunculis unifloris congestis, foliis obovatis.

Scurrula. *Sp. pl.* 110.

Viscum vitice innascens. *Camell. lux.* 3. n. 36. *Pet. gaz.* t. 63 f. 8.

Habitat in China.  $\S$

Frutex parasiticus. Folia opposita, petiolata, ovata, integerrima, glabra, altero latere paulo latiora. Flores axillares, 3—6, propriis pedicellis. Germina inferne attenuata.

americanus, 2. LORANTHUS racemis subramosis aequalibus, foliis ovatis. *Amæn. acad.* 5. p. 396. \*

Loranthus americanus. *Sp. pl.* 1. p. 331. *Jacq. hist.* t. 67.

Loranthus racemis, flore coccineo, baccis nigris. *Vaill. act.* 1722. p. 274.

Lonicera flore coccineo, baccis nigris. *Plum. gen.* 17. ic. 166. f. 1.

Scurrula parasitica, foliis ovatis oppositis; racemis rarioribus alaribus. *Brown. jam.* 197.

Habitat in Americæ arboribus.  $\S$

occidentalis 3. LORANTHUS racemis simplicibus, floribus irregularibus. *Læfl. it.* 187. \* *Amæn. acad.* 5. p. 396. n. 38. \*

Scurrula parasitica, foliis majoribus subrotundis, spicis florum simplicibus alaribus. *Brown. jam.* 197.

*Viscum latioribus & subrotundis foliis, flore purpureo.*  
*Sloan. jam. 168. hist. 2. p. 92. t. 200. f. 2. Raj.*  
*dendr. 52?*

*Habitat in Americæ arboribus. ‡*

4. LORANTHUS floribus aggregato-capitatis.

Loniceriol-  
des.

*Lonicera pedunculis multifloris, involucris pentaphyl-  
 lis, foliis ovato-lanceolatis petiolatis. Fl. zeyl. 83.\**

*Sp. pl. 1. p. 1. 175.*

*Periclymenum surrectum, persicæ foliis, maderaspatanum.*  
*Pluk. alm. 287. t. 212. f. 5.*

*Itti Canni. Rheed. mal. 7. p. -- t. 27.*

*Habitat in Asiæ arboribus. ‡*

*Flores huic sæpius pentandri.*

5. LORANTHUS racemis trichotomis, pedunculis tri-  
 gonis, floribus æqualibus. *Læfl. it. 187.*

*Habitat in Cumanæ arboribus. ‡*

FRANKENIA.

1. FRANKENIA foliis linearibus basi ciliatis.

lævis.

*Frankenia foliis confertis. Roy. lugdh. 452.*

*Frankenia foliis aciformibus congestis. Sauv. monsp. 46.*

*Franca floribus solitariis sessilibus, foliis prismaticis  
 triangularibus Guett. flamp. 2. p. 459.*

*Franca maritima supina saxatilis glauca ericoides sem-  
 pervirens, flore purpureo. Mich. gen. 23. t. 22. f. 1.*

*Lychnis supina maritima, ericæ facie. Raj. angl. 3.  
 p. 38.*

*Anthyllis repens italica, thymi foliis, rubente flore,  
 polygoni facie Till. pis. 46.*

*Calli f. Vermiculari marinæ non dissimilis planta. Bauh.  
 hist. 3. p. 703.*

*Polygonum fruticosum supinum ericoides cinereum,  
 thymi folio, hispanicum. Barr. ic. 714. Bocc. mus. 1.*

*p. 7. t. 11.*

*Polygonum maritimum minus, foliis serpylli. Bauh.  
 p. n. 281.*

*Habitat in Europæ australis maritimis. 24*

2. FRANKENIA caulibus hirsutis, floribus fascicula-  
 tis terminalibus.

*Franca maritima supina multiflora candida, caulibus  
 hirsutis, foliis quasi vermiculatis. Mich. gen. 23. t.*

*22. f. 2.*

*Polygonum creticum*, thymi folio. *Bauh. pin.* 281  
*prodr.* 131.

*Alfina cretica maritima supina*, caule hirsuto, foliis  
quasi vermiculatis. *Tournef. cor.* 45.

*Habitat in Apulia, Creta.*

pulvera-  
lenta,

3. **FRANKENIA** foliis obovatis retusis subtus pulve-  
ratis.

*Frankenia* foliis ovalibus. *Sauv. monsp.* 168.

*Franca maritima quadrifolia supina*, chamæfyces folio  
& facie. *Mich. gen.* 23.

*Anthyllis valentina*. *Clus. hist.* 2. p. 186.

*Anthyllis marina*, chamæfyces similis. *Bauh. pin.* 282.

*Habitat in Sufexiæ, Narbonæ, Italiæ, Apuliæ lito-  
ribus*, ☉

*In ordine Naturali ad XL. referatur.*

## PEPLIS.

Portula

1. **PEPLIS** floribus apetalis. *Fl. lapp.* 128. *Fl. succ.* 291,  
312. *Hall. helv.* 406.

*Glaucoides palustre*, portulacæ folio, flore purpureo.  
*Mich. gen.* 21. t. 18.

*Glaux palustris*, flore striato clauso, foliis portulacæ.  
*T. Petit. gen.* 43. t. 43.

*Glaux aquatica*, folio subrotundo. *Læf. pruf.* 106.  
t. 20.

*Glaux altera*, subrotundo folio. *Vaill. parif.* 80. t. 13.  
f. 5.

*Alfina palustris minor serpillifolia*. *Bauh. pin.* 251.

*Habitat in Europæ inundatis*. ☉

- retandra. 2. **PEPLIS** floribus tetrandris monopetalis. *Jacq. amer.*  
18. t. 180. f. 29. *Amæn. acad.* 5. p. 412. \*

*Amannia hirta*, floribus parvis orbiculatis, floribus  
singularibus ad alas. *Brown. jam.* 145.

*Habitat in Jamaica*. ☉

## D I G Y N I A.

### VELEZIA,

- rigida. 1. **VELEZIA**. *Læf.*

*Silene* foliis subulatis cauli appressis, calycibus rigi-  
dis intermedio longioribus. *Sauv. monsp.* 145.

Lych-

*Lychnis sylvestris minima*, exiguo flore. *Bauh. pin.* 206. *prodr.* 103.

*Lychnis minima rigida cherleri.* *Bauh. hist.* 3. p. 352. *Raj. hist.* 997.

*Lychnis corniculata minor.* *Barr. rar.* 665. t. 1018. 1017. *Bocc. mus.* 2. p. 50. t. 43.

*Habitat in Europa australi.* ☉

*Stamina tenuissima difficillime numerantur; in siccis specimenibus 5 vidi; at in vivis circiter 6. Læfing. Si hujus ordo naturalis sit n. XL, numerus naturalis erit flaminum sex, si vero XLII, erit quinque.*

## ORYZA.

1. *ORYZA.* *Bauh. pin.* 24. *theatr.* 479. *Cam. epit.* 192. *sativa.*

*Dod. pempt.* 509. *Catesb. car.* 1. p. 14. t. 14. *Hort.*

*cliff.* 137. *Mat. med.* 174. *Roy. lugdb.* 58.

*Habitat forte in Æthiopia, colitur in Indiæ paludosis.* ☉

## ATRAPHAXIS.

1. *ATRAPHAXIS ramis spinosis.* *Hort. cliff.* 138. *Roy. spinosa.* *lugdb.* 409.

*Atriplex orientalis*, frutex aculeatus, flore pulchro.

*Tournesf. cor.* 38. *Buxb. cent.* 1. p. 19. t. 30. *Dill.*

*elth.* 47. t. 40. f. 47.

*Habitat in Media ad urbem Hansen locis glareosis. juxta fluvios.* b

2. *ATRAPHAXIS inermis.*

*Atraphaxis inermis*, foliis undulatis. *Hort. cliff.* 137.

*Roy. lugdb.* 409.

*Arbuscula africana repens*, folio ad latera crispo, ad polygonam relata. *Dill. elth.* 36. t. 32. f. 36.

*Habitat in Æthiopia?* b

*undulata.*

## TRIGYNIA.

### FLAGELLARIA.

1. *FLAGELLARIA.* *Fl. zeyl.* 133. \*. *Amœn. acad.* 1. indica. p. 398. *Osborn. it.* 276.

*Palmjuncus lævis.* *Rumph. amb.* 5. p. 120. t. 59. f. 2.

*Panambu-valli.* *Rheed. mal.* 7. p. 99. t. 53. *Raj. suppl.* 573.

*Ha-*

*Habitat in Java, Malabaria, Zeylona. 6*  
*Frutex humanam altitudinem superans. Rami distichi.*  
*Affinitas proxima Calami*

## RUMEX.

\* *Hermaphroditi: valvulis floris granulo notatis.*

- Patientia.** 1. **RUMEX** floribus hermaphroditis: valvulis integerrimis: unica granifera, foliis cordatis. *Mat. med.* 178.  
*Lapathum hortense, folio oblongo. Bauh. pin. 114.*  
*Lapathum sativum. Dod. pempt. 648.*  
*Habitat in Italia. 24*  
*Confer. Lapathum hortense latifolium. Bauh. pin. 115.*
- sanguineus.** 2. **RUMEX** floribus hermaphroditis: valvulis integerrimis: unica granifera, foliis cordato-lanceolatis. *Hort. cliff. 138. Hort. upf. 89. Mat. med. 528. Roy. lugdb. 229.*  
*Lapathum folio acuto rubente. Bauh. pin. 115.*  
*Lapathum rubens. Dod. pempt. 650. Cam. epit. 229.*  
*Habitat in Virginia. 3*  
*Valvula exterior floris grano magno globofo rubro notata, rarius 3 altera grano minimo.*
- crispus.** 3. **RUMEX** floribus hermaphroditis: valvulis integris graniferis, foliis lanceolatis undulatis acutis.  
**Rumex** floribus hermaphroditis: valvulis integerrimis planis, foliis cordato-oblongis. *Hort. cliff. 138. Fl. succ. 294, 314. Roy. lugdb. 229.*  
**Rumex** foliis cordato-oblongis acuminatis integris. *Fl. lapp. 129.*  
*Lapathum folio acuto crispo. Bauh. pin. 115.*  
*Lapathum acutum crispum. Tabern. ic. 436.*  
*Habitat in Europæ humosis. 24*
- verticillatus** 4. **RUMEX** floribus hermaphroditis: valvulis integerrimis: omnibus graniferis, foliis lanceolatis: vaginis cylindricis.  
*Lapathum aquaticum, foliis longis angustis acutis, floribus ad genicula verticillatim congestis. Gron. virg. 39.*  
*Habitat in Virginia. 24*
- Britannica.** 5. **RUMEX** floribus hermaphroditis: valvulis integerrimis:

mis: omnibus graniferis, foliis lanceolatis: vaginis obsoletis. *Mat. med.* 17.

*Rumex aquatica*, calycis foliolis omnibus æqualibus, radice exterius nigra vel flava. *Cold. novch.* 83.

*Lapathum* foliis longis latis vix acuminatis costis caulibusque rubentibus, radice intus crocea. *Gron. virg.* 39.

*Habitat in Virginia.* 24

*In præcedenti stipula cylindrica membranacea fere ad dimidium vaginans internodium, in hac vero non item, sed ut in Europæis. Pedicelli in priori crassiores, in hac capillares; prior magis, spicata, hæc magis paniculata. Plantam Gron. in Fl. virginica habui a Cl. Authore, quæ non rubra erat caule aut costis.*

6. **RUMEX** floribus hermaphroditis: valvulis dentatis: pulcher. subunica granifera, foliis radicalibus panduriformibus.

*Rumex* floribus hermaphroditis, foliis lyratis. *Guett. stamp.* 1. p. 7.

*Lapathum* ramis procumbentibus, seminis involucrio dentato, foliis inferioribus instar fidium. *Morif. hist.* 2. p. 580.

*Lapathum pulchrum bononiense sinuatum. Bauh. hist.* 2 p. 988. *Raj. hist.* 174.

*Habitat in Anglia, Gallia, Italia.* 24

*Folia Caulina lateribus non sinuata, ovato oblonga, integra. Rami floriferi recurvati. Calycis valvule dentatæ, apice obtusæ; harum exterior granulum gerit; reliquæ vero granulum obsoletum habent.*

7. **RUMEX** floribus hermaphroditis: valvulis trifido-ægyptiacus. setaceis: unica granifera. *Hort. ups.* 89.

*Rumex* floribus hermaphroditis, foliis oblongis, valvulis dentato-setaceis longissimis. *Wach. ult.* 391.

*Lapathum ægypticum, capsula seminis alba & crenata. Dill. ult.* 191. t. 158 f. 191.

*Lapathum ægyptiacum annuum, parietariæ foliis, capsula seminis longius barbata. Till. pij.* 43. t. 37. f. 1.

*Habitat in Ægypto.* ☉

8. **RUMEX** floribus hermaphroditis: valvulis dentatis: perficariol-  
omnibus graniferis, foliis lanceolatis. des.

*Habitat in Virginia.* ☉

Planta



*Planta spithamea, ramosissima. Folia lanceolata, petiolata, lævia, undata, integræ: Floris valvulæ dentibus utrinque tribus longis, omnes teçæ granis pallidis magnis.*

- maritimus. 9. RUMEX floribus hermaphroditis: valvulis dentatis graniferis, foliis linearibus. *Lech. scan. 26. It. scan. 248. Fl. svec. 2. n. 313.*

*Lapathum aquaticum, luteolæ folio. Tournef. inst 504.*

*Lapathum aquaticum, angustissimo acuminato folio.*

*Bocc. mus. 2. p. 142. t. 104.*

*Anthoxanthon. Bauh. hist. 2. p. 987.*

*Habitat in Europæ littoribus maritimis.*

- divaricatus. 10. RUMEX floribus hermaphroditis: valvulis dentatis graniferis, foliis cordato oblongis obtusis pubescentibus.

*Lapathum arvense subhirsutum, folio retuso, caule longius brachiato, capsula seminis crenata. Till. pif. 93. t. 37. f. 2.*

*Habitat in Italiâ. ☉*

- acutus. 11. RUMEX floribus hermaphroditis: valvulis dentatis graniferis, foliis cordato-oblongis acuminatis. *Hort. cliff. 138. Mat. med. 176. Fl. svec. 293, 316. Roy. lugdb. 229.*

*Lapathum acutum f. Oxylapathum. Bauh. hist. 2. p. 93. Lob. ic. 284.*

*Lapathum acutum. Raj. hist. 175.*

*Habitat in Europæ succulentis. 24*

- obtusifolius. 12. RUMEX floribus hermaphroditis: valvulis dentatis graniferis, foliis cordato-oblongis obtusiusculis crenulatis.

*Lapathum involucro seminis dentato, foliis obtusis ora crispula. Hall. helv. 172.*

*Lapathum sylvestre, folio subrotundo, seminis involucris dentatis. Moris. hist. 2. p. 580.*

*Lapathum folio minus acuto. Bauh. pin. 111.*

*Lapathum vulgare, folio obtuso. l. B. Raj. hist. 175.*

*Lapathum. Cam. epit. 228.*

*Habitat in Sudermania, Germania, Helvetia, Gallia, Anglia. 24*

\* \* *Hermaphroditi: valvulis floris grano destitutis.*

13. RUMEX floribus hermaphroditis: valvulis dentatis nudis, pedicellis planis reflexis. *Hort. upf.* 90. *bucephalophorus.*  
 Rumex fructibus dentatis calyce reflexis, caput bovinum referentibus. *Hort. cliff.* 139.  
 Acetosa, ocymi folio, neapolitana. *Bauh. pin.* 114.  
 Acetosa, ocymi folio, bucephalophoros. *Col. ecphr.* 1. p. 151. t. 150.  
*Habitat in Italia.* ☉  
*Flores semper terni.*
14. RUMEX floribus hermaphroditis: valvulis integer- aquaticus.  
 rimis nudis, foliis cordatis acutis. *Fl. svec. 2. n.* 315.  
 Lapathum aquaticum, folio cubitali. *Bauh. pin.* 116.  
 Lapathum palustre. *Tabern. ic.* 437.  
 Hippolapathum. *Dalech. hist.* 604. *Cam. epit.* 232.  
 Herba britannica. *Munt. brit. t.* 1.  
*Habitat in Europa ad ripas fluviorum* ☿ *paludum.* 2.
15. RUMEX floribus hermaphroditis: valvulis lævibus, Lunar-  
 caule arboreo, foliis subcordatis. *Vir. cliff.* 32. *Hort.*  
*cliff.* 139. *Roy. lugdb.* 320.  
 Acetosa arborefcens, subrotundo folio. *Pluk. alm.* 8. 1. 252. f. 3.  
 Lunaria magorum arabum. *Bauh. hist. 2. p.* 994?  
*Habitat in Canariis.* ☿  
*Flos tripartitus, nec sexpartitus est.*
16. RUMEX floribus hermaphroditis geminatis: valvu- vesicarius.  
 larum alis omnibus maximis membranaceis reflexis, foliis indivisis. *Hort. cliff.* 130. *Hort. upf.* 90. *Roy. lugdb.* 230.  
 Acetosa americana, foliis longissimis pediculis donatis. *Bauh. pin.* 114. *prodr.* 54. *Morif. hist. 2. p.* 583. f. 5. t. 28. f. 7.  
*Habitat in Africa.* ☉
18. RUMEX floribus hermaphroditis distinctis, valvu- tingitanus.  
 lis cordatis obtusis integerrimis, foliis hastato-ovatis erosus.  
 Acetosa dentata perennis. *Zanon. hist.* 14.  
 Acetosa vesicana tingitana perennis repens, foliis longis sinuatis. *Morif. hist. 2. p.* 583. f. 5. t. 28. f. 8.  
 Lapathum maritimum scetidum. *Bauh. pin.* 116. *prodr.* 56.  
*Habitat in Barbaria, Hispania. Astræmer.* 2.

Cau-

*Caules ramosi. Folia petiolata, ovata, acuminata, subsabara, basi subhaflata, juxta basin erofa, margine fcaherrima & fore inaequalia. Racemi longi, floribus 5. f. 3. propriis pedicellis ad fingulos dentes racemi. Calycis valvulae cordatae, integerrimae, n. de.*

rofeus. 19. RUMEX floribus hermaphroditis diftinctis: valvulae alterius ala maxima membranacea, declinata, foliis erofis. *Roy. lugdb. 230.*

Acetofa aegyptia, rofeo feminis involucro. *Shav. afr. 5. f. 5.*

*Habitat in Aegypto. ☉*

*Seminum valvulae maximae, unciales, orbiculae, reticulatae, margine denticulae, apice & bafi bifida, quod non in R. tingitano.*

fcutatus. 20. RUMEX floribus hermaphroditis, foliis cordato-haftatis. *Vir. cliff. 32. Hort. cliff. 138. Hort. upf. 89. Roy. lugdb. 230.*

Acetofa rotundifolia hortenfis. *Bauh. pin. 114. Morif. hift. 2. p. 583. f. 5. t. 28. f. 9.*

Lapathum androgynum acetofum, foliis fagittatis, ramis divergentibus. *Hall. helv. 170.*

Oxalis rotundifolia. *Dod. pempt. 649.*

β. Acetofa fcutata repens. *Bauh. pin. 114. prodr. 55. Habitat in Helvetia, Galloprovincia, inter acervos lapidum. 24*

digynus. 21. RUMEX floribus hermaphroditis digynis. *Hort. cliff. 138. Fl. fvec. 294, 317. Roy. lugdb. 230.*

*Rumex foliis orbiculatis emarginatis. Fl. lapp. 132.*

Acetofa rotundifolia alpina. *Bauh. pin. 55. prodr. 114. Oed. dan. t. 14.*

Acetofa repens veftmorlandia, cochleariae foliis non-nihil finuatis. *Pluk. alm. 8. t. 252. f. 2.*

Acetofa britannica rotundifolia, fructu compreffo. *Blair. obf. 67. t. 67.*

*Habitat in Alpibus Lapponicis, Helveticis, Wallicis. 24*

*Flos quadrididus: Calyce diphyllus, Corolla dipetala, Staminiibus 6 Stylis 2.*

\* \* \* *Floribus diclinis.*

alpinus. 22. RUMEX floribus hermaphroditis ftérilibus femineisque, valvulis integerrimis nudis, foliis cordatis obtufis rugofis.

Ru.

*Rumex folio pedali cordato, radice repente. Monnier.*  
*obs. 165. Sauv. monsp. 111.*

*Lapathum androgynum obtusifolium, spica confertissima. Hall. helv. 170.*

*Lapathum hortense rotundifolium f. montanum. Bauh. pin. 115.*

*Lapathum rotundifolium. Clus. hist. 2. p. 69.*

*Rhabarbarum monachorum. Mat. med. 178.*

*Habitat in Helvetia, Gallia australi. ♂*

*Statura ♀ magnitudo Rhei.*

22. **RUMEX** floribus androgynis: calycibus femineis spinosis, monophyllis: valvulis exterioribus reflexo-uncinatis. *Hort. upf. 89.*

*Rumex floribus androgynis, calycibus fructus uncinatis. Hort. cliff. 139. Roy. lugdb. 230.*

*Beta cretica, semine aculeato. Bauh. pin. 118. prodr. 57.*

*Beta cretica, semine spinoso. Bauh. hist. 2. p. 962.*

*Habitat in Creta. ☉*

23. **RUMEX** floribus dioicis, foliis oblongis sagittatis: tuberosus, hamis patentibus.

*Acetosa tuberosa radice. Bauh. pin. 114.*

*Oxalis tuberosa. Dod. pempt. 649.*

*Habitat in Italia. 4*

*Affinis nimium R. Acetosa ac si ejus hybrida: sed Folia hamis postice magis hiantia ♀ Stipulae longiores, Radix omnino Filipendulae.*

24. **RUMEX** floribus dioicis, foliis oblongis sagittatis. *Acetosa.*  
*Vir. cliff. 32. Hort. cliff. 130. Hort. upf. 89. Fl. jvec. 295, 318. Fl. lapp. 130. Mat. med. 529. Roy. lugdb. 231.*

*Acetosa pratensis. Bauh. pin. 114.*

*β. Acetosa pratensis, flore albo. Tournes. inst. 502.*

*γ. Oxalis crispata. Bauh. hist. 2. p. 990. Tabern. ic 440.*

*δ. Acetosa montana maxima. Bauh. pin. 114. Scheuch. alp. 129.*

- ε. Acetosa montana, lato ari folio rotundo. Bocc. mus. 165. t. 125.*

*Habitat in Europæ pascuis. δ in Alpibus. 4*

25. **RUMEX** floribus dioicis, foliis lanceolato-hastatis. *Acetosella.*

*Vir. cliff. 32. Hort. cliff. 139. Fl. lapp. 131. Fl.*

*succ. 296, 319. Roy. lugdb. 231. Gron. virg. 152.*

*Linnaei Spec. plant. pars 1.*

*H h*

*Ace-*

*Acetosa arvensis lanceolata. Bauh. pin. 114.*

*Oxalis ovina. Tabern. ic. 440.*

β. *Acetosa lanceolata angustifolia repens. Bauh. pin. 14. prodr. 55.*

γ. *Acetosa arvensis minima non lanceolata. Bauh. pin. 114.*

δ. *Acetosa minor erecta, lobis multifidis. Eccc. mus. 164. t. 126.*

*Habitat in Europæ pascuis. ♂ arvis arenosis. 21*

multifidus. 26. *RUMEX floribus dioicis, foliis hastatis: auriculis palmatis. †*

*Acetosa minor erecta lobis multifidis. Bocc. mus. 2. p. 164. t. 126.*

*Habitat in Alpibus Calabriae, Hetruriæ, Orientis. 21*

aculeatus. 27. *RUMEX floribus dioicis, foliis lanceolatis petiolatis, fructibus reflexis: valvulis ciliatis.*

*Acetosa cretica, semine aculeato. Bauh. pin. 114. prodr. 55. t. 55.*

*Oxalis minor aculeata candiæ. Bauh. hist. 2. p. 991.*

*Habitat in Creta, Hispania. 21*

*Similis R. Acetosellæ, sed Folia basi attenuata absque hamis. Florum semineorum interiores valvule ciliatæ aculeis minimis.*

## SCHEUCHZERIA.

palustris. 1. *SCHEUCHZERIA. Fl. lapp. 133. t. 10. f. 1. Fl.*

*succ. 297, 320. Roy. lugdb. 45. Hall. helv. 258.*

*Gmel. sib. 1. p. 73. Berg. viadr. 67.*

*Juncus floridus minor. Bauh. pin. 12. prodr. 43. theatr. 190. Rudb. elys. 1. p. 110. f. 2.*

*Juncoidi affinis palustris. Scheuch. gram. 336.*

*Gramen junceum aquaticum, semine racemoso. Læf. pruss. 114. t. 28.*

*Habitat in Lapponiæ, Helvetiæ, Borussia, Sueciæ paludosis. 21*

## TRIGLOCHIN.

palustre. 1. *TRIGLOCHIN capsulis trilocularibus sublinearibus.*

*Fl. succ. 298, 321. Act. stockh. 1742. p. 147. t. 6. f. 1. 2. 3.*

*Triglochin fructu tenui. Fl. lapp. 134. Roy. lugdb. 44. Hall. helv. 258.*

Gra-

## HEXANDRIA TRIGYNIA.

483

- Gramen junceum spicatum* f. *Triglochin*. *Bauh. pin.*  
6. *theatr.* 80.  
*Calamagrostis* IV. *Dalech. hist.* 1006.  
β. *Juncago maritima perennis*, bulbosa radice. *Mich.*  
*gen.* 44.  
*Juncago maritima*. *Barr. rar.* 55. t. 271.  
*Habitat in Europæ inundatis uliginosis.* ♂  
2. **TRIGLOCHIN** capsulis sexlocularibus ovatis. *Fl. maritimum.*  
*succ.* 299; 322. *Atl. stockh.* 1742. p. 147. t. 6.  
f. 4. 5.  
*Triglochin* fructu subrotundo. *Fl. lapp.* 135. *Roy.*  
*lugdb.* 45. *Gort. gelr.* 76.  
*Juncago palustris & maritima perennis*, fructu brevior  
quinquecapsulari. *Mich. gen.* 44.  
*Gramen spicatum alternum*. *Bauh. pin.* 6. *theatr.* 82.  
*Habitat in Europæ maritimis.* 24

## MELANTHIUM.

1. **MELANTHIUM** petalis ungviculatis. *virgaticum.*  
*Melanthium* foliis linearibus integerrimis longissimis,  
floribus paniculatis. *Gron. virg.* 59.  
*Asphodelo* affinis floridana, ramoso caule, floribus or-  
nithogali obsoletis. *Pluk. amalth.* 40. t. 434. f. 8.  
*Habitat in Virginia.* 24  
2. **MELANTHIUM** petalis sessilibus. *Amæn. acad.* 2. *sibiricum.*  
p. 349. t. 11. \*  
*Ornithogalum* spicis florum longissimis ramosis. *Gmel.*  
*fib.* 1. p. 45. t. 8.  
*Habitat in Sibiria.* 24  
3. **MELANTHIUM** petalis punctatis, foliis cucullatis. *capense.*  
*Amæn. acad.* 6. *afr.* 12. \*  
*Habitat ad Cap. b. Spei.* 24

## MEDEOLA.

1. **MEDEOLA** foliis verticillatis, ramis inermibus. *virginiana.*  
*Medeola* foliis stellatis lanceolatis, fructu baccato.  
*Gron. virg.* 39. \*  
*Lilium* f. *Martagon pusillum*, floribus minutissime her-  
baceis. *Pluk. alm.* 401. t. 328. f. 4.  
*Habitat in Virginia.* ♂

H h a

2. ME-

asparagol-  
des.

2. **MEDEOLA** foliis pinnatis: foliolis alternis.  
*Asparagus foliis ovato-lanceolatis solitariis. Hort. cliff. 122. Roy. lugdb. 29.*  
*Asparagus africanus scandens, myrti folio. Till. pif. 16. t. 12. f. 1. 2.*  
*Laurus alexandrina ramosa, foliis e summitate caulium prodeuntibus. Herm. lugdb. 679. t. 631.*  
*Habitat in Æthiopia. §*  
 Caules teretes, volubiles, inermes, annui plerumque.  
 Folia alterna, remota, forte, pinnata? Foliola ovato-oblonga, sessilia, nitida, alterna, plerumque septena. Simillima *Asparago* sarmentoso, sed inermis & ultra humanam altitudinem ascendens. Collæ albidae, revolutæ, serotinæ. Flores pedunculati, penduli, solitarii, ex ala foliorum; Differunt folia ista pinnata ab aliis pinnatis foliis, & consideranda ut fasciculata, sed distincta & remota petiolo communi, omnino ut in *Phyllantho Niruri*.

### TRILLIUM.

cernuum.

1. **TRILLIUM** flore pedunculato cernuo.  
*Paris foliis ternis, flore pedunculato nutante. Cold. noveb. 81.*  
*Solanum triphyllum, flore hexapetalo carneo. Catesb. carol. 1. p. 45.*  
*Habitat in Carolina. 24.*

erectum.

2. **TRILLIUM** flore pedunculato erecto.  
*Paris foliis ternis, flore pedunculato erecto. Amoen. acad. 1. p. 154. \**  
*Solanum triphyllum brasilianum. Bauh. pin. 167. prodr. 91. Burf. IX. 12.*  
*Solanum triphyllum canadense. Corn. canad. 166. t. 167.*  
*Solano congener triphyllum canadense. Moris. hist. 3. p. 532. f. 13. t. 3. f. 7.*  
*Habitat in Virginia.*

sessile.

3. **TRILLIUM** flore sessili erecto.  
*Paris foliis ternatis, flore sessili erecto. Gron. virg. 44.*  
*Solanum virginianum triphyllum, flore tripetalo atropurpureo in foliorum sinu, absque pediculo, sessili. Pluk. alm. 352. t. III. f. 6.*

*Solanum triphyllum*, flore hexapetalo: tribus petalis  
purpureis, cæteris viridibus reflexis. *Catesb. car.* 1.  
p. 50. t. 50.

*Habitat in Virginia, Carolina.* 2

## COLCHICUM.

1. COLCHICUM foliis planis lanceolatis erectis. *Hort.* autumnale.  
*cliff.* 140. *Hort. ups.* 90. *Roy. lugdb.* 41. *Sauv.*  
*monsp.* 18. 19. *Bæhm. lips.* 29.

*Colchicum commune.* *Bauh. pin.* 67.

*Colchicum.* *Fuch. hist.* 356. 357.

- β. Colchicum flore pleno. *Bauh. pin.* 67.

*Habitat in Europæ australioris succulentis.* 2

2. COLCHICUM foliis linearibus patentissimis. *Læfl.* montanum.

*Colchicum montanum angustifolium.* *Bauh. pin.* 68.

*Colchicum montanum.* *Clus. hisp.* 266. t. 267. *hist.*

1. p. 200.

*Habitat in Hispania, Helvetia.* 2

*Folia una cum floribus autumnis erumpunt.*

3. COLCHICUM foliis undulatis patentibus. *Hort.* variegatum.  
*cliff.* 140. *Roy. lugdb.* 42.

*Colchicum chionense*, floribus fritillariæ instar tessula-  
tis, foliis undulatis. *Morif. hist.* 2. p. 341. f. 4. t.

3. f. 7.

*Habitat in Chio insula.* 2

## HELONIAS.

1. HELONIAS.

*Helonias foliis radicalibus lanceolatis.* *Amæn. acad.* 3.

p. 12. t. 1. f. 1. \*

*Ephemerum phalangoides virginianum*, flosculis arbu-  
teis bullatis in spicam dispositis. *Pluk. alm.* 135. t.

174. f. 5. *Morif. hist.* 3. p. 606. f. 15. t. 2. f. 1.

*Habitat in Pensylvaniæ paludosis.* 2

bullata.

2. HELONIAS foliis caulinis setaceis.

*Habitat in Pensylvania.* *Barthram.*

*Caulis simplicissimus, bipedalis. Folia alternata s,*  
*sparsa, lineari-setacea, erecta, lævia, margine sca-*  
*bra. Racemus terminalis, simplex, pedunculis flore*  
*longioribus. Flores albi. Simillima Asphodelo, sed*  
*Styli tres recurvati.*

asphodelol-  
des.



## TETRAGYNIA.

## PETIVERIA.

- alliacæ. 1. PETIVERIA floribus hexandris. *Hort. cliff.* 141.  
*Hort. upf.* 91. *Ast. stockh.* 1744. p. 287. t. 7.  
 Petiveria foliis oblongo-ovatis, spicis longioribus terminalibus. *Brown. jam.* 274.  
 Verbenæ aut Scorodonizæ affinis anomala, flore albido calyce aspero, allii odore. *Sloan. jam.* 64. *hist.* 1. p. 172. *Raj. suppl.* 287.  
*Habitat in Jamaica nemorosis.* ♀
- octandra. 2. PETIVERIA floribus octandris.  
 Petiveria foliis, loculis spinosis. *Plum. gen.* 50. ic. 219.  
*Habitat in America meridionali.* Jacquin. ♀  
 Simillima *P. præcedenti*, ut vix, nisi ab exercitato, distinguatur. Differt vero Caule breviori strictiore; Staminiibus octo; Flamentis purpureis, nec albis: Foliis rigidioribus & omnino glabris.

## POLYGYNIA.

## ALISMA.

- Plantago. ▽ 1. ALISMA foliis ovatis acutis, fructibus obtuse trigonis.  
 Alisma fructu obtuso trigono. *Fl. lapp.* 138. *Fl. suec.* 300; 323. *Hort. cliff.* 141. *Gmel. sib.* 1. p. 77.  
 Plantago aquatica latifolia. *Bauh. pin.* 190.  
 Plantago aquatica. *Cam. epit.* 264.  
*Habitat in Europæ aquosis & ad ripas fluviorum, lacuum.* 24.
- flav.:. 2. ALISMA foliis ovatis acutis, pedunculis umbellatis, fructibus globosis.  
 Damasonium maximum, plantaginis folio, flore flavescente, fructu globoso. *Plum. spec.* 7. ic. 115.  
*Habitat in America meridionali.* 24.
- Damasonium. 3. ALISMA foliis cordato-oblongis, floribus hexagynis, capitulis subulatis.  
 Alisma fructu sexcorni. *Hort. cliff.* 141. *Roy. lugdb.* 46. *Sauv. monsp.* 14.  
 Plantago aquatica stellata. *Bauh. pin.* 190.

Plan.

*Plantago aquatica minor altera. Lob. ic. 301.*

*Damaſonium ſtellatum. Dalech. hiſt. 1058.*

*Habitat in Angliæ, Galliæ aquoſis.*

4. **ALISMA** foliis cordatis obtuſis, floribus dodecan- cordifolia-  
dris, feminibus uncinato muricatis.

*Damaſonium ramoſum, folio cordiformi. Vaill. aët.*

*1719. p. 28.*

*Sagittaria virginiana, obtuſiore lato folio, floribus mi-  
noribus albis. Morif. hiſt. 3. p. 618. f. 15. t. 4. f. 6.*

*Ranunculus aquaticus, aſari foliis, ad nodos umbelli-  
ſterus. Plum. ſpec. 7. ic. 234. f. 2.*

*Habitat in America aſtrali & ſeptentrionali. 24*

*Media inter Alifmas & Sagittarias.*

5. **ALISMA** foliis ovatis obtuſis, pedunculis ſolitariis. natans.

*Fl. ſucc. 2. 324.*

*Damaſonium repens, potamogetonis rotundifolli folio.*

*Vaill. aët. 1719. p. 29. t. 4. f. 8.*

*Damaſonium radículas emittens ex geniculis. Vaill.  
pariſ. 46.*

*Ranunculus paluſtris, foliis gramineis & ſubrotundis.*

*Petit. gen. 47.*

*Habitat in Galliæ, Sueciæ ſoffis.*

6. **ALISMA** foliis lineari-lanceolatis, fruſtibus globoſo- Ranuncu-  
ſquarroſis. loides.

*Alifma fruſtu globoſo undique echinato. Hort. cliſt.*

*141. It. gotl. 298. Fl. ſucc. 301; 325. Roy. lugdb.*

*46. Gort. gelr. 77. Sauv. monſp. 14*

*Damaſonium anguſtiſſimo plantaginis folio. Vaill. aët.*

*1719. p. 35.*

*Plantago aquatica humilis anguſtifolia. Bauh. hiſt. 3.*

*p. 738. Lob. ic. 300.*

*Ranunculus aquaticus, plantaginis folio anguſtiſſimo.*

*T. Pet. gar. 41. t. 26. f. 12.*

*Habitat in Gotlandiæ, Belgii, Angliæ, Galliæ ſoffis.*

7. **ALISMA** foliis ſubulatis, Gron. virg. 153. ſubulata,  
*Habitat in Virginia.*

## Classis VII.

HEPTANDRIA  
MONOGYNIA.

## TRIENTALIS.

europæa.

1. TRIENTALIS foliis lanceolatis integerrimis.  
 Trientalis. *Fl. lapp.* 139. *Fl. fvec.* 302.; 326. *Roy. lugdb.*  
 438. *Hall. opusc.* 123. *Kalm. it.* 2. p. 263.  
 Alfine alpina. *Svenkf. fl. sc.* 117.  
 Pyrola, alfine flore, europæa. *Bauh. pin.* 191. *prodr.* 101.  
 Pyrola, alfine flore, minor brasiliiana. *Bauh. prodr.*  
 101. t. (sub *Hermodyli* nomine) 99. *Burf. X.* 107.  
*Habitat in Europæ borealis sylvis & juniperetis.* 2.

## ÆSCULUS.

Hippo Ca-  
stanum,

1. ÆSCULUS floribus heptandris. *Hort. upf.* 92.  
 Æsculus. *Hort. cliff.* 142. *Roy. lugdb.* 463.  
 Castanea folio multifido. *Clus. hist.* 1. p. 7. *Bauh.*  
*pin.* 419.  
*Habitat in Asia septentrionaliore, unde in Europam*  
*1550. §*

Pavia.

2. ÆSCULUS floribus octandris.  
 Pavia. *Boerh. lugdb.* 2. p. 260. t. 260. *Hort. angl.* 54.  
 t. 19. *Hort. cliff.* 143. *Roy. lugdb.* 463. *Trew. chret.*  
 t. 15. *Mill. ic.* 198. *Duham. arb.* 2. p. 98.  
 Saamouna pisonis f. filiquifera brasiliensis arbor, digi-  
 tatis foliis serratis, floribus teucree purpureis. *Pluk.*  
*alm.* 326. t. 56. f. 4.  
*Habitat in Carolina, Brasilia. §*  
*Capsula hujus glabra, præcedentis muricata. Flores*  
*in eodem thyrsu Masculos & Hermaphroditos obser-*  
*vavit Dav. v. Royen; si ita, Polygama erit.*

## DIGYNIA.

## LIMEUM.

africanum.

1. LIMEUM.

Ha-

*Habitat in Æthiopia. 21.*

*Statura Corrigiolæ f. Telephii. Caules prostrati, debiles, spithamæi, angulati, nudi, basi perennantes. Folia alterna, remota, lineari-lanceolata, parva, subpetiolata. Corymbi terminales, solitarii, compositi, nudi, longius pedunculati.*

## TRIGYNIA.

### SAURURUS.

1. SAURURUS foliis cordatis petiolatis, amentis solitariis recurvis. *Hort. ups. 91. cernuus.*

*Saururus foliis profunde cordatis ovato-lanceolatis, spicis solitariis. Hort. cliff. 139. Gron. virg. 40. Roy. lugdb. 8.*

*Serpentaria repens, floribus stamineis spicatis, bryonia nigra folio ampliore pingui. Pluk. alm. 343. t. 117. f. 4.*

*Habitat in Virginia. 21.*

*Amenti Squamæ petiolatæ, albæ; at defloratæ virescentes, pedicellis incurvatis ad rachin, germi incumbens. Stamina vix numerantur, nisi deflorato amento persistens.*

## HEPTAGYNIA.

### SEPTAS.

1. SEPTAS. *Amzn. acad. 6. afr. 13. \* capensis.*

*Doronici? species pumila, auriculæ urfi folio glabro. Pluk. mant. 166. t. 330. f. 9.*

*Trientalis foliis subrotundis crenatis. Roy. lugdb. 538.*

*Habitat ad Cap. b. Spei 24. Dav. v. Royen.*



## Classis VIII.

OCTANDRIA  
MONOGYNIA.

## TROPÆOLUM.

minus.

1. TROPÆOLUM foliis integris, petalis acuminato-setaceis. *Hort. upf.* 93.  
 Tropæolum foliis peltatis orbiculatis. *Hort. cliff.* 145.  
 Cardamindum minus & vulgare T. *Fewill. peruv.* 3.  
*p.* 14. *t.* 8.  
 Nasturtium indicum majus. *Bauh. pin.* 306.  
 Nasturtium indicum. *Lob. ic.* 516. *f.* 2, 1.  
*Habitat in Peru, Limæ innotuit 1580. per Dodonæum.* ☉. 24

majus.

2. TROPÆOLUM foliis subquinelobis, petalis obtusis. *Hort. upf.* 93.  
 Tropæolum foliis peltatis orbiculatis. *Hort. cliff.* 143.  
 Cardamindum ampliori folio & majori flore. T. *Few.*  
*peruv.* 9. *p.* 14. *t.* 8.  
 Acriviola maxima odorata. *Boerh. lugdb.* 1. *p.* 244.  
 Viola inodora scandens, nasturtii sapore, maxima odorata. *Herm. lugdb.* 628. *t.* 629.  
*Habitat in Peru, unde in Europam venit 1684. cura Bewerningii.* ☉. 24  
 Flores ante crepusculum fulminant, observante E. C. *Linnaea.*

peregrinum.

3. TROPÆOLUM foliis palmatis, petalis multifidis.  
 Cardamindum quinquefolii folio, vulgo Malla. *Few.*  
*peruv.* 2. *p.* 756. *t.* 42.  
*Habitat in Peru, nondum mihi visum in Europa.* ☉

## OSBECKIA.

chinensis.

1. OSBECKIA. *Osbeck. it.* 213. *t.* 2. \*  
 Echinophora maderaspatana, sideritis non ferratis nervosis foliis, fructu capsulari caucalidis æmulo. *Pluk. alm.* 132. *t.* 173. *f.* 4.  
 Kadali maderaspatanum minus, capsulis pilosis. *Raj. suppl. app.* 236.

I. A.

*Habitat in India; in honorem Petr. Osbeck, Sveci, qui 1751. plantas Chinæ & Javæ periculoso itinere adiit, legit, examinavit, descripsit, communicavit.*

*Habitus Melastomæ. Caulis erectus, brachiatus, acute tetragonus. Folia angusto-lanceolata, trinervia, opposita, scabra, subsessilia. Flores terminales aliquot, sessiles, cincti foliis 4 flore longioribus patentibus.*

## R H E X I A.

1. RHEXIA foliis sessilibus ferratis, calycibus glabris. *virginica.*

*Gron. virg. 41.*

*Alifanus vegetabilis carolinianus. Pluk. amalth. 8.*

*Lysimachia non papposa virginiana, tuberariæ foliis hirsutis, flore tetrapetalo rubello. Pluk. alm. 235. t. 202. f. 8.*

*Habitat in Virginia.*

*Caulis tetragonus angulis membranaceis. Folia opposita, sublanceolata, internodiis longiora, sessilia, trinervia, pilis vagis hispidiuscula, subserrata serraturis setaceis. Pedunculus terminalis, dichotomus. Flores solitarii ex dichotomia, subsessiles, petalis rubris; Antheris falcatis, luteis.*

2. RHEXIA foliis cillatis.

*mariana.*

*Lysimachia non papposa, terræ marianæ, leptoneuros, flore tetrapetalo rubello, folio & caule hirsutis ferruginea hispidis. Pluk. mant. 123. t. 428. f. 1.*

*Habitat in Marilandia, Brasilia.*

*Calyx adpersus setis apice stellatis.*

3. RHEXIA floribus alternis axillaribus pedunculatis *Acisanthera.* quinquefidis. *Amæn. acad. 5. p. 396. \**

*Acisanthera erecta ramosa, ramulis quadratis, foliolis trinerviis ovatis crenatis oppositis, floribus singulis. Brown. jam. 217. t. 22. f. 1.*

*Habitat in Jamaica. 24*

## O E N O T H E R A.

1. OENOTHERA hirta, foliis supra glabris.

*hirta.*

*Oenothera affurgens hirsuta, floribus solitariis. Brown. jam. 208 ?*

*Onagra frutescens & hirsuta nerii folio, flore magna luteo. Plum. spec. 7. ic. 174. f. 2.*

*Habitat in America calidiore. 5*

2. OE-

- biennis.** 2. **OENOTHERA** foliis ovato-lanceolatis planis, caule muricato-subvillosa. *Vir. clif.* 33. *Hort. upf.* 94. *Gron. virg.* 254. *Roy. lugdb.* 251. *Gort. gelr.* 78. *Kalm. it.* 3. p. 509.  
*Oenothera* foliis ovato-lanceolatis denticulatis, floribus lateralibus in summo caulis. *Hort. clif.* 144.  
*Lyfimachia lutea corniculata.* *Bauh. pin.* 245. 516. \*  
*Morif. hist.* 2. p. 271. f. 3. t. 11. f. 7.  
*Hyoscyamus virginianus.* *Alp. exot.* 325. t. 324.  
*Habitat in Virginia, unde 1614, nunc vulgaris Europæ.* ♂
- parviflora.** 3. **OENOTHERA** foliis ovato-lanceolatis planis, caule lævi subvillosa.  
*Oenothera* foliis lanceolatis, capsulis ventricoso-conicis. *Zin. gætt.* 199.  
*Habitat in America septentrionali.* ♂  
*Fruſtus apex coronatus margine oſtifiſido, nec quadriſido, uti OE. biennis, cui ſimillima. Caulis pilis adſperſus, ſed abſque tuberculis ad eorum baſin. Folia repando ſubdentata, minus mollia. Calyx tubo ſuo quadruplo brevior, infra apicem denticulo notatus, hinc ante explicationem mucrones in hac diſtantes. Petalorum latera diſtancia, dimidio minora præcedentis petalis. Capſulæ quadrivalves.*
- octovalvis.** 4. **OENOTHERA** foliis lanceolato-oblongis acumina-  
tis planis glabris. *Jacq. amer.* 19. *hiſt.* t. 70. †  
*Onagra* foliis perſicariæ anguſtioribus, flore magno luteo. *Plum. ſpec.* 7. ic. 175. f. 1.  
*Habitat in America.*  
*Plumieri figura etiam OE. parviſſoræ reſert.*
- meſſiſſima.** 5. **OENOTHERA** foliis lanceolatis undulatis. *Vir. clif.* 33. *Gron. virg.* 42. *Roy. lugdb.* 251.  
*Oenothera* foliis lineari-lanceolatis dentatis, floribus e medio caule. *Hort. upf.* 144.  
*Onagra bonarienſis villoſa, flore mutabili.* *Dill. elth.* 297. t. 219. f. 286.  
*Habitat in agro Bonarienſi.* ☉
- ſruticeſa.** 6. **OENOTHERA** foliis lanceolatis ſubdentatis, capsulis pedicellatis acutangulis, racemo pedunculato.  
*Oenothera* florum calyce monophyllo, hinc tantum aperto. *Gron. virg.* 42.

*Onagra angustifolia*, caule rubro, flore minore.  
*Tournef. infl.* 302.

*Habitat in Virginia.* 24

*Calyx purpurascens, tetraphyllus, sed rumpitur altero  
superius & unico latere. Tubus filiformis, angustis-  
simus. Capsula angulis 4 acutis compressis. Folia  
rarius denticulata.*

7. **OENOTHERA** foliis lanceolatis obtusis glabris sub- pumila.  
petiolatis, caulibus prostratis, capsulis acutangulis.

*Oenothera foliis radicalibus ovatis, caulinis lanceolatis  
obtusis, capsulis ovatis obtusis. Mill. dict. t.* 188.

*Lyfimachia filiquosa glabra minor mariana, angustio-  
ribus foliis. Pluk. mant.* 123.

*Lyfimachia marilandica parva, foliis angustis acutis.  
Raj. suppl.* 1416.

*Habitat in America septentrionali.* 24

*Caulis palmaris, herbaceus, teres, flexuosus, divari-  
catissimus. Folia elliptico-lanceolata, integerrima.  
Flores axillares, solitarii, erecti, subsessiles. Peta-  
la lutea, obcordata, lineata. Capsulae obovatae, ob-  
tusae, octangulares, angulis quatuor latioribus com-  
pressis, quadrivalvis. Radix perennis.*

## G A U R A.

1. **GAURA.** *Aman. acad.* 3. p. 26. *Afl. holmens.* 1756. biennia.  
*p.* 222. t. 8. \*

*Lyfimachia Chamænerio similis floridana, foliis nigris  
punctis, capsulis carinatis in ramulorum cymis. Pluk.  
amalth.* 139. t. 428. f. 2.

*Habitat in Virginia, Pennsylvania.* ♂

## EPILOBIUM

\* *Staminibus declinatis.*

1. **EPILOBIUM** foliis sparsis lineari-lanceolatis, flori- angustifol-  
um.  
bus inæqualibus.

*Epilobium floribus difformibus, pistillo declinato. Fl.  
svcc.* 304; 327.

*Epilobium foliis lanceolatis integerrimis. Fl. lapp.* 146.  
*Hort. cliff.* 154. *Roy. lugdb.* 250.

*Lyfimachia Chamænerion dicta angustifolia. Bauh.  
pin.* 245.

6. *Ly.*



β. *Lyfimachia Chamænerion dicta latifolia. Bauh. pin.*  
245.

γ. *Lyfimachia Chamænerion dicta alpina. Bauh. pin.* 245.  
*prodr.* 116.

*Habitat in Europa boreali.* 24

*latifolium.* 2. **EPILOBIUM** foliis alternis lanceolato-ovatis, floribus inæqualibus.

*Habitat in Sibiria.* 24

*Differt a præcedenti floribus duplo majoribus, foliis lanceolato-ovatis, alternis nec sparsis, utrinque levissimo tomento mollissimis. Hec & antecedens differunt a sequentibus in non paucis: his enim Flores inæquales petalis integris; Stamina basi conniventia ad Stylum; Pistillum declinatum; Folia alterna; Folia erumpentia revoluta contra ac in reliquis.*

\* \* *Staminibus erectis regularibus, petalis bifidis.*

*hirsutum.* 3. **EPILOBIUM** foliis oppositis lanceolatis ferratis decurrenti-amplexicaulibus. *Hort. cliff.* 145. *Fl. svec.* 305.  
328. *Gron. virg.* 154. *Roy. lugdb.* 251.

*Lyfimachia filiquosa hirsuta, magno flore. Bauh. pin.*  
245.

*Lyfimachia purpurea. Fuchf. hist.* 491.

β. *Lyfimachia filiquosa hirsuta, parvo flore. Bauh. pin.*  
245. *prodr.* 116.

*Habitat in Europæ humidiusculis.* 24

*Planta subhirsuta. Foliorum basjs media altior lateralibus.*

*montanum.* 4. **EPILOBIUM** foliis oppositis ovatis dentatis. *Fl. lapp.*  
147. *Fl. svec.* 306; 329. *Hort. cliff.* 145. *Roy. lugdb.*  
251.

*Lyfimachia filiquosa glabra major. Bauh. pin.* 245.

*Pseudo-Lyfimachium purpureum primum. Dod. pempt.*  
85.

*Habitat in Europæ montosis.*

*Caulis mollissimus. Folia glabra.*

*tetragonum.* 5. **EPILOBIUM** foliis lanceolatis denticulatis: imis oppositis, caule tetragono. *Sauv. monsp.* 75.

*Lyfimachia filiquosa glabra minor. Bauh. pin.* 303.  
*Raj. hist.* 801.

*Lyfimachia minor. Tabern. ic.* 854.

- Habitat in Europa.* 24.  
*Folia tenera macula livida. Summitas; adhuc tenella, nutans.*
6. **EPILOBIUM** foliis oppositis lanceolatis integerrimis, petalis emarginatis, caule erecto. *Fl. suec.* 307; 330. \*  
*Epilobium foliis linearibus. Fl. lapp.* 149. *Roy. lugdb.* 251.  
*Lyfimachia filiquosa glabra angustifolia. Bauh. pin.* 245.
- β. *Epilobium foliis lanceolatis, ramose florens. Fl. lapp.* 148.  
*Habitat in Europæ humidiusculis. β in Alpibus.* 24.
7. **EPILOBIUM** foliis oppositis ovato-lanceolatis integerrimis, filiquis sessilibus, caule repente. *alpinum.*  
*Epilobium foliis ovalibus: superioribus attenuatis. Fl. suec.* 308; 331.  
*Epilobium foliis ovato-oblongis integerrimis. Fl. lapp.* 150.  
*Epilobium, foliis ellipticis obtuse lanceolatis, totum læve. Hall. herb.* 409.  
*Chamænerion alpinum, alines foliis. Scheuch. alp.* 132. 332.  
*Lyfimachia filiquosa nana, prunellæ foliis. Bocc. mus.* 2. p. 161. t. 108.  
*Habitat in Alpibus Helveticis, Lapponicis.* 24  
*Repens est, primo vere florens germinibus sessilibus.*

## MELICOCCA.

1. **MELICOCCA.** *Jacq. hist. t.* 71. *bijuga.*  
*Melicoccus foliis ut plurimum bijugatis ovatis, per pennas alas dispositis. Brown. jam.* 210.  
*Nux americana costa foliorum appendicibus acuta. Pluk. alm.* 265. t. 207. f. 4.  
*Nux americana foliis alatis bifidis. Comm. hort.* 1. p. 183. t. 94.  
*Habitat in America meridionali. †*  
*Obnoxia Cocco hesperidum præ reliquis, ut Sapindus, cui simillima.*

## AMYRIS-

1. **AMYRIS** foliis ternatis acutis. *Elemifera.*  
*Elemifera foliis ternatis. Hort. cliff.* 486.

Fru-

Frutex trifolius resinofus floribus tetrapetalis albis racemosis. *Catech. car.* 2. p. 33. t. 33. f. 3.

Cornus racemosa trifolia & quinquefolia, foliis foraminulatis. *Plum. spec.* 19. ic. 100?

*Habitat in Carolina.* ‡

- maritima. 2. AMYRIS foliis ternatis obtusis. *Jacq. amer.* 19.  
Amyris fruticosus minor, foliis orbiculatis venosis pin-  
nato-ternatis, racemis terminalibus. *Brown. jam.* 209.  
*Habitat in America.* ‡

- toxifera. 3. AMYRIS foliis impari pinnatis.  
Elemifera foliis pinnatis. *Hort. cliff.* 486.  
Toxicodendron foliis alatis, fructu purpureo pyr for-  
mi sparso. *Catech. car.* 1. p. 40. t. 40.  
*Habitat in Carolina.*

- balsamifera. 4. AMYRIS foliis bijugis.  
Arboris arboreus, foliis bijugatis ovatis glabris, race-  
mis laxis terminalibus. *Brown. jam.* 208.  
Lauro affinis terebinthi folio alato, ligno odorato can-  
dido, flore albo. *Sloan. jam.* 137. *hist.* 2. p. 24. t. 168.  
f. 4. *Raj. dendr.* 88.  
Lucinium arbor tilie foliis minoribus, americanum.  
*Pluk. a.m.* 228. t. 201. f. 3.  
*Habitat in Jamaica.* ‡

## GRISLEA.

- secunda. 1. GRISLEA. *Hort. cliff.* 146. *Læfl. it.* 245. \*  
*Habitat in America calidiore.* ‡

## COMBRETUM.

- laxum. 1. COMBRETUM spicis laxis. *Læfl. it.* 308.  
Combretum floribus octandris, spicis laxis secundis.  
*Jacq. amer.* 19.  
Gaura fruticosa scandens, foliis oppositis. *Læfl. it.*  
248. \*  
*Habitat in America meridionali.* ‡

- secundum. 2. COMBRETUM spicis secundis. *Jacq. amer.* 19. †  
*Habitat in America meridionali.* ‡

## ALOPHYLLUS.

- zeylanicus. 1. ALOPHYLLUS. *Fl. zeyl.* 140. \*  
*Habitat in Zeylona.* ‡

## XIMENIA.

1. XIMENIA foliis oblongis, pedunculis multifloris. americana.  
*Jacq. amer. 19. hist. t. 177. f. 31. Hort. cliff. 1193.*  
*Ximania aculeata, flore villosa, fructu luteo. Plum.*  
*gen. 6. ic. 261. f. 1.*  
*Habitat in America. ♀*  
*Folia ovato-oblonga, alterna, integerrima, petiolata.*  
*Spina ad singulam foliorum axillam. Racemus multi-*  
*florus, inter spinam & ramum.*
2. XIMENIA foliis ovatis, pedunculis unifloris. inermis.  
*Amyris arborescens, foliis ovatis glabris: vetustiori-*  
*bus confertis, petiolis submarginatis, floribus solita-*  
*riis. Brown. jam. 209.*  
*Habitat in Jamaica. ♀*

## MIMUSOPS.

1. MIMUSOPS foliis alternis remotis. *Fl. Zeyl. 138. \* Elengi.*  
*Flos cuspidum. Rumph. amb. 2. p. 189. t. 63*  
*Kauki indorum. Pluk. alm. 203. Breyn. cent. 20. t. 8.*  
*Elengi. Rhed. mal. 1. p. 34. t. 20.*  
*Habitat in India. ♀*
2. MIMUSOPS foliis confertis. *Fl. Zeyl. 137. \* Kauki.*  
*Metrolderops macassarensis. Rumph. amb. 3. p. 19.*  
*t. 8.*  
*Habitat in India. ♀*

## JAMBOLIFERA.

1. JAMBOLIFERA. *Fl. Zeyl. 139. \* peduncu-*  
*Jambulones. Bauh. pin. 460. lata.*  
*Habitat in India. ♀*

## SANTALUM.

1. SANTALUM. *Mat. med. 183. album.*  
*Santalum verum. Breyn. ic. 94. t. 5. f. 1.*  
*Santalum album. Bauh. pin. 392. Rumph. amb. 2. p.*  
*42. t. 11.*  
*Habitat in India. ♀*

## MEMECYLON.

1. MEMECYLON foliis ovatis. *Fl. Zeyl. 136. \* capitella-*  
*I i Cor- tum.*

*Cornus sylvestris*, foliis croceum colorem tingentibus,  
flosculis ad foliorum alas globosis. *Burm. zeyl.* 76.  
t. 30.

*Habitat in Zeylona.* ‡

## LAWSONIA.

inermis.

1. *LAWSONIA* ramis inermibus. *Fl. zeyl.* 135. *Gron. orient.* 47.

*Ligustrum ægyptiacum latifolium.* *Bauh. pin.* 476.

*Alhenna* f. *Henna arabum.* *Walth. hort.* 3. t. 4.

*Pontaletice.* *Rheed. mal.* 4. p. 117. t. 57.

*Alcanna arabum.* *Bell. itin.* 35.

*Cyprus*, *Henna*, *Alcanna.* *Rauw. itin.* 60. t. 60.

*Habitat in India, Ægypto.* ‡

*Affinis maxime sequenti; ut forte cultura mitior facta.*

sp. n. 106a

2. *LAWSONIA* ramis spinosis. *Fl. zeyl.* 134. *Hasselt.* it. 464. \*

*Rhamnus malabaricus.* *Mail anschi. Pluk. alm.* 28. 220. f. 1.

*Cyprus Rumph. amb.* 4. p. 42 t. 17.

*Mail-anschi. Rheed. mal.* 1. p. 73. t. 40.

*Habitat in India.* ‡

## VACCINIUM.

\* *Foliis annotinis deciduis.*

Myrtillus.

1. *VACCINIUM* pedunculis unifloris, foliis serratis ovatis deciduis, caule angulato. *Fl. lapp.* 143. *Fl. suec.* 313; 333. *Mat. med.* 184. *Hort. cliff.* 148. *Roy. lugdb.* 239. *Hall. helv.* 419.

*Vitis idæa* foliis oblongis crenatis, fructu nigricante. *Bauh. pin.* 470.

*Myrtillus germanica* & *Vitis idæa.* *Dalech. hist.* 191; 192.

*Habitat in Europæ borealis sylvis umbrosis.* ‡

stamineum.

2. *VACCINIUM* pedunculis solitariis unifloris, aptheris corolla longioribus, foliis oblongis integerrimis. *Vaccinium* staminibus corolla longioribus. *Gron. rug.* 43.

*Vitis idæa americana*, longiori mucronato folio & crenato, floribus urceolatis racemosis. *Pluk. alm.* 331. t. 239. f. 3.

Ha-

*Habitat in America septentrionali.* ♀

*Frutex.* Folia oblonga, integerrima. Pedunculi axillares, solitarii, uniflori, filiformes, corolla longiores. Corollæ quinquefidæ, campanulato-patentes. Antherae 10. Stylus flore longior.

3. **VACCINIUM** pedunculis unifloris, foliis integerrimis ovalibus basi ciliatis. uliginosum.

*Vaccinium* foliis ovalibus integerrimis deciduis. *Fl. fvec.* 312; 332.

*Vaccinium* foliis annuis integerrimis. *Fl. lapp.* 142. *Hall. helv.* 414.

*Vitis idæa* foliis subrotundis exalbidis. *Bauh. pin.* 470.

*Vitis idæa* 11. *Clus. hist.* 1. p. 61. t. 62.

*Habitat in Sueciæ borealibus & alpinis, uliginosis.* ♀

4. **VACCINIUM** pedunculis simplicibus, foliis integerrimis ovatis subtus tomentosis. album.

*Habitat in Pensylvania.* *Kalm.* ♀

*Frutex.* Folia ovata, subtus tomentosa. Flores duo vel tres ad apices ramorum: pedunculis aggregatis, brevissimis, nudis.

5. **VACCINIUM** pedunculis, simplicissimis unifloris, foliis ovatis mucronatis glabris integerrimis. mucronatum.

*Habitat in America septentrionali.* *Kalm.* ♀

*Frutex.* Ramis teretibus, glaucis. Folia elliptica, utrinque glabra, integerrima, mucronata, petiolis canaliculatis insidentia. Pedunculi uniflori, foliis paulo breviores.

6. **VACCINIUM** floribus corymbosis ovatis, foliis oblongis acuminatis integerrimis. corymbosum.

*Habitat in America septentrionali.* *Kalm.* ♀

*Frutex.* Folia oblonga, acuta, nervis subtus villosis. Corymbi ex gemmaceo involucre, sessiles. Corollæ cylindraceo-ovata, longiores, quam in reliquis. Stamina decem.

7. **VACCINIUM** racemis filiformibus foliosis: foliis oblongis integerrimis. frondosum.

*Vaccinium* foliis ovatis integris deciduis, racemis foliosis. *Giron. virg.* 155. \*

*Habitat in America septentrionali.* ♀

*Frutex.* Folia oblonga, integerrima. Racemi infra folia, bracteis oblongis, parvis. Pedunculi partialiter bracteis

*filis longiores, instructi bracteolis duabus filiformibus.*  
*Corolla ovato-campanulata. Stamina 19.*

*ligustrinum.* 8. **VACCINIUM** racemis nudis, caule fruticoso, foliis crenulatis oblongis.

*Habitat in Pennsylvania. Kalm. h*

*Flores in Racemis alternis, numerosis, destitutis bracteis, s. foliolis, qua nota facile distinguitur.*

*Aristostaphylos.*

9. **VACCINIUM** floribus racemosis, foliis crenulatis ovatis acutis, caule arboreo.

*Vitis idæa orientalis maxima, cerasi folio, flore variegato. Tournef. cor. 42. it. 2. p. 223. t. 223.*

β. *Vitis idæa cappadocica maxima, mespili folio, flore variegato. Tournef. cor. 42.*

*Habitat in Cappadocia. h*

*Racemi foliis longiores, infra folia sui, foliolis ovatis instructi.*

\* \* *Foliis sempervirentibus.*

*Vitis idæa.* 10. **VACCINIUM** racemis terminalibus nutantibus, foliis obovatis revolutis integerrimis subtus punctatis.

*Vaccinium foliis obverse ovatis perennantibus. Fl. lapp. 145. Fl. suec. 314, 334. Mat. med. 185. Hort. cliff. 148. Roy. lugdb. 239. Hall. helv. 414. Gort. getr. 81.*

*Vitis idæa foliis subrotundis non crenatis, baccis rubris. Bauh. pin. 470. Oed. dan. t. 40.*

*Habitat in Europæ frigidioris sylvis macris. h*

*Oxycoccus.* 11. **VACCINIUM** foliis integerrimis revolutis ovatis, caulibus repentibus filiformibus nudis.

*Vaccinium ramis filiformibus repentibus, foliis ovatis perennantibus. Fl. lapp. 145.*

*Vaccinium ramis filiformibus, foliis ovatis perennantibus, pedunculis simplicibus stipula duplici. Fl. suec. 315; 335. Mat. med. 186.*

*Vitis idæa palustris. Bauh. pin. 471.*

*Vaccinia palustris. Dod. pempt. 770. Lob. ic. 109.*

*Habitat in Europæ paludibus sphagno repletis. 24*

*Racemus brevissimus, pedunculis longissimis: bracteis 2 alternis; Corolla revoluta.*

*hispidulum.* 12. **VACCINIUM** foliis integerrimis revolutis ovatis, caulibus repentibus filiformibus hispidis. *Kalm. it. 3.*

*p. 37.*

*Vitis idæa palustris americana*, oblongis splendentibus foliis, fructu grandiore rubro plurimis intus acinis referto. *Pluk. alm.* 392. t. 320. f. 6.

*Vitis idæa palustris virginiana*, fructu majore. *Raj. hist.* 685.

*Habitat in Americæ septentrionalis paludosis.* 24.

*Structura antecedentis*, sed *majora omnia*, & *Caulis squamis setaceis imbricatus.*

## ERICA.

\* *Antheris biaristatis.*

1. **ERICA** antheris bicornibus inclusis, corollis inæqualibus campanulatis mediocribus, foliis oppositis sagittatis. *vulgaris.*

*Erica* foliis quadrifariam imbricatis triquetris glabris erectis, corollis inæqualibus calyce brevioribus. *Hort. cliff.* 146. *Fl. suec.* 309, 336. *Roy. lugdb.* 442.

*Erica vulgaris glabra.* *Bauh. pin.* 485. *Fl. lapp.* 141.

2. *Erica myricæ folio hirsuto.* *Bauh. pin.* 485.

*Erica vulgaris hirsuta.* *Raj. angl.* 2. p. 471.

*Habitat in Europæ campestribus sterilibus, frequens.* 5

*Comparativa*: mediocris, longior, brevior, corolla respicit proportionem ad calycem.

2. **ERICA** antheris bicornibus exsertis, corollis globosis umbellatis, foliis ternis acerosis glabris. *umbellata.*

*Erica* foliis acerosis glabris ternis, corollis ovatis staminibus brevioribus terminalibus. *Læfl. it.* 138.

*Habitat in Lusitania.* 5

*Fruticulus habitu Ericæ vulgaris.* Folia terna, acerosa, brevia, glabra, subtus linea alba notata. Umbellulæ nullæ absque involucrio. Calyx compositus. Corolla palide cærulea, angulata.

3. **ERICA** antheris bicornibus inclusis, corollis campanulatis mediocribus secundis, foliis ternis triquetris patulis. *herbacea.*

*Erica procumbens herbacea.* *Bauh. pin.* 486.

*Erica coris folio* 8. *Clus. hist.* 1. p. 44.

*Habitat in Europa australiori.*

4. **ERICA** antheris bicornibus inclusis, corollis ovatis cinereis racemosis, foliis ternis glabris linearibus.



*Erica* foliis linearibus ternis, floribus globoso oblongis laxè spicatis. *Gutt. stamp.* 2. p. 110.

*Erica* foliis acerosis glabris ternis, corollis oblongo-ovatis staminibus longioribus verticillato-racemosis. *Læfl. it.* 137. \*

*Erica* humilis, cortice cinereo arbuti flore. *Bauh. pin.* 486.

*Erica* coris folio 6. *Clus. hist.* 1. p. 47. *Ord. dan.* 1. 38. *Habitat in Europa media, Oriente.* ‡

scoparia.

5. *ERICA* antheris bicornibus inclusis, corollis campanulatis longioribus, foliis ternis patentibus, ramis albis.

*Erica* erecta, foliis ternis, floribus ad alas sessilibus verticillatis. *Sauv. monsp.* 170.

*Erica* major scoparia, foliis deciduis. *Bauh. pin.* 485.

*Erica* scoparia, flosculis herbaceis. *Lob. ic.* 2. p. 215. *Sauv. monsp.* 46.

*Habitat* Montpelii, in Hispania, & Europa australi. ‡

castra.

6. *ERICA* antheris bicornibus inclusis, corollis subovatis mediocribus solitariis, foliis ternis, caule arboreo.

*Erica* foliis lineari-subulatis ternis, corollis ovatis, calycibus acutis patulis. *Hort. cliff.* 148.

*Habitat in Æthiopia.* ‡

viridi purpurea.

7. *ERICA* antheris bicornibus inclusis, corollis ovaris longioribus, racemis corymbosis secundis, foliis ternis.

*Erica* foliis lanceolatis oppositis imbricatis, floribus uno versu racemosis. *Hort. cliff.* 148. *Sauv. mansp.* 170.

*Erica* major, floribus ex herbaceo purpureis. *Bauh. pin.* 485.

*Erica* corios folio 3. *Clus. hist.* 1. p. 42.

*Habitat in Lusitania.* ‡

Tetralix.

8. *ERICA* antheris bicornibus inclusis corollis subglobosis confertis folio longioribus, foliis quaternis ciliatis patentibus.

*Erica* foliis subulatis ciliatis quaternis, corollis globoso ovatis terminalibus confertis. *Hort. cliff.* 148. *H. fr. c.* 310, 337. *Roy. lugdb.* 442.

*Erica* ex rubro nigricans scoparia. *Bauh. pin.* 486.

*Habitat in Europæ borealis paludibus cespitosis.* ‡

arborea.

9. *ERICA* antheris bicornibus inclusis, corollis campanu-

nulatis longioribus, foliis quaternis patentissimis, caule subatboreo tomento o.

*Erica erecta*. foliis quaternis, floribus spicato-racemosis. *Sauv. mon* p. 170. 46.

*Erica* foliis acerosis linearibus patentissimis, corollis ovatis, staminibus brevibus. *Hort. cliff.* 47. *Roy. lugdb.* 442.

*Erica maxima alba*. *Bauh. pin.* 485.

*Habitat in Europa australi.* ♀

10. *ERICA* foliis in summitate quinis, caule procumbente. *Segu. ver.* 280. purpurea-  
fcens.

*Erica procumbens dilute purpurea*. *Bauh. pin.* 486. *Raj. hist.* 1715.

*Erica* foliis corios, flore purpureo dilutionis coloris. *Bauh. hist.* 1. p. 358.

*Erica coris folio VII.* *Clus. hist.* 1. p. 43.

*Habitat in Europa australi.* ♀

\* \* *Antheris muticis obtusis emarginatis.*

11. *ERICA* antheris simplicibus inclusis, corollis ovatis ciliatis irregularibus, floribus terno-racemosis, foliis ternis ciliatis.

*Erica* foliis ovatis ciliatis ternis, corollis ovatis apice tubulosis irregularibus verticillato-racemosis. *Less. it.* 138. \*

*Erica hirsuta anglica*. *Bauh. pin.* 602.

*Erica XII.* *Clus. hist.* 1. p. 46.

*Habitat in Lusitania.* ♀

*Planta suffruticosa, bipedalis. Folia terna, patentia, sessilia, ovato-oblonga, acuta, margine reflexa, ciliata. Racemus terminalis, terno verticillatus. Corolla magna, ovata, ore contracto inaequali.*

12. *ERICA* antheris bifidis exsertis, calycibus duplicatis imbricata, coloratis acutis corolla longioribus, foliis ternis laevibus.

*Habitat in Aethiopia.* ♀

*Calyx albus est.*

13. *ERICA* antheris simplicibus bifidis exsertis, corollis cylindricis, foliis quinis patentibus. multiflora.

*Erica*, foliis corios, multiflora. *Bauh. hist.* 1. p. 356.

*Raj. angl.* 3. p. 471. \*

*Erica maxima purpurascens, longioribus foliis. Bauh. pin. 485.*

*Erica juniperifolia dense fruticans narbonensis. Lob. hist. 620. Garid. aix. 160. t. 32. Sauv. monsp. 46.*

*Habitat in Aeglia, G. Narbonensi, Oriente. §*

carnea.

14. *ERICA* antheris simplicibus exsertis, corollis ovatis sublongioribus, foliis quaternis triangularibus patentibus glabris.

*Erica procumbens, ternis foliis, carnea. Bauh. pin. 486.*

*Erica coris folio IX. Clus. hist. 1. p. 44.*

*Habitat in Pannoniæ, Helvetiæ, Baldi, Tridentini arenosis. §*

bruniades.

15. *ERICA* antheris bifidis exsertis, corollis globosis calyce lanato testis, foliis ternis subciliatis patentibus.

*Erica promont. b. spei floribus albidis, staminibus rubris. Pet. gaz. 5. t. 2. f. 9.*

*Eriocephalus bruniades ericaformis monomopatenensis, capitulis globulorum instar interius cavis & densa lanugine testis. Pluk. mant. 69. t. 347. f. 9.*

*Frutex africanus aromaticus. Seb. mus. 2. p. 64. t. 63. f. 7.*

*Habitat in Æthiopia. §*

*Folia linearia, remota, patula, pilis varis adpersa Calyces alba lana.*

capitata.

16. *ERICA* antheris bifidis exsertis, corollis globosis, calycibus lanatis coloratis, foliis ternis hirtis.

*Erica carnea promont. cap. spei foliis & floribus villosis. Pet. gaz. 5. t. 2. f. 10.*

*Erica africana, calyce lanuginoso ex viridi luteo, floribus concoloribus exiguis lanugine obfatis. Seb. mus. 1. p. 30. t. 20. f. 1.*

*Erica capitata nodiflora, globulis lanugine ex flavo-virescente obductis Raj. dendr. 98.*

*Habitat in Æthiopia. §*

*Dignoscitur floribus globosis, lanugine luteo-viridi obductis.*

plukenetii.

17. *ERICA* antheris bifidis exsertis, corollis subcylindricis subtruncatis longioribus, calycibus simplicibus, foliis ternis linearibus.

Cha-

*Chamæpithys æthiopica*, foliis læte virentibus, flore oblongo phœniceo plus quam eleganti s. plusquene-  
tii. *Pluk. mant.* 45. t. 344. f. 3. *Seb. mus.* 2. t. 25.  
f. 5.

*Habitat in Æthiopia.* ♀

*Folia terna, linearia, conferta.* Flores pedunculati, mutantes. Calyx corolla dimidio brevior, subulatus, carinatus. Corolla cylindrica, longa, ore lævissime quadrifido. Stamina fere membranacea, linearia, obtusa, corollæ duplo longiora.

18. *ERICA* antheris bifidis exsertis, corollis clavatis coccinea, quadrifidis, calycibus imbricatis, foliis ternis glabris.

*Erica africana angustifolia*, floribus oblongis tubulosis dependentibus coccineis, cum longissimis filamentis concoloribus. *Raj. dendr.* 98. *Seb. mus.* 1. p. 32. t. 21. f. 4.

*Habitat in Æthiopia.* ♀

*Statura E. plukenetii, sed Caulis tomentosus. Folia linearia, brevia. Pedunculi cassiores, breviores. Calyx imbricatus squamis octo, ovatis, per paria oppositis, nec intimis quaternis. Corolla præcedentis, sed profundius divisa. Stamina prioris, corolla duplo longiora, nec Antheris distinguendis a filamentis.*

19. *ERICA* antheris bifidis exsertis, corollis clavato-cylindricis curvis solitariis, foliis ternis glabris.

*Erica foliis linearibus quaternis oppositis villosis, corollis longissimis solitariis.* *Hort. cliff.* 148.

*Erica fruticosa capensis.* *Seb. mus.* 1. p. 32. t. 20. f. 4.

*Habitat in Æthiopia.* ♀

20. *ERICA* antheris bifidis inclusis, corollis clavato-tubiflora, cylindricis incurvatis pubescentibus acutis, foliis quaternis ciliatis.

*Erica spicata, floribus oblongis ex carneo purpureis.* *Pluk. mant.* 67. t. 346. f. 9.

*Erica africana frutescens.* *Seb. mus.* 2. p. 20. t. 19. f. 5.

*Habitat in Æthiopia.* ♀

*Folia linearia, lævia: summa ciliata. Flores terminales, solitarii, sessiles. Corollæ purpureæ, acuminatæ, pubescentes, multoties calyce longiores.*

21. *ERICA* antheris bifidis inclusis, corollis subclavatis cerinthodes, hirtis, floribus aggregatis sessilibus, foliis quaternis ciliatis.

*Erica africana*, flore purpureo. *Barth. ast.* 2. p. 57.

*Erica coris folio hispido*, *cerinthoides africana*. *Breyn. cent.* 25. t. 33. *Seb. mus.* 2. t. 22. f. 4. *U t.* 34. f. 6.

*Habitat in Æthiopia.* h

*Frutex ramis compositis. Folia quaterna, oblonga, convexa, levia, subius fulco exarata, spinulis ciliata. Flores magni, ad latera aggesti in capitulum sessiles. Calyx hirtus pilis albis. Corolla, late sanguinea, pilis albis hirta, ore obsolete quadrifida.*

abietina.

22. *ERICA* antheris bifidis inclusis, corollis ventricoso-cylindricis glabris, floribus lateralibus, foliis quaternis ciliatis.

*Erica foliis subulatis glabris quinis pluribusque verticillatis, floribus longissimis terminalibus confertis. Hort. cliff.* 148.

*Erica africana*, abieris folio longiore & tenuiore, floribus oblongis saturate rubris. *Raj. dendr.* 95. *Seb. mus.* 1. p. 31. t. 21. f. 1.

*Habitat in Æthiopia.* h

*Frutex determinate ramosus. Folia quaterna, oblongo-subulata, levia, canaliculata, acuta. Flores laterales, pedunculis solitariis ex alis foliorum. Calyx foliolis ovatis, purpureis. Corollæ longæ, similes E. curvisloræ, sed rectæ, ventricosæ, glaberrimæ obtusæ.*

pentaphylla.

23. *ERICA* antheris bifidis inclusis, corollis ovatis acutis, foliis quinis lævibus.

*Erica africana frutescens, juniperi folio, flore urceolari brevissimo. Seb. mus.* 1. p. 32. t. 21. f. 2.

*Habitat in Æthiopia.* h

pubescens.

24. *ERICA* antheris bifidis inclusis, corollis subovatis pubescentibus, foliis quaternis linearibus hispidis.

*Habitat in Æthiopia.* h

parviflora.

25. *ERICA* antheris bifidis inclusis, corollis subrotundis longitudine calycis, foliis quaternis ciliatis.

*Habitat ad Cap. b. spel.* h

*Folia linearia, incurvata, hispida. Flores inter folia terminalia conniventia, foliis breviores.*

phyiodes.

26. *ERICA* antheris bifidis inclusis, corollis ovatis inflatis glabris, foliis quaternis linearibus glabris obtusis.

Ha-

*Habitat in Æthiopia. h*

*Folia patula, lato-linearia, carinata. Flores ad apices ramorum quasi umbellati. Calyx ovatus, glaber, coloratus, corolla triplo brevior.*

27. *ERICA* antheris bifidis inclusis, corollis inflatis *halicacaba*. maximis, foliis ternis linearibus lævibus. *Amæn. acad. 6. afr. 15. \**

*Habitat in Æthiopia. h*

*Frutex alius: Ramis purpurascens; Ramulis submentosis, albis. Folia tria, confertissima, lævia, margine scabra. Pedunculi foliis breviores. Corolla purpurascens, glabra, ab omnibus distincta magnitudine & facie calycis Ihyfalidis annuæ.*

28. *ERICA* antheris bifidis inclusis, corollis subovatis *calycina*. calyces coloratos æquantibus, foliis ternis lævibus. *Erica africana humilis, flore albo urceolari, coris folio, calyce amplo. Seb. mus. 2. p. 13. t. 11. f. 7.*

*Habitat in Æthiopia. h*

*Calyx & Bractæe albæ. Corollæ luteæ.*

29. *ERICA* antheris bifidis inclusis, corollis ovatis *tenuifolia*. calyces coloratos æquantibus, foliis oppositis linearibus appressis.

*Erica africana tenuifolia, unedonis flore, foliis longioribus cauli adpressis. Pluk. mant. 68. t. 346. f. 11.*

*Habitat in Æthiopia. h*

30. *ERICA* antheris bifidis inclusis, corollis ovatis subfessilibus, foliis quinis obtusis ciliatis. *empetrifolia*.

*Erica africana, coridis folio, pilosa, flosculis minutissimis purpureis inter ramulos dispersis. Pluk. mant. 68.*

*Habitat in Æthiopia. h*

31. *ERICA* antheris bifidis inclusis, corollis globosis *pilulifera*, glabris quaternis, foliis quaternis setaceis patulis.

*Habitat in Æthiopia. h*

32. *ERICA* antheris bifidis inclusis, corollis ovatis, *corifolia*. calycibus planis, bracteis coloratis, foliis ternis glabris.

*Erica antheris bifidis exsertis, corollis globosis medio-cribus, pedunculis triphyllis, foliis quaternis. Syst. nat. 2. p. 355.*

Eri-

*Erica africana* glabra fruticosa, coris folio, arbuti flore dilute purpureo. *Seb. mus.* 1. p. 32. t. 21. f. 3.

*Erica rosmarini* foliis incanis, æthiopica, floribus arbutis magno calyce inclusis. *Pluk. mant.* 68. t. 346. f. 5.

*Erica capensis*, coridis folio, flore rubello, *Pct. gaz.* 7. t. 3. f. 7.

*Habitat in Æthiopia.* ♀

*Caulis fruticosus, compositus. Folia ternaria, linearia, glabra. Flores terminales s. umbellulae, sessiles. Bracteae in singulo pedunculo tres, alternæ, ovatae, purpureae, magnitudine 3 facie folioli calycini. Calyx patens, foliolis ovatis, purpureis. Corolla ovata, purpurea.*

triflora.

33. *ERICA* antheris bifidis inclusis, corollis subglobosis subvillosis ternis, pedunculis triphyllis, foliis ternis.

*Habitat in Æthiopia.* ♀

gnapholodes.

34. *ERICA* antheris bifidis inclusis, corollis globosis ternis, calycibus duplicibus obtusis.

*Habitat in Æthiopia.* ♀

planifolia.

35. *ERICA* antheris bifidis inclusis, corollis campanulatis, foliis quaternis ciliatis patentibus.

*Erica africana* hirsuta, thymi foliis ternis. *Pluk. mant.* 69. t. 347. f. 1.

*Habitat in Æthiopia.* ♀

paniculata.

36. *ERICA* antheris bifidis inclusis, corollis campanulatis subumbellatis, foliis quaternis linearibus lævibus.

*Erica tenuifolia*, flosculis suave rubellis. *Pluk. alm.* 136. t. 279. f. 2.

*Erica africana*, angustissimis foliis, flosculis in capitulum congestis. *Seb. mus.* 2. p. 46. t. 44. f. 4.

*Habitat in Æthiopia.* ♀

*Frutex. Folia linearia, erecta, fere lævia. Flores parvi, purpurei, uti etiam calyces 3 pedunculi minimi, adeo copiosi, ut totam tegant plantam.*

spumosa.

37. *ERICA* antheris bifidis inclusis, calycibus trifloris: squamis inflatis inflexis, foliis ternis lævibus. *Amman. acad.* 6. afr. 14.

*Erica africana*, flore rubro plano. *Seb. mus.* 2. p. 11. t. 9. f. 10. mala.

*Habitat in Æthiopia.* ♀

*Æthiopicas has omnes ad me misit inspiciendas ex Herbario Oldenlandiano, egregii thesauri, de re herbaria optime meritus, possessor C. Burmannus.*

\* \* \* *obscura.*

28. *ERICA* racemo terminali, foliis alternis subtus tomentosis. dabocil.

*Erica cantabrica*, flore maximo, foliis myrti subtus incanis. *Tournef. inst.* 603.

*Erica* f. *Dabeci hibernis*. *Raj. dendr.* 98.

*Erica hibernica*, foliis myrti pilosis subtus incanis. *Pet. gar.* 42. t. 27. f. 4.

*Vaccinium cantabricum*. *Hudson. angl.* 143.

*Habitat in Hiberniæ Gallovidia in montibus Mayo, solo cespitoso.* §

*Statura* § *magnitudo Andromedæ polifoliæ*. *Caules fruticosi, pilosi, ferruginei. Folia alterna, Andromedæ, lanceolata, subpetiolata, lateribus revoluta, subtus albo tomentosa, supra pilis rarissimis adspersa. Rami breviores. Racemus longus, terminalis. Calyx quadripartitus, minimus. Capsula erecta, quadrilocularis, quadrivalvis, magna. Flores non vidi, quos describant autoptæ.*

## DAPHNE.

\* *Floribus lateralibus.*

1. *DAPHNE* floribus sessilibus ternis caulinis, foliis lanceolatis deciduis. *Mezereum.*

*Daphne floribus sessilibus infra folia elliptico-lanceolata.*

*Fl. lapp.* 140. *Fl. suec.* 311, 338. *Mat. med.* 179.

*Hort. cliff.* 147. *Roy. lugdb.* 201.

*Laureola folio deciduo, flore purpureo, officinis Laureola femina.* *Bauh. pin.* 462.

*Daphnoides.* *Cam. epit.* 937.

*Habitat in Europæ borealis sylvis.* §

2. *DAPHNE* floribus sessilibus axillaribus, foliis lanceolatis, caulibus simplicissimis. *Thymelæa.*

*Daphne floribus tetrandris secundum caules simplicissimos.* *Sauv. monsp.* 56.

*Thymelæa foliis polygalæ glabris.* *Bauh. pin.* 463.

*Thymelæa alpina glabra, flosculis subluteis ad foliorum ortum sessilibus.* *Pluk. alm.* 266. t. 229. f. 2.

Sa-



*Sanamunda viridis vel glabra. Bauh. prodr. 160.*

*Sanamunda glabra. Bauh. hist. 1. p. 592.*

*Habitat in Hispania & Montspeli in Horto. De. 5*

villosa.

3. *DAPHNE* foliis sessilibus lateralibus solitariis, foliis lanceolatis planis ciliatis pilosis confertis.

*Thymelæa villosa minor lusitanica, polygoni folio.*

*Tournef. inst. 594.*

*Habitat in Lusitania, Hispania, Afræner. 5*

*Frutex ramis alternis. Folia polygoni, lanceolata, vix petiolata, utrinque pilis albis remotis: paucioribus in pagina superiore, pluribus ad margines. Ex alis rudimenta plurima foliosa ramulorum, unde quasi verticillata. Flores angusti, parvi, foliis breviores.*

Tarton-raira.

4. *DAPHNE* floribus sessilibus aggregatis axillaribus, foliis ovatis utrinque pubescentibus nervosis. *Gron. orient. 125.*

*Thymelæa foliis candicantibus & serici instar mollibus.*

*Bauh. pin. 463.*

*Tarton-Raire galloprovinciæ monspeliensium. Lob. ic. 371.*

*Habitat in Galloprovincia. 5*

alpina.

5. *DAPHNE* floribus sessilibus aggregatis lateralibus, foliis lanceolatis obtusiusculis subtus tomentosis.

*Daphnes aliud rarum, foliis supremis hirsutis. Gen. fasc. 6. t. 3. f. 7.*

*Thymelæa cantabrica juniperifolia, ramulis procumbentibus. Tournef. inst. 595.*

*Thymelæa floribus inter folia, folio utrinque hirsuto.*

*Hall. helv. 187. Sauv. monp. 57.*

*Chamelæa alpina, folio inferne incano. Bauh. pin. 1462.*

*Chamelæa alpina incana. Lob. ic. 376.*

- β. *Chamelæa sabaudica, folio utrinque incano, flore albo. Raj. hist. 1583.*

*Thymelæa, incana, mezerei folio & facie, surculis admodum fragilibus. Pluk. alm. 366. t. 229. f. 3.*

*Habitat in Alpibus Helvetiæ, Genevæ, Italiæ, Austriæ. 5*

laureola.

6. *DAPHNE* racemis axillaribus, foliis lanceolatis glabris.

*Daph-*

*Daphne racemis lateralibus, foliis lanceolatis integris.*  
*Hort. cliff. 147. Hort. upf. 94. Mat. med. 180. Sauv.*  
*monsp. 57.*

*Laureola sempervirens, flore viridi, quibusdam Lau-*  
*reola mas. Bauh. pin. 662.*

*Laureola. Dod. pempt. 365.*

*Habitat in Anglia, Helvetia, Gallia, Baldo. ‡*

7. *DAPHNE* pedunculis lateralibus bifloris, foliis lanceolato-ovatis. pontica.

*Thymelæa pontica, citrei foliis. Tournesf. itin. 2. p.*  
*180. t. 180.*

*Habitat in Ponto. ‡*

\* \* *Floribus terminalibus.*

8. *DAPHNE* capitulo terminali pedunculato, foliis oppositis oblongo-ovatis glabris. *India.*

*positis oblongo-ovatis glabris. Osh. it. 246.*

*Habitat in China. ‡*

*Frutex parvus. Folia opposita, oblongo ovata, integerrima, glabra. Pedunculus terminalis, brevissimus, apice gerens flores 6. ad 8. sessiles.*

9. *DAPHNE* floribus congestis terminalibus sessilibus, *Cneorum.*  
*foliis lanceolatis nudis.*

*Daphne humifusa, foliis oblongis flores sessiles terminales subcingentibus. Sauv. monsp. 57. Duham. arb.*  
*2. t. 94.*

*Thymelææ affinis facie externa. Bauh. pin. 463.*

*Cneorum. Matth. hist. 46. Clus. hist. 89.*

*Habitat in Helvetia, Hungaria, Pyrenæis, Baldo, L.*  
*B. Munchhausen. ‡*

10. *DAPHNE* panicula terminali, foliis lineari-lanceolatis acuminatis.

*Daphne foliis lanceolatis basi angustioribus, racemo nudo terminali. Sauv. monsp. 56.*

*Daphne floribus racemosis, foliis lineari-lanceolatis acuminatis integris. Guett. stamp. 2. p. 427.*

*Thymelæa foliis lini. Bauh. pin. 463.*

*Thymelæa. Clus. hist. 1. p. 87. Cam. epit. 974.*

*Habitat in Hispania, Italia, G. Narbonensi. ‡*

11. *DAPHNE* floribus terminalibus pedunculatis, foliis sparsis linearibus patentibus mucronatis.

11. *THY-*

*Thymelæa capitata lanuginosa, foliis creberrimis minimis aculeatis. Burm. afr. 134. t. 49. f. 1.*  
*Habitat in Æthiopia. h*

## DIRCA.

- palustris. 1. *DIRCA. Amæn. acad. 3. p. 12. t. 1. f. 7. Kalm. it. 3. p. 88. Duham. arb. 1. t. 212.*  
*Thymelæa floribus albis primo vere erumpentibus, foliis oblongis acuminatis, viminibus & cortice valde tenacibus. Gron. virg. 155.*  
*Habitat in Virginie paludosis. h*

## GNIDIA.

- pinifolia. 1. *GNIDIA foliis sparsis lineari subulatis, floribus verticillatis, floribus aggregatis terminalibus.*  
*Rapunculus foliis nervosis linearibus, floribus argenteis non galeatis. Burm. afr. 112. t. 41. f. 3.*  
*Habitat in Æthiopia. h*
- tomentosa. 2. *GNIDIA foliis alternis ovato-lanceolatis.*  
*Habitat in Æthiopia. h*  
*Flores tomentosi, capitati, terminales.*
- oppositifolia. 3. *GNIDIA foliis oppositis lanceolatis.*  
*Habitat in Æthiopia. h*  
*Frutex ramis rectis. Folia opposita, lanceolata, glabra, integerrima. Flores terminales, conferti, infundibuliformes, tubo villosi, petalis emarginatis.*

## STELLERA.

- Passerina. 1. *STELLERA foliis linearibus, floribus quadrifidis.*  
*Amæn. acad. 1. p. 400. Sauv. mosp. 69.*  
*Passerina foliis linearibus, floribus laxè spicatis alternis in foliorum binorum alis sessilibus. Guett. Jlamp. 25.*  
*Dalib. parif. 118.*  
*Passerina. Bauh. hist. 3. p. 456.*  
*Lithospermum, liniaræ folio, germanicum. Bauh. pin. 259.*  
*Linaria alterna botryoides montana. Column. ecphr. 1. p. 82. t. 80.*  
*Habitat in Italiæ, Galliæ, Germaniæ campis aridis apricis.*

2. STELL-

2. **STELLERA** foliis lanceolatis, floribus quinquesi- Chamæjasme  
dis. *Amæn. acad.* 1. p. 400. \*  
Chamæjasme radice mandragoræ. *Amm. ruth.* 16. t. 2.  
*Habitat in* Sibiria. *Gmelin* 24

## PASSERINA.

1. **PASSERINA** foliis linearibus convexis quadrifariam filiformis.  
imbricatis, ramis tomentosis.  
*Passerina* foliis linearibus. *Hort. cliff.* 246. t. 11. \*  
*Roy. lugdb.* 208.  
*Thymelæa æthiopica* fruticosa, foliis in longum striatis,  
furculis valde tomentosis. *Pluk. alm.* 180.  
*Thymelæa æthiopica*, passerinæ foliis. *Breyn. cent.* [10.  
t. 6.  
*Habitat in* Æthiopia. §
2. **PASSERINA** foliis carnosis extus glabris, caulibus hirsuta.  
tomentosis.  
*Thymelæa tomentosa*, foliis sedî minoris. *Bauh. pin.*  
461.  
*Sanamunda* 3. *Clus. hist.* 1. p. 89. *Breyn. cent.* t. 19.  
*Sesamoides parvum dalechampi.* *Bauh. hist.* 1. p. 595.  
*Habitat in* Galloprovincia, Italia, Oriente. §
3. **PASSERINA** foliis ovatis tomentosis, floralibus sericea,  
quaternis, caule hirsuto, filamentis bifurcatis. *Amæn.*  
*acad.* 4 p. 313. \*  
*Thymelæa sericea*, foliis oblongis, floribus tubulosis  
angustissimis. *Burm. afric.* 135. t. 49. f. 2.  
*Habitat in* Æthiopia. §
4. **PASSERINA** foliis lanceolatis glabris acuminatis ad- dodecandra.  
pressis, floribus dodecandris sessilibus, staminibus  
quatuor inclusis. *Amæn. acad.* 4 p. 271. \*  
*Thymelæa* foliis oppositis cruciatis. *Burm. afr.* 127.  
t. 147. f. 1, 2.  
*Habitat in* Æthiopia. §
5. **PASSERINA** foliis ovatis glabris acutis, floribus ob- lavigata.  
tusis. *Amæn. acad.* 4 p. 312. \*  
*Thymelæa* foliis planis acutis, coma & floribus purpu-  
reis. *Burm. afric.* 137. t. 49. f. 2.  
*Habitat in* Æthiopia. §
7. **PASSERINA** foliis lanceolato-linearibus glabris, flo- capitata.  
*Linnaei Spec. plant. pars I.* K k ribus

ribus capitatis: receptaculo incrassato. *Amæn. acad.*  
6. afr. 16. \*

*Thymelæa* foliis linearibus alternis, floribus ex uno petiolo copiosis. *Burm. afric.* 133. t. 48. f. 3.

*Habitat ad Cap. b. Spei.* ‡

7. *PASSERINA* foliis lanceolatis subciliatis erectis, ramis nudis.

*Passerina* foliis lanceolatis. *Hort. cliff.* 146. *Roy. lugdb.* 208. *Gron. orient.* 126.

*Thymelæa* foliis chamæleæ minoribus hirsutis. *Bauh. pin.* 463.

*Thymelæa* foliis oblongis acutis ad oras fimbriatis. *Burm. afric.* 129. t. 47. f. 2.

*Sanamunda* 1. *Clus. hist.* 1. p. 88.

*Erica africana*, rusci folio. *Seb. mus.* 2. p. 15. t. 12. f. 9.

*Habitat in Æthiopia, Hispania, Oriente.* ‡

8. *PASSERINA* foliis oppositis linearibus, floribus solitariis terminalibus, ramis glabris.

*Thymelæa* foliis triquetris cruciatim oppositis, flore sericeo. *Burm. afr.* 132. t. 48. f. 2.

*Thymelæa* ramosa, linearibus foliis angustis, flore solitario. *Burm. afric.* 131. t. 48. f. 1.

*Habitat in Æthiopia.* ‡

## LACHNEA.

eriecephala. 1. *LACHNEA* capitulis solitariis lanatis, foliis quadrifariam imbricatis.

*Habitat in Æthiopia.* ‡

Folia *trigona gibba*. Capitula *terminalia*, *solitaria*, *tomentosa*, *cincta lana candida*.

conglomerata. 2. *LACHNEA* capitulis confertis, foliis laxis.

*Thymelæa africana*, polii angustioribus foliis, floribus in capitulis summo caule conglomeratis. *Pluk. mant.* 179. *Raj. dendr.* 54.

*Sanamunda* 3. *Clusii* affinis, polii folio. *Breyn. Cent.* 18. t. 7.

*Habitat ad Cap. b. Spei.* ‡

## BÆCKEA.

frutescens. 1. *BÆCKEA*. *Osbeck. it.* 231. t. 1. \*

Ha-

*Habitat in China.* † *D. D. Abrah. Bæck S. R. M. Sociæ Archiater, & Amicus noster vere sincerus, e cujus horto feci purissimas rariores stirpes obtinui-mus, etiam hanc primus nobiscum communicavit.*

*Frutex habitu abrotani, ramis virgatis: Ramulis oppositis, brevibus, simplicibus. Folia opposita, linearia, acuta, glabra, integerrima. Flores axillares, solitarii: pedunculo rudo longitudine floris, foliis longe brevior. Perianthium infundibuliforme, quinque-dentatum. Petala 5, subrotunda, calyci inserta. Filamenta 8, quorum 6 paria, & 2 solitaria, brevissima, inflexa. Antheræ subovatae, parvæ. Germen subrotundum. Stylus filiformis, corolla brevior. Stigma capitatum. Chinenfibus Tjongina. Osbeck.*

## DIGYNIA.

### GALENIA.

- I. *GALENIA. Hort. cliff. 150. Roy. lugdb. 209. africana.*  
*Sherardia. Pont. epist. 14.*  
*Kali lignosum, flore muscoso, rosmarini folio. Bocc. musf. 150. t. 110.*  
*Atriplex africana lignosa frutescens, rosmarini foliis. Till. pisf. 20. t. 15.*  
*Habitat in Africa. †*

### WEINMANNIA.

- I. *WEINMANNIA.*  
*Windmannia fruticosa, foliis subrotundis serratis per pinnata. pinnae cordato alatas, racemis terminalibus, pinnis & ramis oppositis. Brown. jam. 212.*  
*Habitat in Jamaica. †*

### MOEHRINGIA.

- I. *MOEHRINGIA.*  
*Alfina montana, capillaceo folio. Bauh. pin. 251. muscosa. Scheuch. alp. 503. t. 6. Pluk. alm. 23. t. 75. f. 1.*  
*Alfina saxifraga angustifolia minima montana. Col. ecphr. 1. p. 292. t. 290.*  
 β. *Alfina tenuifolia muscosa. Bauh. pin. 251. Moris. blesf. 10. hist. 2. p. 551. f. 5. t. 23. f. 12.*

*Habitat in Alpibus Helvetiæ, Italiæ, Austriæ, Mos-  
peli ad fontes & scaturigines.*

## TRIGYNIA. POLYGONUM

\* *Atraphaxoides caule frutescente.*

- frutescens.* 1. **POLYGONUM** caule fruticoso, calycinis foliolis  
duobus reflexis. *Hort. upf.* 95.  
*Atraphaxis inermis, foliis planis. Hort. cliff.* 138. *Roy.*  
*lugdb.* 209.  
*Lapathum orientale, frutex humilis, flore pulchro.*  
*Tournef. cor.* 38.  
*Lapathum dauricum montanum fruticans, ramis late*  
*iparlis. Amm. ruth.* 227.  
*Habitat in Sibiria, Dauria. §*

\* \* *Bistorta spica unica.*

- Bistorta.* 2. **POLYGONUM** caule simplicissimo monostachyo,  
foliis ovatis in petiolum decurrentibus. *Mat. med.* 188.  
*Bistorta foliis ovato-oblongis acuminatis Hort. cliff.*  
*150. Hort. upf.* 95. *Roy. lugdb.* 217.  
*Bistorta major, radice magis intorta. Bauh. pin.* 192.  
*Bistorta. Cam. epit.* 683.  
*Habitat in montibus Helvetiæ, Austriæ, Galliæ. ¶*
- viviparum.* 3. **POLYGONUM** caule simplicissimo monostachyo,  
foliis lanceolatis.  
*Bistorta foliis lanceolatis. Fl. lapp.* 152. *Fl. Suec.* 321;  
*340. Hort. cliff.* 150.  
*Bistorta montana minor, radice intorta inodora, flo-*  
*sculis in spicæ cacumine albis sterilibus, inferna spi-*  
*cæ parte tuberculis proliferis turbinatis puniceis fe-*  
*cunda. Amm. ruth.* 169.  
*Bistorta alpina media. Bauh. pin.* 192. *Oed. dan. t.* 13.  
*Bistorta alpina minor. Bauh. pin.* 192,  
*Bistorta alpina minima. Hall. opusc.* 234.  
*Habitat in Europæ subalpinis pascuis duris. ¶*

\* \* \* *Perficaria pistillo bifido.*

- virginiana-*  
*num.* 4. **POLYGONUM** floribus pentandris semidigynis,  
corollis quadrifidis inæqualibus, foliis ovatis.  
Per-

- Perficaria floribus pentandris digynis, corolla quadrifida inæquali. *Wach. ultr.* 258.
- Perficaria florum staminibus quinis, stylo duplici, corolla quadrifida inæquali. *Hort. cliff.* 42. *Gron. virg.* 43. *Roy. lugdb.* 216.
- Perficaria frutescens maculosa virginiana, flore albo. *Morif. hist.* 2. p. 589. *Raj. hist.* 183.
- Habitat in Virginia.* 2.
5. POLYGONUM floribus pentandris semidigynis, staminibus corollæ regulari æqualibus. *Lapathifolium.*
- Perficaria floribus pentandris digynis, corolla regulari staminibus æquali. *Wach. ultr.* 257.
- Perficaria florum staminibus quinis semidigynis, stylo bifido corollæ regulari æquantibus. *Hort. cliff.* 42.
- Perficaria major, lapathi foliis, calyce floris purpureo. *Tournef. inst.* 510. *Raj. suppl.* 119.
- Perficaria Hydropiper. *Lob. ic.* 315.
- Habitat in Gallia.*
6. POLYGONUM floribus pentandris semidigynis, amphibium. staminibus corolla longioribus.
- Perficaria florum staminibus quinis corollam superantibus, stylo bifido. *Hort. cliff.* 41. *Roy. lugdb.* 216.
- Perficaria floribus pentandris digynis, corolla staminibus brevior. *Fl. suec.* 318; 341.
- Potamogeton salicis folio. *Bauh. pin.* 193.
- Habitat in Europa.*
7. POLYGONUM foliis crenatis. † *seriatum.*
- Polygonum folio oblongo crenato. *Shaw. afr.* 489.
- Habitat in Mauritania.*
8. POLYGONUM floribus pentandris trigynis, foliis ocreatum. lanceolatis.
- Perficaria spicis longis numerosis, vaginis integris, floribus pentandris trigynis. *Gmel.*
- Habitat in Sibiria.* *D. Gmelin.*
- Stipulæ sunt vaginae integræ, ventricosæ, læves, ampliores multo quam caulis. Folia lanceolata, revoluta. Flores racemosi.
9. POLYGONUM floribus hexandris, semidigynis, Hydropiper foliis lanceolatis, stipulis submuticis.



*Polygonum foliis lanceolatis, floribus hexandris semidigynis. Mat. med. 187.*

*Perficaria floribus hexandris semidigynis. Fl. suec. 320; 343.*

*Perficaria florum staminibus senis, stylo bifido. Hort. cliff. 46. Roy. lugdb. 216.*

*Perficaria urens f. Hydropiper. Bauh. pin. 101.*

*Hydropiper. Dod. pempt. 607.*

*Habitat in Europæ subhumidis. ☉*

*Perficaria. 10. POLYGONUM floribus hexandris digynis, spicis ovato-oblongis, foliis lanceolatis, stipulis ciliatis.*

*Perficaria floribus hexandris digynis. Fl. suec. 319; 342*

*Perficaria florum staminibus senis, stylo duplici. Hort. cliff. 42. Gron. virg. 157. Roy. lugdb. 216.*

*Perficaria mitis. Bauh. hist. 3. p. 779. Fl. lapp. 71.*

*Perficaria mitis maculosa & non maculosa. Bauh. pin. 101.*

*Perficaria foliis angustissimis pene glabris, spinis interruptis rarissimis, vaginis ciliaribus. Hall. helv. 181. gatt. 29.*

*β. Perficaria angustifolia. Bauh. pin. 101. prodr. 43.*

*γ. Perficaria foliis subtus tomentosis. Hall. helv. 181. gatt. 28.*

*Habitat in Europæ cultis. ☉*

*Folia subtus tomentosa.*

*barbatum. 11. POLYGONUM floribus hexandris trigynis, spicis virgatis, stipulistruncatis setaceo-ciliatis, foliis lanceolatis.*

*Perficaria maderaspatana, longiore folio, hirsuta. Pluk. alm. 288. t. 10. f. 7?*

*Perficaria foliis ovatis glabris. Kraschen. comm. petr. 1747. p. 375. t. 13?*

*Perficaria procumbens longissima angustissima non maculosa, spica longiori laxiori & graciliore. Sloan. jam. 48. hist. 1. p. 17. t. 3. f. 1.*

*Velutia-modela-mucu. Rheed. mal. 12. p. 145. t. 77. (non 76.).*

*Habitat in China.*

*Caulis herbaceus, rufus. Stipulæ vaginales laxæ, pilosæ, truncatæ, ciliatæ, setis validis longitudine ferme ipsius stipulæ. Spicæ virgatæ, longæ: floribus remotis, tribus ex singula bractæa turbinata. Folia lanceolata.*

12. **POLYGONUM** floribus heptandris digynis, foliis ovatis, caule erecto, stipulis hirtis hypocrateriformibus. orientale.

*Perficaria* florum staminibus sex pluribusque, stylo duplici. *Hort. cliff.* 42. *Hort. upf.* 56. *Roy. lugdb.* 216.

*Parficaria* foliis ovato-lanceolatis acutis, foliis pentandris, caule erecto. *Mill. dict.* t. 201.

*Perficaria orientalis*, nicotianæ folio, calyce florum purpureo. *Tournef. cor.* 38. *Comm. rar.* 43. t. 43.

*Schovanna-modela muccu.* *Rheed. mal.* 12. p. 147. t. 76. (non 77.).

*Habitat in Oriente, India.* ☉

*Indica longe hirsutior est, quam orientalis.*

13. **POLYGONUM** floribus octandris digynis, pedunculis hispidis, foliis lanceolatis, stipulis muticis. pennsylvanicum.

*Habitat in Pennsylvania.* *Kalm.*

*Facies* ♂ *habitus* *Pol. Perficariæ*, sed majora strictiora omnia. *Caulis* ramosus. angulatus. *Folia* lanceolata, acuminata carina subtus admodum scabra. *Stipulae* truncatae, muticae. *Pedunculi* scabri, setis rigidiusculis apice gluten secernentibus. *Spicae* subracemosa, flosculis pedicellatis.

\* \* \* \* *Polygonum foliis indivisis, floribus octandris.*

14. **POLYGONUM** floribus octandris trigynis axillaribus, foliis ovali-lanceolatis sempervirentibus, caule suffrutescente. maritimum.

*Polygonum maritimum latifolium.* *Bauh. pin.* 281.

*Polygonum maritimum.* *Cam. epit.* 691.

*Habitat in littoribus M. mediterranei, Orientis, Virginiae.* ☿

*Affine* nimis *P. aviculari*, ut vix distinguatur, etiam floribus quaternis ♂ caule hinc planiore; sed majora durioraque omnia. *Folia* ovata, coriacea, sempervirentia, exsiccatione carulescentia. *Caulis* brevior, durus, perennans.

15. **POLYGONUM** floribus octandris trigynis axillaribus, foliis lanceolatis, caule procumbente herbaceo. aviculare.

*Polygonum caule procumbente herbaceo, foliis lanceolatis integerrimis.* *Mat. med.* 186.

*Polygonum.* *Hort. cliff.* 150. *Fl. succ.* 322; 339. *Gronov. arg.* 44. *Roy. lugdb.* 215. *Hall. helv.* 182.

*Polygonum latifolium.* *Bauh. pin.* 281.

- β. *Polygonum brevi angustoque folio. Bauh. pin. 281.*  
 γ. *Polygonum oblongo angusto folio. Bauh. pin. 281. Fl. lapp. 153.*  
 δ. *Polygonum angustis foliis, calycibus purpurascens. Dill. app. 65.*  
 ε. *Polygonum erectum humile, foliis orbis. Dill. app. 65.*  
*Habitat in Europæ cultis ruderalis. ☉*  
 erectum. 16. **POLYGONUM** floribus octandris trigynis axillaribus, foliis ovalibus, caule erecto herbaceo.  
*Habitat in Philadelphia, enata ex seminibus D. Kal-mii. ☉*  
 articulatum. 17. **POLYGONUM** floribus octandris trigynis, spicis articulatis paniculatis, stipulis vaginalibus truncatis.  
*Habitat in Canada. Kalm. ☉*  
*Radix fibrosa, parva. Caulis pedalis, ramis superne alternis; subdivisus in paniculam floriferam. Stipulæ solitariae, vaginales, superne latiores, integræ, truncatæ, nec fissæ. Folia linearia, glabra. Paniculam terminant spicæ filiformes, angustissimæ, imbricatæ bracteis vaginalibus, turbinatis, truncatis: ex sinu singulo oritur Pedunculus solitarius, capillaris, basi articulatus, coloratus, uniflorus. Corollæ incarnata, quinquepartita, fundo laxiore quam in P. aviculari, staminibus octo, Pistillo stylis 3.*  
 divaricatum 18. **POLYGONUM** floribus octandris trigynis racemosis, foliis lanceolatis, caule divaricato patulo.  
*Helxine foliis lanceolatis, caule diffuso. Hort. ups. 96.*  
*Fagopyrum orientale ramosum & multiflorum, persicariæ folio. Tournef. cor. 39. Buxb. cent. 2. p. 31. t. 31.*  
*Persicaria alpina, folio nigricante, floribus albis. B. All. pedem. 41. t. 8.*  
*Persicaria alpina altera saxatilis, foliis durioribus acutis. Bocc. mus. 2. p. 108. t. 83.*  
*Habitat in Sibiria, Corsica. 24*  
 \* \* \* \* \* *Helxine foliis subcordatis.*  
 chinense. 19. **POLYGONUM** floribus octandris trigynis, pedunculis scabris, foliis ovatis.  
*Habitat in India, China.*

*Caulis flexuosus. Folia ovato-cordata, f. lato-ovata, glabra utrinque. Pedunculi hispidi, apice sæpius trifidi. Flores capitati. Stipulæ vaginales, late, integerrimæ nec ciliatæ, ovatæ. Flores in capitulis subrotundis, pauci, terminales.*

20. **POLYGONUM** foliis triangularibus, caule aculeato, stipulis perfoliato-foliosis patentibus subrotundis. *perfoliatum*

*Habitat in India. Burmannus.*

*Similis P. sagittato* ♂ *arifolio. Caulis angulatus aculeis parvis recurvis, ut etiam Petioli. Stipulæ non vaginantes, sed patentes, foliaceæ, ovatæ f. subrotundæ, integerrimæ. Folia triangularia, basi parum sinuata, latiora quam longa, interdum petiata. Pedunculi ad apicem caulis, pauciflori, aculeati.*

21. **POLYGONUM** foliis sagittatis, caule aculeato. *Helxine caule aculeato, foliis sagittatis. Gron. virg. 44. sagittatum, Cold. noveb. 92.*

*Helxine caule erecto aculeis reflexis exasperato. Hort. cliff. 151. t. 12. \* Roy. lugdb. 211.*

*Fagotritico similis, angustiori folio, convolvuli modo scandens, caule spinulis deflexis densius obsito. Pluk. mant. 74. t. 394. f. 5.*

*Planta posterior e novo belgio. Laët. amer. 73. t. 74. Habitat in Virginie, Marilandie madidis.*

22. **POLYGONUM** foliis hastatis, caule aculeato. *Fagotritico similis spinosa scandens, ari folio latiore, floridana. Pluk. amalth. 87. t. 398. f. 3. arifolium.*

*Planta prior e novo belgio. Laët. amer. 73.*

*Habitat in Virginia, Florida.*

23. **POLYGONUM** foliis cordato-sagittatis, caule inermi erecto, seminibus subdentatis. *tataricum.*

*Helxine caule erectiusculo inermi, foliis cordato-sagittatis, seminibus subdentatis. Hort. upf. 96.*

*Helxine caule erecto inermi, foliis cordato-sagittatis, seminum angulis dentatis. Hort. cliff. 151.*

*Fagotriticum sibiricum. Aët. stockh. 1744. p. 117. t. 4.*

*Habitat in Tataria. ☉*

- Fagopyrum*. 24. **POLYGONUM** foliis cordato - sagittatis, caule erectiusculo inermi, seminum angulis æqualibus. Polygonum caule erectiusculo inermi ramofo, foliis cordato-sagittatis, feminibus lævibus. *Mat. med.* 188. Helxine caule erectiusculo inermi, foliis cordato - sagittatis, feminibus integerrimis. *Hort. upf.* 96. Helxine caule erectiusculo inermi, foliis cordato - sagittatis. *Hort. cliff.* 151. *Fl. suec.* 324, 345. *Roy. lugdb.* 217. *Erysimum cereale*, folio hederaceo. *Bauh. pin.* 27. *Ocimum cereale*. *Tabern. hist.* 276. *Habitat in Asia.* ☉
- Convolvulus*. 25. **POLYGONUM** foliis cordatis, caule volubili, floribus planiusculis. Helxine caule volubili. *Fl. lapp.* 134. *Fl. sve.* 323, 344. *Hort. cliff.* 150. *Gron. virg.* 157. *Roy. lugdb.* 217. *Convolvulus minor*, semine triangulo. *Bauh. pin.* 295. *Convolvulum nigrum*. *Dod. pempt.* 396. *Habitat in Europæ agris.* ☉ *Caules striati. Racemi ex alis. Flores etiam ad basin pedunculi communis. Semina calyce triquetro testâ.*
- dumetorum*. 26. **POLYGONUM** foliis cordatis, caule volubili lævi, floribus carinato - alatis. *Fagopyrum prælongum dumetorum*, feminibus alatis duplici more dispositis. *Dill. app.* 60. *Fagopyrum scandens altissimum dumetorum*, feminibus tribus alis pellucidis. *Rupp. jen.* 99. *Hall. helv.* 173. *Fagopyrum sylvaticum scandens*, flore foliaceo. *Pont. anth.* 265. *Fagopyrum majus scandens*. *Vaill. parif.* 52. *Habitat in Europæ australioris sylvis umbrosis.* ☉ *Folia cordata lobis baseos rotundatis. Caulis non striatus. Panicula e racemo composito, alternatim bifario.*
- scandens*. 27. **POLYGONUM** foliis cordatis, caule erecto scandente, petiolis basi subtus poro pertusis. Polygonum foliis cordatis, caule volubili, floribus carinatis. *Syst. nat.* 1. p. 364.

*Fagopyrum scandens americanum maximum. Tournef. inst. 511.*

*Fagopyrum scandens, caule rubente, semine nigro. Gron. virg. 44. Cold. noveb. 93.*

*Fagotriticum volubile majus virginianum. Pluk. alm. 143. t. 177. f. 7.*

*Fagopyrum scandens f. Volubilis nigra major, flore & fructu membranaceis compressis. Sloan. jam. 46. hist. 1. p. 138. t. 90. f. 1.*

*Habitat in America. ☉*

## COCOLOBA.

### 1. COCCOLOBA foliis cordato-subrotundis nitidis. *Uvisera.*

*Coccoloba foliis subrotundis integris nitidis planis, racemis fructuum cernuis. Jacq. amer. 19. hist. t. 73.*

*Coccolobis foliis crassis orbiculatis sinu aperto. Brown. jam. 208.*

*Polygonum foliis subrotundis, caule arboreo, fructibus baccatis. Sp. pl. 1. p. 365.*

*Uvisera foliis subrotundis amplissimis. Hort. cliff. 487.*

*Uvisera litorea, foliis amplioribus fere orbiculatis crassis, americana. Pluk. alm. 394. t. 236. f. 7.*

*Guajabara racemosa, foliis coriaceis subrotundis. Plum. ic. t. 445.*

*Populus americana rotundifolia. Bath. pin. 430.*

*Prunus maritima racemosa, folio subrotundo glabro, fructu minore purpureo. Sloan jam. 183. hist. 2. p. 129. t. 220. f. 3. Raj. dendr. 40. Catesb. car. 2. p. 96. t. 96.*

*Habitat in America. ☿*

### 2. COCCOLOBA foliis orbiculatis pubescentibus. *pubescens.*

*Coccoloba grandifolia foliis subrotundis integris hirsutis rugosis. Jac. amer. 19.*

*Coccolobis arborea, foliis orbiculatis. Brown. jam. 210.*

*Scortea arbor americana, amplissimis foliis averfa parte nervis extantibus. Pluk. phyt. 222. f. 8.*

*Habitat in America. ☿*

### 3. COCCOLOBA foliis lanceolato-ovatis. *punctata.*

*Coccoloba foliis ovali-oblongis acuminatis planis. Jacq. amer. 19.*

*Coccolobis foliis oblongo-ovatis venosis, uvis minoribus punctatis. Brown. jam. 240.*

*Uvi-*

Uvifera arbor americana, fructu aromatico punctato.

*Pluk. alm.* 394. t. 237. f. 4.

*Habitat in America.* §

- excoriata. 4. COCCOLOBA foliis ovatis, ramis quasi excorticatis.  
Coccoloba foliis ovato-oblongis acuminatis rugosis.  
*Jacq. amer.* 19.

Coccolobis montana major arborea, foliis subrotundis, cortice lævi. *Brown. jam.* 210.

Guajabara alia racemosa, foliis oblongis. *Plum. ic.* 146. f. 1?

Arbor indica, glycyrrhizæ foliis subrotundis, floribus in prælongam spicam adactis. *Pluk. amalth.* 22. t. 363. f. 4?

*Habitat in America.* §

- semulifolia. 5. COCCOLOBA foliis ovatis membranaceis. *Anan. acad.* 5. p. 397. \*

Coccolobis frutescens, foliis subrotundis, fructu minore trigono. *Brown. jam.* 210. t. 14. f. 3.

*Habitat in Jamaica.* §

## PAULLINIA.

- asiatica. 1. PAULLINIA foliis ternatis, caule aculeato, cirrhis nullis. *Fl. zeyl.* 143. \*

Chamælæa trifolia aculeata. *Burm. zeyl.* 58. t. 24.

Frutex baccifer indicus spinosus trifolius, floribus spicatis. *Pluk. alm.* 202. t. 95. f. 5.

Kaka-toddali. *Rheed. mal.* 5. p. 81. t. 41.

*Habitat in India.* §

- Seriana. 2. PAULLINIA foliis ternatis, petiolis teretiusculis, foliolis ovato-oblongis.

Paullinia foliis ternatis, foliolis crenatis, pedunculis cirriferis. *Hort. cliff.* 152.

Seriana scandens triphylla & racemosa. *Plum. gen.* 34. ic. 113. f. 2.

*Habitat in America calidiore.* §

- Curura. 3. PAULLINIA foliis ternatis, petiolis marginatis, foliolis cuneiformibus obtusis subdentatis.

Paullinia foliis ternatis, foliolis obtusis vix denticulatis glabris desinentibus in petiolum proprium.

*Hort. cliff.* 151. \*

Curura scandens triphylla. *Plum. gen.* 34. ic. 111. f. 2.

*Habitat in America calidiore.* §

4. PAUL-

4. **PAULLINIA** foliis biternatis, petiolis marginatis, fo- mexicana.  
lioli. ovatis integris.  
**Paullinia** foliis ternato-decompositis, caule aculeato.  
*Hort. cliff.* 152.  
**Seriana** scandens enneaphylla & racemosa. *Plum. gen.*  
34. *ic.* 113. *f.* 1.  
**Quauhmecati.** *Hern. mex.* 289.  
*Habitat in America calidiore.* b
5. **PAULLINIA** foliis biternatis, foliolis ovatis subfi. curassavica.  
nuatis.  
**Paullinia** foliis decompositis ternatis, caule inermi.  
*Hort. cliff.* 152. \*  
**Cururu** scandens enneaphylla, fructu racemoso rubro.  
*Plum. gen.* 34. *ic.* 111. *f.* 1.  
**Cordis indi** folio & facie frutescens curassavica latifolia.  
*Pluk. alm.* 120. *t.* 168. *f.* 6.  
*Habitat in Curassao.* b
6. **PAULLINIA** foliis triternatis.  
**Paullinia** foliis supradecompositis. *Hort. cliff.* 152. \* polyphylla.  
**Seriana** frutescens polyphylla & racemosa. *Plum. gen.*  
34. *ic.* 112.  
**Cordis indi** folio & facie frutescens portoricensis.  
*Pluk. alm.* 120. *t.* 168. *f.* 5.  
*Habitat in America calidiore.* b  
**Cirrus axillaris; Flores paniculati.**
7. **PAULLINIA** foliis pinnatis: foliolis incis, petiolis pinnata.  
marginatis. *Hort. cliff.* 52. \* *Roy. lugdb.* 464.  
**Cururu** scandens pentaphylla. *Plum. gen.* 34.  
**Clematis** pentaphylla, pediculis alatis, fructu racemoso  
tricocco & coccineo. *Plum. amer.* 76. *t.* 91.  
**Pisum cordatum** non vesicarium. *Sloan. jam.* 111.  
**Cururu** -ape. *Marcgr. bras.* 22. *Pis. bras.* 114. *Raj.*  
*hist.* 1347.  
*Habitat in Brasilia, Jamaica, Domingo.* b  
*Foliola oblongo-ovata, profunde 3 remote serrata.*

## CARDIOSPERMUM.

1. **CARDIOSPERMUM** foliis lævibus. *Hort. cliff.* 150. *Halicac-*  
*Hort. upf.* 97. *Fl. Zeyl.* 142. *Roy. lugdb.* 464. *bum.*  
**Pisum** vesicarium, fructu nigro: alba macula notato.  
*Bauh. pin.* 743.

Hali-



*Halicacabum. Rumph. amb. 6. p. 60. t. 24. f. 2.*

*Halicacabus pefegrinus. Dod. pempt. 455.*

β. *Corindum ampliore folio, fructu maximo. Tournef. infl. 431.*

γ. *Corindum fructu & folio minori. Tournef. infl. 431.*

*Habitat in Indiis. ☉*

Corindum. 2. **CARDIOSPERMUM** foliis subtus tomentosis.

*Cardiospermum villosum, foliis incisus obtulis, petiolis brevibus. Mill. dict.*

*Habitat in Brasilia.*

*Similis C. Halicacabo, sed Capsula & Folia subtus tomentosa. Pedunculi inter cirrhos 10! f. 11 floribus, a casu fructificationum persistentes, rigidi: Capsula arctiores.*

## SAPINDUS.

Saponaria. 1. **SAPINDUS** foliis impari-pinnatis, caule intermi.  
*Sapindus. Hort. cliff. 152. Mat. med. 532. Roy. lugdb. 464.*

*Sapindus foliis oblongis vix petiolatis per costam ample alatam. Brown. jam. 206.*

*Saponaria. Rumph. amb. 2. p. 134*

*Prunifera racemosa, folio alato costa media membranulis utrinque exstantibus donata. Sloan. jam. 184. hist. 2. p. 131.*

*Nux americana, foliis alatis bifidis. Comm. hort. 1. p. 183. t. 94.*

*Nuciprunifera arbor americana, fructu saponario orbiculato monococco nigro. Pluk. alm. 255. t. 217. f. 7.*

*Habitat in Indiis. ☿*

trifoliata. 2. **SAPINDUS** foliis ternatis.

*Poerinii f. Vercœpcelongi. Rheed. mal. 4. p. 43. t.*

*Habitat in Malabaria. ☿*

spinofus. 3. **SAPINDUS** foliis abrupte pinnatis, caule spinosissimo.

*Sapindus fruticosus, caudice & ramis spinosissimis, foliis ovatis pinnatis. Brown. jam. 207. t. 20. f. 2.*

*Habitat in Jamaica. ☿*

## TETRAGYNIA.

## PARIS

## 1. PARIS.

quadrifolia.

Paris foliis quaternis. *Fl. lapp.* 155. *Fl. fvec.* 325; 346.*Hort. cliff.* 153. *Roy. lugdb.* 461. *Hall. helv.* 412.Solanum quadrifolium bacciferum. *Bauh. pin.* 167.Aconitum salutiferum. *Tabern. hist.* 720.*Habitat in Europæ nemoribus.* 24

## ADOXA.

1. ADOXA. *Hort. cliff.* 152. *Fl. fvec.* 326, 347. \* Moschatellina.*Roy. lugdb.* 252.Moschatella. *Cord. hist.* 172.Ranunculus nemorosus Moschatellina dictus. *Bauh. pin.* 178.Fumaria bulbosa f. tuberosa minima. *Tabern. ic.* 39.*Habitat in Europæ nemoribus.* 24*Huc relata secundum florem terminalem ex principis**Phil. bot.* 178.

## ELATINE.

1. ELATINE foliis oppositis. *Fl. lapp.* 156. *Fl. fvec.* Hydropiper 327; 348. *Gron. virg.* 158. *Roy. lugdb.* 452.Hydropiper. *Buxb. cent.* 3. p. 35. t. 37. f. 3.

Alismastrum serpillifolium, flore albo tetrapetalo.

*Vaill. parif.* 5. t. 2. f. 2.β. Alismastrum serpillifolium, flore roseo tripetalo. *Vaill. parif.* 5 t. 2. f. 1.*Habitat in Europæ inundatis.* ☉2. ELATINE foliis verticillatis. *Fl. fvec.* 2. n. 349. Alismastrum.*Roy. lugdb.* 452. *Sauv. monsp.* 164. *Boem. lips.* 127.

Elatine foliis emerfis linaribus, immerfis capillaceis.

*Sauv. act. monsp.* 1743. p. 50.Ericoides facie pinastellæ. *Rupp. gen.* 90. \*Equisetum palustre, linariæ scopariæ folio. *Bauh. pin.* 15.Alismastrum galii folio. *Vaill. parif.* 6. t. 1. f. 5.β. Alismastrum gratiolæ folio. *Tournef. inst.* 244 *Raj. suppl.* 502.*Habitat Abœ, Lipsiæ, Parisiis, Monspeli, in fossis.*

Clas-

## Classis IX.

ENNEANDRIA  
MONOGYNIA.

## LAURUS.

Cinnamomum.

1. LAURUS foliis trinerviis ovato-oblongis nervis versus apicem evanescentibus.
- Fl. zeyl.*
145. \*
- Nat. med.*
- 190.

Laurus foliis oblongo-ovatis nitidis planis. *Hort. cliff.* 154.Cinnamomum, foliis latis ovatis, frugiferum. *Burm. zeyl.* 62. t. 27.Cinnamomum f. *Canella zeylanica*. *Bauh. pin.* 408.*Cassia cinnamomea*. *Herm. lugdb.* 129. t. 655.*Habitat in Zeylona; in Martinia*. *Jacquin.* §

Cassia.

2. LAURUS foliis triplinerviis lanceolatis.
- Fl. zeyl.*
- 146.
- Nat. med.*
- 191.

Cinnamomum perpetuo florens, folio tenuiore acuto. *Burm. zeyl.* 63. t. 28.Cinnamomum f. *Canella malabarica* f. *javanensis*. *Bauh. pin.* 409.*Carna*. *Rheed. mal.* 1. p. 107. t. 59.*Habitat in Malabaria, Sumatra, Java.* §

Camphora.

3. LAURUS foliis triplinerviis lanceolato-ovatis.
- Nat. med.*
- 192.

Laurus foliis ovatis utrinque acuminatis trinerviis nitidis, petiolis laxis. *Hort. cliff.* 154.Laurus Camphorifera. *Kämpf. jap.*Camphora officinarum. *Bauh. pin.* 500.Arbor camphorifera japonica. *Breyn. prodr.* 2. p. 16. ic. 16. t. 2.*Habitat in Japonia.* §

Chloroxylon.

4. LAURUS foliis trinerviis ovatis coriaceis: nervis apicem attingentibus.

Chloroxylon foliis ovatis glabris rigidis trinerviis, floribus singularibus. *Brown. jam.* 187. t. 7. f. 1.*Habitat in Jamaica.* §

5. LAU-

5. LAURUS foliis venosis lanceolatis perennantibus, floribus quadrifidis dioicis. *Hort. cliff.* 155. *Hort. upf.* 98. *Mat. med.* 194. *Roy. lugdb.* 98.  
*Laurus vulgaris.* *Bauh. pin.* 460.  
*Laurus.* *Cam. epit.* 60.  
*Habitat in Italia, Græcia.* ♀
6. LAURUS foliis venosis lanceolatis perennantibus planis, ramulis tuberculatis cicatricibus, floribus racemosis. *Hort. cliff.* 154. *Gron. virg.* 159.  
*Laurus indica.* *Pluk. alm.* 210. t. 301. f. 1.  
*Laurus latifolia indica.* *Barr. rar.* 123. t. 877.  
*Cinnamomum sylvestre americanum.* *Seb. thes.* 2. p. 90. t. 84. f. 6.  
*Habitat in Virginia.* ♀
7. LAURUS foliis venosis ovatis coriaceis perennantibus, floribus corymbosis. *Persea.*  
*Persea.* *Clus. hist.* 1. p. 2.  
*Persea americana.* *Bauh. pin.* 441.  
*Pyro similis fructus in nova hispania, nucleo magno.* *Bauh. pin.* 439.  
*Prunifera arbor fructu maximo pyriformi viridi, pericarpio esculento butyraceo nucleum unicum maximum nullo officulo tectum cingente.* *Sloan jam.* 132. *hist.* 2. p. 132. t. 222. f. 2. *Raj. dendr.* 48.  
*Arbor americana, amplissimis pergamenis foliis superficie nitidissima, fructu pyriformi crustaceo cortice coriato.* *Pluk. alm.* 39. t. 267. f. 1.  
*Habitat in America calida.* ♀
8. LAURUS foliis venosis lanceolatis, calycibus fructus baccatis. *Hort. cliff.* 154. *Gron. virg.* 46. *Roy. lugdb.* 226. *Borbonia.*  
*Laurus carolinensis, foliis acuminatis, baccis cæruleis, pedicellis longis rubris insidentibus.* *Catesb. car.* 1. p. 63. t. 63.  
*Borbonia fructu oblongo nigro, calyce coccineo.* *Plum. gen.* 4. ic. 60.  
*Habitat in Carolina, Virginia.* ♀
9. LAURUS foliis venosis oblongis acuminatis annuis, ramis supra axillaribus. *anivialis.*  
*Laurus foliis lanceolatis enervibus annuis.* *Gron. virg.* 159.

*Laurus foliis enervibus ovatis utrinque acutis. Gron. virg. 46.*

*Cornus foliis salicis laureæ acuminatis, floribus albis, fructu sassafras. Catesb. car. 2. p. 28. t. 28?*

*Habitat in Virginia ad ripas rivulorum. ‡*

*Pedunculi hujus fructiferi colorati sunt.*

**Benzoinæ**

10. **LAURUS** foliis enerviis ovatis utrinque acutis integris annuis. *Hort. cliff. 154. Mat. med. 195. Gron. virg. 46. Roy. lugdb. 226.*

*Arbor virginiana, citreæ vel limonii folio, Benzoinum fundens. Comm. hort. 1. p. 189. t. 97.*

*Arbor virginiana, pichaminis folio, baccata Benzoinum redolens. Pluk. alm. 42. t. 139. f. 3, 4.*

*Habitat in Virginia. ‡*

**Sassafras.**

11. **LAURUS** foliis integris trilobisque. *Hort. cliff. 154. Mat. med. 193. Gron. virg. 46. Roy. lugdb. 227. Cold. noveb. 94. Kalm. it. 2. p. 270, 434.*

*Cornus mas odorata, folio trifido: margine plano, Sassafras dicta. Pluk. alm. 120. t. 222. f. 6. Catesb. car. 1. p. 55. t. 55.*

*Sassafras arbor, ex florida, ficulneo folio. Bauh. pin. 431.*

β. *Cornus mas f. Sassafras laurinis foliis indivisis. Pluk. amalth. 66.*

*Habitat in Virginia, Carolina, Florida. ‡*

## TINUS.

**occidentalis.**

1. **TINUS,**

*Volkameria arborea, foliis oblongo-ovatis alternis superne glabris, subtus subvillofis & nervosis, spicis ramosis & terminalibus. Brown. jam. 214. t. 21. f. 1.*

*Baccifera arbor calyculata, foliis laurinis, fructu racemoso esculento subrotundo monopireno pallide luteo. Sloan. jam. 165. hist. 2. p. 86. t. 198. f. 2. Raj. dend. 49.*

*Habitat in Jamaica. ‡*

## CASSYTHA.

**californica.**

1. **CASSYTHA.** *Osborn. it. 243. \* Jacq. amer. 20. hist. 1. 79. Sp. pl. 1. p. 35.*

*Cuscuta altera f. major. Camell. luc. 1. n. 1. Pet. gaz. 77. t. 49. f. 12.*

Cu-

*Cuscuta baccifera barbadensium. Pluk. alm. 126. t. 172. f. 2.*

*Cuscuta. Rumph. amb. 5. p. 491. t. 184. f. 4.*

*Acatia-valli. Rheed. mal. 7. p. 83. t. 44. Raj. suppl. 551.*

*Habitat in India.*

*Simillima Cuscutæ; relata ad Triandriam ab Osbeckio, ad Enneandriam a Jacquinio; mihi non floruit.*

## TRIGYNIA.

### RHEUM.

1. RHEUM foliis glabris, petiolis subfulcatis.

Rhaponticum.

*Rheum foliis glabris. Hort. upf. 98. Mat. med. 198.*

*Rheum. Hort. cliff. 155.*

*Rhaponticum. Alp. rhapont. 1. t. 1.*

*Rhaponticum folio lapathi majoris glabro. Buzh. pin. 116.*

*Habitat in Thracia, Scythia, Monte Aureo. 2.*

*Petoli subius fulcati, quod non in sequenti specie.*

2. RHEUM foliis subvillosis undulatis petiolis æqualibus. *undulatum.*

*Amzn. acad. 3. p. 212. t. 4. \**

*Rheum foliis subvillosis. Hort. upf. 98. Mat. med. 197.*

*Rhabarbarum folio oblongo crispo undulato, flabellis*

*sparsis. Geoffr. mat. 2 p. 125.*

*Rhabarbarum sinense, folio crispo, flagellis rarioribus*

*& minoribus. Amm. herb. 206.*

*Rhabarbarum folio longiori hirsuto crispo, florum thyr-*

*so longiori & tenuiori. Amm. ruth. 9.*

*Acetosa montana, folio cubitali oblongiore crispo,*

*floribus in subviridi luteolis. Messersschmidii. Amm.*

*ruth. 226.*

*Habitat in China, Sibiria. 2.*

3. RHEUM foliis palmatis acuminatis.

*palmatum.*

*Habitat in China ad murum. a. Dav. Gortero. 2.*

*Gemma vernans non rubescens, sed flavescens. Folia*

*scabriuscula. Foliorum lacinia oblongæ acutiusculæ.*

4. RHEUM foliis sublobatis obtusissimis glaberrimis ar-

*compactum.*

*gute denticulatis glabris.*

*Rheum foliis cordatis glabris marginibus sinuatis, spi-*

*cis divisis nutantibus. Mill. dict. t. 218*

*Habitat in Tataria, China.* 24

*Folia, magis quam reliqua, coriacea f. compacta: Lobis rotundatis, obsoletioribus, crenatis margine cartilagineo acutis denticulis, utrinque glaberrimis, venis robustioribus. Paniculae rami nutantes.*

Ribes.

5. RHEUM foliis granulatis, petiolis æqualibus. *Gron. orient.* 130.

Lapathum orientale, aspero & verrucoso folio, Ribes arabibus dictum. *Dill. elth.* 191. t. 158. f. 192.

Lapathum orientale tomentosum rotundifolium, Ribes arabum dictum. *Breyn. E. N. C. cent.* 7. p. 7.

Lapathum orientale asperum, folio subrotundo, fructu magno purpureo. *Pocock. orient.* 189. t. 84.

Ribes arabum. *Rauw. it.* 266. 282.

Ribes arabum, foliis petasitidis. *Bauh. pin.* 455.

*Habitat in Persia, Libano, Carmelo.* 24

*Flores hujus speciei non vidi. Facies juadet hujus generis esse.*

## HEXAGYNIA.

### BUTOMUS.


umbellatus. 1. BUTOMUS. *Fl. lapp.* 159. *Fl. suec.* 328, 350. *Roy. lugdb.* 45. *Hall. helv.* 298.

*Juncus floribus major. Bauh. pin.* 112.

*Sedo affinis Juncoides umbellata palustris. Moris. hist.* 3. p. 468. f. 12. t. 5. f. 1.

*Gladiolus aquatilis. Dod. pempt.* 600.

*Habitat in Europæ aquosis.* 24



## Classis X.

# DECANDRIA

## MONOGYNIA.

### SOPHORA.

1. SOPHORA foliis pinnatis: foliolis numerosis oblongis villosis, caule herbaceo. alopecuroides.  
*Sophora. Hort. cliff. 156.*  
*Ervum orientale alopecuroides perenne, fructu longissimo. Tournef. cor. 27. Dill. elth. 136.*  
*Glycyrrhiza siliquis nodosis quasi articulatis. Buxb. cent. 3. p. 25. t. 46.*  
*Habitat in Oriente. 2.*
2. SOPHORA foliis pinnatis: foliolis numerosis subrotundis tomentosis.  
*Sophora tomentosa, foliolis subrotundis. Fl. zeyl. 163.*  
*Indigophora foliis tomentosis. Hort. cliff. 487.*  
*Colutea zeylanica argentea tota. Herm. lugdb. 169. t. 171. Raj. hist. 1720.*  
*Habitat in Zeylona. 5*
3. SOPHORA foliis pinnatis: foliolis numerosis subcordatis. occidentalis.  
*Galega fruticosa, foliis subrotundis pinnatis, spicis simplicibus terminalibus. Brown. jam. 289. t. 31. f. 1.*  
*Coraliodendron foliis pseudo-acaciæ subtus tomentosis, flore luteo. Plum. spec. 21. ic. 103.*  
*Arbori coral affinis non spinosa, fraxini folio rotundiore, foliis & ramulis pubescentibus. Sloan. jam. 144. hist. 2. p. 40. t. 107. f. 3.*  
*Habitat in America.*  
*Differt a Soph. tomentosa parum admodum. Folia minus tomentosa, & fere subtus tantum subto- mentosa.*
4. SOPHORA foliis pinnatis: foliolis septenis glabris. heptaphylla.  
*Sophora glabra, foliolis septenis. Fl. zeyl. 164.*  
*Fruticulus sinensis, sennæ sylvestris folio angustiore,*



nodosa siliqua rostro longiore donata. *Pluk. amalth.*

18. t. 451. f. 10 ?

Anticholerica. *Rumph. amb.* 4. p. 60. t. 22.

Habitat in India, §

- Genistoides. 5. *SOPHORA* foliis ternatis sessilibus: foliolis linearibus.

Genistra africana, foliis galli. *Old. afr.* 31 -

Habitat ad Cap. b. Spei.

- tinctoria. 6. *SOPHORA* foliis ternatis subsessilibus: foliolis obovatis glabris.

Cytisus foliis fere sessilibus, calycibus bractea triplici auctis. *Gron. virg.* 82.

Cytisus procumbens americanus, flore luteo, ramossimus, qui Anil suppeditat. *Pluk. alm.* 129. t. 86. f. 2. *Ehret. t.* 1. f. 3.

Habitat in Barbados, Virginia.

- lupinoides. 7. *SOPHORA* foliis ternatis petiolatis: foliolis ovalibus pilosis.

*Sophora* foliis ternatis, spica verticillata. *Amæn. acad.* 2. p. 350. \*

Habitat in Camtschatca, G. Demidoff.

- biflora. 8. *SOPHORA* foliis simplicibus obovatis subtomentosis, pedunculis bifloris.

*Crotalaria* foliis simplicibus ovatis villosis, petiolis simplicissimis, ramis teretibus. *Hort. cliff.* 357. \* *Roy. lugdb.* 374.

*Crotalaria* arbor africana, styracis folio molli incano. *Pluk. alm.* 122. t. 185. f. 2.

*Genista* arborescens africana, styracis folio. *Herm. lugdb.* 270. t. 271. *Seb. mus.* 2. t. 99. f. 3.

*Myrto-Genista* capitis b. Spei. *Breyn. cent.* t. 29.

Habitat in Æthiopia. §

## ANAGYRIS.

foetida.

1. *ANAGYRIS* *Clus. list.* 1. p. 93.

*Anagyris* floribus lateralibus. *Roy. lugdb.* 371.

*Anagyris* foetida. *Bauh. pin.* 391.

Habitat in Italia, Sicilia, Hispania montibus. §

## CERCIS

Siliquastrum.

1. *CERCIS* foliis cordato-orbiculatis glabris. *Hort. cliff.*

156.

156. *Hort. upf.* 99. *Roy. lugdb.* 463. *Gron. orient.*  
131.

*Siliqua sylvestris rotundifolia. Bauh. pin.* 402.

*Arbor judæ. Dod. pempt.* 786.

*Habitat in Italia, Hispania, Narbona, Oriente.* ‡

2. *CERCIS foliis cordatis pubescentibus. Hort. cliff. canadensis.*

156. *Hort. upf.* 99. *Roy. lugdb.* 463. *Gron. virg.*

47.

*Ceratia agrestis virginiana, folio rotundo minori. Raj.*

*dendr.* 100.

*Habitat in Virginia.* ‡

## BAUHINIA.

1. *BAUHINIA caule cirrhifero. Læst. it.* 218.

*scandens*

*Clematis indica, folio bifido, flore fructuque carens,*

*arbores transcendens. Raj. suppl.* 328.

*Folium linguæ. Rumph. amb.* 5. p. 1. t. 1.

*Naga-mu-valli. Rheed. mal.* 8. p. 57. t. 29.

*Habitat in Malabaria, Amboina, Cumana.* ‡

2. *BAUHINIA caule aculeato. Hort. cliff.* 156. t. 14. *aculeata.*

*Roy. lugdb.* 463. *Læst. it.* 219.

*Bauhinia aculeata, folio rotundo & emarginato. Plum.*

*gen.* 23. *ic.* 44. *f.* 1.

*Habitat in America calidiore.* ‡

3. *BAUHINIA foliis ovatis: lobis divaricatis.*

*divaricata.*

*Bauhinia foliis quinquenerviis: laciniis acuminatis re-*

*motissimis. Hort. cliff.* 156. t. 15. *Roy. lugdb.* 464.

*Bauhinia non aculeata, folio ampliori & bicorni.*

*Plum. gen.* 23. *ic.* 44. *f.* 2.

4. *Bauhinia foliis ovato-cordatis lobis longissimis paral-*

*lelis. Mill. dict.* t. 61.

*Habitat in America.* ‡

5. *BAUHINIA foliis ovatis: lobis parallelis.*

*ungulata.*

*Bauhinia foliis ovatis oblongis recta linea bifidis. Hort.*

*cliff.* 157.

*Habitat in America.* ‡

6. *BAUHINIA foliis cordatis: lobis coadunatis obtusis.*

*variegata.*

*Arbor f. Thomæ f. Asitira. Zanon. hist.* 26. t. 15.

*Chovanna-mandaru. Rheed. mal.* 1. p. 57. t. 32. *Raj.*

*hist.* 1751.

- Habitat in Malabariæ, Maderæ arenosis.* ‡  
*Corollæ patentes: Petalis ovatis, acutis, roseis, luteo purpureoque variegatis.*
- purpurea. 6. *BAUHINIA* foliis subcordatis bipartitis rotundatis subtus tomentosis.  
*Chovanna-mandaru. Rheed. mal. 1. p. 59. t. 33.*  
*Habitat in Indiæ arenosis.* ‡  
*Flores patentes, purpurascentes: Petalis lanceolatis, distantibus.*
- tomentosa. 7. *BAUHINIA* foliis cordatis: lobis semiorbiculatis tomentosis.  
*Bauhinia foliis cordato subrotundis: laciniis rotundatis.*  
*Hort. cliff. 157. Hort. upf. 99. Fl. zeyl. 147. Roy. lugdb. 464.*  
*Bauhinia foliis subrotundis, flore flavescente striato.*  
*Burm. zeyl. 44. f. 18.*  
*Mandaru maderaspatense, foliis firmioribus bisulcis glabritie splendentibus. Pluk. alm. 240. t. 44. f. 6.*  
*Chanschena-pou. Rheed. mal. 1. p. 63. t. 35.*  
*Habitat in India.* ‡  
*Corollæ campanulatæ, albo-flavescentes: Petalis ob-ovatis.*
- acuminata. 8. *BAUHINIA* foliis ovatis: lobis acuminatis semiovatis.  
*Bauhinia inermis, foliis cordatis semibifidis: laciniis acuminato-ovatis erectis dehiscentibus. Hort. cliff. 157. Fl. zeyl. 148. \**  
*Bauhinia foliis bilobis, spicis laxis terminalibus. Brown. jam. 286.*  
*Sennæ spurix aut Asphalto affinis arbor filiquosa, foliis bifidis. Sloan. jam. 150. hist. 1. p. 51.*  
*Velutta-mandaru. Rheed. mal. 1. p. 61. t. 34. Raj. hist. 1751.*  
*Habitat in Indiis.* ‡  
*Corollæ campanulatæ, albæ: Petalis obtusis. Stamina diadelpha in B. hac & B. divaricata; contrario ac in Papilionaceis modo.*

## PARKINSONIA.

- aculeata. 1. *PARKINSONIA. Hort. cliff. 157. t. 13. Hort. upf. 99. Roy. lugdb. 465. Brown. jam. 222. Jacq. hist. 1. 80.*

Par-

*Parkinsonia aculeata*, foliis minutis uni costæ adnexis.

*Plum. gen.* 25.

*Parkinsonia aculeata*, foliis mimosæ uni costæ affixis.

*Walth. hort.* 6. t. 13.

*Habitat in America calidiore.* ‡

## HYMENÆA.

1. *HYMENÆA*. *Hort. cliff.* 484. *Hort. upf.* 303. *Mat. Courbaril.*  
*med.* 515. *Brown. jam.* 221.

*Courbaril bifolia*, fructu pyramidato. *Plum. gen.* 49.

*ceratia diphyllas anteguanæ*, ricini majoris fructu nigro, siliqua grandi incluso. *Pluk. alm.* 96. t. 82. f. 2.

*Arbor siliquosa ex qua Gummi Elemi.* *Bauh. pin.* 404.

*Itaiba.* *Pis. bras.* 123.

*Habitat in America meridionali.* ‡

## CASSIA.

\* *Sennæ.*

1. *CASSIA* foliis conjugis, stipulis cordato-lanceolatis. *diphylla.*

*Habitat in India.* ☉

*Frutex Caule tereti. Folia conjugata s. duo in petio-  
lo brevi, semiorbiculata, obtusa, striata. Stipulæ  
cordato-lanceolatæ, totos ramos obtegentes.*

2. *CASSIA* foliis bijugis subovatis, glandulis duabus *absus.*  
subulatis inter infima. *Fl. Zeyl.* 153. \*

*Senna quadrifolia*, siliqua plana hirsuta, flore aureo  
sanguineo. *Burm. Zeyl.* 312. t. 97.

*Senna exigua maderaspatana tetraphylla siliquifera gla-  
bra, florum pediculis ad exortum foliorum propen-  
dentibus.* *Pluk. alm.* 341. t. 60. f. 1.

*Loto affinis ægyptiaca.* *Bauh. pin.* 332.

*Absus.* *Alp. ægypt.* 97.

*Habitat in India, Ægypto.* ☉

3. *CASSIA* foliis bijugis ovato-oblongis acuminatis, *viminea.*  
glandula oblonga inter infima, spinis subpetiolaribus  
obsoletis tridentatis. *Amæn. acad.* 5. p. 397. \* *Læfl.*  
*it.* 232.

*Cassia viminea*, foliis ovatis acuminatis bijugatis, ra-  
cemis laxis alaribus, siliquis brevioribus compressis.  
*Brown. jam.* 223.

*Habitat in Jamaica.* ‡

Tagera.

4. CASSIA foliis trijugis, glandula petiolari, stipulis ciliatis cordatis acuminatis.

Cassia 6, Senna spuria tetraphylla arborescens, filiquis tenuibus longissimis pendulis. *Amm. herb.* 603. n. 33?

Tagera. *Rheed. mal.* 2. p. 103. t. 52? *Raj. hist.* 1743. *Habitat in India.*

Fruticulus procumbens: Ramis filiformibus. Petioli brevissimi; Foliola quatuor, semiorbiculata, venosa, quorum duo exteriora majora. Stipulae cordatae f. cordato-lanceolatae, margine pilosae.

Tera.

5. CASSIA foliis trijugis obovatis: exterioribus majoribus; glandula subulata inter inferiora quatuor. *Roy. lugdb.* 468. *Fl. zeyl.* 152. *Hort. upf.* 100. *Mill. dict.* t. 82.

Cassia filiqua quadrangulata. *Dill. elth.* 72. t. 63. f. 73.

8. Cassia humilis, filiquis foeni graeci. *Plum. spec.* 18. t. 76. f. 2.

Senna orientalis hexaphylla, Tala zeylonensium. *Herm. lugdb.* 557. *Raj. hist.* 1743.

Senna orientalis hexaphylla, filiquis longis incurvis. *Pluk. alm.* 342.

Galega indica minor hexaphyllos amanni. *Raj. hist.* 911.

*Habitat in India.* ☉

Caules ♂ Petioli scabri. Stipulae lineares, acuminatae, pilosae. Glandulae acuminatae, apice fuscae, inter singula paria 2 inferiora, non vero inter extremum par sitae.

Varietas β differt filiqua recurva. An species distincta?

bica-psu-laris.

6. CASSIA foliis trijugis obovatis glabris: interioribus rotundioribus minoribus: glandula interjecta globosa. *Hort. cliff.* 159. *Hort. upf.* 100. *Roy. lugdb.* 468.

Cassia hexaphylla, filiqua bica-psulari. *Plum. spec.* 18. t. 76. f. 1.

*Habitat in India.* ☿

Arborescens est, ♂ glaberrima. Foliola interiora breviora. Sliqua teretiuscula, coriacea, loculo duplici tereti.

emarginata.

7. CASSIA foliis trijugis ovatis rotundatis emarginatis aequalibus. *Hort. cliff.* 159. *Roy. lugdb.* 467.

Cassia minor hexaphylla fruticosa, sennae foliis. *Scan. jam.*

*jam.* 146. *hist.* 2. p. 44. t. 180. f. 1. 2. 3. 4. *Raj.*  
*dendr.* 110.

*Habitat in Caribæis.* ☿

8. CASSIA foliis trijugis ovatis obtusiusculis.

obtusifolia.

*Cassia foetida*, foliis sennæ italicæ. *Dill. clh.* 71. t. 62.  
f. 72.

*Cassia minor* herbacea plerumque hexaphylla, folio  
obtus. *Sloan. jam.* 148. *hist.* 2. p. 47. t. 180. f. 5.

*Gallinaria rotundifolia.* *Rumph. amb.* 5. p. 283. t. 97.  
f. 2.

*Habitat in Cuba.* ☉

9. CASSIA foliis trijugis quadrijugisve-sexjugis subovatis.

Senna.

*Hort. cliff.* 159. *Mat. med.* 200. *Roy. lugdb.* 468.

*Senna Alexandrina* f. foliis acutis. *Bauh. pin.* 397.

*Morif. hist.* 2. p. 201. f. 2. t. 24. f. 1.

13. *Senna italica* f. foliis obtusis. *Bauh. pin.* 397. *Morif.*  
*hist.* 2. p. 200. f. 2. t. 24. f. 2.

*Senna. Dod. pempt.* 361.

*Habitat in Aëgypto.* ☉

10. CASSIA foliis quadrijugis ovato-lanceolatis re- falcata.  
trofalcatis: glandula baseos petiolorum. *Hort. cliff.*

159. \*

*Habitat in America.* ☉

11. CASSIA foliis quinquejugis ovatis - lanceolatis occidenta-  
litate margine scabris: exterioribus majoribus; glandula  
baseos petiolorum

*Cassia foliolis* quatuor parium ovato-lanceolatis, glan-  
dula baseos petiolorum. *Hort. cliff.* 159. *Roy. lugdb.*  
467.

*Senna occidentalis*, odore opii viroso, orobi panno-  
nici foliis mucronatis, glabra. *Comm. hort.* 1. p. 51.  
t. 26.

*Habitat in Jamaica.*

*Caulis sesquipedalis*, punctis vagis scaber, exaratus  
deorsum a singulo petiolo sulcis duobus. — Folia su-  
periora foliolis quinque parium, ovato-lanceolatis,  
glabris, margine scabris, acuminatis, exterioribus sen-  
sim majoribus, fixidis. Racemus terminalis. Flo-  
res lutei absque macula. Folia impuberis plantæ  
quinquejuga; adultæ tantum trijuga.

12. CAS-

- pilosa.** 12. CASSIA foliis quinquejugis eglandulatis, stipulis semicordatis acuminatis, caule stricto piloso. Cassia suffruticosa erecta hirsuta, floribus singularibus ad alas. *Brown. jam.* 224.  
*Habitat in Jamaica.*  
Caulis erectiusculus, fruticulosus. Ramis erectis, undique valde pilosis. Folia glabra. Stipulae semicordatae, acuminatae, ciliatae. Folia levia. Pedunculi erecti, gemini, uniflori, laeves. Siliquae oblongae, compressae. Flores circiter pentandri.
- planifolia.** 13. CASSIA foliis quinquejugis ovato-lanceolatis glabris, glandula baseos petiolorum. *Roy. lugdb.* 468.  
Cassia siliquis planis. *Plum. spec.* 18. t. 77.  
*Habitat in America calidiore.*
- fistula.** 14. CASSIA foliis quinquejugis ovatis acuminatis glabris, petiolis eglandulatis. *Fl. zeyl.* 149. \* *Mat. med.* 199. *Hasseltq. in.* 468.  
Cassia foliolis quinque parium lanceolatis: extimis minoribus. *Hort. cliff.* 158.  
Cassia fistula alexandrina. *Bauh. pin.* 405. *Comm. hort.* 1. p. 215. t. 110. *Rumph. amb.* 2. p. 83. t. 21.  
*Conna. Rheed. mal.* 1. p. 37. t. 21.  
*Habitat in India, Aegypto.* 5
- biflora.** 15. CASSIA foliis sejugis oblongiusculis glabris inferioribus minoribus, glandula subulata inter infima, pedicellis subbifloris. *Hort. cliff.* 159. \*  
Cassia biflora. *Amæn. acad.* 5. p. 397. \*  
Cassia fruticosa, foliis minoribus obovatis sexjugatis, floribus geminis. s. bigeminatis, racemis alaribus. *Brown. jam.* 223.  
Cassia minor arborescens, siliquis planis articulatis. *Plum. spec.* 18. ic. 78. f. 1.  
*Habitat in America.* 5
- hirsuta.** 16. CASSIA foliis sejugis ovatis acuminatis lanatis. *Hort. cliff.* 159. *Roy. lugdb.* 467.  
Cassia americana foetida, foliis amplioribus villosis. *Tournef. inst.* 519.  
Senna occidentalis, odore opii viroso, foliis orobi pannonicici mucronatis hirsutis, *Herm. lugdb.* 356.  
*Habitat in America.*

17. CASSIA foliis septemjugis, floribus pentandris, cau- serpens.  
libus filiformibus prostratis herbaceis.  
Cassia herbacea tenuissima procumbens, floribus singu-  
laribus ad alas. *Brown. jam.* 225.  
*Habitat in Jamaica.* ☉  
Caules *filiformes*, *herbacei prostrati*, *rarius ad basin*  
*ramosi, glabri.* Foliola *oblonga, striata, angustata.*  
Stipulæ *subulatæ.* Pedunculi *axillares, solitarii, lon-*  
*gitudine foliorum, nudi, horizontales.* Siliquæ *ere-*  
*ctæ, rhombeo-oblongæ; ad angulum rectum cum pe-*  
*dunculo positæ.*
18. CASSIA foliis septemjugis lanceolatis: extimis *ligustrina.*  
minoribus, glandula baseos petiolorum. *Hort. cliff.*  
199. *Hort. upf.* 100. *Gron. virg.* 47. *Roy. lugdb.*  
467.  
Cassia ligustri folio. *Plum. spec.* 10. *Dill. elth.* 350. t.  
259. f. 338.  
Cassia bahamensis, pinnis foliorum mucronatis angu-  
stis, calyce floris non reflexo. *Mart. cent.* 21. t. 21.  
*Habitat in Virginia, Bahama.* ☿  
*Folia glabra, subtus obsolete pubescentia.*
19. CASSIA foliis octojugis ovali-oblongis: interiori- alata.  
bus minoribus, petiolis eglandulosis, stipulis patulis.  
*Hort. cliff.* 158. *Hort. upf.* 100. *Roy. lugdb.* 467.  
Cassia sylvestris foetida, filiquis alatis. *Plum. spec.* 18.  
Faba dulcis *Mer. surin.* 58. t. 58.  
Herpetica. *Rumph. amb.* 7. p. 35. t. 18.  
*Habitat in America calidiore.* ☿
20. CASSIA foliis octojugis ovato-oblongis æquali- marilandica.  
bus, glandula baseos petiolorum. *Hort. cliff.* 159.  
*Hort. upf.* 100. *Roy. lugdb.* 467.  
Cassia mimosæ foliis, filiqua hirsuta. *Dill. elth.* 351. t.  
260. f. 339.  
Cassia marilandica, pinnis foliorum oblongis, calyce  
floris reflexo. *Mart. cent.* 23. t. 23.  
*Habitat in Virginia, Marilandia.* ☿  
*Caulis ☿ folia adpersa pilis raris.*
21. CASSIA foliis novemjugis oblongis, glandula tenuissima.  
subulata inter infima. *Roy. lugdb.* 478.  
Senna spuria frutescens, filiquis tenuissimis. *Houft. mff.*  
*Habitat in Havana.* ☿



- Sophera.** 22. CASSIA foliis decemjugis lanceolatis, glandula  
baseos oblonga. *Fl. zeyl.* 150. \*  
Senna vigintifolia filiquis teretibus. *Burm. zeyl.* 213.  
t. 98.  
Galegæ affinis. *Bauh. pin.* 352.  
Gallinaria acutifolia. *Rumph. amb.* 5. p. 283. t. 97. f. 1.  
Ponnam-Tongera. *Rheed. mal.* 2. p. 101. t. 52.  
*Habitat in Indiæ umbrosis.*
- auriculata.** 23. CASSIA foliis duodecim-jugis obtusis mucronatis,  
glandulis subulatis pluribus, stipulis reniformibus  
barbatis. *Fl. zeyl.* 151. \*  
Senna spuria maderaspatana, caule ad foliorum alas  
geminatis subrotundis appendiculis auriculato. *Pluk.*  
*alm.* 341. t. 314. f. 4. *Raj. dendr.* 110.  
*Habitat in India.*
- javanica.** 24. CASSIA foliis duodecim-jugis oblongis obtusis  
glabris, glandula nulla.  
Cassia fistula javanica, flore carneo. *Comm. hort.* 1. p.  
217. t. III.  
Cassia foliis plurimis oblongis, flore rubello, filiquis  
maximis crassioribus trinerviis. *Brown. jam.* 223.  
Cassia fistula brasiliensis. *Bauh. pin.* 403.  
Cassia nigra s. fistulosa secunda. *Sloan. jam.* 146. *hist.*  
2. p. 44.  
*Habitat in Indiis.*  
*Flores mihi non protulit.*
- \* \* Chamæcristæ foliolis numerosis.
- Chamæcrista.** 25. CASSIA foliis multijugis, glandula petioli pedi-  
cellata, stipulis ensiformibus. *Hort. upf.* 101.  
Cassia foliolis plurium parium linearibus, stipulis su-  
bulatis. *Hort. cliff.* 158. *Gron. virg.* 47.  
Cassia suffruticosa erecta, foliolis linearibus plurimis  
pinnatis, floribus singularibus vel geminis sparsis.  
*Brown. jam.* 225.  
Chamæcrista pavonis major. *Comm. hort.* 1. p. 53. t. 37.  
*Habitat in Jamaica, Barbados, Virginia.* ☉  
Foliola 20. Pedicellus spatio supra petiolum egredi-  
ens. Petala duo superiora macula purpurea. An-  
theræ rubræ
- glandulosa.** 26. CASSIA foliis multijugis multiglandulatis, stipulis  
subulatis.

*Chamæcrista pavonis americana*, filiqua multiplici.

*Breyn. cent. 64. t. 24.*

*Habitat in Jamaica.*

*Caules suffruticosi; ramis nudiusculis. Foliola lanceolata petiolo, inter singula paria, glandula pedicellata notato. Stipulae subulatae. Pedunculi axillares foliorum, gemini, uniflori, folio breviores. Siliquae orobi. Flores circiter hexandri antheris duabus longissimis.*

27. *CASSIA* foliis multijugis linearibus: glandula basios petiolorum obsoleta, stipulis setaceis. *Fl. zeyl. des. 154. \**

*Habitat in Zeylona.*

*Petioli absque glandula. Caulis non flexuosus. Stipulae lanceolatae, seta terminatae.*

28. *CASSIA* foliis multijugis, stipulis dimidiato-cordatis. *flexuosa.*

*Senna spuria occidentalis, filiqua singulari, foliis herbæ mimosæ. Herm. lugdb. 558. Sloan. jam. 150. Raj. suppl. 18.*

*Chamæcrista pavonis brasiliiana, filiqua singulari. Breyn. cent. 65. t. 23.*

*Habitat in Brasilia. ☉*

29. *CASSIA* foliis multijugis, floribus pentandris, nictitans. *caule erecto.*

*Cassia calycibus acutis, floribus pentandris. Hort. cliff. 497. t. 36. Hort. upf. 101.*

*Senna spuria virginiana, mimosæ foliis, floribus parvis nictitantibus. Pluk. alm. 341. t. 314. f. 3.*

*Amcena incesta. Rumph. amb. 6. p. 147. t. 61. f. 1.*

*Habitat in Virginia. ☉*

*Glandula petiolorum pedicellata ♂ fusca. Pedunculi sæpius tripartiti ♂ triflori, spatio supra alas enati.*

30. *CASSIA* foliis multijugis, caule procumbente. *procum.*

*Cassia americana procumbens herbacea, mimosæ foliis, floribus parvis filiquis angustis planis. Comm. petrop. t. 11. \ bens;*

*Chamæcrista mariana, flore minore. Pet. succ. 242.*

*n. 40.*

*Habitat in Indiis, Virginiae apricis, siccis. ☉*

## POINCIANA.

- elata.** 1. POINCIANA caule inermi. *Amæn. acad.* 4. p. 313. \*  
*Poinciana* foliis bipinnatis: foliolis ovato-oblongis, caule inermi. *Hort. cliff.* 158. *Fl. zeyl.* 159.  
*Habitat in Indiis.* ‡  
*Foliola integerrima, minuta. Calyces coriacei. Flamenta atropurpurea, longissima, basi villosa.*
- bi-juga.** 2. POINCIANA aculeis solitariis, foliolis emarginatis. *Jacq. amer.* 20.  
*Frutex pavonius, Crista pavonis. Breyn. cent.* 61. t. 22. *Raj. hist.* 981.  
*Acacia gloriosa jamaicensis, foliis minoribus subrotundis, spinis ad genicula simplicibus, floribus flavo-purpureis. Pluk. alm.* 5.  
*Crista pavonis. Rumph. amb.* 4. p. 53. t. 20.  
*Habitat in Indiis.* ‡
- pulcherri-ma.** 3. POINCIANA aculeis geminis. *Hort. upf.* 101.  
*Acacia orientalis gloriosa, coluteæ foliis, ad genicula spinis gemellis aculeata. Pluk. alm.* 5.  
*Flos pavonis. Mer. sur.* 45. t. 45. ?  
*Tsietti mandaru. Rheed. mal.* 6. p. 1. t. 1.  
*Habitat in Indiis.* ‡

## CÆSALPINIA.

- brasiliensis.** 1. CÆSALPINIA caule foliisque inermibus.  
*Cæsalpinia* foliolis obovatis integerrimis. *Mat. med.* 205.  
*Cæsalpinia arborea inermis, foliis minoribus: paribus bipinnatis, ligno kermesino: Brasiletto. Brown. jam.* 227.  
*Cæsalpinia* foliolis duplicato-pinnatis, foliolis emarginatis, floribus decandris. *Mill. dict.*  
*Pseudo-Santalum croceum. Sloan. jam.* 213. *hist.* 2. p. 184. *Catesb. car.* 2. p. 51. t. 51.  
*Habitat in Carolina, jamaica, Brasilia. ‡*  
*Brasiletto officinis dicta.*
- erecta.** 2. CÆSALPINIA caule aculeato, foliolis ovatis, floribus pentandris.  
*Cæsalpinia* foliis duplicato-pinnatis, foliolis ovatis integerrimis, floribus pentandris. *Mill. dict.*

Cæ-

*Cæsalpina polyphylla aculeis horrida. Plum. gen. 26. t. 68.*

*Habitat in Jamaica. ‡*

3. *CÆSALPINIA* caule aculeato, foliolis obcordatis vesicaria, subrotundis.

*Cæsalpina spinosa, foliis minoribus obcordatis bipinnatis. Brown. jam. 227.*

*Senna spuria arborea spinosa, foliis alatis ramosis f. decompositis flore luteo, siliquis brevibus sulcatis nigris, sabinae odore. Sloan. jam. 149. hist. 2. p. 50. t. 181. f. 2, 3. bona. Raj. dendr. 11.*

*Habitat in Jamaica. ‡*

4. *CÆSALPINIA* caule aculeato, foliolis oblongis inæquilateribus emarginatis. *Fl. zeyl. 158. \* Mat. med. 203.*

*Ligno brasiliiano simile. Bauh. pin. 393.*

*Lignum Sappan. Rumph. amb. 4. p. 56. t. 21. Meist. 11. 90.*

*Tiam-pangam. Rheed. mal. 6. p. 3. t. 2.*

*Habitat in Indiis. ‡*

## GUILANDINA.

1. *GUILANDINA* aculeata, pinnis ovatis, foliolis aculeis solitariis. *Bonduc.*

*Guilandina caule fructuque aculeatis. Hort. cliff. 158.*

*Bonduc vulgare majus polyphyllum. Plum. gen. 25.*

*Acacia gloriosa, lentisci folio. spinosa, flore spicato luteo, siliqua magna muricata. Pluk. alm. 4. t. 2. f. 2.*

*Lobus echinatus, fructu flavo, foliis rotundioribus. Sloan. jam. 144. hist. 2. p. 40.*

*Frutex globulorum. Rumph. amb. 5. p. 89. t. 48.*

*Habitat in Indiis. ‡*

2. *GUILANDINA* aculeata, pinnis oblongo-ovatis, foliolis aculeis geminis. *Bonducella.*

*Guilandina aculeata, foliolis ovalibus acuminatis. Fl. zeyl. 156. \* Hort. upf. 101.*

*Bonduc vulgare minus polyphyllum. Plum. gen. 25.*

*Crista Pavonis, glycyrrhizæ folio, minor repens spinosissima, flore luteo spicato minimo, siliqua latissima echinata, semine rotundo cinereo. Breyn. prol. 3. app. 33. t. 28.*

*Linnæi Spec. plant. pars I.*

*M m •*

*Glo-*

Globuli majores. *Rumph. amb. 5. p. 93. t. 49. f. 1.*  
 Lobus echinatus fructu cæsius, foliis longioribus. *Sloan. jam. 144. hist. 2. p. 41.*

*Habitat in Indiis. §*

- Nuga. 3. **GUILANDINA** caule inermi, foliis petiolo primario subtus aculeis geminis. †

Nugæ silvarum. *Rumph. amb. 5. p. 94. t. 50.*

*Habitat in Amboina. §*

*Legumina levia sunt.*

- Moringa. 4. **GUILANDINA** inermis, foliis subbipinnatis: foliolis inferioribus ternatis. *Fl. zeyl. 155. \* Mat. med. 202.*

Lignum peregrinum aquam cæruleam reddens. *Bauh. pin. 416.*

Moringha zeylanica, foliorum pinnis pinnatis, flore majore, fructu anguloso. *Burm. zeyl. 162. t. 75.*

Morungu. *Rheed. mal. 6. p. 19. t. - Rumph. amb. 1. p. 184. t. 74, 75.*

*Habitat in Zeylona, America, Ægypto. §*

*Semina ex Asia tribus membranis longitudinaliter alata fuere, ex Africa vero membranis his carere.*

*Hæc dioica secundum Rumphium; an etiam præcedentes?*

- dioica 5. **GUILANDINA** inermis, foliis bipinnatis: basi apiceque simpliciter pinnatis. *Gen. plant. 2. 51.*

Bonduc canadense polyphyllum non spinosum, mas & femina. *Duham. arb. 1. p. 108. t. 42.*

*Habitat in Canada. §*

## GUAJACUM.

- officinale. 1. **GUAJACUM** foliolis bijugis obtusis.

Guajacum foliis pinnatis: foliolis quaternis obtusis.

*Hort. cliff. 187. Mat. med. 207.*

Guajacum foliis fere impetiolatis bijugis obovatis & leniter radiatis, pinnis & ramulis dichotomis. *Brown.*

*jam. 225.*

Guajacum flore cæruleo, fructu subrotundo. *Plum. gen. 39.*

Guajacum jamaicense, foliis velut muria conditis spissius virentibus, flore subcæruleo. *Pluk. alm. 180. t. 35. f. 4.*

Guajacum magna matrice. *Bauh. pin. 448.*

Pru

Pruno vel Evonymo affinis arbor, folio alato buxæ subrotundo. *Sloan. jam. 186. hist. 2. p. 133. t. 222.*

*f. 3. 4. 5. 6.*

Arbor ligni Sancti vel Guajacum. *Seb. thes. 1. p. 86. t. 53. f. 2.*

3. Guajacum jamaicense, lentisci subrotundis foliis late virentibus, flore albo. *Pluk. alm. 180. t. 35. f. 3.*

*Habitat in Hispaniola, Jamaica. §*

4. GUAJACUM foliolis multijugis obtusis.

*Sandum;*

Guajacum foliis pinnatis, foliolis obverse ovatis integerrimis. *Roy. lugdb. 208.*

Guajacum flore cæruleo fimbriato, fructu tetragono. *Plum. gen. 39.*

Guajacum americanum, lentisci folio. *Comm. hort. 1. p. 171. t. 88.*

Guajacum propemodum sine matrice. *Bauh. pin. 448.*  
Jasminum vulgo americanum f. Euonymo affinis occidentalis, alatis rusci foliis, nucifera, cortice adg. nacula fungoso. *Pluk. alm. 139. t. 94. f. 4.*

*Habitat in Americis insula f. Johannis de Porto Rico. §*

5. GUAJACUM foliolis multijugis acutis.

*afrum.*

Guilandinoides. *Hort. cliff. 439.*

Acacia africana, quæ Acaciæ similis, foliis myrti parvis aculeatis pinnatis, flore coccineo tetrapetaloides.

*Walth. hort. 2. t. 2.*

Afra arbor acaciæ similis, foliis myrti aculeatis splendentibus. *Boerh. lugdb. 2. p. 57.*

*Habitat in Æthiopia, China. §*

*Flores & fructus non vidi, hinc de genere dubius.*

## CYNOMETRA.

1. CYNOMETRA trunco florifero. *Act. upf. 1741. p. cæuliflora.*  
*79. Fl. zeyl. 166.*

Cynomorium. *Rumph. amb. 1. p. 163. t. 62.*

*Habitat in India. §*

2. CYNOMETRA ramis floriferis. *Act. upf. 1741. p. ramiflora.*  
*79. Fl. zeyl. 167.*

Cynomorium sylvestre. *Rumph. amb. 1. p. 164. t. 63.*

*Iripa. Rheed. mal. 4. p. 65. t. 31.*

*Habitat in India. §*

## ANACARDIUM.

- occidentale. 1. ANACARDIUM. *Hort. cliff.* 161. *Fl. zeyl.* 165.  
*Mat. med.* 206. *Hort. upf.* 102. *Roy. lugdb.* 463.  
*Brown. jam.* 227.  
 Anacardii alia species. *Bauh. pin.* 512.  
 Pomitera f. potius Prunifera indica, nuce reniformi.  
*Catesb. car.* 3. p. 9. t. 9.  
 Acaiou. *Pif. bras.* 58. *mant.* 193.  
 Castuvium. *Rumph. amb.* 1. p. 177. t. 69.  
 Casthou. *Mer. surin.* 16. t. 16.  
 Kapa-mava. *Rheed. mal.* 3. p. 65. t. 54.  
 Habitat in Indiis. §

## SWIETENIA.

- Mahagoni. 1. SWIETENIA. *Jacqu. amer.* 20. †  
 Cedrela foliis pinnatis, floribus sparsis, ligno odorato. *Brown. jam.* 158.  
 Arbor foliis pinnatis nullo impari: nervo ad latus.  
*Catesb. car.* 2. p. 81. t. 81.  
 Habitat in America meridionali: §  
 Affinitate proxima jungitur Cedrelæ.

## DICTAMNUS.

- albus. 1. DICTAMNUS. *Hort. cliff.* 161. *Hort. upf.* 102.  
*Mat. med.* 208. *Roy. lugdb.* 463. *Sauv. monsp.* 232.  
 Dictamnus albus vulgo Fraxinella. *Bauh. pin.* 222.  
 Fraxinella. *Renal. spec.* 122. t. 121.  
 Habitat in Germania, Gallia, Italia. ‡  
 Ordo naturalis Rutæ communis.

## RUTA.

- graveolens. 1. RUTA foliis decompositis, petalis laceris, floribus lateralibus quadrifidis. *Hort. cliff.* 145. *Hort. upf.* 102. *Mat. med.* 209. *Roy. lugdb.* 461.  
 Ruta sylvestris major. *Bauh. pin.* 336. *Morif. hist.* 2. p. 507. f. 5. t. 14. f. 3.  
 β. Ruta chalepensis tenuifolia, florum petalis villis scattentibus. *Morif. hist.* 2. p. 509.  
 Ruta hortensis altera. *Bauh. pin.* 336. *Raj. hist.* 874.  
 γ. Ruta hortensis latifolia. *Bauh. pin.* 336.  
 Ruta graveolens hortensis. *Dod. pempt.* 119.  
 δ. Ruta hortensis latifolia arbusculæ similis. *Boerh. lugdb.* 1. p. 260.  
 Ru-

*Ruta africana maxima. Schwer. hort. 24.*

*Habitat in Europæ australis, Alexandria, Mauritania sterilibus. ‡*

*Flos primarius quinquesfidus, reliqui quadrifidi. Phil. Bot. 178. Paniculae rami primi bipartiti; reliqui flores alterni.*

*Confer: RUTA sylvestris minor. Bauh. pin. 336.*

*Cam. epit. 495. Ruta montana legitima. Clus. hist. 2. p. 136. app. alt. fig. structura eadem, facie distincta, descripta Læsl. it. 140.*

2. *RUTA foliis terminatis sessilibus.*

patavina.

*Pseudo-Ruta patavina trifolia, floribus luteis umbellatis. Mich. gen. 22. t. 19.*

*Habitat prope Patavium.*

3. *RUTA foliis simplicibus indivisis.*

linifolia.

*Ruta foliis simplicibus solitariis. Hort. cliff. 146.*

*Ruta sylvestris linifolia hispanica. Bocc. mus. 2. p. 82. t. 73. Barr. ic. 1186.*

β. *Ruta montana, foliis integris subrotundis. Buxb. cent. 2. p. 30. t. 28. f. 2.*

*Ruta orientalis, linariæ folio, flore parvo. Tournef. cor. 19. Buxb. cent. 2. p. 30. t. 28. f. 1.*

*Habitat in Hispania; β. in Rodastro; γ. in Media. 24*

*Paniculae rami laterales bifidi, triflori, flore intermedio sessili. Petala 5, sessilia, integra, ovata. Filamenta basi pilosa arcuata includunt receptaculum poris melliferis pertusum. Pericarpia punctis elevatis.*

## TOLUIFERA.

1. *TOLUIFERA. Mat. med. 201.*

Balsamum.

*Balsamum Tolutanum, foliis ceratiæ similibus. Bauh. pin. 401.*

*Habitat in America prope Carthagenam. ‡*

## HÆMATOXYLUM.

1. *HÆMATOXYLUM. Hort. cliff. 160. Roy. lugdb. 465.*

Campechianum.

*Lignum campechianum, species quædam. Sloan. jam. 213. hist. 2. p. 183. t. 10. f. 1. 2. 3. 4. Catesb. car. 3. p. 66. t. 66.*

*Habitat in Campeche Americæ. ‡*



## ADENANTHERA.

- payonina. 1. *ADENANTHERA* foliis decompositis utrinque glabris. *Roy. lugdb.* 462. *Vir. cliff.* 36. *Fl. zeyl.* 160. \*  
*Poinciana* foliis duplicato-pinnatis: foliolis alternis.  
*Hort. cliff.* 158.  
*Corallaria parvifolia.* *Rumph. amb.* 3. p. 173. t. 109.  
*Mandsiadi.* *Rheed. mal.* 6. p. 25. t. 14. *Raj. hist.* 1752.  
*Habitat in India.* ‡
- Falcataria. 2. *ADENANTHERA* foliis decompositis subtus tomentosis.  
*Clypearia alba.* *Rumph. amb.* 3. p. 176. t. 111.  
*Habitat in India.* ‡

## MELIA.

- Azedarach. 1. *MELIA* foliis bipinnatis. *Fl. zeyl.* 162. \* *Gron. orient.* 133.  
*Melia* foliis decompositis. *Hort. cliff.* 161. *Roy. lugdb.* 462.  
*Azedarach.* *Dod. pempt.* 848.  
*Arbor fraxini folio, flore cæruleo.* *Bauh. pin.* 415.  
*Pseudo-Sycomorus.* *Cam. epit.* 181.
- sempervirens 2. *Azedarach sempervirens* & *florens.* *Tournef. inst.* 616.  
*Azadirachta indica, foliis ramosis minoribus, flore albo subcæruleo purpurascente majore.* *Comm. hort.* 1. p. 147. t. 76.  
*Habitat in Syria, β. in Zeylona.* ‡
- Azadirachta* 2. *MELIA* foliis pinnatis. *Hort. cliff.* 161. *Fl. zeyl.* 161.  
\* *Roy. lugdb.* 462.  
*Olea malabarica, fraxini folio.* *Pluk. alm.* 269. t. 247. f. 1.  
*Azedarach foliis falcato-serratis.* *Burm. zeyl.* 40. t. 15.  
*Azadirachta indica, folio fraxini.* *Breyn. ic.* 21. t. 15.  
*Arbor indica, fraxino similis, oleæ fructu.* *Bauh. pin.* 416.  
*Aria Bepou.* *Rheed. mal.* 4. p. 107. t. 52.  
*Habitat in India.* ‡

## TRICHILIA.

- hirs. 1. *TRICHILIA* foliis pinnatis subhirsutis, foliolis obovatis.  
*Trichilia subhirsuta, foliis pinnatis ovatis, racenis alaribus.* *Brown. jam.* 278.  
2. *Evonymus* caudice non ramofo, folio alato, fructu rotundo tripyreno. *Sloan. jam.* 171. *hist.* 2. p. 103. t. 210. f. 2, 3. *Raj. dendr.* 70.  
*Habitat in Jamaica.* ‡

2. **TRICHILIA** foliis ternatis. *Jacq. hist. t. 82.* trifoliata.  
*Trichilia Halesia. Læff. it. 188. \**  
*Habitat in America meridionali. ♀*
3. **TRICHILIA** foliis pinnatis glabris, floribus octandris. *Guara.*  
*Trichilia foliis oblongo-ovatis pinnatis nitidis, racemis*  
*laxis. Brown. jam. 279.*  
*Melia guara floribus octandris. Jacq. amer. 20.*  
*Guidonia, nucis juglandis foliis, major. Plum. gen.*  
*4. ic. 147. f. 2.*  
*Habitat in Brasilia aliisque indiæ occidentalis. ♀*  
*Calyx planiusculus. Petala 4, linearia. Nectarium lon-*  
*gitudine corollæ; intra ejus os Filamenta cum anthe-*  
*ris brevissimis. Receptaculum subcylindricum, coro-*  
*natum margine glanduloso. Stigma orbiculatum. Fru-*  
*ctus ignotus Flores octandri.*

## ZYGOPHYLLUM.

1. **ZYGOPHYLLUM** foliis petiolatis: foliolis obova- *Fabago.*  
 tis, caule herbaceo.  
*Zygophyllum capsulis prismaticeo pentahædri. Hort.*  
*sp. 103. Sp. pl. 1. p. 385. Gron. orient. 132.*  
*Zygophyllum foliis petiolatis. Hort. cliff. 160. Roy.*  
*lugdb. 460.*  
*Capparis portulacæ folio. Bauh. pin. 480.*  
*Capparis Fabago. Dod. pempt. 741.*  
*Habitat in Syria, Mauritania. 24*  
*Folia ovata, petiolata. Petala obtusa.*
2. **ZYGOPHYLLUM** foliis petiolatis: foliolis linea- *coccineum.*  
 ribus carnosiss.  
*Zygophyllum capsulis cylindricis. Sp. pl. 1. p. 386.*  
*Fabago arabica teretifolia, flore coccineo. Shaw. afr.*  
*231. f. 231.*  
*Habitat in Africa.*  
*Distincta a Z. albo floribus rubris, Capsulis oblongis.*
3. **ZYGOPHYLLUM** foliis petiolatis: foliolis clava- *album.*  
 tis carnosiss. *Linn. dec. 1. t. 8. \**  
*Habitat in Ægypto. Roque. ♀*
4. **ZYGOPHYLLUM** foliis subpetiolatis: foliolis ob- *Mergfana.*  
 ovatis, caule fruticoso.  
*Zygophyllum capsularum angulis compresso-mem-*  
*branaceis. Sp. pl. 1. p. 385.*  
 M m 4 Fa-

*Fabago triphylla* & *tetraphylla*, flore tetrapetalo, fructu membranaceo quadrangulari. *Burm. afr.* 7. t. 3. f. 2.

*Fabago capensis frutescens major.* *Dill. elth.* 142. t. 116. f. 141.

*Planta africana frutescens*, portulacæ foliis, Morgfani fyrorum, ex brevi pediculo binis. *Pluk. amalh.* 173. t. 429. f. 4.

*Habitat in Æthiopia.* §

*Folia obverse ovata, subpetiolata; nec perfecte sessilia. Petala acuta.*

sessilifolium. 5. **ZYGOPHYLLUM** foliis sessilibus: foliolis lanceolato-ovalibus margine scabris, caule fruticoso.

*Zygophyllum capsulis globoso-depressis.* *Sp. pl.* 1. p. 385.

*Zygophyllum foliis sessilibus.* *Hort. eliff.* 160. *Roy. lugdb.* 460.

*Fabago capensis frutescens minor.* *Dill. elth.* 142. t. 116. f. 142.

*Fabago humilis quadrifolia glabra*, flore albedo, fructu rotundo. *Burm. afr.* 4. t. 2. f. 1.

*Fabago africana arborescens*, flore sulphureo, fructu rotundo. *Comm. rar.* 10. t. 10.

β. *Zygophyllum fulvum capsulis ovatis acutis.* *Sp. pl.* 1. p. 386.

*Fabago flore luteo, petalorum unguibus rubris, fructu sulcato acuto, oblongo.* *Burm. afr.* 6. t. 3. f. 1.

*Habitat in Æthiopia.* §

*Folia sessilia: Foliola ovali-lanceolata, margine cartilagineo crenata s. scabra. Caulis angulatus. Pedunculi solitarii, foliis longiores. Petala crenata.*

spinosum. 6. **ZYGOPHYLLUM** foliis sessilibus: foliolis linearibus, caule suffruticoso.

*Zygophyllum caule aculeato.* *Sp. pl.* 1. p. 386.

*Fabago tenuifolia spinosa*, fructu rotundo. *Burm. afr.* 5. t. 5.

*Habitat in Æthiopia.* §

*Folia linearia, acuta.*

æstivans. 7. **ZYGOPHYLLUM** foliis sessilibus: foliolis obovatis retusis.

*Habitat in Surinamo. Rolander; sed impubes perit.*

*Caules herbacei, pedales, laves, diffus, teretiusculi, late-*

*latere superiore plani. Folia opposita, conjugata, sessilia: Foliola obovata, avenia, retusq. Stipulæ quinæ, reflexæ; duabus inter folia a latere superiore; unica inter folia a latere inferiore; unica inter foliorum utrumque par.*

## Q VASSIA.

## 1. QVASSIA.

amara.

*Habitat in Surinamo C. Dahlberg. ♂*

## FAGONIA.

## 1. FAGONIA spinosa, foliolis lanceolatis planis læceticis.

*vibus. Hort. upf. 103.*

*Fagonia spinosa. Hort. cliff. 160.*

*Trifolium spinosum creticum. Bauh. pin. 330. prodr.*

*142. Clus. hist. 2. p. 212.*

*Habitat in Creta. ☉*

## 2. FAGONIA inermis.

hispanica.

*Fagonia hispanica non spinosa. Tournef. inst. 265.*

*Habitat in Hispania. ♂*

## 3. FAGONIA spinosa, foliolis linearibus convexis. arabica.

*Shaw. afr. 229.*

*Fagonia arabica; longissimis aculeis armata. Shaw.*

*afr. 229.*

*Habitat in Arabia.*

## TRIBULUS.

1. TRIBULUS foliolis subquadrijugis: exterioribus maximis majoribus. *Lxfl. it. 208.*

*Tribulus foliolis trium parium pinnatis. Hort. cliff. 160.*

*Tribulus foliis senis pinnatis: extimismajoribus, floribus singularibus. Brown. jam. 220. t. 132. f. 3.*

*Tribulus terrestris major, flore maximo odorato.*

*Sloan. jam. 90. hist. 1. p. 209. t. 132. f. 1. Raj. suppl. 650.*

*Tribulus terrestris fructu turbinato, foliis lanuginosis.*

*Plum. spec. 7. ic. 254. f. 1.*

*Habitat in Jamaica aridis. ☉*

*Fructus decaspermus, levis f. muticus.*

## 2. TRIBULUS foliolis quinquejugatis subæqualibus, lanuginosis seminibus bicornibus.

M m 5

Tri-

*Tribulus foliolis* quinque parium. *Fl. zeyl.* 168. \*

*Tribulus terrestris zeylanicus*, cauliculis villosis, fructu acuminato. *Burm. zeyl.* 226. t. 106. f. 1.

*Habitat in Zeylona.*

terrestris. 3. *TRIBULUS* foliolis sexjugatis subæqualibus, seminibus quadricornibus *Hort. cliff.* 160. *Hort. upf.* 103.

*Roy. lugdb.* 460. *Sauv. monsp.* 227. *Brown. jam.* 220.

*Tribulus terrestris*, ciceris folio, fructu aculeato. *Bauh. pin.* 250.

*Tribulus terrestris.* *Lob. ic.* 2. p. 84.

*Tribulus terrestris minor incanus hispanicus.* *Barr. rar.* 54. t. 558.

*Habitat in Europa australi ad semitas.* ☉

Cistoides. 4. *TRIBULUS* foliolis octojugatis subæqualibus.

*Tribulus foliis octo parium pinnatis.* *Roy. lugdb.* 460.

*Tribulus terrestris major curassavicus.* *Herm. par.* 236. t. 236.

*Tribulus terrestris americanus*, argemones flore flavo. *Pluk. alm.* 374. t. 64. *Raj. suppl.* 649.

*Habitat in America calidiore.*

## THRYALLIS.

brasiliensis. 1. *THRYALLIS.*

Fruticescens herba pisonis. *Marcgr. bras.* 79. f. 3.

*Habitat in Brasilia.* †

Fruticulus ramis teretibus, articulatis. Folia opposita, petiolata, ovata, integerrima. *Stipulæ setaceæ.*

Racemus terminalis, e dichotomia ramorum, simplex, pedalis: Bracteis setaceis, brevissimis; Pedicellis filiformibus, flore longioribus. Flores parvi lutei. Fructus tricocci.

## LIMONIA.

acidissim a 1. *LIMONIA.*

*Limonia malus sylvestris zeylanica*, fructu pumilo.

*Burm. zeyl.* 143.

*Schinus foliis pinnatis*: petiolo marginato articulato, spinis axillaribus solitariis. *Fl. zeyl.* 175. \*

*Anisifolium.* *Rumph. amb.* 2. t. 43.

Catu Tsiern Naregam. *Rheed. mal.* 4. p. t. 14.

*Habitat in India.* †

## MONOTROPA.

1. MONOTROPA floribus lateralibus octandris; ter. Hypopithys  
 minali decandro. *Fl. succ.* 329; 351. *Phil. bot.* §.  
 178.  
 Orobanche quæ Hypopithys dici potest. *Bauh. pin.*  
 88. *prodr.* 31.  
 Orobanche Hypopithys lutea. *Mentz. pug.* 3. f. 5.  
 Orobanche flore brevior duplici, verbasculi odore.  
*Morif. hist.* 3. p. 504. f. 12. t. 16. f. 20.  
 Orobanche verbasculi odore. *Plot. oxf.* 146. t. 9. f. 6.  
*Pluk. alm.* 273. t. 209. f. 5.  
*Habitat in Sveciæ, Germaniæ, Angliæ, Canadæ syl-*  
*vis Parasitica radicum.*
2. MONOTROPA caule unifloro decandro. uniflora.  
 Monotropa flore nutante. *Grøn. virg.* 41.  
 Orobanche monanthos virginiana, flore majore penta-  
 petalo. *Morif. hist.* 3. p. 502. f. 12. t. 16. f. 5. *mal.*  
 Orobanche virginiana, flore pentapetalo cernuo. *Pluk.*  
*alm.* 273. t. 209. f. 2. *Catesb. car.* 1. p. 36. t. 36. *bon.*  
*Habitat in Marilandia, Virginia, Canada.*

## JUSSIÆA.

1. JUSSIÆA repens, floribus pentapetalis decandris, repens.  
 pedunculis folio longioribus. *Fl. Zeyl.* 169. \*  
 Nir-Carambu. *Rheed. mal.* 2. p. 99. t. 51. *Raj. hist.*  
 1510.  
*Habitat in India.*
2. JUSSIÆA erecta, floribus pentapetalis, pedunculis peruviana.  
 foliosis.  
 Onagra laurifolia, flore amplo pentapetalo. *Few. peruv.*  
 716. t. 11.  
*Habitat in Lima.*
3. JUSSIÆA erecta villosa, floribus pentapetalis decan- pubescens.  
 dris sessilibus. *Mill. dict. Lxfl. it.* 282. n. 201.  
 Lyfimachia lutea erecta non papposa major, foliis hir-  
 sutis, fructu caryophylloide. *Sloan. jam.* 85.  
*Habitat in America.*
4. JUSSIÆA erecta villosa, floribus tetrapetalis octan- suffruticosa.  
 dris pedunculatis.  
 Ludwigia capsulis oblongis uncialibus. *Roy. lugdb.*  
 252.

Ly-

*Lyfimachia indica* non papposa, flore luteo minimo, siliquis caryophyllum aromaticum æmulantibus.  
*Herm. lugdb.* 396.

Carambu. *Rheed. mal.* 2. p. 55. t. 49. *Raj. hist.* 1510.  
*Habitat in India.*

erecta. 5. *JUSSLÆA* erecta glabra, floribus tetrapetalis octandris sessilibus. *Fl. zeyl.* 170. *Hort. upf.* 103.

*Ludwigia* capsulis oblongis. *Hort. cliff.* 491.

*Onagra persicæ* foliis amplioribus, parvo flore luteo.  
*Plum. spec.* 7. t. 175. f. 2.

*Lyfimachia lutea* non papposa erecta, foliis glabris, fructu caryophylloide. *Sloan. jam.* 85. *hist.* 1. p. 37. t. 11 f. 1.

*Jasminum catalanicum*, flore luteo. *Seb. thes.* 1. p. 42. t. 24. f. 3.

• *Herba vitiliginum.* *Rumph. amb.* 6. p. 49. t. 21. f. 1.

*Habitat in America* ☿ forte in *Virginia.* ☉

## QUISQUALIS.

indica. 1. *QVISQUALIS.*

*Ovis qualis.* *Rumph. amb.* 5. p. 71. t. 38.

*Habitat in India.* ☿

Rami teretes, pubescentes. Folia opposita, petiolata, cordata s. ovata, integerrima. Stipulæ nullæ. Spica terminalis ☿ axillares pedunculatæ Bracteis ovato-oblongis, oppositis, Floribusque oppositis statura *Daphnes*, calyce filiformi extus pubescente.

## DAIS.

cotinifolia. 1. *DAIS.* *D. v. Royen.*

*Habitat ad Cap. b. Spei.* ☿

Folia opposita, obovata, integerrima, glabra, petiolata. Flores fasciculo aggregati, terminales, pubescentes: Involucro gemmaceo, quadrivalvi. Affinis *Pasferinis.*

## BUCIDA.

Buceras. 1. *BUCIDA.* *Aman. acad.* 5. p. 397. \*

*Buceras* ramulis flexuosis tenuioribus, foliis obovatis confertis, spicis plurimis terminalibus. *Brown. jam.* 221. t. 23. f. 1.

*Mangle julifera*, foliis subrotundis versus summum latissimis confertim nascentibus, cortice ad coriaden-

deſignanda utili. *Sloan. jam. 156. hiſt. 2. p. 67. t. 189.*

*f. 3. Raj. dendr. 116.*

*Habitat in Jamaica. †*

*Affinis Rhizophoræ.*

## COPAIFERA.

## 1. COPAIFERA.

*officinalis.*

*Copaiva. Mat. med. 513. Jacq. amer. 21. hiſt. t. 86.*

*Coapoiba. Marcgr. braſ. 131. Piſ. braſ. 118. Raj. hiſt. 1593.*

*Habitat in Braſilia, Antillis. †*

## SAMYDA.

## 1. SAMYDA floribus decandris, foliis ovato-oblongis parviflora. utrinque glabris.

*Samyda parviflora. Læſt. it. 260.*

*Samyda foliis ovatis cum acumine, fructibus plurimis minoribus confertis. Brown. jam. 217.*

*Caffearia floribus decandris. Jacq. amer. 21. hiſt. t. 85.*

*Arbor baccifera foliis oblongis acuminatis, floribus confertim ex alis foliorum erumpentibus, fructu minimo croceo. Sloan. jam. 137. hiſt. 2. p. 108. t. 211. f. 2. Raj. dendr. 108.*

*Habitat in America. †*

## 2. SAMYDA floribus octandris, foliis cordatis glabris. nitida.

*Samyda foliis nitidis cordatis læviſſime crenatis rudimentis mollibus rubentibus, racemis tenuioribus alaribus. Brown. jam. 217. t. 23. f. 3. Guilandina.*

*Habitat in America. †*

## 3. SAMYDA floribus octandris, ramis spinosis. spinosa.

*Caffearia aculeata, floribus octandris. Jacq. amer. 21.*

*Guidonia, aurantii foliis, aculeata. Plum. gen. 4. ic. 147. f. 1.*

*Habitat in America.*

## 4. SAMYDA floribus dodecandris, foliis obovatis sub-pubescentibus tomentosis.

*Samyda foliis ovatis villoſis, floribus confertis, fasciculis ſparſis. Brown. jam. 218. ?*

*Frutex baccifer folio oblongo integro, flore pentapetalo pallide luteo odoratiſſimo. Sloan. jam. 173. hiſt. 2. p. 109. ? Raj. dendr. 75. ?*

*Habitat in America.*

5. SA



- ferrulata.* 5. **SAMYDA** floribus dodecandris, foliis ovato-oblongis ferrulatis.  
*Samyda floribus dodecandris. Jacq. amer. 21.*  
*Guidonia ulmi folio, flore roseo niveo. Plum. gen. 4.*  
*ic. 146. f. 2.*  
*Habitat in America. ♀*

## MELASTOMA.

- Acinodendron.* 1. **MELASTOMA** foliis denticulatis subtrinerviis ovatis acutis.  
*Melastoma foliis ovato-lanceolatis crenatis nervis quinque longitudinalibus: extimis obsoletioribus. Hort. cliff. 162.*  
*Acinodendrum americanum pentaneuron: foliis crassius hirsutis ad ambitum rarioribus ferris. Pluk. mac. 4. t. 159. f. 1.*  
*Grossularia alia, plantaginis folio, fructu rariore violaceo. Plum. ic. 142. f. 2.*  
*Grossulariæ fructu arbor maxima non spinosa, malabathri folio maximo inodoro, flore racemoso albo. Sloan. jam. 164. hist. 2. p. 84 t. 196. f. 1. Ray. densif. 26.*  
*Habitat in America calidiore. ♀*
- Grossularioides.* 2. **MELASTOMA** foliis denticulatis triplinerviis ovatis acuminatis.  
*Arbor americana, latiore & acuminato folio trinervio, utrinque glabro & margine leviter crenato. Pluk. alm. 40. t. 249. f. 1.*  
*Habitat in Surinamo D. Bartsch.*
- scabrofolia.* 3. **MELASTOMA** foliis denticulatis quinquenerviis cordatis scabris subtus tomentosis, ramis tomentoso villosis.  
*Melastoma subhirsuta, foliis cordatis scabris minutissime denticulatis reticulatis, racemis minoribus alaribus. Brown. jam. 219. t. 24. f. 3.*  
*Habitat in Jamaica. ♀*
- sessilifolia.* 4. **MELASTOMA** foliis integerrimis trinerviis spatulatis sessilibus subtus subtomentosis.  
*Melastoma foliis amplioribus per petiolum recurrentibus & contractis, fasciculis florum sparsis. Brown. jam. 219. t. 24. f. 1, 2.*

Arbor americana, foliis a concurfu nervorum ima parte longius productis, subtus lanugine candicante tomentosis. *Pluk. phyt.* 249. f. 2.

*Habitat in Jamaica.* §

5. MELASTOMA foliis denticulatis quinquenerviis hirta. ovato-lanceolatis, caule hispido.

Grossularia plantaginis folio angustiore hirsuto. *Phen. spec.* 18. ic. 141.

Grossulariæ fructu non spinosa, malabathri foliis longa & rufa lanugine hirsutis, fructu majore cæruleo. *Sloan. jam.* 165. *hist.* 2. p. 85. t. 197. f. 2. *Raj. dendr.* 74.

Arbuscula jamaicensis quinquenervis, minutissime dentatis foliis & caule pubescentibus. *Pluk. alm.* 40. t. 264. f. 1. vel potius. 265. f. 1.

*Habitat in America meridionali* §

*Flores in foliorum alis breviter pedunculati.*

6. MELASTOMA foliis integerrimis trinerviis oblongo-ovatis subtus tomentosis, racemis brachiatis, spicis bipartitis. holosericea.

Melastoma foliis integerrimis ovato-lanceolatis subtus sericeis: nervis ante basin coadunatis. *Hort. cliff.* 162.

Acinodendrum americanum, ampliore folio trinervi, inferius alba lanugine incano maximo utrinque glabro. *Pluk. mant.* 4. t. 250. f. 2.

Arbor racemosa brasiliiana, folio malabathri. *Breyn. cent.* 3. t. 2. 4.

*Muiwa. Maregr. bras.* 117.

*Habitat in Brasilia, Jamaica, Surinamo.* §

Arbor Ramis tomentosis. Folia petiolata, trinervia excepto marginali nervo, reticulata, subtus tomentosa uti petioli. Racemi terminales e dichotomia brachiati: spicis lateralibus conjugatis secundis, flavis.

7. MELASTOMA foliis integerrimis quinquenerviis malabathri-lanceolato-ovatis scabris. *Fl. zeyl.* 171. ca.

Melastoma quinquenervia hirta major, capitulis sericeis villosis. *Burm. zeyl.* 155. t. 73.

*Kadali. Rheed. mal.* 4. p. 87. 42.

*Habitat in India.* §

8. MELASTOMA foliis integerrimis quinquenerviis lanceolato-ova-

ovato-oblongis læviusculis acuminatis: margine lævibus.

*Melastoma fruticosa minor.* foliis tenuibus ovatis, racemis terminalibus. *Brown. jam.* 219.

*Grossulariæ* fructu arbor non spinosa, malabathri folio maximo glabro & splendente. *Sloan. jam.* 165. *hist.* 2. p. 85. t. 197. f. 1. *Raj. dendr.* 26.

*Grossularia plantaginis* folio lato, fructu minimo cæruleo. *Plum. spec.* 18. t. 140. †

*Habitat in America.* ‡

*Folia tenuiora* ♀ minus consistentia. *Racemi terminales compositi Floribus minutis.*

discolor. 9. **MELASTOMA** foliis integerrimis quinquenerviis oblongo-ovatis glabris margine lævibus. *Jacq. hist.* t. 84.

*Grossularia plantaginis* folio apicibus candidis & falcatis. *Plum. ic.* t. 42. f. 1.

*Grossularia* fructu non spinosa, malabathri foliis subtus niveis, fructu racemoso umbellæ modo disposito. *Sloan. jam.* 165. *hist.* 2. p. 86. t. 198. f. 1.

Arbor americana quinquenervia, comantibus flosculis, foliis amplissimis glabris prona parte albicantibus.

*Pluk. alm.* 40. t. 264. f. 4.

*Habitat in America.* ‡

aspera. 10. **MELASTOMA** foliis integerrimis trinerviis lanceolatis scabris. *Fl. zeyl.* 172.

*Fragarius ruber.* *Rumph. amb.* 4. p. 135. t. 71.

*Katou-Kadali.* *Rheed. mal.* 4. p. 91. t. 43. *Raj. hist.* p. 1493.

*Habitat in India.* ‡

œandra. 11. **MELASTOMA** foliis integerrimis trinerviis ovatis glabris margine hispidis. *Fl. zeyl.* 173. \* *Osb. it.* 213.

*Habitat in India.* ‡

crispata. 12. **MELASTOMA** foliis integerrimis quinquenerviis quaternis, ramis crispatis. †

*Funis murænarum.* *Rumph. amb.* 5. p. 66. t. 35.

*Habitat in Amboina.* ‡

## KALMIA.

latifolia. 1. **KALMIA** foliis ovatis, corymbis terminalibus  
*Amen.*

*Amæn. acad.* 3. p. 13. \* *Kalm. it.* 2. p. 43. *Æ* 3. p. 148.

*Andromeda* foliis ovatis obtusis, corollis corymbosis infundibuliformibus, genitalibus declinatis. *Gron. virg.* 160.

*Chamæ Daphne* foliis tini, floribus bullatis. *Catesb. car.* 2. p. 98. t. 98.

*Ledum* floribus bullatis confertis in summis caulibus. *Trew. chret. t.* 38. f. 1.

*Cistus Chamæ Rhododendros mariana* laurifolia, floribus expansis summo ramulo in umbellam plurimis. *Pluk. alm.* 49. t. 379. f. 6.

*Habitat in Marilandia, Virginia, Pensylvania.* *h*

2. *KALMIA* foliis lanceolatis, corymbis lateralibus. angustifolia. *Amæn. acad.* 3. p. 14. *Kalm. it.* 3. p. 147.

*Azalea* foliis lanceolatis integerrimis non nervosis glabris, corymbis terminalibus. *Gron. virg.* 21.

*Chamæ Daphne* sempervirens, foliis oblongis angustis, foliorum fasciculis oppositis. *Catesb. car.* 3. p. 17. t. 17. f. 1.

*Ledum* floribus bullatis fasciculatis ex alis oppositis foliorum. *Trew. chret. t.* 38. f. 2.

*Cistus sempervirens* laurifolia, floribus eleganter bullatis. *Pluk. alm.* 106. t. 161. f. 3.

*Anonyma.* *Cold. noveb.* 100.

*Habitat in Pensylvania, Nova Cæsarea, Noveboraco* *h*

## L E D U M.

1. *LEDUM.* *Fl. lapp.* 160. *Fl. suec.* 2. n. 352. \* palustre.

*Ledum* foliis linearibus subtus hirsutis, floribus corymbosis. *Fl. suec.* 1. n. 341; 352. *Mat. m. d.* 210.

*Cistus Ledon* foliis rosmarini ferrugineis. *Bauh. pin.* 467.

*Roimarinum* sylvestre. *Cam. epit.* 546.

*Habitat in Europæ septentrionalis, paludibus uliginosis.* *h*

## R H O D O R A.

1. *RHODORA.* † canadensis.

*Chamærhododendros.* *Duham. sem. app.* 10. t. 27. f. 2.

*Habitat in Canada.* *h*

*Plantam non vidi; fide Cl. Duhamelii assumpsi.*

N n

RHO-

## RHODODENDRON.

- ferrugineum.** 1. RHODODENDRON foliis glabris subtus leprosis, corollis infundibuliformibus.  
*Azalæa maculis ferrugineis subtus adpersa, floribus decandris. Lemonier. Sauv. monsp. 57.*  
*Ledum foliis glabris, flore tubuloso. Hall. helv. 417.*  
*Ledum alpinum, foliis ferrea rubigine nigricantibus. Bauh. pin. 468. Raj. hist. 1005.*  
*Evonymus theophrasti. Dalech. hist. 27.*  
*Montana allobrogum lentiscifolia. Lob. ic. 366.*  
*Habitat in Alpibus Helveticis, Allobrogicis, Pyrenæis. ‡*
- dauricum.** 2. RHODODENDRON foliis glabris utrinque nudis, corollis rotatis.  
*Chamærhododendros folio glabro majusculo, amplo flore roseo. Amm. ruth. 181. t. 21.*  
*Habitat in Dauria. ‡*
- hirsutum.** 3. RHODODENDRON foliis ciliatis nudis, corollis infundibuliformibus.  
*Ledum foliis glabris ciliatis ovatis, flore tubuloso. Hall. helv. 418.*  
*Ledum alpinum hirsutum. Bauh. pin. 468. Raj. hist. 1005.*  
*Ledum alpinum. Clus. hist. 1. p. 82.*  
*Balsamum alpinum Lob. ic. 468.*  
*Habitat in Alpibus Helveticis, Austriacis, Stiriacis. ‡*
- Chamæcistus.** 4. RHODODENDRON foliis ciliatis, corollis rotatis.  
*Ledum foliis serpylli ad margines cili inftar pilosis, flore purpureo. Mich. gen. 225. t. 106.*  
*Chamæcistus hirsuta. Bauh. pin. 466.*  
*Chamæcistus VIII. Clus. hist. 1. p. 76.*  
*Cistus pumilus montis baldi. Bauh. hist. 2. p. 19. Raj. hist. 1005.*  
*Cistus Chamærhododendros foliis confertis ferrea rubigine nitentibus margine pilosis. Pluk. alm. 106. t. 23. f. 4.*  
*Habitat in Baldo ☿ prope Salisburgum. ‡*
- ponticum.** 5. RHODODENDRON foliis nitidis lanceolatis utrinque glabris, racemis terminalibus.  
*Chamærhododendros pontica maxima, folio lauroce-rasi. Tournef. cor. 42. it. 2. p. 99.*

*Habitat in Oriente, Gibraltariæ subhumidis umbrosis.*

*Alstræmer. 5*

Folia *Pruni-Laurocerasi*, perennia, acuta: Petiolis brevibus, glabris. Racemus terminalis, longitudine pedicellorum s. foliorum: Pedicellis alternis, nudis. Calyx minutissimus. Corolla campanulata, magnitudine *Nerii*, purpurea. Stamina declinata, longitudine corollæ.

6. RHODODENDRON foliis nitidis ovalibus obtusis maximum, venosis margine acuto reflexo, pedunculis unifloris.

Ledum laurocerasi folio. *Amæn. acad. 2. p. 200.*

Kalmia foliis lanceolato-ovatis nitidis subtus ferrugineis, corymbis terminalibus. *Mill. dict. 1. 229.*

Chamærhododendros, lauri folio, sempervirens, floribus bullatis corymbosis. *Catesb. car. 3. p. 17. t. 17. f. 2.*

*Habitat in Virginia. Collinson. 5*

Rami & Folia magnitudine & consistentia Pudi Laurocerasi dicti, venis reticulatis inscripta, margine reflexa.

## ANDROMEDA.

1. ANDROMEDA pedunculis solitariis lateralibus, corollis campanulatis, foliis oppositis imbricatis obtusis revolutis. tetragona.

Andromeda foliis quadrifariam imbricatis obtusis. *Fl. svec. 338; 356.*

Andromeda, foliis quadrifariam imbricatis obtusis, ex alis florens. *Fl. lapp. 166. t. 1. f. 4. Hort. cliff. 163.*

*Habitat in Alpibus Lapponicis. 24*

2. ANDROMEDA pedunculis solitariis terminalibus, hypnoides, corollis campanulatis, foliis confertis subulatis. *Oed. dan. t. 10.*

Andromeda foliis aciformibus confertis. *Fl. lapp. 165. t. 1. f. 3. Fl. svec. 337; 355.*

*Habitat in Alpibus Lapponicis. 24*

3. ANDROMEDA pedunculis aggregatis, corollis ovatis, foliis sparsis linearibus obtusis planis. cerulea.

Andromeda foliis linearibus obtusis sparsis. *Fl. lapp. 165. t. 1. f. 5. Fl. svec. 336, 354. Hort. cliff. 163.*

Andromeda rarior norvegica. *Burser. Amoen. acad. 1. p. 332. Oed. dan. t. 57.*

*Erica folio abietis, flore arbuti. Buxb. cent. 4. p. 26.*  
t. 53.

*Habitat in Lapponiæ Alpibus. §*

*mariana.*

4. ANDROMEDA pedunculis aggregatis, corollis cylindricis, foliis alternis ovatis integerrimis.

*Andromeda foliis ovatis, pedunculis fasciculatis, capsulis pentagonis apice dehiscentibus. Gron. virg. 49.*

*Arbuscula mariana, brevioribus evonymi foliis pallide virentibus, floribus arbutis ex eodem nodo plurimis, spicatum uno versu erumpentibus. Pluk. mant.*

25. t. 448.

*Habitat in Virginia. §*

*Polifolia.*

5. ANDROMEDA pedunculis aggregatis, corollis ovatis, foliis alternis lanceolatis revolutis.

*Andromeda foliis alternis lanceolatis margine reflexis.*

*Fl. lapp. 163. t. 1. f. 3. Fl. suec. 335; 353. Hort. cliff. 163. Roy. lugdb. 440. Hall. helv. 416. opusc. 140.*

*Polifolia. Buxb. act. 2. p. 345. cent. 5. p. 28. t. 49. f. 1.*

*Vitis idææ affinis polifolia montana. Oed. dan. t. 54.*

*Erica humilis, rosmarini foliis, unedonis flore, capsula cistoide. Pluk. alm. 136. t. 175. f. 3.*

*Habitat in Europæ frigidioris paludibus turfosis. §*

*paniculata.*

6. ANDROMEDA racemis secundis nudis paniculatis, corollis subcylindricis, foliis alternis oblongis crenulatis.

*Andromeda foliis ovatis acutis crenulatis planis alternis, floribus racemosis. Hort. cliff. 162. Gron. virg. 48.*

*Frutex foliis ferratis, floribus longioribus spicatis subviridibus, spica pentagona. Catesb. car. 2. p. 43.*

t. 43.

*Vitis idææ americana, longiori mucronato & crenato folio, floribus urceolatis racemosis. Pluk. alm. 391.*

*Habitat in Virginia.*

*racemosa.*

7. ANDROMEDA racemis secundis bracteatis, corollis gibbo-cylindricis, foliis alternis oblongis ferratis.

*Habitat in Pennsylvania. Kalm.*

*Hæc differt a priori: 1. Racemis minus paniculatis. 2.*

*Bractea sub singulo flore lineari-lanceolata, stricla, flore longiore, viridi, decidua; in præcedenti nulla bractea. 3. Capsulis retinentibus stylum, quem antecedens dimittit. 4. foliis magis ferratis.*

8. AN-

8. **ANDROMEDA** racemis secundis nudis, corollis arborea. rutundo-ovatis.  
*Andromeda* arborea, foliis oblongo-ovatis integerri-  
 mis, floribus paniculatis nutantibus, racemis simpli-  
 cissimis. *Gron. virg.* 48.  
*Frutex* foliis oblongis acuminatis, floribus spicatis u-  
 no versu dispositis *Catesb. car.* 1. p. 71. t. 71.  
*Habitat in Virginia, Carolina.* ♀
9. **ANDROMEDA** racemis secundis foliaceis, corol- calyculata.  
 lis subcylindricis, foliis alternis lanceolatis obtusis  
 punctatis. *Fl. suec.* 2. n. 357. *Gort. ingr.* 64.  
*Cistus* Ledon f. *Andromeda*, floribus monopetalis par-  
 vis albis tubulosis spicatum in summis ramulis dis-  
 positus, foliis & facie vitis idææ, capsula minima  
 sicca quinquepartita. *Gron. virg.* 21?  
*Chamae Daphne.* *Buxb. act.* 1. p. 241. t. 8. f. 1.  
*Habitat in Virginia, Canada, Sibiria, Ingria.* ♀  
*Frutex foliis ovali-lanceolatis, coriaceis, myricæ confi-*  
*sentia, obtusis, margine recurvis, adpersis punctis*  
*vix conspicuis, callosis, ab utraque parte. Racemi*  
*ex ramis propriis, foliolis ovalibus alternis, in quo-*  
*rum ala flos solitarius, pedicellatus. Calyx a basi*  
*tegitur duobus foliolis ovatis. Flores ad unum latus*  
*nutant. Stamina 10, antheris oblongis.*

## EPIGÆA.

1. **EPIGÆA.** *Amæn. acad.* 3. p. 17. \* repens.  
*Memecylum.* *Mich. gen.* 13.  
*Arbutus* foliis ovatis integris, petiollis laxis longitudine  
 foliorum. *Gron. virg.* 49.  
*Pyrolæ* affinis repens fruticosa, foliis rigidis scabritie  
 exasperatis, flore pentapetaloide fistuloso. *Pluk. alm.*  
 309. t. 107. f. 1. *Raj. suppl.* 596.  
*Habitat in Virginie, Canadæ pinetis.* ♀

## GAULTHERIA.

1. **GAULTHERIA.** *Kalm. Amæn. acad.* 3. p. 14. \* procumbens.  
*Dukam arb.* 1. p. 286. t. 113.  
*Vitis* idæa canadensis, pyrolæ folio. *Tournef. inst.*  
 608.  
*Anonyma* pedunculis arcuatis. *Cold. noveb.* 98.  
*Habitat in Canadæ sterilibus arenosis.* ♀



## A R B U T U S.

Unedo,

1. **ARBUTUS** caule arboreo, foliis glabris ferratis, baccis polyspermis. *Hort. cliff.* 163. *Roy. lugdb.* 440. *Sauv. monsp.* 59. *Gron. orient.* 137.  
*Arbutus folio serrato. Bauh. pin.* 460. *Mill. dict. t.* 48. f. 1, 2.  
*Arbutus. Cam. epit.* 168.  
*Habitat in Europæ australis, Hiberniæ occidentalis, Orientis, sylvis.* ‡  
*Baccæ polyspermæ, tuberculis seminum exasperatæ.*

Andrachne.

2. **ARBUTUS** caule arboreo, foliis glabris integerrimis, baccis polyspermis.  
*Arbutus folio non serrato. Bauh. pin.* 460.  
*Andrachne theophrasti. Clus. hist. i. p.* 48.  
*Habitat in Oriente.* ‡  
*Cortice non scabro & Foliis integerrimis ab A. Unedine differt.*

acadiensis,

3. **ARBUTUS** caulibus procumbentibus, foliis ovatis subserratis, floribus sparsis, baccis polyspermis.  
*Vitis idæa acadiensis, foliis alaterni. Tournes. inst.* 608.  
*Habitat in Acadia.* ‡

alpina,

4. **ARBUTUS** caulibus procumbentibus, foliis rugosis ferratis. *Fl. lapp.* 161. *Fl. suec.* 340; 359. *Hort. cliff.* 163. *Roy. lugdb.* 440. *Hall. helv.* 415.  
*Vitis idæa foliis oblongis albicantibus. Bauh. pin.* 470.  
*Vitis idæa. Clus. hist. i. p.* 61.  
*Habitat in Alpibus Laponiæ, Helvetiæ, Sibiricæ, Angliæ.* ‡

Uva ursi.

5. **ARBUTUS** caulibus procumbentibus, foliis integerrimis. *Fl. lapp.* 162. t. 6. f. 3. *Fl. suec.* 339; 358. *Mat. med.* 211. *Hort. cliff.* 163. *Roy. lugdb.* 440. *Hall. helv.* 415.  
*Uva ursi. Clus. hist. i. p.* 63. *Oed. dan. t.* 33.  
*Radix idæa putata & Uva ursi. Bauh. hist. i. p.* 524.  
*Arbutus nqveboracensibus Bearberry. Cold. nov.* 104.  
*Habitat in Europa frigida, Canada.* ‡

## C L E T H R A .

alnifolia.

1. **CLETHRA.** *Gron. virg.* 47. *Duham. arb. i. p.* 176. t. 71.  
*Alnifolia americana serrata, floribus pentapetalis albis*  
in

in spicam dispositis, *Pluk. alm.* 18. t. 115. f. 1. *Catesb. car.* 1. p. 66. t. 66.

*Habitat in Carolina, Virginia, Pennsylvania. ‡*

## P Y R O L A.

1. PYROLA staminibus adscendentibus, pistillo decli- rotundifo-  
nato. *Fl. succ.* 330; 360. *Mat. med.* 212. *Amæn. acad.* 11a.  
1. p. 156.  
Pyrola staminibus & pistillis declinatis. *Fl. lapp.* 169.  
Pyrola foliis integerrimis. *Vir. cliff.* 37. *Roy, lugdb.*  
440.  
Pyrola foliis subrotundis, scapo racemoso. *Hort. cliff.*  
162.  
Pyrola rotundifolia major. *Bauh. pin.* 191.  
Pyrola noveboracensis. *Cold. noveb.* 99.  
*Habitat in Europa septentrionaliore, Virginia, Brasi-*  
*lia. 2*
2. PYROLA floribus racemosis dispersis, staminibus pi- minor.  
stillisque rectis. *It. gotl.* 206. *Fl. succ.* 331; 361.  
Pyrola foliis subrotundis, scapo undique racemoso,  
genitalibus rectis. *Aët. stockh.* 1741. p. 192.  
Pyrola foliis obiter serratis subrotundis racemosis, flo-  
ribus tuba erecta. *Hall. helv.* 420. *opusc.* 82.  
Pyrola minor. *Riv. pent.* 149. *Raj. angl.* 3. p. 363.  
*Habitat in Europa frigidior. 2*
3. PYROLA racemo unilaterali. *Fl. succ.* 332; 362. secunda.  
Pyrola floribus uno versu sparsis. *Fl. lapp.* 168.  
Pyrola folio mucronato ferrato. *Bauh. pin.* 181.  
Pyrola 2 tenerior. *Clus. pan.* 506.  
*Habitat in Europæ borealis sylvis. ‡*
4. PYROLA pedunculis subumbellatis. *Fl. succ.* 333; umbellata.  
363.  
Pyrola frutescens, arbuti flore. *Bauh. pin.* 191.  
Pyrola 3. fruticans. *Clus. pan.* 507.  
*Habitat in Europæ, Asiæ & Americæ septentrionalis*  
*sylvis. ‡*
5. PYROLA pedunculis bifloris. maculata;  
Pyrola petiolis apice bifloris vel trifloris. *Gron. virg.*  
48.  
Pyrola marilandica minor, folio macronato arbuti.  
*Pet. mxf.* 675.

*Pyrola mariana*, arbuti foliis angustioribus, trifoliata ad medium nervium linea alba utrinque per longitudinem discurrente. *Pluk. mant.* 157. t. 349. f. 4.  
*Habitat in Americæ septentrionalis sylvis.* §

uniflora.

6. *PYROLA* scapo unifloro. *Fl. lapp.* 167. *Fl. svec.* 334; 364. *Hall. helv.* 420.

*Pyrola rotundifolia minor.* *Bauh. pin.* 191.

Singulari flore ampliore. *Morif. hist.* 3. p. 505. f. 12. t. 10. f. 2.

*Pyrola* IV. *Oed. dan.* t. 8. 1

*Habitat in Europæ borealis sylvis.* 2

## DIGYNIA

### ROYENA.

lucida.

1. *ROYENA* foliis ovatis scabriusculis. *Hort. cliff.* 149. *Roy. lugdb.* 411.

*Staphylodendron africanum* sempervirens, foliis splendentibus. *Comm. hort.* 1. p. 187. t. 96.

*Staphylodendrum africanum*, folio singulari lucido. *Herm. parad.* 232. t. 232.

*Pistacia africana.* *Pluk. alm.* 298. t. 63. f. 4. t. 317. f. 5. *Habitat ad Cap. b. Spei.* §

glabra.

2. *ROYENA* foliis lanceolatis glabris. *Roy. prodr.* 441. *Royena* foliis lanceolatis. *Hort. cliff.* 149.

*Vitis idæa æthiopica*, buxi minoris folio, floribus albis. *Comm. hort.* 1. p. 125. t. 65.

*Vitis idæa æthiopica*, myrti foliis, flosculis dependentibus. *Pluk. alm.* 391. t. 321. f. 4.

*Habitat ad Cap. b. Spei.* §

hirsuta.

3. *ROYENA* foliis lanceolatis hirsutis. *Roy. lugdb.* 441. *Arbutus* foliis lanceolatis integerrimis hirsutis. *Hort. cliff.* 163.

*Staphylodendrum africanum*, folio lanuginoso rosmarini latiore. *Hoerh. lugdb.* 2. p. 235.

*Habitat ad Cap. b. Spei.* §

## HYDRANGÆA.

arborescens.

1. *HYDRANGÆA.* *Gron. virg.* 50. *Duham. arb.* 1. p. 298. t. 118.

Ha-

*Habitat in Virginia. ‡*

*Frutex ex ordine Arborescentium XIX, foliis oppositis, floribus in cymam digestis.*

CUNONIA.

1. CUNONIA.

*capensis.*

*Oosterdykia floribus spicatis pentapetalis, foliis oblongis subincanis serratis. Burm. afr. 259. t. 96.*

*Arbustula, arbuti alatis foliis, africana. Pluk. alm. 45. t. 191. f. 4.*

*Habitat ad Cap. b. Spei. ‡*

*Folia opposita, pinnata cum impari sessili, nec longiore. Foliola sæpe septem, lanceolata, glabra, serrata. Glandula maxima, spatulae forma, ovata, integerrima, compressa, foliacea, petiolo ancipiti, tum e singula foliorum ala, tum major inter pedunculos. Racemi terminales, bini, simplices: pedicelli ex singulis punctis sparsis plures, uniflori.*

CHRYSOSPLENIUM.

1. CHRYSOSPLENIUM foliis alternis. *Fl. suec. 317; alternifolium. 365. lt. scan. 16.*

*Chrysosplenium. Hort. cliff. 149. Roy. lugdb. 209.*

*Chrysosplenium foliis amplioribus auriculatis. Rudb. lapp. 97. Fl. lapp. 151.*

*Chrysosplenium foliis pediculis oblongis insidentibus. Tourn. f. hist. 146.*

*Saxifraga aurea, foliis pediculis oblongis insidentibus. Raj. hist. 206.*

*Sedum palustre luteum majus, foliis pediculis longis insidentibus. Moris. hist. 3. p. 477. f. 12. t. 8. f. 8.*

*Habitat in Sveciæ, Germaniæ, Angliæ opacis humen- tibus. 24*

*Flos primarius Decandrus, reliqui Octandri. Phil. Bot. 178.*

2. CHRYSOSPLENIUM foliis oppositis. *Sauv. mon. p. oppositifolium. 128.*

*Chrysosplenium foliis amplioribus auriculatis. Tournef. infl. 146.*

*Chrysosplenium l. Saxifraga aurea. Tabern. hist. 1224.*

*Saxifraga aurea. Dod. pempt. 316. Lob. hist. 336. ic. 312.*

*Saxifraga rotundifolia aurea. Bauh. pin. 309.*

*Saxifraga romanorum. Dalech. hist. 1114.*

*Sedum palustre luteum, foliis subrotundis sessilibus.*

*Morif. hist. 3. p. 477. f. 12. t. 8. f. 7.*

*Alchimilla rotundifolia aurea hirsuta. Herm. lugdb. 14.*

*Habitat in Belgio, Anglia, Canada, locis umbrosis humentibus.*

*Affinis primæ præcedenti.*

## SAXIFRAGA.

\* *Foliis indivisis, Caule subnudo.*

**Cotyledon.** 1. **SAXIFRAGA** foliis radicatis aggregatis lingulatis cartilagineo-ferratis, caule paniculato.

*Saxifraga foliis ligulatis radicalibus margine cartilagineo acute ferratis, floribus paniculatis. Fl. suec. 356; 366.*

*Saxifraga foliorum marginibus cartilagine crenatis. Hort. cliff. 168. Roy. lugdb. 453.*

*Saxifraga foliis radicalibus in orbem positis: ferraturis cartilagineis Fl. lapp. 177. t. 2. f. 2.*

*Sanicula montana crenata, folio longiore, pediculo folioso. Pluk. alm. 331. t. 222. f. 1.*

*Sedum ferratum album bicornè, marginibus argenteis. Morif. hist. 3. p. 478. f. 12. t. 9. f. 19. 20.*

*Cotyledon media, foliis oblongis, ferratis. Bauh. pin. 285.*

β. *Cotyledon minor, foliis subrotundis ferratis. Bauh. pin. 285. prodr. 133.*

γ. *Saxifraga, sedi folio, flore albo, multiflora. Tournef. inst. 252.*

*Sedum ferratum, flore albo, multiflorum. Dod. pempt. 113.*

δ. *Saxifraga, sedi folio, pyrenaica ferrata. Tournef. inst. 252.*

ε. *Cotyledon pyramidale, lato crenato & retuso folio, polyanthos. Bocc. mus. 2. p. 109. t. 86.*

*Saxifraga sedi folio angustiore ferrato. Segu. ver. 448. t. 9. f. 1*

*Habitat in Alpibus Europæ. 24*

mutata.

**SAXIFRAGA** foliis radicalibus aggregatis lingulatis cartilagineo-ferratis, caule racemoso.

*Saxifraga foliorum limbo cartilagineo integro, spica longa, floribus purpureo croceis. Hall. helv. 402. n.*

*9. t. 2. Seguier. ver. 3. p. 199. Burf. XVI. 98.*

*112.*

*Habitat in Alpibus Helvetiæ, Italiæ.*

*Hybrida quasi apparet hæc planta S. Cotyledoni affinis, sed distincta. Folia Radicalia aggregata, lingulata, cartilagine serrata, versus basin ciliato-villosa. Caulis firmior, simplex foliis copiosioribus, magis viscidus. Racemi alterni, breves. Flores quadruplo majores: Petalis lanceolatis, longis, diversissimis, croceis nec albis.*

3. **SAXIFRAGA** foliis lanceolatis denticulatis, caule nudo paniculato, floribus subcapitatis. pennsylvanica.

*Saxifraga foliis radicalibus lanceolatis denticulatis, caule subnudo piloso ramoso, floribus confertis capitatis. Gron. virg. 49.*

*Saxifraga noveboracensis. Cold. noveb. 105.*

*Saxifraga pennsylvanica, floribus muscosis recemosis.*

*Dill. elth. 337. t. 253. f. 328.*

*Sanicula virginiana alba, folio oblongo mucronato.*

*Pluk. alm. 331. t. 59. f. 1. & t. 222. f. 5.*

*Habitat in Virginia, Pennsylvania, Canada.*

4. **SAXIFRAGA** foliis lanceolatis obtusis pilosis, caule nudo bifloro. androsacea.

*Saxifraga caule paucifloro, foliis subhirsutis lingulatis & tridentatis. Hall. helv. 405.*

*Sedum alpinum tertium. Col. ecphr. 2. p. 66. t. 67. f. ultim.*

*Saxifraga alpina, habitu androsaces, villosa. Hall. norib. 1736. t. 1. f. 3.*

*Habitat in Sibiria, Helvetia.*

5. **SAXIFRAGA** foliis linearibus subtus punctatis aggregatis recurvatis, caule nudo, multifloro. caesia.

*Saxifraga rosulis densissimis, foliis crassis glaucis repandis subtus sulcatis. Hall. helv. 402.*

*Saxifraga alpina minima, foliis caësis deorsum incurvis. Tournef. inst. 253. Segu. veron. 449. t. 9. f. 2.*

*Sedum alpinum album, foliis compactis. Bauh. pin. 284. Moris. hist. 3. f. 12. t. 7. f. 32.*

*Sedum alpinum minimum, foliis cinereis, flore candido. Scheuch. alp. 341; 141. t. 21. f. 1.*

*Habitat in Alpibus Helveticis, Austriacis, Pyrenaicis, Baldo.*

*Folia pagina superioris duplici ordine punctorum perforatorum evadunt albida.*

- burseriana.* 6. **SAXIFRAGA** foliis aggregatis imbricatis subulatis lævibus, caule subnudo unifloro.  
*Saxifraga alpina*, foliis glaucis acutis, monanthos, caule folioso. *Segu. ver.* 3. p. 201. t. 5. f. 2.  
*Sedum alpinum*, *saxifragæ* albæ flore. *Bauh. pin.* 284. *Burf. XVI:* 6.  
*Habitat in Taurero Raftadienfi, montibus Lessanenſibus.* 24.  
*Facies Sedi minoris. Folia juniperina, dense imbricata, acuta, carinata, glabra. Caulis vix digitalis, rufescente hispidus f. glutinosus, foliolis 3 f. 4 alternis, linearibus. Flos magnitudine Saxifragæ granulatæ: venis incarnatis.*
- sedoides.* 7. **SAXIFRAGA** foliis caulinis aggregatis alternis oppositisque sublanœolatis, flore pedunculato.  
*Saxifraga alpina minima*, foliis ligulatis in orbem actis, flore ochroleuco. *Segu. veron.* 450. t. 9. f. 4.  
*Habitat in Alpibus Tridentinis, Sibiricis.* 24
- bryoides.* 8. **SAXIFRAGA** foliis ciliatis inflexis imbricatis, caule nudiusculo unifloro.  
*Saxifraga rosulis densissimis*, foliolis ciliatis, caule unifloro. *Hall. helv.* 403.  
*Saxifraga pyrenaica minima lutea musco similis. Tournes. inst.* 253.  
*Sedum muscosum. Bauh. hist.* 3. p. 695. *Scheuch. alp.* 142. t. 21. f. 2.  
*Sedum ejusdem alpinum quartum. Col. ecphr.* 2. p. 66. t. 67. f. 1, 2.  
*Habitat in Alpibus Helveticis, pyrenaicis. Folia spinoso-ciliata, licet cilia sæpius obtegantur foliis imbricatis.*
- bronchialis.* 9. **SAXIFRAGA** foliis imbricatis subulatis ciliatis spinosis, caule subnudo multifloro.  
*Habitat in Sibiria.*  
*Stolones oblongi confertim imbricati: Foliis planis, subulatis, mucronato-spinosis, spinoso-ciliatis. Caulis altus, nudiusculus: foliis aliquot minimis mucronatis margine lævibus. Panicula parva.*
- stellaris.* 10. **SAXIFRAGA** foliis ferratis, caule nudo ramoso, petalis acuminatis.

*Saxifraga foliis lanceolatis dentato-ferratis, caule nudo simplici. Fl. fvec. 335; 367.*

*Saxifraga caule nudo simplici, foliis lanceolatis dentatis, petalis acutis. Fl. Lapp. 175.*

*Sanicula myosotis, floribus albicantibus fere umbellatis. Pluk. alm. 331. t. 58. f. 2. & t. 222. f. 4.*

*Sedum montanum hirsutum, mucronato & dentato folio, flore albo guttato. Morif. hist. 3. p. 478. f. 12. 1. 9. f. 13.*

*Sedum alpinum IV. Oed. dan. t. 23.*

*Habitat in Alpibus Spitsbergenfibus, Lapponicis, Helveticis, Styriacis, Westmorlandicis.*

*Flores inferi.*

11. **SAXIFRAGA** foliis ovalibus retusis obsolete serratis petiolatis, caule nudo, panícula conglomerata: *crassifolia.*  
*Saxifraga foliis ovalibus crenulatis, caulibus nudis. Gmel.*

*Habitat in Alpibus Sibiriae. 24.*

*Radix crassitie fere digiti. Folia bina, palmaria, coriacea, ovalia, retusa, repando-crenata, glabra, foliatione involuta. Gemmae maximae foliis 2 f. 3, ovatis, concavis, quorum unum scapo adherens attollitur. Scapus nudus, pedalis, crassitie digiti, subpaniculatus ramis pedatis. Flores superi, submutantes, campanulati: petala obovata, rubro-purpurea.*

12. **SAXIFRAGA** foliis obovatis crenatis subsessilibus, nivalis, caule nudo, floribus congestis.

*Saxifraga foliis subovatis crenatis, caule nudo, floribus capitatis. Fl. fvec. 354; 368.*

*Saxifraga caule nudo simplici, foliis elliptico-subrotundis crenatis, floribus capitatis. Fl. Lapp. 176. t. 2. f. 5. 6.*

*Saxifraga foliis cordato-ovalibus crenatis, corolla alba, caule hirsuto aphylo. Gron. virg. 160.*

*Saxifraga foliis oblongo-rotundis dentatis, floribus compactis. Raj. angl. 3. p. 354. t. 16. f. 1.*

*Sempervivum minus dentatum. Mart. Spitzb. 43. t. F. f. A. Pluk. phyt. 222. f. 2.*

*Sedum III. Oed. dan. t. 28.*

*Habitat in summis Alpibus Spitzbergenfibus, Lapponicis, Arvonicis, Virginia, Canada. 24*



- punctata.** 13. *SAXIFRAGA* foliis subrotundis dentatis longius petiolatis, caule nudo.  
*Sedum bicornne ferratum*, pallidiore folio rotundiore, floribus punctatis. *Morif. hist.* 3. p. 478. f. 12. t. 9. f. 17.  
*Habitat in Sibiria.* 24  
*Folia subrotunda, dentibus incisa, ut fere lobata absque ora cartilaginea, minime retusa, dentibus acutis; Petioli laeves foliis triplo longiores, superne sensim dilatati in folium. Flores minores quam in S. Geo, fere inferi.*
- umbrosa.** 14. *SAXIFRAGA* foliis obovatis subretusis cartilagineo-crenatis, caule nudo paniculato.  
*Saxifraga* foliis cuneiformibus retusis radicalibus sinuato-ferratis margine acutis, caule, paniculato. *Hort. ups.* 108.  
*Geum folio subrotundo minori, pistillo floris rubro. Magn. hort.* 88. t. 8. *Mill. ic.* 141. f. 2.  
*Habitat in Montibus apud Cantabrös.* 24  
*Affinis nimium S. Geo; Folia basi minime cordata, crenis angulatis; petiolis ciliato-villosis. Flores inferi.*
- cuneifolia.** 15. *SAXIFRAGA* foliis cuneiformibus obtusissimis repandis, caule nudo paniculato.  
*Cotyledon altera olim matthioli. Bauh. hist.* 3. p. 684.  
*Cotyledon aut Sedi species quædam. Gesn. fasc.* 19. t. 12. f. 37. bene.  
*Habitat in Alpibus Strizæ, Gotthardo. Gérard;*  
*Simillima præcedenti, sed Foliorum margo non crenatus, at obtusissime repandus, ut 3 folia longe minor. Flores inferi.*
- hirsuta.** 16. *SAXIFRAGA* foliis cordato-ovalibus retusis cartilagineo-crenatis, caule nudo paniculato.  
*Geum folio circinato acute crenato, pistillo floris rubro. Magn. hort.* 87. t. 87.  
*Habitat in Pyrenæis.* 24  
*Affinis nimium sequenti, sed folia ovalia, nec reniformia. Flores inferi.*
- Geum.** 17. *SAXIFRAGA* foliis reniformibus dentatis, caule nudo paniculato.  
*Sanicula montana rotundifolia minor. Bauh. pin* 243.

*Sedum montanum rotundifolium minus album non guttatum. Moris. hist. 3. p. 478. f. 12. t. 9 f. 12.*

*Habitat in Alpibus Europæ. 24*

*Flores inferi.*

\* \* Foliis indivisis, Caule folioso.

18. **SAXIFRAGA** foliis caulinis ovatis oppositis imbricatis: summis ciliatis. *Fl. suec. 359; 369.* oppositifolia.

*Saxifraga, foliis ovatis quadrangulo imbricatis, rariis procumbentibus. Fl. lapp. 179. t. 2. f. 1. Hall. helv. 403.*

*Sedum alpinum ericoides purpurascens. Bauh. pin. 284. prodr. 132. Moris. hist. 3. p. 480. f. 12. t. 10. f. 36.*

*Habitat in rupibus Alpium Spitzbergenium, Laponicarum, Pyrenaicarum, Helveticarum. 24*

*Caules filiformes, dependentes, repentes. Folia ovata, quadrifariam imbricata opposita, sessilia, ciliata. Flos terminalis, solitarius, sessilis.*

19. **SAXIFRAGA** foliis caulinis lanceolatis alternis aspera. spinoso-ciliatis, caule inferne procumbente.

*Saxifraga foliis simplicibus ciliatis, caule folioso ramoso. Hall. helv. 403.*

*Saxifraga sedi foliis crenatis asperis. T. Scheuch. it. 2. p. 140. t. 20. f. 3.*

*Sedum alpinum, foliis crenatis asperis. Bauh. pin. 284. prodr. 132. Gesn. fasc. 22. t. 6. f. 27.*

*Sedum alpinum hispidum fere spinosum, flore pallido. Bauh. hist. 3. p. 695. Moris. hist. 3. p. 479. f. 12. t. 10. f. 25.*

*Sedum minimum alpinum villosum alterum. Park. theatr. 738.*

*Habitat in Alpibus Helveticis.*

20. **SAXIFRAGA** foliis caulinis linearibus alternis ciliatis: radicalibus aggregatis. autumnalis.

*Saxifraga foliorum margine ciliari, floribus luteis maculosis. Hall. helv. 399.*

*Saxifraga angustifolia autumnalis, flore luteo guttato, foliis florum magis acuminatis. Breyn. cent. 106. t. 48.*

*Sedum alpinum, floribus luteis maculosis. Bauh. pin. 284.*

*Sedum angustifolium autumnale, flore luteo guttato. Moris. hist. 3. p. 477. f. 12. t. 8. f. 6.*

Ha

*Habitat in humentibus Borussiae, Angliae, Helvetiae.*  
*Calyx corolla longe brevior & viridis est. Flores inferi.*

- Hirculus.** 21. **SAXIFRAGA** foliis caulinis lanceolatis alternis nudis inermibus, caule erecto. *Fl. suec. 2. n. 370.*  
*Saxifraga petalis latissimis luteis lineatis. Hall. herb. 399. t. 8.*  
*Geum palustre luteum bicorne, nardi celticae similis. Moris. hist. 3. p. 477.*  
*Chamaecistus fruticulosus, foliis nardi celticae. Bauh. pin. 466.*  
*Hirculus qui Chamaecisti genus. Clus. cur. 5.*  
*Habitat in Svecia, Helvetia, Lapponia, Sibiria.*  
*Petala ad basin Lacuna nectarifera ex rima bivahri apicibus acutis instructa. Flores inferi.*
- ovoides.** 22. **SAXIFRAGA** foliis caulinis lineari-subulatis sparsis nudis inermibus, caulibus decumbentibus.  
*Saxifraga foliis linearibus sparsis glabris. Fl. suec. 357; 371.*  
*Saxifraga foliis subulatis sparsis. Fl. lapp. 178.*  
*Sedum alpinum, flore pallido. Bauh. pin. 284. Moris. hist. 3. p. 477. f. 12. t. 6. f. 3.*  
*Sedum minus VI. Clus. hist. 2. p. 60.*  
*Habitat in Alpibus Lapponicis, Styriacis, Westmoriandicis, Baldo.*
- rotundifolia.** 23. **SAXIFRAGA** foliis caulinis reniformibus dentatis petiolatis, caule paniculato.  
*Saxifraga foliis reniformibus acute crenatis, caule ramoso folioso. Hort. cliff. 167. Roy. lugdb. 453. Hall. herb. 400.*  
*Sanicula montana rotundifolia major. Bauh. pin. 243.*  
*Sanicula alpina. Cam. epit. 764. Gesn. fasc. 19. t. 10. f. 25.*  
*Habitat in Alpibus Helveticis, Austriacis. 24*  
*Flores inferi.*

\* \* \* Foliis lobatis, Caule erecto.

- granulata.** 24. **SAXIFRAGA** foliis caulinis reniformibus lobatis, caule ramoso, radice granulata. *Hort. cliff. 167. Fl. suec. 350; 372. Niat. med. 215. Roy. lugdb. 453. Hall. herb. 401.*

Saxi-

*Saxifraga rotundifolia alba. Bauh. pin. 309.*

*Saxifraga alba. Dod. pempt. 316.*

*Saxifraga IV. Cam. epit. 719.*

*Habitat in Europæ apricis. 24*

25. **SAXIFRAGA** foliis caulinis palmato-lobatis: cau- bulbifera.  
linis sessilibus, caule ramoso bulbifero.

*Saxifraga bulbosa altera bulbifera montana. Col. ecphr.*

1. p. 318. t. 317.

*Saxifraga ad folia bulbos gerens. Bauh. pin. 309.*

*Sedum bicornne rotundifolium erectum, radice granu-  
lata, & ad caulem tubercula proferens. Moris. hist.*

3. p. 474. f. 12. t. 9. f. 24.

*Habitat in Italiæ pratis saxosis umbrosis. 24*

*Varietas potius præcedentis, quam species. Segu. ver.*

3. p. 207.

26. **SAXIFRAGA** foliis caulinis palmatis petiolatis, cornu.  
caule simplicissimo unifloro bulbifero. *Oed. dan. t.*  
*22.*

*Saxifraga foliis palma is, caule simplici unifloro. Fl.*

*lapp. 172. t. 2. f. 4. Hort. cliff. 167. Fl. suec. 351. 373.*

*Habitat in Alpibus Lapponicis frequens. 24*

*Flos parum nutat. Folia caulina glabra, magis pal-  
mata & germen supra receptaculum, magis quam in  
antecedente, ut distinctissima a priore planta. Vi-  
detur tamen affinis sequenti, licet & ab hac di-  
stincta.*

27. **SAXIFRAGA** foliis reniformibus palmatis pilosis, sibirica.  
caule pedunculisque filiformibus.

*Habitat in Sibiria.*

*Radiculæ capillares. Folia radicalia petiolata, reni-  
formia, septemloba f. palmata: lobis ovatis, pilis ra-  
ris adpersa, petiolis hirsutis. Caulis filiformis, ad-  
scendens, sempipedalis, debilis, Foliis 2. f. 3. sessilibus,  
palmatis. Pedunculi terminales & laterales longif-  
simi, filiformes, bifidi, nudi. Flos majusculus, al-  
bus, inferus.*

28. **SAXIFRAGA** foliis caulinis palmatis: summo flo- rivularis.  
rali ovato, caule simplici subbifloro.

*Saxifraga foliis radicalibus quinquelobis, florali ovato.*

*Fl. lapp. 174. t. 2. f. 7. Fl. suec. 352, 374.*

*Habitat ad Alpium Lapponicarum latera & rivulos.*

*Linnei Spec. plant. pars 1.*

Oo

29. SA-

- groenlandica. 29. *SAXIFRAGA* foliis caulinis palmato-multifidis sparsis: laciniis acutis, caule erecto.  
*Saxifraga tridactylites groenlandica*, caulibus valde foliosis. *Dill. elih.* 337. t. 353. f. 329.  
*Habitat in Groenlandia., forte etiam in Pyrenæis & Helveticis Alpibus.*
- geranioides. 30. *SAXIFRAGA* foliis radicalibus reniformibus quinquelobis multifidis, caulinis linearibus, caule subnudo ramoso. *Amæn. acad.* 4. p. 271. \*  
*Habitat in Pyrenæis. Burf. & VI. 86.*
- ajugifolia. 31. *SAXIFRAGA* foliis radicalibus palmato quinquepartitis, caulinis linearibus indivisis, caulibus adscendentibus multifloris. *Amæn. acad.* 4. p. 271. \*  
*Habitat in Galloprovinciæ montibus.*
- tridactylites 32. *SAXIFRAGA* foliis caulinis cuneiformibus trifidis alternis, caule erecto ramoso. *Fl. suec.* 353; 375.  
*Saxifraga foliis trifidis basi angustis, caule erecto. Roy. lugdb.* 457. *Sauv. monsp.* 208.  
*Saxifraga foliis omnibus trilobis basi angustis, caule erecto. Fl. lapp.* 173. *Hort. cliff.* 168.  
*Sedum tridactylites tectorum. Bauh. pin.* 285.  
*Paronychia altera. Dod. pempt.* 113.  
*Habitat in Europæ arenosis, & Alpibus. ☉*
- petræa. 33. *SAXIFRAGA* foliis caulinis palmato-tripartitis: laciniis subtrifidis, caule ramosissimo laxo.  
*Saxifraga alba petræa. Pon. bald.* 339. f. 337. \*  
*Saxifraga aizoides alpina, trifido folio, major alba. Pluk. alm.* 331. t. 222. f. 3.  
*Sedum tridactylites alpinum majus album. Bauh. pin.* 284. *Morif. hist.* 3. p. 479. f. 12. t. 9. f. 28. \*  
*Habitat in Alpibus Norvegicis, Pyrenæis, Baldi.*  
*Affinis nimium S. tridactyliti, etiam viscida, sed Caulis ramosissimus. Folia lata, palmata, tripartita, incisa. Corollæ quadruplo-majores.*
- caespitosa. 34. *SAXIFRAGA* foliis radicalibus aggregatis linearibus integris, trifidisque, caule erecto subnudo subbifloro. *Fl. suec.* 2. n. 376. \*  
*Saxifraga foliis simplicibus & trifidis, caule pæncaphyllo paucifloro. Hall. helv.* 405. *opus.* 291. t. 1. f. 1.

*Saxifraga* foliis partim integris partim trifidis. *Segu.*  
ver. 451. t. 9. f. 4.

*Sedum tridactylites* alpinum minus. *Bauh. pin.* 284.  
prodr. 131.

*Sedum Moschatella* alpina lutea. *Gesn. fasc.* 25. t. 6.  
f. 31.

*Habitat in Alpibus Lapponicis, Helveticis, Tridenti-*  
*nis, Montpelii.* 2.

\* \* \* Foliis lobatis ; Caulibus procumbentibus.

35. *SAXIFRAGA* foliis caulinis cordatis trilobis inter- cymbalaria.  
grisque, caulibus procumbentibus.

*Saxifraga* exigua, foliis cymbalariae. *Buxb. cent.* 2. p.  
40. t. 45. f. 2.

*Habitat in Oriente.*

36. *SAXIFRAGA* foliis caulinis ovatis lobatis, caule hederacea.  
filiformi flaccido.

*Saxifraga* cretica annua minima, hederaceo folio.  
*Tournef. cor.* 18.

*Habitat in Creta.*

37. *SAXIFRAGA* foliis caulinis linearibus integris tri- *Hypnoides.*  
fidisve, stolonibus procumbentibus, caule erecto  
nudiusculo.

*Saxifraga* procumbens, foliis linearibus integris trifidis-  
que. *Hort. cliff.* 168. *Roy. lugdb.* 453. *Sauv. monsp.*  
208. *Gort. gelr.* 248.

*Saxifraga* foliis omnibus angustis petiolatis apice triden-  
tatis, caule folioso ramoso viscido. *Hall. helv.* 404.

*Sedum* alpinum, trifido folio. *Bauh. pin.* 284. *Morif.*  
*hist.* 3. p. 479. f. 12. t. 9. f. 26.

38. *Saxifraga* procumbens, foliis linearibus integris trifidis  
& quinquefidis. *Roy. lugdb.* 454. *Sauv. monsp.* 208.

*Habitat in Alpibus Helveticis, Austriacis, Pyrenaicis,*  
*Westmorlandicis.* 2.

38. *SAXIFRAGA* foliis caulinis cuneiformibus apice adscendens.  
dentatis, caule adscendente subviloso.

*Sedum tridactylites* alpinum, caule folioso. *Bauh. pin.*  
284.

*Sedum tridactylites* alpinum majus. 3. *Bauh. prodr.*  
131. *Burj. XVI.* 89.

*Habitat in Pyrenæis, Baldo, Taurero Rastadiensi.* 2.

## TIARELLA.

- cordifolia. 1. **TIARELLA** foliis cordatis. *Amæn. acad.* 3. p. 17.  
*Mitella* scapo nudo. *Hort. cliff.* 167. \* *Roy. lugdb.*  
 459. *Gron. virg.* 160.  
*Cortusa americana*, flore spicato, petalis integris. *Herm.*  
*par.* 129. t. 129.  
*Habitat in America & Asia septentrionali.* 24
- trifoliata. 2. **TIARELLA** foliis ternatis. *Amæn. acad.* 3. p. 17.  
*Mitella* foliis ternatis. *Amæn. acad.* 2. p. 351.  
*Habitat in Asia boreali.* *G. Demidoff.* 24

## MITELLA.

- diphylla. 1. **MITELLA** scapo diphyлло. *Hort. cliff.* 167. \* *Roy.*  
*lugdb.* 459. *Cold. noveb.* 96.  
*Cortusa americana altera*, floribus minutim fimbriatis.  
*Mentz. pug.* t. 10.  
*Cortusa americana*, spicato flore, petalis fimbriatis.  
*Herm. par.* 130.  
*Sanicula* f. *Cortusa indica*, flore spicato fimbriato.  
*Dodart. mem.* 209.  
*Habitat in America septentrionali.*
- nuda. 2. **MITELLA** scapo nudo. *Amæn. acad.* 2. p. 352.  
*Habitat in Asia boreali.* 24

## SCLERANTHUS.

- annuus. 1. **SCLERANTHUS** calycibus fructus patulis. *Fl.*  
*svæc.* 348; 377.  
*Scleranthus.* *Hort. cliff.* 166. *Roy. lugdb.* 215.  
*Polygonum*, gramineo folio, majus erectum. *Bauh.*  
*pin.* 281.  
*Polygonum*, angustissimo & gramineo folio, minus  
 repens. *Bauh. pin.* 281. (*sed mixtum*)  
*Polygonum minus alterum.* *Tabern. hist.* 1217.  
*Habitat in Europæ arvis arenosis.* ☉
- perennis. 2. **SCLERANTHUS** calycibus fructus clausis. *Fl. svæc.*  
 349; 378.  
*Knavel incanum*, flore majore, perenne. *Raj. angl.* 3.  
*p.* 160. t. 5. f. 1.  
*Polygonum minus polycarpon.* *Tabern. hist.* 1215.  
 Alchi-

*Alchimilla* gramineo folio, majori flore. *Vaill. parif.*

4. t. 1. f. 5.

*Habitat in Europæ campis apricis arenosis.* 24

3. **SCLERANTHUS** calycibus fructus patentissimis polycarpes.

spinosis, caule subvillosa. *Amæn. acad.* 4. p. 313.

*Polycarpus. Dalch. hist.* 444.

*Polygonum montanum. vermiculata foliis. Bauh. pin.*  
281.

*Vermiculata nova planta. Col. ecphr.* 1. p. 295. 2. 294.

*Habitat Montpellii & in Italia.* ☉

## GYPSOPHILA.

1. **GYPSOPHILA** foliis mucronatis recurvatis, flo- aggregatis.  
ribus aggregatis. *Amæn. acad.* 3. p. 23.

*Saponaria* calycibus pentaphyllis, floribus aggregatis,  
foliis mucronatis canaliculatis recurvis. *Hort. upf.*  
107.

*Caryophyllus saxatilis, ericæ foliis, umbellatis corym-*  
*bis. Bauh. pin.* 211. *prodr.* 105. *Raj. hist.* 1033.

*Magn. monsp.* 53. t. 5.

*Habitat in monte Lupi Galliæ, Montpellii, Pyre-*  
*næis.* 24

2. **GYPSOPHILA** foliis lanceolatis, petalis emarginatis, repens.  
staminibus pistillo brevioribus. *Amæn. acad.* 3. p. 23.

*Saponaria* caule simplici, foliis subulatis planis, ex  
alis ramulosa. *Hort. cliff.* 166.

*Saponaria* foliis glaucis pulposis angustis heteromallis.  
*Hall helv.* 380. n. 6. *Rupp. j. n.* 117.

*Alfina* orientalis altissima, gramineo folio, flore albo.  
*Buxb. cent.* 3 p. 32. t. 60.

*Alfina* angustifolia caryophylloides multiflora glabra  
purpurascens, radice atragaliti. *Pluk. alm.* 22. t. 75.  
f. 2.

*Caryophyllus saxatilis, foliis gramineis, minor. Bauh.*  
*pin.* 211. *Burf. XI.* 126.

*Habitat in Sibiria, Austria, Helvetia, Pyrenæorum*  
*montibus.* 24

*Corollæ albæ, extus incarnatæ, emarginatæ; antheræ*  
*rufescentes polline violaceo.*

3. **GYPSOPHILA** foliis lanceolatis lævibus, caulibus prostratis.  
diffusis, pistillis corolla campanulata longioribus.

O O 3

Ha-



*Habitat in Alpibus Europæis? 24*

Caulis plures, diffusi, læves, teretes, subpedales, ad genicula rufescentes. Folia lanceolata, glabra. Panicula brachiata, trichotoma: pedunculo intermedio simpliciore ad triangulum recto. Corolla alba: Petalis obtusis, patentibus, apice canaliculatis. Stamina corolla dimidio breviora. Antheræ flavæ. Scylli corolla paullo longiores. Simillima *G. reptans*, sed differt Radice vix repente; Caulis geniculis minus purpurascens. Calycis foliolis non carinatis. Corollis minoribus albis, nec subrubentibus. Antheris flavis. Stylis divergentibus; petalo longioribus. Foliis minus carnosiss.

fastigiata. 4. *GYPHOPHILA* foliis lanceolato-linearibus obsolete triquetris lævibus obtusis secundis. *Amæn. acad.* 3. p. 23.

Saponaria calycibus pentaphyllis, corymbis fastigiatis, foliis linearibus, caule adscendente. *L. gotl.* 282. *Fl. suec.* 346.; 379. *Sauv. monsp.* 145.

Saponaria petalis ovatis, foliis glaucis pulposiss linearibus. *Hall. jen.* 117. t. 2. f. 1.

Caryophyllus saxatilis, floribus graminæ, umbellatis corymbis. *Bauh. pin.* 211.

Polygonum majus erectum angustifolium, floribus candidis. *Mentz. pug. t. 2. f. 2.*

*Habitat in Gotlandiæ, Borussia, Helvetiæ rupibus. 24*

Caulis in hac, non in antecedente, ante florescentiam procumbit, hinc folia secunda.

altissima. 3. *GYPHOPHILA* foliis lanceolatis subtrinerviis rectis.

Saponaria calycibus pentaphyllis, corymbis fastigiatis, foliis lanceolatis, caule adscendente. *Hort. upl.* 107.

*Habitat in Sibiria. 24*

Differt a sequente magnitudine dupla vel quadrupla omnium partium; Structura eadem, tamen diversa species. Caulis erectus, sesquipedalis. Folia longitudine 3 latitudine digiti.

Struthium. 6. *GYPHOPHILA* foliis linearibus; axillis confertis teretibus. *J. est. it.* 73.

Saponaria caule simplici, foliis linearibus ex alis foliorum confertis teretibus. *Hort. cliff.* 166. *Rey. jugdh.* 444.

*Saponaria lychnidis folio*, flosculis albis. *Bauh. pin.* 206.

*Kali vermiculatum*, albo globoso flore. *Barr. rar.* 64. t. 119. *Bocc. mus.* 2 p. t. 122.

*Habitat in Hispania.* 24

7. *GYPSOPHILA* foliis lanceolatis subtrinerviis tomentosa, mentosis, caule pubescente. *Amæn. acad.* 4. p. 271.\*

*Linum sylvestre latifolium*, flore albicante. *Barr. rar.* 670. t. 1002.

*Habitat in Hispania.* 24

8. *GYPSOPHILA* foliis lanceolatis scabris, floribus paniculatis, dioicis, corollis revolutis. *Amæn. acad.* 3. p. 23.

*Alsine frutescens*, caryophylli folio, flore parvo albo. *Gerb. tanaïf.* 15.

*Habitat in Sibiria, Tataria desertis sabulosis, Tawrow, Bielogrod.* 24

*Folia margine scabra, acuta. Stamina in fundo floris sterilia latent. Petala revoluta, minora quam in congeneribus. Styli longiores corolla, incurvi. Mas mihi non visus.*

9. *GYPSOPHILA* foliis ovato-lanceolatis semiamplexi- perfoliatis, caulibus.

*Saponaria foliis lanceolatis, calycibus campanulatis angulatis. Hort. cliff.* 165.

*Spergula multiflora*, foliis inferioribus saponariae, superioribus behen similibus. *Dill. elth.* 368. t. 276. f. 357.

*Habitat in Hispania & Oriente.*

10. *GYPSOPHILA* foliis lineari-lanceolatis planis, caule dichotomo, pedunculis bifloris, petalis emarginatis. *Amæn. acad.* 3. p. 24.

*Saponaria caule dichotomo, foliis subulatis planis. Hort. cliff.* 445. *Roy. lugdb.* 445.

*Saponaria caule brachiato, capillaceo folio, calyce campanulato, floribus sparsis. Sauv. monsp.* 145.

*Tunica minima. Dalech. hist.* 1191.

*Habitat Monspelii & forte in Belgio.* 24

11. *GYPSOPHILA* foliis linearibus planis, calycibus muralis, aphyllis, caule dichotomo, petalis crenatis. *Amæn. acad.* 3. p. 23.

- Saponaria calycibus pentaphyllis, corollis crenato-emarginatis, foliis subulatis planis. Fl. suec. 347; 382.*  
*Saponaria foliis linearibus. Fl. lapp. 171.*  
*Caryophyllus minimus muralis. Bauh. pin. 211.*  
*Lychnis parva palustris, foliis acutis lanceolatis, flosculis purpureis. Menz. pug. t. 7. f. 4.*  
*Habitat in Svecia, Germania, Helvetia ad vias. ☉*  
*Similis G. saxifragæ, at caret squamis calycinis.*  
*saxifraga. 12. GYPSOPHILA foliis linearibus, calycibus angulatis: squamis duabus. corollis crenatis.*  
*Dianthus saxifragus. Sp. pl. 1. p. 413.*  
*Dianthus radice repente, ramis decumbentibus, foliis subulatis. Roy. lugdb. 444.*  
*Dianthus calyce lato & brevissimo. Hall. helv. 381.*  
*Caryophyllus saxifragus strigosior f. Caryophyllus sylvestris, flore minimo. Bauh. pin. 211.*  
*Lychnis pumila caryophyllata, flore rubello, italica Hæmorrhoidalis aldrovando. Barr. rar. 64. t. 998.*  
*Habitat in Austria, Helvetia, Gallia. 4*

## SAPONARIA.

- officinalis. 1. SAPONARIA calycibus cylindricis, foliis ovato-lanceolatis. Hort. cliff. 165. Hort. upf. 106. Mat. med. 214. Roy. lugdb. 414. Gron. virg. 160. Gort. getr. 245.*  
*Saponaria major lævis. Bauh. pin. 206.*  
*Saponaria vulgaris. Cam. epit. 152.*  
*hybrida. 2. Saponaria concava anglica. Bauh. pin. 206. Morij. hist. 2. p. 548. f. 5. t. 22. f. 52.*  
*Gentiana folio convoluto. Bauh. hist. 3. p. 521.*  
*Habitat in Europa media. 2.*  
*Hybrida varietas singularis, instar alterius Pelorice Antirrhini, qua ex Saponaria transit in monopetalam Gentianam, reperta in Spineis, Lichbarrow, Northamptonshire Angliæ, a Gerardo, amplius ibi non occurrit teste Mortonio, forte etiamnum alicubi in Hortis Curiosorum asservata, quæ semine non propagatur.*  
*cretica 2. SAPONARIA calycibus striatis quinquangularibus, caule erecto subdichotomo, foliis subulatis.*  
*Saxifraga altera. Alp. exot. 292. t. 291.*  
*Habitat in Cretæ aridis.*  
*Caulis brachiatus f. dichotomus, teres, villosus viscidus.*  
 284

*angustus, pedalis. Folia lineari-subulata, levia. Pedunculi axillares, solitarii, uniflori, striati. Flos erectus. Calyx quinquangularis angulis striis tribus, monophyllus, oblongus, acutus, inter dentes membranaceus. Petala parva, integra.*

3. SAPONARIA calycibus pyramidatis quinquangularibus, foliis ovatis acuminatis sessilibus. *Hort. cliff. 166. Hort. upf. 107. Roy. lugdb. 443. Guett. stamp. 287. Sauv. monsp. 153.* Vaccaria.

*Lychnis fr. getum rubra, foliis perfoliatis. Bauh. pin. 204. Vaccaria Dod. pempt 104.*

*Habitat inter segetes Galliae, Germaniae, Helvetiae, Orientis. ☉*

4. SAPONARIA calycibus cylindricis villosis, caulis dichotomis procumbentibus. Ocymoides.

*Saponaria minor quibusdam. Bauh. hist. 3. p. 344.*

*Lychnis vel Ocymoides repens montanum. Bauh. pin. 206.*

*Habitat in Helvetiae, Italiae, Monspeli petrosis, umbrosis. 24*

5. SAPONARIA calycibus cylindricis villosis, caule dichotomo erecto patulo. *Hort. upf. 106. n. 2.* orientalis.

*Saponaria caule dichotomo, foliis lanceolatis petiolatis.*

*floribus solitariis pedunculatis. Roy. lugdb. 445.*

*Lychnis orientalis annua supina, antirrhini folio, flore minimo purpurascens. Tournef. cor. 25. Dill. elth. 205. t. 167. f. 204.*

*Habitat in Oriente. ☉*

*Similis praecedenti; huic Folia lineari-lanceolata, illi lanceolato ovata. Calyces fructus ovati, adpersi punctis elevatis piliferis. Petala acute emarginata absque coronula.*

6. SAPONARIA calycibus teretibus, corollis coronatis, floribus subumbellatis, foliis sublinearibus canaliculatis. lutea.

*Lychnis floribus umbellatis ochroleucis, petalis ovatis, filamentis nigris. All. pædem. 29. t. 5. f. 2.*

- β. Globularia lutea montana. *Col. ecphr. 1. p. 152. t. 153.*

*Bellis montana, globoso luteo flore. Bauh. pin. 262.*

*Lychnis lutea montana, globulariae capite & facie. Barr. ic. 498.*

*Lychnis rubra*, globulariæ capitulo. *Boeck. mus.* 2. p. 75. t. 62. f. 1.

*Habitat in Alpitus Exilliarum, Fenestrellarum, Canisio.* 24

## DIANTHUS.

\* *Flores aggregati.*

barbatus. 1. **DIANTHUS** floribus aggregatis fasciculatis: squamis calycinis ovato subulatis tubum æquantibus foliis lanceolatis. *Hort. cliff.* 155. *Hort. upf.* 165. *Roy. lugdb.* 444. *Sauv. monsp.* 144.

*Caryophyllus hortensis barbatus latifolius.* *Bauh. pin.* 208.

*Thyrsis. Reneal. spec.* 47.

β. *Caryophyllus barbatus hortensis angustifolius.* *Bauh. pin.* 209.

*Armerius flos alter.* *Dod. pempt.* 176.

*Habitat in - - - 24*

carthusiano- 2. **DIANTHUS** floribus subaggregatis: squamis calycinis ovatis aristatis tubum subæquantibus, foliis linearibus trinerviis. *Hort. upf.* 105.

*Dianthus floribus aggregatis, squamis calycinis lanceolatis, corollis crenatis.* *Guert. flamp.* 284.

*Caryophyllus sylvestris vulgaris latifolius.* *Bauh. pin.* 209.

*Caryophyllus arvensis, calyculo florum numerofo.* *Læf. pruss.* 37. f. 7.

β. *Caryophyllus sylvestris, flore rubro plurimo de summo caule prodeunte.* *Segu. ver.* 434. t. 8. f. 2.

*Habitat in Germaniæ, Italiæ, Siciliæ sterilibus apricis.* 24

*Differt a D. barbato Foliis dimidio angustioribus, rigidioribus, trinerviis nec uninerviis. Caule parum scabro, nec levissimo. Petalis distantibus, supra villosis, nec glabris. Pistillis tubo longioribus.*

Armeria. 3. **DIANTHUS** floribus aggregatis fasciculatis: squamis calycinis lanceolatis villosis tubum æquantibus. *Hort. cliff.* 165. *Fl. suec.* 345. 381. *It. gotl.* 301. *Roy. lugdb.* 443. *Sauv. monsp.* 144.

*Caryophyllus barbatus sylvestris.* *Bauh. pin.* 208.

*Armeria sylvestris altera.* *Lob. ic.* 448.

*He-*

*Habitat in sterilibus* Gotlandiæ, Germaniæ, Galliæ, Italiæ. ☉

*Petala acuminata, uno alterove dente notata.*

4. DIANTHUS floribus aggregatis capitatis: squamis prolifera. calycinis ovatis obtusis muticis tubum superantibus.

*Dianthus floribus aggregatis, capitulo magno, squamis calycinis ovatis obtusis magnis. Hort. upf. 106. Sauv.*

*monsp. 144.*

*Caryophyllus sylvestris prolifera. Bauh. pin. 209. Segu. veron. 26. t. 7. f. 1.*

*Caryophyllus sylvestris annuus, multis capsulis simul junctis donatus. Morif. hist. 2. p. 563.*

*Habitat in Germaniæ & australioris Europæ pascuis sterilibus.* ☉

5. DIANTHUS floribus solitariis, squamis calycinis diminutis, octonis florem superantibus.

*Caryophyllo prolifero affinis, unico ex quolibet capitulo flore. Bauh. pin. 219.*

*Caryophyllus sylvestris minimus. Tabern. hist. 290.*

*Habitat in Germania.* ☉

*Nata D. prolifera, eique nimis affinis, an pro Specie agnoscenda? Folia angustiora. Caulis ramosus. Flores terminales simplices, nec aggregati. Calyx paribus 4 squamarum, quarum interiores per gradus majores obtusiores. Corolla brevissima, vix extra tubum prominens; adeoque fructificatio D. prolifera, sed non aggregata, verum simplex.*

\* \* Flores solitarii.

6. DIANTHUS floribus solitariis: squamis calycinis Caryophyllis subovatis brevissimis, corollis crenatis. *Hort. cliff. 164. Hort. upf. 104. Mat. med. 213. Roy. lugdb. 443. Sauv. monsp. 143.*

7. Caryophyllus hortensis simplex, flore majore. *Bauh. coronarius. pin. 208.*

8. Caryophyllus altis major. *Bauh. pin. 207.*

9. Caryophyllus maximus ruber & variegatus. *Bauh. pin. 209.*

10. Caryophyllus - - flore pleno ex squamis calycinis imbricatis. longissime imbricatis. *Hort. cliff. l. c.*

*Caryophyllus spicam frumenti referens. E. N. C. cent. 3. p. 368. t. 9. Hort. cliff. 164. Phil. bot. 3.*

- inodorus*. 1. *Tunica angustifolia procumbens, petalis serratis. Hall. helv. 382.*  
*Caryophyllus sylvestris biflorus. Bauh. pin. 209. prodr. 104.*  
*Caryophyllus sylvestris, flore rubro inodoro, calyce oblongo cum brevibus unguibus. Segg. ver. 425. t. 7. f. 3.*  
*Habitat in Italia, Rajus; in Alpibus Helveticis, sed angustior. 24.*  
*Coronarius a. cum reliquis γ. β. δ. — forte ex 1. inodoro prognatus est.*  
*Varietas δ. maxime singularis est.*
- beltoides*. 7. **DIANTHUS** floribus solitariis: squamis calycinis lanceolatis binis, corollis crenatis. *Hort. cliff. 164. Fl. fvec. 342; 382. Sauv. monsp. 143.*  
*Caryophyllus minor repens nostras. Raj. hist. 988. Dill. elth. 402.*  
*Caryophyllus simplex supinus latifolius. Bauh. pin. 209.*  
*Betonica coronaria f. Caryophyllus minor, folio viridi nigricante, repens. Bauh. hist. 3. p. 329.*  
*Habitat in Europæ pratis. 24.*
- glauca*. 8. **DIANTHUS** floribus subfoliariis squamis calycinis lanceolatis quaternis brevibus, corollis crenatis. *Hort. cliff. 164 Hort. upf. 104.*  
*Tunica ramosior, flore candido cum corolla purpurea. Dill. elth. 400. t. 298. f. 348.*  
*Habitat in Anglia. 24.*
- chinensis*. 9. **DIANTHUS** floribus solitariis: squamis calycinis subulatis patulis tubum æquantibus, corollis dentatis. *Hort. cliff. 164 Hort. upf. 104. \* Roy. lugdh. 443.*  
*Caryophyllus sinensis supinus, leucoji folio, flore unico, Tournef. art. 1705. p. 348. f. 5. Mill. ic. 81. f. 2.*  
*Habitat in China. ☉*
- monspessulanus*. 10. **DIANTHUS** floribus solitariis: squamis calycinis subulatis tubum æquantibus, petalis multifidis. *Amern. acad. 4. p. 313. \**  
*Habitat Montpelii, Veronæ. 24.*

11. **DIANTHUS** floribus paniculatis : squamis calycinis brevibus acuminatis, corollis multifido-capillaribus, caule erecto. *Aman. acad.* 4. p. 272. \* *Fl. svec.* 2. p. 383. *superbus.*  
*Caryophyllus simplex alter, flore laciniato odoratissimo. Bauh. pin.* 210.  
*Caryophyllus sylvestris VI. Clus. hist.* 1. p. 284.  
*Habitat in Gallia, Germania, Dania, Kemi.* ♂
12. **DIANTHUS** floribus solitariis : squamis calycinis subovatis brevissimis, corollis multifidis fauce pubescentibus. *plumarius.*  
*Dianthus floribus solitariis, petalis multifidis basi canaliculatis. Hort. upf.* 105. \* *Sauv. monsp.* 143.  
*Dianthus floribus solitariis, corollis lacero partitis, squamis calycinis ovatis acutis. Fl. svec.* 244.  
*Dianthus petalis multifidis. Fl. lapp.* 170. *Hort. cliff.* 174. *Roy. lugdb.* 443.  
*Caryophyllus sylvestris, floribus lamuginosis hirsutis. Bauh. pin.* 210.  
*Caryophyllus sylvestris V. species alia. Clus. hist.* 1. p. 284.  
*Habitat in Europæ & Canadæ pascuis nemorosis.*  
*Folia rore cæsis glauca.*  
 \* \* \* *Caule unifloro herbaceo.*
13. **DIANTHUS** caulibus subunifloris : squamis calycinis ovatis obtusis, corollis multifidis, foliis linearibus. *It. gotl.* 318. *Fl. svec.* 343; 384. \* *arenarius.*  
*Dianthus caule simplici unifloro. Monnier. obs.* 152.  
*Dianthus foliis brevibus, squamis muticis, caule unifloro. Sauv. meth.* 143.  
*Caryophyllus sylvestris humilis, flore unico. Bauh. pin.* 209.  
*Caryophyllus sylvestris 1. Clus. hist.* 1. p. 282. *Dill. elth.* 402.  
*Armerius flos tertius. Dod. pempt.* 176.  
*Habitat in Europæ frigidioris arena mobili.* 2.  
*Affinitatem hujus summam cum præcedente docet Petalorum faux; Petala magis oblonga, divisa & lacera ultra medium disci (illi vero tantum multifida) basi macula livida acuta, pilis purpurascensibus adspersa.*



alpinus. 14. **DIANTHUS** caule unifloro, corollis crenatis: squamis calycinis exterioribus tubum æquantibus, foliis linearibus obtusis.

Caryophyllus pumilus latifolius. *Bauh. pin. 209. prodr. 104. Burf. XI. 95.*

Caryophyllus sylvestris, flore magno inodoro hirsuto. *Bauh. pin. 209.*

Caryophyllus sylvestris 2. *Clus. hist. 1. p. 283. f. 1. bon.*

*Habitat in Stiria, Austria. 24*

*Radix lignosa, exserens caules plures. Caulis longitudine digiti, articulis tribus constans. Folia caulis linearia, obtusiuscula, plana, obtusa, vel versus terram, ubi cespites constituunt, lanceolato-linearia. Calycis squamæ exteriores longitudine tubi calycis: interiores dimidio breviores. Corolla magna. Petala calycis longitudine, rotundata, dentibus brevissimis numerosis crenata, fauce parum villosa.*

virginicus. 15. **DIANTHUS** caule subunifloro, corollis crenatis: squamis calycinis brevissimis, foliis subulatis.

Caryophyllus sylvestris repens multiflorus. *Bauh. pin. 209; Burf. XI. 99.*

β. 1 unica rupestris, folio cæcio molli, flore carneo. *Dill. elth. 401. t. 298. f. 385.*

*Habitat Monspelli. 24*

*Radix perennis. Folia radicalia, copiosa, erecta, in cespitem imbricata, acuminata, facie Statice ferme. Caules aliquot, spithamæ, 4 articulis simplicissimis constantes, foliis minoribus & summis minimis ex vagina perfoliata ortis. Calycis squamæ latæ, ovata, acutæ, brevissimæ: paribus remotis. Corollæ limbus dimidio brevior tubo calycis, rotundatus, crenatus. Flos unicus in caule, interdum & aliter ex ala folii supremi.*

\* \* \* \* *Frutescentes.*

arbores. 16. **DIANTHUS** caule fruticoso, foliis subulatis.

Caryophyllus creticus arboreus, juniperi folio. *Tournef. cor. 23.*

Caryophyllus arboreus creticus. *Bauh. pin. 208. prodr. 104.*

Betonica coronaria arborea cretica. *Bauh. hist. 3. p. 328.*

*Habitat in Creta. 6*

17. **DIANTHUS** caule fruticoso, foliis lanceolatis. fruticosus.  
*Caryophyllus græcus arboreus, leucoj folio peramaro.* *Tournef. cor. 23. itin. 1. p. 183. t. 9.*  
*Habitat in Græcia. 5*

## TRIGYNIA.

### CUCUBALUS.

1. **CUCUBALUS** calycibus campanulatis; petalis distinctibus, pericarpis coloratis, ramis divaricatis. bacciferus.  
*Cucubalus caule ramoso, floribus trigynis, fructu colorato globoso.* *Hort. cliff. 170. Hort. ups. 109. Roy. lugdb. 448. Sauv. monsp. 152.*  
*Alfine scandens baccifera.* *Bauh. pin. 250.*  
*Alfine repens.* *Dod. pempt. 403.*  
*Habitat in Tataria, Germaniæ, Galliæ, Italiæ nemoribus, sepibus. 2*
2. **CUCUBALUS** calycibus subglobosis glabris reticulato-venosis, capsulis trilocularibus, corollis subnudis. Behen.  
*Fl. sic. 360; 385.*  
*Cucubalus floribus trigynis, calycibus glabris, capsulis trilocularibus.* *Hort. cliff. 176. Roy. lugdb. 448.*  
*Cucubalus calycibus ovatis hermaphroditis.* *Fl. lapp. 180.*  
*Lychnis sylvestris, quæ Behen album vulgo.* *Bauh. pin. 205.*  
*Papaver spumeum.* *Lob. ic. 340.*
- β. **Lychnis** maritima repens. *It. gotl. 197.*  
*Habitat in Europæ septentrionalis pratis siccis. 2*  
*Varietas Feminea in Horto upsaliensi frequens. Hæc Hermaphrodito minor. Calyces exacte ovati, magis obscuri, nec oblongo-ovati. Corolla minor. Stamina corolla dimidio breviora terminata tuberculo absque antheris. Styli 3. f. 5. corolla longiores, declinati. Planta utraque fertilis.*
3. **CUCUBALUS** foliis obovatis carnosiss. *Roy. lugdb. Fabarius. 448.*  
*Lychnis maritima saxatilis, folio anacampserotis.* *Tournef. cor. 24. Boert. lugdb. 1. p. 214.*

Behen album f. Polemonium saxatile, fabariæ folio, ficulum. *Bocc. mus.* 133. t. 92.

*Habitat in Sicilia.*

- viscosus. 4. CUCUBALUS floribus lateralibus undique decumbentibus, caule indiviso, foliis basi reflexis. *Fl. suec.* 2. n. 386. \*

Cucubalus foliis amplexicaulibus, floribus verticillatis, pedunculis erectis. *Roy. lugdb.* 448?

Lychnis montana viscosa noctiflora hirsuta latifolia alba, floribus absque appendicibus. *Till. pis.* 105.

Lychnis orientalis maxima, buglossi folio undulato. *Tournef. cor.* 24. itin. 2. p. 361. t. 361.

Lychnis major noctiflora drubensis. *Raj. hist.* 995?

*Habitat in Svecia, Italia, Anglia, Ararat.* ♂.

- stellatus. 5. CUCUBALUS foliis quaternis. *Hort. upf.* 110.

Drypis foliis quaternis. *Cold. noveb.* 106.

Silene foliis quaternis. *Gron. virg.* 50.

Lychnis caryophyllæus virginianus, gentianæ foliis glabris quatuor ex singulis geniculis caulem amplexantibus, flore amplo fimbriato. *Raj. hist.* 1895.

*Habitat in Virginia, Canada.*

Petala, absque corona, quadrifida, lobis bipartitis.

- egyptiacus. 6. CUCUBALUS floribus erectis, petalis emarginatis retroflexis: utrinque denticulo notatis. *Hassletq.*

*Habitat in Agypto.*

- fibricus. 7. CUCUBALUS petalis emarginatis, floribus subverticillatis: verticillis umbellatis aphyllis.

Viscago foliis imis petiolatis ovatis, floribus paniculatis, petalis emarginatis. *Hall. gert.* 1. p. 150. fig. bona.

*Habitat in desertis Tacorow; inter Woroniz & Bielgrad.*

*Stamina flore longiora.*

- tataricus. 8. CUCUBALUS petalis bipartitis, floribus secundis decumbentibus, pedunculis oppositis solitariis erectis, caule simplicissimæ. *Hort. upf.* 111. \*. *Gort. ingr.* 68.

Lychn.

*Lychnis septentrionalium*, foliis hyssopi, floribus uno versu positis albis. *Hall. gert.* 33. \*

*Habitat in Tartaria, Russia.* 24

9. **CUCUBALUS** calycibus clavatis, petalis semibifidis, panicula dichotoma erecta, genitalibus declinatis, caule incano. *italicus.*

*Habitat in Italia.* ♂

Folia lanceolata, obtusiuscula, subnuda. Caulis, praesertim inferne, tomento vix conspicuo incanus. Panicula dichotoma, floribus pedunculatis, erectis. Calyces erecti, clavati, exarati sulcis decem. Petala semibifida, latiuscula, alba, subtus plumbea. Stamina alba.

10. **CUCUBALUS** petalis bipartitis, floribus paniculatis, staminibus longis, foliis lanceolato-ovatis. *Hort. upl.* III. \* *catholicus.*

*Silene foliis ovatis utrinque acutis, caule paniculato, floribus nutantibus tenuissimis. Roy. lugdb.* 447.

*Lychnis viscosa altissima, flore mucoso, ocymastri facie. Cupan. cath.* 110.

- β. *Lychnis nocturna non viscosa, herbaceo flore. Dill. elth.* 425. t. 316. f. 408.

*Habitat in Italia, Sicilia.* 24

*Calyces fusci, nitidi.*

11. **CUCUBALUS** petalis semibifidis, panicula brachiata dichotoma, caule foliisque holosericeis: radicalibus spatulatis. *mollissimus.*

*Lychnis maritima pulverulenta, folio carnosio. Tournef. inst.* 338.

*Habitat in Italiae maritimis.* 24. ♀

Caules plures, rigiduli, erecti, pedales, subtomentosi, mollissimi: Ramis inferne alternis, brevioribus. Caules his immixti steriles dimidio breviores, erecti, apice confertim foliosi. Folia inferiora spatulata, reliqua lanceolata; omnia subcarnosa, mollissima, tomentosa, obtusiuscula, apice subreflexa, patentia. Panicula brachiata: ramis dichotomis. Calyces obtuse dentati, clavati, erecti, mollissimi. Flores albi: Laminis calyce brevioribus, semibifidis. Stamina longitudine calycis. Styli & colla longiores.

- Oites.** 12. **CUCUBALUS** floribus dioicis, petalis linearibus indivisis. *Hort. cliff.* 272. *Roy. lugdb.* 445.  
*Lychnis viscosa*, flore muscoso. *Bauh. pin.* 206.  
*Lychnis sylvestris sesamoides minor*, flore muscoso.  
*Mentz. pug.* t. 3. f. 2.  
*Sesamoides magnum falmanticum.* *Clus. hist.* 1. p. 295.  
*Habitat in Silesia, Austria, Vallesia, Anglia, Gallia, Sibiria.* 24
- reflexus.** 13. **CUCUBALUS** floribus spicatis alternis secundis subsessilibus, corollis obsoletis nudis. *Hort. upf.* 112. \*  
*Lychnis sylvestris alba*, spica reflexa. *Magn. monsp.* 171.  
t. 170. *Raj. hist.* 996. n. 12.  
*Habitat Montpelii.* 24  
*Confer. Silene mutabilis, cujus an sola varietas?*

## S I L E N E.

\* *Floribus lateralibus solitariis.*

- anglica.** 1. **SILENE** hirsuta, petalis integerrimis, floribus erectis, fructibus reflexis pedunculatis alternis. *Hort. upf.* 113.  
*Viscago cerasii foliis*, vasculis pendulis anglica. *Dill. elth.* 417. t. 309. f. 398.  
*Lychnis sylvestris hirsuta annua*, flore minore albo.  
*Vaill. parif.* 121. t. 16. f. 12.  
*Habitat in Anglia, Gallia.* ☉  
*Petala in hac non crenata, in sequente ☿ emarginata ☿ crenata. Folia inferiora obovato-lanceolata versus basin ciliata. Calyces non pilosi, sed angulis muricatis: aculeis reflexis vix conspicuis.*
- lufitanica.** 2. **SILENE** hirsuta, petalis dentatis, floribus erectis, fructibus divaricato-reflexis alternis. *Hort. upf.* 113.  
*Hort. cliff.* 172.  
*Silene floribus alternis*, petalorum limbis integris crenatis. *Hort. cliff.* 172.  
*Viscago hirsuta lufitanica*, stellato flore. *Dill. elth.* 420. t. 311. f. 401.  
*Habitat in Lusitania.* ☉  
*Calyces fructus horizontaliter reflexi, valde pilosi.*  
*Corolla indivisa, crenata, incarnata.*

3. **SILENE** petalis integerrimis subrotundis, fructibus erectis alternis. *Hort. cliff.* 171. *Hort. upf.* 113. *Roy. lugdb.* 446. quinqueval-  
nera.
- Silene spica incurva uno versu florida, foliis imis spatulatis. Sauv. monsp.* 146.
- Lychnis sylvestris lanuginosa minor. Bauh. pin.* 206.
- Lychnis hirsuta, flore eleganter variegato. Raj. hist.* 997.
- Lychnis hirsuta minor, flore variegato. Dod. mem.* 4. p. 291.
- Habitat in Hispania, Lusatia, Italia, Gallia. ☉*
- Calyces fructus erecti, pilosi, visciidi. Petala obtusa, vix emarginata, integerrima, purpurea margine pallido.*
4. **SILENE** floribus spicatis alternis secundis sessilibus, petalis bifidis. nocturna.
- Silene foliis lanceolatis, caule ramoso & floribus secundis erectis, calycibus hirsutis. Roy. lugdb.* 447.
- Vilcago hirta noctiflora, floribus obsoletis spicatis. Dill. elth.* 420. t. 310. f. 400.
- Lychnis sylvestris hirsuta elatior spicata, lini colore. Barr. ic.* 10. t. 27. f. 1.
- Lychnis sylvestris nocturna pilosa, floribus unius ordinis dilute purpureis. Cup. cath.* 119. *Raj. suppl.* 480.
- Lychnis segetum meridionalium annua hirta, floribus albis uno versu dispositis. Moris. hist.* 2. p. 346. f. 5. t. 36. f. 7.
- Habitat in Hispania, Monspeli, Pennsylvania. ☉*
- Folia inferiora obovata, concava, rigida. Flores sessiles, ex bracteis oppositis alterni, alternatim florentes. Petala alba, bifida, majuscula, subius virefcentia, nec post anthesin rubra. Similis Cucubalo reflexo, sed Caule ramosiore altiore, Petalis extra calycem patentibus.*
5. **SILENE** floribus subspicatis alternis secundis, petalis indivisis, fructibus erectis
- Silene hirsuta. foliis sessilibus oblongo-cuneiformibus obtusis, calycibus striatis, fructibus erectis alternis. Dalib. paris.* 129.

- Odites.** 12. **CUCUBALUS** floribus dioicis, petalis linearibus indivisis. *Hort. cliff.* 272. *Roy. lugdb.* 445.  
*Lychnis viscosa*, flore muscoso. *Bauh. pin.* 206.  
*Lychnis sylvestris sesamoides minor*, flore muscoso.  
*Mentz. pug. t. 3. f. 2.*  
*Sesamoides magnum salmanticum.* *Clus. hist. r. p.* 295.  
*Habitat in Silesia, Austria, Vallesia, Anglia, Gallia, Sibiria.* 24
- reflexus.** 13. **CUCUBALUS** floribus spicatis alternis secundis subseffilibus, corollis obsoletis nudis. *Hort. upf.* 112. \*  
*Lychnis sylvestris alba*, spica reflexa. *Magn. monsp.* 171.  
*t. 170. Raj. hist.* 996. n. 12.  
*Habitat Monspeli.* 24  
*Confer. Silene mutabilis, cujus an sola varietas?*

## S I L E N E.

\* *Floribus lateralibus solitariis.*

- anglica.** 1. **SILENE** hirsuta, petalis integerrimis, floribus erectis, fructibus reflexis pedunculatis alternis. *Hort. upf.* 113.  
*Viscago cerastii foliis, vasculis pendulis anglica.* *Dill. elth.* 417. t. 309. f. 398.  
*Lychnis sylvestris hirsuta annua*, flore minore albo.  
*Vaill. parif.* 121. t. 16. f. 12.  
*Habitat in Anglia, Gallia.* ☉  
*Petala in hac non crenata, in sequente ☿ emarginata ☿ crenata. Folia inferiora obovato-lanceolata versus basin ciliata. Calyces non pilosi, sed angulis muricatis: aculeis reflexis vix conspicuis.*
- lufitanica.** 2. **SILENE** hirsuta, petalis dentatis, floribus erectis, fructibus divaricato-reflexis alternis. *Hort. upf.* 113.  
*Hort. cliff.* 172.  
*Silene floribus alternis, petalorum limbis integris crenatis.* *Hort. cliff.* 172.  
*Viscago hirsuta lufitanica, stellato flore.* *Dill. elth.* 420. t. 311. f. 401.  
*Habitat in Lusitania.* ☉  
*Calyces fructus horizontaliter reflexi, valde pilosi.*  
*Corolla indivisa, crenata, incarnata.*

3. **SILENE** petalis integerrimis subrotundis, fructibus erectis alternis. *Hort. cliff.* 171. *Hort. upf.* 113. *Roy. lugdb.* 446. quinqueval-  
nera.

*Silene* spica incurva uno versu florida, foliis imis spatulatis. *Saur. monsp.* 146.

*Lychnis sylvestris lanuginosa minor.* *Bauh. pin.* 206.

*Lychnis hirsuta*, flore eleganter variegato. *Raj. hist.* 997.

*Lychnis hirsuta minor*, flore variegato. *Dod. mem.* 4. p. 291.

*Habitat in Hispania, Lusatia, Italia, Gallia.* ○

*Calyces fructus erecti, pilosi, visci.* *Petala obtusa, vix emarginata, integerrima, purpurea margine pallido.*

4. **SILENE** floribus spicatis alternis secundis sessilibus, nocturna. petalis bifidis.

*Silene* foliis lanceolatis, caule ramofo & floribus secundis erectis, calycibus hirsutis. *Roy. lugdb.* 447.

*Viscago hirta noctiflora*, floribus obsoletis spicatis. *Dill. elth.* 420. t. 310. f. 400.

*Lychnis sylvestris hirsuta elatior spicata*, limbo colore. *Barr. ic.* 10. t. 27. f. 1.

*Lychnis sylvestris nocturna pilosa*, floribus unius ordinis dilute purpureis. *Cup. cath.* 119. *Raj. suppl.* 480.

*Lychnis segetum meridionalium annua hirta*, floribus albis uno versu dispositis. *Morif. hist.* 2. p. 346. f.

5. t. 36. f. 7.

*Habitat in Hispania, Montpelii, Pennsylvania.* ○

*Folia inferiora obovata, concava, rigida.* Flores sessiles, ex bracteis oppositis alterni, alternatim florentes. *Petala alba, bifida, majuscula, subtus virescens, nec post anthesin rubra.* *Similis Cucubalo reflexo, sed Caule ramofoiore altiore, Petalis extra calycem patentibus.*

5. **SILENE** floribus subspicatis alternis secundis, petalis indivisis, fructibus erectis

*Silene hirtula*, foliis sessilibus oblongo-cuneiformibus obtusis, calycibus striatis, fructibus erectis alternis. *Dalib. paris.* 129.

P P 2

Lychn



*Lychnis sylvestris hirsuta annua, flore minore carnea.*  
*Vaill. parif. 121. t. 16. f. 12.*

*Viscago hirta gallica, flore parvo carneo, petalis integris.* *Dill. elth. 419. t. 310. f. 399.*

*Habitat in Gallia.* ☉

*cerastoides.* 6. **SILENE** petalis bipartitis, fructificationibus erectis subfessilibus, calycibus subpilosis subfessilibus.

*Silene hirsuta, petalis emarginatis, fructibus erectis alternis hirsutis fessilibus.* *Sp. pl. 1. p. 417.*

*Silene rigidula.* *Amæn. acad. 4. p. 313. \**

*Viscago cerastii foliis, vasculis erectis fessilibus.* *Dill. elth. 416. t. 309. f. 307.*

*Habitat in Europa australi.* ☉

\* \* *Floribus lateralibus confertis.*

*mutabilis.* 7. **SILENE** petalis bifidis, calycibus angulatis pedunculatis, foliis lanceolato-linearibus. *Amæn. acad. 4. p. 314. \**

*Lychnis flore albo minimo.* *Raj. hist. 906.*

*Lychnis arvensis minor anglica.* *Raj. hist. 1004.*

*Habitat in Europa australi.* ☉

*Affinis Cucubalo reflexo, cum qua conjungit Raj. hist. 664, sed sata utraque diversam sese mihi ostendebat.*

*nutans.* 8. **SILENE** petalis bifidis, floribus lateralibus secundis cernuis, panicula nutante.

*Silene foliis lanceolatis, caule paniculato, floribus nutantibus, calyce striato, corolla involuta.* *Hort. clif. 171. Fl. svec. 366; 588. \**

*Lychnis montana viscosa alba latifolia.* *Bauh. pin. 205.*

*Habitat in Europæ borealis pratis aridis.* 24

*amœna.* 9. **SILENE** petalis bifidis: coronula subcoadunata, floribus secundis, pedunculis oppositis trifloris, ramis alternis.

*Silene petalis bifidis coronatis: corolla coadunata, calycibus erectis raris pilosis.* *Hort. upf. 114.*

*Lychnis maritima repens.* *Bauh. pin. 205.*

*Lychnis maritima anglicana.* *Bauh. hist. 3. p. 357.*

*Lychnis perennis angustifolia marina anglica procumbens.* *Morif. hist. 2. p. 535. f. 5. t. 20. f. 2.*

*Habitat in Tataria; in Angliæ meridionalis litoris.* 24

*Caules diffusi, leviusculi, adscendentes. Rami alterni, divaricati, breviores. Folia supra glabra. Racemus secundus: pedunculis oppositis, trifloris. Calyces angulis 10 purpurascentibus villosis. Petala alba, semibifida: Nectarii lamellis bifidis, patulis, stria-alba.*

10. *SILENE* petalis bifidis, caule fruticoso, foliis lato-lanceolatis, panicula trichotoma. fruticosa.

*Silene caule folioso fruticoso, foliis lineari-lanceolatis acutis glabris, capsulis ovatis Vir. cliff. 38. Hort. cliff. 171. n. 4. Hort. upf. 111. n. 2. Roy. lugdb. 446.*

*Lychnis myrtifolia behen albo similis. Bauh. pin. 205.*

*Ocymoides fruticosum. Cam. hort. 109. t. 33.*

*Saponaria frutescens, acutis foliis, ex sicilia. Boer. fl. 58. t. 30.*

*Habitat in Sicilia, Germania. ‡*

11. *SILENE* foliis suborbiculatis carnosissimis hirsutis, racemo secundo. crassifolia.

*Habitat ad Cap. b. Spei.*

*Caulis bipedalis, firmus, hirsutus: Ramis alternis. Folia orbiculata s. obovata, obtusissima, Portulaca, crassa, carnosissima, utrinque pilosa: vix extimi pollicis magnitudine. Racemus terminalis, secundus, Flores 9 s. 12, triflori colore; calycibus hirsutis; bracteis ovatis. Petala emarginata, rite coronata. Pistilla tria.*

12. *SILENE* petalis semibifidis, foliis ovatis scabriusculis, panicula elongata subaphylla. viridiflora.

*Silene foliis lanceolatis, caule paniculato nudo, floribus erectis. Hort. cliff. 171. Roy. lugdb. 446.*

*Lychnis ocymastri facie, flore viridi. Herm. parad. 199. t. 199.*

*Habitat in Lusitania. ♂*

*Planta pedalis & ultra. Caules erecti, teretes, pubescentes, geniculis incrassatis. Folia lanceolato-ovata, pubescentia, viridia, acuta. Rami alterni. Panicula brachiata pedunculis dichotomis, paucifloris. Calyces nutantes, cylindrici, angulis decem viscosi, uti pedunculi. Flores virido-virescentes: petalis semibifidis.*

P. p.

2

*semibifidis: lobis recurvatis, linearibus. Stamina longitudine tubi; Pistilla longitudine petalorum.*

*gigantea.*

13. **SILENE** foliis radicalibus cochleariformibus obtusissimis, caule subverticillato.

*Silene* foliis obverse ovatis crassis, limbis corollæ bifidis a sole revolutis. *Wach. ultr.* 391.

*Lychnis græca*, sed arborescens folio & facie, flore albo. *T. Walth. hort.* 32. t. 11.

*Habitat in Africa.* ♂, ♀. h

*Folia prioris anni, maxima. Caulis posterioris anni, bipedalis, simplex: articulis inferioribus brevibus, villosis; superioribus longissimis, nudis, viscoso-glutinosus. Folia inferiora obovata, carnosæ, fragilia, concava, pubescentia, sessilia. Pedunculi communes oppositi, horizontaliter patentes, brevissimi, dichotomi. Calyces pubescentes Corollæ virides corona obsoleta, ut fere nulla quasi Cucubali generis, sed tamen aliquat. Flores noctu tantum explicati.*

*bupleurei-  
des.*

14. **SILENE** petalis bifidis, floribus pedunculatis oppositis bractea brevioribus, foliis lanceolatis acutis glabris.

*Silene* caule folioso herbaceo, foliis lanceolatis acutis glabris, calycibus erectis. *Hort. cliff.* 171. *Roy. lugdb.* 446.

*Lychnis orientalis*, bupleuri folio. *Tournef. cor.* 2. ic. 2. p. 139. t. 154.

*Habitat in Persia.*

\* \* \* *Floribus ex dichotomia caulis.*

*Conoidea.*

15. **SILENE** calycibus fructus globosis acuminatis striis triginta, foliis glabris, petalis integris. *Hort. ups.* 110. *Sauv. monsp.* 146.

*Cucubalus calycibus conicis striatis. Hort. cliff.* 171. *Roy. lugdb.* 448.

*Lychnis sylvestris latifolia*, calycibus turgidis striatis. *Bauh. pin.* 205.

*Lychnis sylvestris.* 2. *Clus. hist.* 1. p. 288.

*Habitat inter segetes Hispaniæ.* ☉

*conica.*

16. **SILENE** calycibus fructus conicis striis triginta, foliis mollibus, petalis bifidis. *Hort. ups.* 110.

*Lychnis sylvestris angustifolia*, calycibus turgidis striatis. *Bauh. pin.* 205.

*Lychn-*

*Lychnis glabra annua, foliis oblongis mucronatis, flore amplo rubello, capsula oblonga pyramidalis. Moris. hist. 2. p. 542. f. 5. t. 36. f. 6.*

*Habitat in Hispania, Galloprovincia, Oriente. ☉*

17. *SILENE calycibus glabris ovatis venoso-reticulatis, Behen, capsulis trilocularibus.*

*Lychnis vesicaria cretica, parvo flore purpurascens.*

*Dill. elth. 427. t. 317. f. 409.*

*Lychnis cretica, parvo flore, calyce striato purpurascens. Tournesf. cor. 24.*

*Habitat in Creta. ☉*

*Differt a Cucubalo Behen corona corollæ, petalorum laminis vix  $\frac{1}{2}$  longitudine calycis, radice annua, corollis purpurascens, antheris albis. Calyx obtusus. Flores nutantes, pedunculo e dichotomia, flore ipso brevior.*

18. *SILENE petalis emarginatis, calycibus glabris reticulato-venosis acuminatis pedunculo longioribus, caule dichotomo stricto. Aman. acad. 4. p. 314. \**

*Silene caule subdichotomo, floribus erectis subsessilibus, foliis lanceolatis piloso-viscidis. Aman. acad. 1. p. 158.*

*Habitat in Hispania & Tolosæ. ☉*

19. *SILENE calycibus fructiferis pendulis inflatis: angulis decem scabris. Hort. upf. 109. \**

*Cucubalus floribus trigynis erectis, fructibus pendulis, calycibus striis decem crispis. Hort. cliff. 170.*

- β. Cucubalus floribus trigynis erectis, calycibus fructus pendulis angulosis. Hort. cliff. 171.*

*Viscago hirsuta sicula, lychnidis aquaticæ facie, supina. Dill. elth. 421. t. 312. f. 404.*

*Habitat in Creta & Sicilia. ☉*

20. *SILENE calycibus decemangularibus: dentibus tum æquantibus, caule dichotomo.*

*Cucubalus calycibus fructus ovatis erectis: angulis decem: alternis venosis. Hort. upf. 109. \* Fl. suec.*

*2. n. 389. Sauv. monsp. 146.*

*Lychnis noctiflora. Bqnh. pin. 205. Raj. hist. 994.*

*Ocymoides noctiflorum. Cam. hort. 109. t. 34.*

*Habitat in Suecia, Germania. ☉*

*Calyces admodum viscerati dentibus longioribus, quam in reliquis.*

virginica.

21. **SILENE** calycibus floris cylindricis villosis, panicula dichotoma.

*Lychnis flore simplici specioso coccineo, foliis oblongis acuminatis adversis, caule viscoso. Gron. virg. 161.*

*Habitat in Virginia. 24*

*Planta polymorpha. Flores facie Lychnis dioicæ albæ, at saturate rubri; Calyces angustiores quam in congeneribus, hirsuti. Stamina cum stylis extra florem prominentia. Caulis nunc simplex digiti longitudine, terminatus floribus in fasciculum collectis, nunc pedalis apice trichomus, pedunculo intermedio semper unifloro. Folia oblonga s. lato-lanceolata, ut in Gomphrena globosa.*

anthirrhina.

22. **SILENE**, foliis lanceolatis subciliatis, pedunculis trifidis, petalis emarginatis, calycibus ovatis. *Roy. lugdb. 447.*

*Silene petalis bilobis coronatis, floribus erectis, foliis subciliatis. Hort. ups. 114.*

*Silene corymbo dichotomo, floribus pedunculatis, ramis alternis erectis, foliis lanceolatis integerrimis. Gron. virg. 50.*

*Viscago americana noctiflora, anthirrhini folio. Dill. elth. 442. t. 313. f. 403.*

*Habitat in Virginia, Carolina. ☉*

rubella.

23. **SILENE** erecta lævis, calycibus subglobosis glabris venosis, corollis inapertis. *Hort. ups. 112.*

*Viscago lusitanica, flore rubello vix conspicuo. Dill. elth. 324. t. 314. f. 406.*

*Habitat in Lusitania, Oriente. ☉*

inaperta.

24. **SILENE** caule dichotomo paniculato, calycibus lævibus, petalis brevissimis emarginatis, foliis glabris lanceolatis.

*Viscago lævis, inaperta flore. Dill. elth. 424. t. 315. f. 407.*

*Habitat in Europa australi. ☉*

*Stamina sæpe 5. ☉ corolla absque limbo.*

portensis.

25. **SILENE** caule dichotomo paniculato, calycibus stri-

statiatis, petalis bifidis, folijs linearibus.

*Habitat in Lusitania. Læfl. ☉*

*Habitus S. inapertæ, at Folia angustiora & Calycæ longitudinaliter striati. Planta spithamæa, brachiata, dichotoma. Folia linearia, acuta, glabra. Pedunculi filiformes, angustissimi, longi, erecti, e dichotomia caulis uniflori. Calyces clavati, longi, purpurascens, striis alternis albis. Stamina longitudine fere corollæ. An Silenes inapertæ sola varietas? Aman. acad. 3. p. 409.*

26. *SILENE erecta lævis, calycibus erectis decangulis creta. sulcatis; petalis bifidis. Hort. upf. 112.\**

*Viscago foliis inferioribus bellidis, superioribus tunicæ, calyce strictiore & turgidiore. Dill. elth. 422. t. 314. f. 404, 405.*

*Lychnis viscosa, foliis inferioribus bellidi similibus, flore minimo carneo s. rubro. Magn. hort. 126.*

*Habitat in Creta. ☉*

*Caulis dichotomo-paniculatus, patens, filiformis: articulis viscidis. Calyces glabri. Flores amænæ purpurei petalis semibifidis, erecto-patulis.*

27. *SILENE petalis bifidis, caule dichotomo, floribus axillaribus sessilibus, folijs glabris. Muscipula.*

*Silene floribus tristylis, folijs lanceolatis sessilibus, capsulis trilocularibus, floribus ternis. Sauv. monsp. 145.*

*Lychnis sylvestris viscosa rubra alterna. Bauh. pin. 205.*

*Lychnis sylvestris. 3. Clus. hist. 1. p. 289.*

*Habitat in Hispania, G. Narbonensi.*

28. *SILENE foliis fasciculatis setaceis: ramorum flo- rentium oppositis. Roy. lugdb. 447.*

*Lychnis sylvestris, plurimis foliolis simul junctis. Bauh. pin. 205.*

*Lychnis sylvestris VIII. Clus. hist. 1. p. 290.*

*Habitat in Pannonia, Austria, Bohemia. 24*

29. *SILENE floribus fasciculatis fastigiatis, folijs superioribus cordatis glabris. Hort. upf. 110.*

*Silene foliis lanceolato-ovatis glabris, floribus terminalibus fasciculatis fastigiatis. Hort. cliff. 172. Roy. lugdb. 446. Sauv. monsp. 153.*

*Lychnis viscosa purpurea latifolia laevis. Bauh. pin. 205.*

*Armerius flos quartus. Dod. pempt. 176.*

*Habitat in Anglia, Gallia. ☉*

*quadrifida.* 30. *SILENE* petalis quadrilobis, caule dichotomo, floribus pedunculatis, foliis glabris recurvis. *Leyf. hal. 383.*

*Cucubalus quadrifidus. Sp. pl. 1. p. 415.*

*Lychnis alpina, foliis angustis reflexis, petalis quadripartitis. Segu. veron. 3. p. 186. t. 5. f. 1.*

*Lychnis saxatilis, linariae folio, glabra, flore albo, petalis tricuspidatis, femine limbo fimbriato. Till. pis. 106.*

*Lychnis viscosa angustifolia major. Bauh. pin. 205.*

*Lychnis sylvestris X. Clus. hist. 1. p. 291.*

*Caryophyllus minimus humilis alter exoticus. Lob. 445.*

*Habitat in Italiae, Austriae montibus. ☉*

*rupestris.* 31. *SILENE* floribus erectis, petalis emarginatis, calycibus teretibus, foliis lanceolatis. *Fl. svec. 367; 390. R. W. goth. 144. \**

*Silene floribus erectis laxè distantibus, caule dichotomo. Fl. lapp. 183.*

*Alfina alpina glabra. Bauh. pin. 251, prodr. 118. Oed. dan. t. 4. Raj. hist. 1003, 1031.*

*Auricula muris alpina glabra. Bauh. hist. 3. p. 360.*

β. *Caryophyllus holosteus alpinus gramineus. Bauh. pin. 210, prodr. 104. Burf. XI. 117.*

*Habitat in montibus aridis Sveciae, Helvetiae. ♂*

*Varietas β. foliis linearibus in umbra nata.*

*Saxifraga.* 32. *SILENE* caulibus unifloris, pedunculis longitudine caulis, foliis glabris, floribus hermaphroditis femineisque.

*Caryophyllus saxifragus. Bauh. pin. 211.*

*Saxifraga antiquorum quibusdam. Bauh. hist. 3. p. 338.*

*Lychnis minor saxifraga. Segu. ver. 431. t. 6. f. 1.*

*Habitat in montibus cretaceis Galliae, Italiae. ♀*

*Radix lignosa, ramosa. Caules plures, digitales, glabri. Folia opposita, linearia, acuta, glabra, per 4 ad 6 oppositiones. Pedunculus terminalis, filiformis longitudine caulis; raro pedunculus alter lateralis. Flos unicus. Calyx clavatus, glaber. Petala bifida limbo calyce brevior, subius rubro.*

23. **SILENE** caulibus subunifloris decumbentibus, foliis lanceolatis tomentosis longitudine calycis. *vallesia.*  
*Lychnis pumila saxatilis alpina*, flore carneo, folio crasso molli tomentoso. *Amæn. acad.* 1. p. 158.  
*Lychnis maritima* pinguis e corfica. *Boec. mus.* 1. 34?  
*cui folia latiora breviora.*  
*Habitat in Alpibus Vallefiacis, Valdensibus. Allioni.*

34. **SILENE** acaulis. *Fl. lapp.* 184. *Fl. fvec.* 368; 387. *acaulis.*  
*Hall. herb.* 373.  
*Lychnis alpina pumila*, folio gramineo. *Bauh. pin.* 206. *Dill. elth.* 206. 1. 167. f. 206. *Oed. dan.* 1. 21.  
*Muscus alpinus*, lychnidis flore. *Bauh. hist.* 3. p. 767.  
*Habitat in Alpibus Lapponicis, Austriacis, Helveticis, pyrenæis.* 24.  
*Herba Bryo similis. Petala emarginata.*

## S T E L L A R I A.

1. **STELLARIA** foliis cordatis petiolatis, panicula pedunculis ramosis. *nemorum.*  
*dunculis ramosis. Fl. fvec.* 2. n. 391. \*  
*Alfne altissima nemorum. Bauh. pin.* 250.  
*Alfne montana hederacea maxima. Col. ecphr.* 1. p. 289. 1. 290.  
 β. *Alfne montana latifolia*, flore laciniato. *Bauh. pin.* 250. *Rupp. jen.* 109.  
*Habitat in Europæ nemoribus.*
2. **STELLARIA** foliis ovatis sessilibus, caule dichotomo, floribus solitariis, pedunculis fructiferis reflexis. *dichotoma.*  
*Habitat in Alpibus Helveticis, Sibiricis.* ☉  
*Rami divaricati, dichotomi. Pedunculi fructiferi toti reflexi.*
3. **STELLARIA** foliis lanceolatis serrulatis, petalis quinquepartitis. *radiana.*  
*Alfne saxatilis*, angusto & oblongo salicis folio, flore albo tenuissime laciniato. *Amn. ruth.* 83. 1. 10.  
*Habitat in Sibiria.*
4. **STELLARIA** foliis lanceolatis serrulatis, petalis bifidis. *Holosteæ.*  
*Alfne foliis lanceolatis. Hort. cliff.* 172. *Fl. fvec.* 371; 392.  
*Caryophyllus holosteus arvensis*, flore majore. *Bauh. pin.* 210.



- Gramen. *Fuchf. hist.* 136.  
*Habitat in Europæ nemoribus.*
- graminea. 5. **STELLARIA** foliis linearibus integerrimis, floribus paniculatis.  
 Ailne foliis linearibus. *Fl. lapp.* 189; 393. *Fl. suec.* 372. *Hort. cliff.* 172. *Roy. lugdb.* 449.  
 Caryophyllus arvensis glaber, flore minore. *Bauh. pin.* 210.  
 Gramen floridum minus. *Tabern. ic.* 232.  
 β. Ailne, folio gramineo angustiore, palustris. *Dill. app.* 69.
7. **Caryophyllus holosteus alpinus angustifolius.** *Bauh. pin.* 210. *prodr.* 104. n. 6. *Burf. XI.* 2.  
*Habitat in siccis juniperetis sepibus testis Europæ.* 4
- Cerastoides. 6. **STELLARIA** foliis oblongis, pedunculis subbifloris.  
*Fl. suec.* 2. n. 394. \*  
*Habitat in Alpibus Lapponicis.*
- biflora. 7. **STELLARIA** foliis subulatis, scapis subbifloris, petalis emarginatis, germinibus oblongis, calycibus striatis. *Fl. suec.* 2. n. 395. \*  
 Sagina ramis erectis bifloris. *Fl. lapp.* 158. *Oed. dan.* t. 12.  
 Mœhringia scapis bifloris. *Fl. suec.* 1. n. 316. *Amæn. acad.* 2 p. 264.  
 Ailne polygonoides, foliis brevibus, flore albo. *Segu. ver.* 3. p. 177. t. 4. f. 1.  
 Arenaria foliis subulatis, calycibus striatis, germinibus oblongis, floribus corymbosis. *Amæn. acad.* 1. p. 158. \*  
*Habitat in Alpibus Lapponicis.* 4  
 Planta parva, facie Saginæ. Folia radicalia subulata, in acervum congesta. Caules digitales, filiformes, maxima parte nudi: superne bifidi in 2. pedunculos subjecto pari foliorum: altero pedunculo, in medio, foliolis duobus.
- Arenaria. 8. **STELLARIA** caule erecto bifido, ramis alternis, petalis emarginatis. *Sl. pl.* 1. p. 11. 26.  
*Habitat in Hispania.* | ☉  
 Radix fibrosa. Caulis erectus, teres, spithameus, pilis patulis pubescens & subglutinosus. Rami alterni, longitudine fere caulis. Folia sessilia, spatulata, apice

*pice ovato, basin versus linearia angustiora, subtus  
 & margine pilosa, supra vero vix manifeste. Pri-  
 mus flos e dichotomia caulis: reliqui ex alis ramorum  
 alterni. Omnes majusculi. Pedunculi recti, foliis  
 longiores. Calyx viscosus uti tota planta, apicibus  
 purpurascens, uti etiam Folia floralia. Corolla cam-  
 panulata, calyce longior, alba, obtusissima petalis  
 vix manifeste emarginatis. Stamina alba, longitu-  
 dine calycis. Antheræ cerulescentes. Germen glo-  
 bosum. Styli tres, longitudine flaminum.*

## ARENARIA.

1. **ARENARIA** foliis ovatis acutis carnosiss. *Fl. fvec. peploides.*  
 375; 396. *Fl. lapp. 149. It. al. 151.*  
*Alpine litoralis, foliis portulacæ. Bauh. pin. 251. Læf.*  
*pruff. 12. t. 2.*  
*Telephium maritimum, portulacæ folio. Buxb. act. 3.*  
*p. 271.*  
*Habitat ad litora maris Europæi borealia. 24*
2. **ARENARIA** foliis ovatis carinatis recurvis quadri- *tetraquetra.*  
 fariam imbricatis. *Bauh. pin. 211. prodr. 125. Burf.*  
*XI. 135.*  
*Habitat in Pyrenæis. 24*  
*Herba omnino Gypsophila aggregatæ, sed Pedunculus*  
*uniflorus, terminalis.*
3. **ARENARIA** foliis ovatis obtusis, pedunculo laterali *lateriflora.*  
 bifloro.  
*Habitat in Sibiria.*  
*Caulis digitalis, filiformis, simplicissimus. Folia ovata,*  
*obtusa, subpetiolata, glabra. Pedunculus axillaris u-*  
*nicus, longus, bifidus. Corolla calyce longior.*
4. **ARENARIA** foliis ovatis acutis petiolatis nervosis. *trinervia.*  
*Hort. cliff. 173. Fl. fvec. 374; 397.\* Roy. lugdb. 471.*  
*Alpine plantaginis folio. Bauh. hist. 3. p. 364.*  
*Habitat in Europæ sylvis. 0*
5. **ARENARIA** foliis ovatis nervosis sessilibus acutis, *multicaulis.*  
 corollis calyce majoribus. *Aman. acad. 1. p. 162. n.*  
*155.*  
*Alpine foliis lanceolatis, petalis integris calyce maio-*  
*ribus. Hall. helv. 390. t. 6. f. 3.*  
*Habitat in Alpibus Helvetiæ, Pyrenæis.*

serpyllifolia

6. *ARENARIA* foliis subovatis acutis sessilibus, corollis calyce brevioribus. *Fl. suec.* 373; 398. *Hort. cliff.* 173. *Roy. lugdb.* 451.  
*Alfina minor multicaulis. Bauh. pin.* 251.  
*Alfina minor. Fuchf. hist.* 23.  
*Habitat in Europæ sylvis glareosis. ☉*

montana.

7. *ARENARIA* foliis lanceolato-linearibus scabris; caulibus sterilibus longissimis procumbentibus. *Amxn. acad.* 4. p. 272. \*  
*Alfina foliis linearibus acuminatis, petalis florum integris calyce duplo longioribus. Monn. opf.* 127.  
*Habitat in Galliæ australis montibus.*

rubra.

8. *ARENARIA* foliis filiformibus, stipulis membranaceis vaginantibus. *Fl. suec.* 376; 399.

campestris.

- α. *Alfina*, *Spergula* facie, minor s. *Spergula* minor, subcæruleo flore. *Bauh. pin.* 251. *prodr.* 119.  
*Polygonum foliis gramineis, spergulae capitulis. Læf. pruss.* 203. t. 63.

marina.

- β. *Arenaria* foliis linearibus longitudine internodiorum. *Hort. cliff.* 173. *Gron. virg.* 161. *Roy. lugdb.* 451.  
*Spergula marina nostras. Ray. hist.* 1034.  
*Alfina spergulae facie media. Bauh. pin.* 251.  
*Habitat α in Europæ arenosis collibus, β. in litoribus marinis. ☉*  
*Stamina in campestri certe 10. sunt: in maritima prope Aboam stamina tantum 5 observavit D. Kalm.*

media.

9. *ARENARIA* foliis linearibus carnosiss, stipulis membranaceis.  
*Arenaria teretifolia verna, flore albo, semine limbo foliaceo cincto. Rupp. jen.* 2. p. 113.  
*Spergula annua, semine foliaceo nigro circulo membranaceo albo cincto. Dill. giff.* 46. *E. N. C. cent.* 5. f. 6. p. 275. t. 4. *Morif. blas.* 228.  
*Alfina Spergulae facie minima, seminibus marginatis. T. Vaill. parif.* 8.  
*Habitat in Germania, Gallia.*  
*Simillima A. rubrae maritimæ, sed minus ramosa, caulibus magis adscendentibus, tota pubescenti-viscosa. Folia carnosissima, linearia, acutiuscula, supra depressiuscula, subtus convexa. Flores albi, petalis calyce vix majoribus. Semina margine membranaceo cincta. Planta vernalis ☉ filia spuria A. rubrae, ut fere*

*varietas, quod imprimis ex stipulis scariofo-membranaceis constat.*

10. ARENARIA foliis semicylindricis carnosis obtusis, *bavaria.*  
petalis lanceolatis, pedunculis terminalibus subbinatis. *Amæn. acad. 4. p. 315. \**

Alfina alpina, foliis teretibus obtusis, flore albo. *Segu. veron. 1. p. 429.*

Saxifraga bavarica. *Raj. hist. 1033.*

Habitat in Bavaria, Baldo. 24

11. ARENARIA foliis subulatis, caulibus paniculatis, axillis calycum foliolis ovatis.

Arenaria foliis subulatis, calycinis laciniis membrana ipsis latiore auctis obtusis. *Guett. stamp. 2. p. 281.*

Spergula foliis aciformibus densis, ramis ramosis subnudis. *Sauv. monsp. 45.*

Alfina glabra, tenuissimis foliis, floribus albis. *Herm. par. 12. t. 12.*

Alfina saxatilis & multiflora, capillaceo folio. *Vaill. paris. 7. t. 2. f. 3. Hall. opusc. 113.*

Alfina saxatilis, laricis folio, minor & minori flore. *T. Magn. monsp. 11. t. 11.*

Alfina caryophylloide tenuifolia, flore albo punctato. *Pluk. alm. 22. t. 7. f. 3.*

Anthyllis lychnitis annua. *Barr. ic. 580.*

Habitat in Germania, Helvetia, Gallia, Sibiria. 24

12. ARENARIA foliis subulatis, caule paniculato, pedunculis simplicibus, capsulis erectis, petalis calyce brevioribus.

Arenaria foliis subulatis, calycinis laciniis membrana ipsis angustiore auctis acutissimis. *Guett. stamp. 2. p. 282.*

Alfina tenuifolia. *Bauh. pin. 3. p. 364. Vaill. paris. 7. t. 3. f. 1. Segu. ver. 418. t. 6. f. 2.*

Habitat in Helvetia, Gallia, Anglia, Italia. 24

Habet in flore stamina decem.

13. ARENARIA foliis setaceis, caule superne nudius. laricifolia, sculo, calycibus subhirsutis.

Alfina foliis fasciculatis, petiolis simplicibus, calyce hirsuto. *Hall. helv. 388.*

Alfina alpina, junceo folio. *Bauh. pin. 251. prodr. 118.*

Lychn

*Lychnoides*, juniperi folio, perennis. *Vaill. parif.* 121.

*Habitat in montosis Helvetiæ, Genevæ, Parisiorum, Monspeli.* 24

striata.

14. *ARENARIA* foliis linearibus erectis oppressis, calycibus oblongis striatis. *Amæn. acad.* 4 p. 315. \*

*Auricula muris pulchro flore, folio tenuissimo. Bauh. pin.* 3. p. 361.

*Caryophyllus faxatilis, polygoni minoris folio & facie. Burf. XI.* 129.

*Habitat in Alpibus Austriæ & Vallis augustæ.* 24

elliptica.

15. *ARENARIA* foliis ovatis ciliatis acutis.

*Alsine, serpylli folio, multicaulis & multiflora. Segu. ver.* 1. p. 421. t. 5. f. 2.

*Habitat in Alpibus, Ræticiis, monte Abrahami.*

hispida.

16. *ARENARIA* foliis subulatis subtus hispidis.

*Habitat Monspeli in monte calcaris.*

*Habitus Spergulæ. Caules simplices, pilis raris adspersi. Folia opposita, subulata, plana, subtus pilis raris hispida. Panicula primum dichotoma, mox alternis pedunculis ramosa.*

grandiflora.

17. *ARENARIA* foliis subulatis strictis: radicalibus confertis, caulibus unifloris.

*Alsine uniflora & grandiflora, foliis acuminatis, petalis integris. Allion. pedem.* 49. t. 10. f. 1.

*Habitat in Alpibus Valdensium, Cenisii.* 24

### CHERLERIA.

Sedoides.

1. *CHERLERIA. Hall. helv.* 391. t. 6. *opusc.* 300. t. 1. f. 3. *Segu. ver.* 3. p. 180. t. 4. f. 3.

*Lychnis alpina, muscosis foliis densius stipatis, floribus parvis, calyce duriore. Pluk. alm.* 233. t. 42. f. 8.

*Sedum montanum perpusillum luteum. Morif. hist.* 3. p. 471. f. 12. t. 6. f. 14.

*Sedum montanum perpusillum, luteolis floribus. Park. theatr.* 738.

*Habitat in Alpibus Helvetiæ, Valesiæ, Gotthardo.* 24

### GARIDELLA.

*Nigella-frum.*

1. *GARIDELLA. Hort. cliff.* 170. *Hort. ups.* 108. *Roy. lugdb.* 481.

*Garidella* foliis tenuissime divis. *Garid. prov.* 203.  
t. 39.

*Nigellastrum* raris & fœniculaceis foliis. *Magn. hort.*  
143. t. 143.

*Nigella cretica*, folio fœniculi. C. B. *Morif. hist.* 3.  
p. 516: f. 12. t. 18. f. 6.

*Habitat in Galloprovincia.* ☉

## MALPIGHIA.

1. *MALPIGHIA* foliis ovatis integerrimis glabris, pedunculis umbellatis. *Hort. cliff.* 169. *Hort. upf.* 108.  
*Roy. lugdb.* 459. *Mill. dict.* t. 181. f. 2.

*Malpighia fruticosa erecta*, foliis nitidis ovatis acuminatis, floribus umbellatis, ramulis gracilibus. *Brown. jam.* 230.

*Cerasus jamaicensis*, fructu tetrapyreno. *Comm. hort.*  
1. p. 145. t. 75.

*Arbor baccifera*, folio subrotundo, fructu cerasino sulcato rubro polypyreno: ossiculis canaliculatis. *Sloan. jam.* 172. *hist.* 2. p. 106. t. 207. f. 2. *Raj. dendr.* 74.

*Habitat in Jamaica, Brasilia, Surinamo, Curassao:* ☿

2. *MALPIGHIA* foliis ovatis integerrimis glabris, pedunculis unifloris.

*Malpighia* foliis ovatis integerrimis glabris. *Hort. upf.* 108.

*Malpighia fruticosa erecta*, ramulis gracilibus patetibus, floribus solitariis. *Brown. jam.* 230.

*Malpighia mali punici facie.* *Pluk. gen.* 46. t. 166. f. 2.

*Cerasus americana*, myrti conjugatis foliis, fructu acerbo tetrapyreno. *Pluk. alm.* 94. t. 157. f. 7.

*Habitat in America meridionali.* ☿

3. *MALPIGHIA* foliis lanceolatis integerrimis glabris, nitida, spicis lateralibus.

*Malpighia* foliis oblongo-ovatis utrinque nitidis, racemis axillaribus. *Jacq. amer.* 21.

*Habitat in America.* ☿

4. *MALPIGHIA* foliis oblongo-ovatis: setis decumbentibus rigidis, pedunculis unioris aggregatis. *Mill. dict.* t. 181. f. 1.

*Malpighia latifolia*, folio subtus spinoso. *Plum. gen.* 46. t. 167. f. 1.

*Linnaei Spec. plant. pars I.*

Qq

Me-

*Mespilus americana*, folio lato subtus spinoso, fructu rubro. *Tournef. inst.* 642

Arbor baccifera, folio oblongo subtilissimis spinis subtus obsito, fructu cerasino fulcato polypyreno: of-  
ficulis canaliculatis. *Sloan. jam.* 172. *hist.* 2. p. 106.  
t. 207. f. 3. *Raj. dendr.* 74.

*Habitat in America calidiore.* ♀

Folia ovato-oblonga, integerrima, subpetiolata, ad-  
pressa, imprimis subtus, setis rigidis adpressis pungentius  
geminis: altera anteriore. Stipulae utrinque 2. Flo-  
res axillares utrinque 4, pedunculis unifloris, geni-  
culo instructis.

angustifolia. 5. **MALPIGHIA** foliis linear-lanceolatis: setis utrin-  
que decumbentibus rigidis, pedunculis umbellatis  
*Mill. dict.* *Jacq. amer.* 21.

*Malpighia angustifolia*, folio subtus spinoso. *Plur.*  
*gen.* 46.

*Habitat in America meridionali.* ♀

crassifolia. 6. **MALPIGHIA** foliis ovatis integerrimis subtus to-  
mentosis, racemis terminalibus.

*Malpighia arborea*, floribus spicatis, foliis ovatis ac-  
cuminatis. *Brown. jam.* 230.

*Malpighia arborea*, foliis subrotundis alternis inferne  
sublanuginosis, spicis crassis compositis termina-  
libus. *Brown. jam.* 231.

*Malpighia latifolia*, cortice sanguineo. *Barr. equi.*  
72 ?

*Tilia* affinis *laurifolia*, arbuti floribus albis racemosis  
odoratis, fructu pentagono. *Sloan. jam.* 136. *hist.* 2.  
p. 20. t. 163. f. 1.

*Habitat in America calidiore.* ♀

Arbor, foliis ovatis palmaribus crassis subpetiolatis in-  
tegerrimis, supra pubescentibus, subtus tomentosis, se-  
pius alternis. Racemi terminales longi tomentosi.

verbascifolia. 7. **MALPIGHIA** foliis lanceolato-ovatis tomentosis in-  
tegerrimis, racemis terminalibus.

*Malpighia humilis*, verbasci folio & facie, caule cras-  
sissimo. *Barr. equin.* 71.

Baccifera arbor calyculata, foliis laurinis, fructu ra-  
cemoso esculento subrotundo monoppyreno pallide  
luteo. *Sloan. hist.* 2. t. 198. f. 2.

*Habitat in America calidiore.* ♀

*Folia lanceolato-ovata, in petiolos desinentia, pedalis, villosa, subtus densissimo tomento vestita. Racemi terminales, longi, villosi.*

8. MALPIGHIA foliis lanceolatis dentato spinosis sub-aquifolia.  
tus hispidis.

Malpighia angustis & acuminatis aquifolii foliis. *Plum.*  
*gen. 46. ic. 168. f. 1.*

*Habitat in America calidiore. h*

9. MALPIGHIA foliis subovatis dentato-spinosis. coccigera.

Malpighia humilis, ilicis cocciglandiferæ foliis. *Plum.*  
*gen. 46. ic. 168. f. 2.*

*Habitat in America calidiore. h*

## BANISTERIA.

1. BANISTERIA foliis sinuato-angulosis. angulosa.

Acer scandens, folio anguloso. *Plum. spec. 18.*

Clematis anguloso folio, aceris fructu. *Plum. amer.*

*77. t. 92. Raj. suppl. 328.*

*Habitat in America calidiore. h*

2. BANISTERIA foliis ovatis, spicis lateralibus, semi-purpurea.  
nibus erectis.

Banisteria foliis pinnatis: foliolis ovatis, spicis lateralibus, seminibus erectis. *Mill. dict.*

Acer scandens, pseudoacaciæ folio, flore purpurascente. *Plum. spec. 18. ic. 15.*

*Habitat in America meridionali. h*

3. BANISTERIA foliis ovato-oblongis rigidis, racemis terminalibus. laurifolia. *Mill. dict.*

Acer scandens, foliis laurinis. *Sloan. jam. 137. hist. 2. p. 26. Raj. dendr. 94.*

*Habitat in Jamaica. h*

4. BANISTERIA foliis ovato-oblongis acuminatis, benghalensis.  
racemis lateralibus, seminibus patentibus. *Fl. zeyl. 176.*

Acer scandens, foliis citri, flore cæruleo spicato. *Plum. spec. 18. ic. 14.*

Acer benghalensis, laurinis foliis; fructu tergemino.

*Pluk. alm. 7. t. 3. f. 1.*

*Habitat in Indiis. h*



- dichotoma. 5. **BANISTERIA** foliis ovatis, racemis dichotomis  
*Banisteria* foliis ovatis, ramis dichotomis, seminibus  
 extrorsum tenuioribus, introrsum lacinulam emittentibus. *Hort. cliff.* 169. \*  
*Acer* scandens, convolvuli folio, flore ex aureo cinereo. *Plum. spec.* 18. ic. 13.  
*Habitat in America calidiore.* ♀
- fulgens. 6. **BANISTERIA** foliis subovatis subtus tomentosis,  
 racemis brachiatis, pedunculis umbellatis.  
*Banisteria* foliis ovatis, ramis ramosis, seminibus extrorsum tenuioribus, introrsum lacinulam emittentibus. *Hort. cliff.* 169. \*  
*Banisteria* foliis orbiculatis, petiolis biglandulis, seminibus unialatis rugosis, racemis subumbellatis alaribus. *Brown. jam.* 231.  
*Acer* scandens minus, apocyni facie, folio subrotundo. *Sloan. jam.* 138. *hist.* 2. p. 27. t. 162. f. 2. *Kandendr.* 94.  
*Triopteris americana* scandens, fructu fulgente majore aureo. *Pluk. mant.* 185.  
*Habitat in America.* ♀
- brachiata. 7. **BANISTERIA** foliis subovatis, ramis brachiatis, seminibus introrsum angustioribus.  
*Banisteria* foliis subovatis, ramis ramosis, seminibus introrsum tenuioribus nullam lacinulam emittentibus. *Hort. cliff.* 169. \*  
*Banisteria* foliis ovatis acuminatis, floribus laxo spicatis, ramis diffusis scandentibus. *Mill. dist.*  
*Habitat in America.* ♀

## TRIOPTERIS.

- Jamaicensis. 1. **TRIOPTERIS.** *Hort. cliff.* 169. \*  
*Acer* scandens tricoccus, foliis citri, flore luteo minore. *Plum. ic.* 9. t. 16?  
*Hiræa reclinata.* *Jacq. amer.* 21?  
*Habitat in America calidiore.* ♀

## ERYTHROXYLON.

- areolatum. 1. **ERYTHROXYLON.** *Amæn. acad.* 5. p. 397. \*  
*Erythroxylon* foliis ellipticis lineis binis longitudinalibus

libus subtus notatis, fasciculis florum sparsis. *Brown.*

*jam.* 278. t. 38. f. 2.

*Habitat in Jamaica.* ♀

# PENTAGYNIA.

## A V E R R H O A.

1. AVERRHOA caudice nudo fructificante, pomis oblongis obtusangulis. *Fl. Zeyl.* 177. \* Bilimbi.  
*Blinbungum teres. Rumph. amb.* 1. p. 118. t. 36.  
*Bilimbi. Rheed. mal.* 3. p. 55. t. 45. 46.  
*Habitat in India.* ♀
2. AVERRHOA axillis foliorum fructificantibus, pomis oblongis acutangulis. *Fl. Zeyl.* 178. \* Carambola.  
*Mala goënsia, fructu octangulati pomi vulgari magnitudine. Bauh. pin.* 433.  
*Prunum stellatum. Rumph. amb.* 1. p. 115. t. 35.  
*Tamara Tonga s. Carambolas. Rheed. mal.* 3. p. 51. t. 43. 44.  
*Habitat in India.* ♀
3. AVERRHOA ramis nudis fructificantibus, pomis subrotundis. *Fl. Zeyl.* 179. acida.  
*Neli-poli. Rheed. mal.* 3. p. 57. t. 47. 48. *Raj. hist.* 1450.  
*Cheramela. Rumph. amb.* 7. p. 34. t. 33. f. 2.  
*Habitat in India.* ♀

## S P O N D I A S.

1. SPONDIAS petiolis communibus compressis. *Læf.* purpurea.  
*it.* 209.  
*Spondias diffusa, foliis plurimis minoribus pinnatis, penna compressa fulcata, floribus præcocibus. Brown. jam.* 228.  
*Myrobalanus minor, folio fraxini alato, fructu purpureo officulo magno fibroso. Sloan. jam.* 182. *hist.* 2. p. 126. t. 219. f. 3, 4, 5. *Raj. dendr.* 43.  
*Habitat in India occidentali.* ♀
2. SPONDIAS foliolis nitidis. lutea.  
*Spondias. Hort. cliff.* 484. *Hobo. Læf. it.* 209. \*  
*Spondias foliolis plurimis pinnatis ovatis, racemisternalibus, cortice interno rubente. Brown. jam.* 229.  

Q q 3.      Mon-

Monbin arbor, folio fraxini, flore luteo racemoso.  
*Num. gen.* 44.

Myrobalanus folio fraxini alato, fructu luteo, officulo magno fibroso. *Sloan. jam.* 181. *list.* 2. p. 125. t. 219. f. 1. 2. *Raj. dendr.* 43.

*Prunus americana*, *Mer. surin.* 13. t. 13. bona.

*Prunus brasiliensis*, fructu racemoso ligno intus pro officulo. *Raj. hist.* 1154.

Acaja & Ibametara. *Marcgr. bras.* 129.

3. *Spondias* foliis paucioribus pinnatis ovatis nitidis, *Brown. jam.* 229.

*Habitat in America meridionali.* §

## COTYLEDON.

orbiculata. 1. **COTYLEDON** foliis subrotundis planis integerrimis. *Hort. cliff.* 276. *Roy. lugdb.* 454.

*Cotyledon africana* frutescens incanum, orbiculatis foliis. *Herm. lugdb.* 349. t. 551. *Mori. hist.* 3. p. 474. f. 12. t. 7. f. 39.

*Habitat ad Cap. b. Spei.* §

spuria. 2. **COTYLEDON** foliis alternis spatulatis carnosiss integerrimis.

*Cotyledon africana* frutescens, folio longo & angusto, flore flavescente. *Comm. rar.* 23. t. 23. *Burm. afr. t.* 18, 19. f. 1. & t. 22. f. 1.

*Cotyledon africana*, foliis depressis cruciatis. *Walt. hort.* 16.

*Habitat ad Cap. b. Spei.* §

Simillima *C. orbiculata*; differt saltem foliis oblongis & digito similibus, sed, quod præcipuum, alternis. Fructificatio adeo *C. orbiculata*, ut de matre dubium esse nequeat.

hemisphærica. 3. **COTYLEDON** foliis semiglobosis. *Hort. cliff.* 176. *Roy. lugdb.* 454.

*Cotyledon capensis*, folio semiglobato. *Dill. elth.* 118. t. 95. f. 111.

*Habitat in Æthiopia.* §

ferrata. 4. **COTYLEDON** foliis ovalibus crenatis, caule spicato.

*Cotyledon* foliis radicalibus lanceolatis crenatis: caulinis subulatis, *Hort. cliff.* 497. *Roy. lugdb.* 454.

*Coty-*

*Cotyledon cretica*, folio oblongo fimbriato. *Dill. elth.*

113. t. 95. f. 112.

*Habitat in Creta, Sioiria.*

5. *COTYLEDON* foliis oblongis spinoso-mucronatis, spinosa, caule spicato.

*Habitat in Sibiria. D. Gmelin.*

*Habitus speciei 3. Folia integra spina setacea terminata.*

*Flores quinquepartiti.*

6. *COTYLEDON* foliis cucullato-peltatis serrato-dentatis alternis, caule ramoso, floribus erectis. *Gron. orient.* 141.

*Cotyledon* foliis peltatis. *Vir. cliff.* 39. *Roy. lugdb.* 454.

*Cotyledon* flore luteo, radice repente. *Dod. mem.* 265. *repens.* t. 73.

*Cotyledon* radice tuberosa longa repente. *Morif. præl.* 257. *Raj. hist.* 1878.

*Cotyledon* luteum umbilicatum spicatum, radice repente majus. *Morif. hist.* 3. p. 471.

*Umbilicus repens. Cam. epit.* 858.

- β. *Cotyledon vera*, radice tuberosa. *Bauh. hist.* 5. p. 683. *tuberosa.*

*Cotyledon major. Bauh. pin.* 285. *Raj. hist.* 1878.

*Cotyledon umbilicus veneris. Clus. hist.* 2. p. 63.

*Sedum* luteum murale spicatum, folio umbilicato rotundo. *Morif. hist.* 3. p. 470. f. 12. t. 10. f. 4.

*Habitat in Lusitania, Hispania, Anglia, Judæa.* 24

7. *COTYLEDON* foliis trifidis, floribus quadrifidis. *laciniata.*

*Hort. cliff.* 175. *Roy. lugdb.* 454.

*Cotyledon* asra, folio crasso lato laciniato, flosculo aureo. *Boerh. lugdb.* 1. p. 288. t. 288.

*Telephium africanum*, angustiori folio, flore aurantiaco. *Pluk. alm.* 362. t. 228. f. 3.

*Telephium indicum. Bont. jav.* 132.

*Planta anatis. Rumph. amb.* 5. p. 275. t. 95.

*Habitat in Ægypto, India.* 5

8. *COTYLEDON* foliis oblongis subteretibus, floribus fasciculatis. *hispanica.* *Læfl. it.* t. 1.

*Cotyledon maritima*, fedi folio, flore carneo, fibrosa radice. *Tournef. inst.* 90.

*Cotyledon hispanica*, fedi teretifolio folio, flore umbellato rubro, fibrosa radice. *Minuart. sched.* 3.

Qq 4

Co.

Cotyledon palustris. sedi folio, floribus rubris longioribus. *Shaw. afr. 177.*

β. Cotyledon palustris, sedi folio, floribus luteis brevioribus. *Shaw. afr. 178. f. 178.*

*Habitat in Africa, Oriente, Hispania. ♂*

*Radix fibrosa. Caulis simplex, teres, erectiusculus, digitalis. Folia oblonga, teretiuscula, obtusiuscula, sessilia, subpilosa; supra planiuscula rufo-punctata. Flores fasciculati corymbo terminali. Corollæ infundibuliformes: extus rufescentes, subpilosæ. Limbo quinquepartito, plano, acuto, supra incarnato striis tribus purpureis apice purpureo. Filamenta crassiuscula, prominula. Antheræ minime. Calyx brevis. Facies herbæ est Sedi albi.*

## S E D U M.

\* *Planifolia.*

verticillatum. 1. SEDUM foliis quaternis. *Amæn. acad. 2. p. 253. t. 4. f. 14. \**

Sedum aparines facie, tauromenitanum. *Raj. extr. 283. Morif. hist. 3. p. 473.*

*Habitat in Europa maxime australi, & Sibiria. ♀*

Telephium. 2. SEDUM foliis planiusculis serratis, corymbo folioso, caule erecto.

Sedum foliis planiusculis patentibus serratis, corymbo terminali. *Hort. cliff. 176. Fl. suec. 386; 400. Mus. med. 217. Roy. ludb. 456.*

Telephium vulgare. *Bauh. pin. 287.*

album. Telephium album. *Fuchf. hist. 800.*

purpureum. β. Telephium purpureum majus. *Bauh. pin. 287.*

γ. Telephium purpureum minus. *Bauh. pin. 287. prodr. 133.*

δ. Telephium latifolium peregrinum. *Bauh. pin. 287.*

maximum. ε. Sedum foliis ovalibus serratis, umbellis in alis foliorum. *Hall. helv. 35.*

Telephium hispanicum 1. *Clusf. hist. 2. p. 66.*

Anacampseros maxima. *Bauh. hist. 3. p. 682.*

*Habitat in Europæ siccissimis. ♀*

Anacampseros. 3. SEDUM foliis cuneiformibus integerrimis, caulibus decumbentibus, floribus corymbosis.

Sedum foliis spatulatis indivisis, umbella terminali. *Sauv. monsp. 77.*

Te-

*Telephium repens*, folio deciduo. *Bauh. pin.* 287.

*Telephium* VI. *Cepæa plancii*. *Clus. hist.* 2. p. 67.

*Anacampteros minor*, rotundiore folio, sempervirens.

*Bauh. hist.* 3. p. 682.

*Habitat in* - - - 24

*Folia hyeme decidua.*

4. *SEDUM* foliis lanceolatis ferratis planis, caule ere- Alzoon.  
cto, cyma sessili terminali.

*Sedum* foliis planis ferratis, corymbo folioso. *Hort.*  
*upf.* 116.

*Anacampteros* flore flavo. *Amm. ruth.* 96. t. 11.

*Habitat in* Sibiria. 5

5. *SEDUM* foliis cuneiformibus concavis subdentatis hybridum.  
aggregatis, ramis repentibus, cyma terminali. *Hort.*  
*upf.* 116.

*Anacampteros minor repens*, flore purpureo. *Buxb.*  
*cent.* 1. p. 33. t. 61. f. 2.

*Sedum minus repens*, chamædryos foliis, e singulis  
foliorum alis radices agens. *Amm. ruth.* 93.

*Habitat in* Tataria ad radices montium uralensium, Tra-  
pezunti. 5

6. *SEDUM* foliis radicalibus fasciculatis spatulato-lan- libanoti-  
ceolatis, caule subnudo simplicissimo. cum.

*Habitat in* Palæstina. *Hasselquist.* 24

*Facies sempervivi montani.* Folia Radicalia fascicula-  
ta, spatulato-lanceolata, acuta, glabra, integerrima.  
Caulis ad latus radices, adscendens, semipedalis,  
simplicissimus, inferne adpersus foliis linearibus, a-  
cutis. Racemus terminalis, longitudine caulis: Pe-  
dicellis sparsis, sæpius bifloris. Capsulæ non diver-  
gentes, calyce cinctæ.

7. *SEDUM* foliis planiusculis angulatis, floribus late stellatum.  
ralibus sessilibus solitariis. *Hort. cliff.* 176. *Roy. lugdb.*  
456.

*Sedum echinatum vel stellatum*, flore albo. *Bauh. hist.*  
3. p. 680.

*Sempervivum tertium.* *Col. phytob.* 32. t. 11.

*Cotyledon stellata.* *Bauh. pin.* 295.

*Habitat in* Italia, Gallia, Helvetia.

8. *SEDUM* foliis planis, caule ramoso, floribus pani- *Cepæa.*  
cu-  
Q q 5

culatis. *Hort. cliff.* 176. \* *Roy. lugdb.* 456. *Sauv. monsp.* 76.

*Sedum annuum album*, oblongo portulacæ minoris folio. *Morif. hist.* 3. p. 473. f. 12. t. 7. f. 37.

*Cepæa. Bauh. pin.* 288. *Clus. hist.* 2. p. 68.

*Habitat* Monspelii, Genevæ, Halæ ad lacum salsum. ☉

\* \* *Teretifolia.*

dasyphyllum.

9. *SEDUM* foliis oppositis ovatis obtusis carnosis, caule infirmo, floribus sparsis.

*Sedum* foliis cordato-ovatis compressis sæpius oppositis, floribus sparsis. *Sauv. monsp.* 8.

*Sedum* foliis semiglobosis subovatis sessilibus quadrifariam imbricatis. *Wachend. ultr.* 391.

*Sedum parvum*, folio circinato, flore albo. *Bauh. hist.* 3. p. 691.

*Sedum minus*, folio circinato. *Bauh. pin.* 283. *Morif. hist.* 3. p. 473. f. 12. t. 7. f. 35.

*Aizoon dasyphyllum. Dalech. hist.* 1133.

*Habitat* in Helvetia, Lusitania, Hispania, Italia. ☉

reflexum.

10. *SEDUM* foliis subulatis sparsis basi solutis: inferioribus recurvatis. *Fl. suec.* 2. p. 1296.

*Sedum minus luteum*, folio acuto. *Bauh. pin.* 283.

β. *Sedum minus luteum* ramulis reflexis. *Bauh. pin.* 283.

*Habitat* in Europa ad radices montium. 24.

*Folia viridia*, tenuius subulata, inferiora recurvata uti hamus.

rupestre.

11. *SEDUM* foliis subulatis quinquefariam confertis basi solutis, floribus cymosis. *Hort. cliff.* 176. l. gotl. 178. *Fl. suec.* 388. *Roy. lugdb.* 466. *Gort. gelr.* 268.

*Sedum rupestre repens*, foliis compressis. *Dill. elth.* 343. t. 256. f. 332.

*Habitat* ad radices montium Europæ. 24.

*Folia glauca*, crassius subulata, quinquefariam imbricata.

hispanicum.

12. *SEDUM* foliis teretibus acutis: radicalibus fasciculatis, cyma pubescente. *Amoen. acad.* 4. p. 273. \*

*Sedum hispanicum*, folio glauco acuto, flore albido. *Dill. elth.* 342. t. 256. f. 332.

*Habitat* in Hispania. 24.

13. SE-

13. **SEDUM** foliis oblongis obtusis teretiusculis sessili- album.  
bus patentibus, cyma ramosa. *Hort. cliff.* 177. *Fl.*  
*svec.* 387; 402. *Roy. lugdb.* 457. *Hall. helv.* 398.  
*Sedum minus teretifolium album.* *Bauh. pin.* 283.  
*Vermicularis* f. *Illecebra major.* *Lob. hist.* 205.  
*Habitat in Europæ petris.*
14. **SEDUM** foliis fusiformibus subdepressis: infimis rubens.  
quaternis, cyma subquadrifida, floribus pentandris,  
staminibus reflexis.  
*Sedum foliis oblonge ovatis, floribus in summum cau-*  
*lem congestis, calyce rubente.* *Hall. helv.* 396.  
*Sedum minimum montanum non acre, flore purpura-*  
*sciente parvo, semine stellato.* *Raj. hist.* 692.  
*Sedum saxatile, atrorubentibus floribus.* *Bauh. pin.* 283.  
*prodr.* 132. *Scheuch. it.* 1. p. 48. t. 6. f. 3, 4. *Segu.*  
*ver.* 3. p. 207.
15. *Tillea erecta, floribus lateralibus, foliis obtusis.* *Sp.*  
*pl.* 1. p. 129.  
*Tillea erecta.* *Hort. ups.* 24. *Sauv. monsp.* 129.  
*Sedum annuum minimum stellatum rubrum.* *Magn.*  
*monsp.* 233. t. 237.
16. *Sedum arvense, flore rubente.* *Bauh. pin.* 283. *prodr.*  
132. *Segu. ver.* 1. p. 461.
17. *Sedum minus teretifolium alterum.* *Bauh. pin.* 283.
18. *Sedum minus teretifolium luteum.* *Bauh. pin.* 283.  
*Habitat in Europa australi.* ☉  
*Planta (an unica a — ?) mea. Caulis digitalis,*  
*simplex pilis raris viscosis. Folia sparsa, patentia,*  
*oblonga, obtusa, carnosae: inferiora quaternae. Cyma*  
*tri-f. quadrifida ramis recurvis. Flores sessiles.*  
*Corolla alba, valde mucronata carina villosa pur-*  
*purascens. Stamina 5, mox recurvata. Germina*  
*subpilosa, Fructus purpurascens. Crassulam dice-*  
*rem ob stamina; sed Petala 5 omnino Sedi & ha-*  
*bitus.*
15. **SEDUM** foliis subovatis adnato-sessilibus gibbis ere- acre.  
tiusculis alternis, cyma trifida. *Hort. cliff.* 177. *Fl.*  
*svec.* 389; 403. *Mus. med.* 218. *Roy. lugdb.* 456.  
*Sempervivum minus vermiculatum acre.* *Bauh. pin.*  
283.  
*Illecebra* f. *Sempervivum tertium.* *Dod. pempt.* 129.  
*Habitat in Europæ campis siccissimis sterilissimis.* 24.  
16. SE-



- sexangulare. 16. *SEDUM* foliis subovatis adnato-fessilibus gibbise-  
rectiusculis sexfariam imbricatis. *Fl. suec.* 390; 404.\*  
Sempervivum minus vermiculatum insipidum. *Bauh.*  
*pin.* 284 ?  
Sempervivum minimum. *Cam. epit.* 856.  
*Habitat in Europæ borealis campis apricis siccis.* 24  
*Differt a præcedenti, cui simillimum, Foliis sexfariam*  
*oblique imbricatis; Ramis nunquam copiosis, ut cespiti-*  
*tem constituent; Floribus in singulo ramo cymæ raro*  
*ultra tres.*
- annuum. 17. *SEDUM* caule erecto solitario annuo, foliis ovatis  
fessilibus gibbis alternis, cyma recurva. *Fl. suec.* 391;  
405.\*  
Sedum minimum non acre, flore albo. *Raj. angl.* 3.  
*p.* 270. *t.* 12. *f.* 2. *figura tenuis.*  
*Habitat in Europa boreali.* ☉
- villosum. 18. *SEDUM* caule erecto, foliis planiusculis, peduncu-  
lisque subpilosis.  
Sedum palustre subhirsutum purpureum. *Bauh. pin.* 283.  
Sedum minus 111. f. palustre. *Clus. hist.* 2. *p.* 59.  
Sedum palustre 11. *Oed. dan. t.* 24.  
*Habitat in pratis paludosis Germaniæ, Angliæ, Gal-*  
*liæ, Pyrenæis.*

## PENTHORUM.

- sedoides. 1. *PENTHORUM.* *Gron. virg.* 51. \* *Aët. upf.* 1744  
*p.* 12. *t.* 2.  
*Habitat in Virginia.* 24. ☉

## OXALIS.

\* *Scapo radicali.*

- Acetosella. 1. *OXALIS* scapo unifloro, foliis ternatis, radice squa-  
moso-articulata. *Hort. cliff.* 175. *Fl. suec.* 385; 406.  
*Roy. lugdb.* 458. *Mat. med.* 216.  
*Oxalis* foliis ternatis: scapo unifloro. *Fl. lapp.* 194.  
*Trifolium acetosum* vulgare. *Bauh. pin.* 330.  
*Trifolium acetosum.* *Dod. pempt.* 578.  
β. *Oxys* flore subcæruleo. *Tournef. inst.* 88.  
γ. *Oxys* flore purpurascente. *Tournef. inst.* 88.  
*Habitat in Europæ borealis sylvis.* 24

2. OXALIS scapo unifloro, foliis ternatis indivisis ciliatis. *purpurea.*

*Oxalis* scapo unifloro, foliis ternatis, radice bulbosa.

*Hort. cliff.* 175. *Roy. lugdb.* 458. *Sp. pl.* 1. p. 433.

*Ehret. pict.* 1. 10. f. 2.

*Oxalidi* affinis planta bulbosa africana, flore purpureo magno. *Breyn. cent.* 102. t. 46.

*Oxys* bulbosa africana rotundifolia, caulibus & floribus purpureis amplis. *Comm. hort.* 1. p. 41. t. 21.

*Oxys* bulbosa trifolia hirsuta, flore albo. *Burm. afr.* 67. t. 27. f. 3.

*Oxys* africana bulbosa trifolia, flore luteo magna.

*Seb. mus.* 1. p. 37. t. 22. f. 10.

*Habitat in* Æthiopia. 24.

## 3. OXALIS scapo unifloro, foliis ternatis semibifidis: longiora, lobis lanceolatis.

*Habitat in* Virginia.

*Caulis nudus, simplex, squamis aliquot adpersus, & radice per terram exit terminatus Umbella ex foliis novem ☽ flore unico. Folia petiolata, ternata: foliolis sessilibus, plus quam semibifidis: lobis lanceolatis. Pedunculus uniflorus, in medio stipularum pari notatus. Flos unicus, longus.*

4. OXALIS scapo unifloro foliis ternatis bipartitis. *flava.*

*Oxalis* bulbosa, angustis digitatis foliis, flore solitario luteo. *Burm. afr.* 68. t. 27. f. 4.

*Oxys* africana bulbosa, floribus amplis luteis, foliis minimis hirsutis. *Raj. suppl.* 549.

*Oxalis* africana bulbosa, foliis simpliciter decaphyllis, flore flavo. *Burm. afr.* 69. t. 27. f. 5.

*Oxalis* bulbosa, simplici caule ad summum folioso, flore luteo. *Burm. afr.* 81. t. 30. f. 1.

*Habitat in* Æthiopia. 24.

5. OXALIS scapo umbellifero, foliis ternatis obcordatis, calycibus apice callosis. *violacea*

*Oxalis* caule aphylllo, flore purpureo, radice tuberosa rotunda. *Gron. virg.* 161.

*Oxys* purpurea virginiana, radice lilii more nucleata.

*Pluk. alm.* 274. t. 102. f. 4.

*Habitat in* Virginia. Canada. 24.

*Affinis sequenti, sed huic Foliola parum biloba, illi semibifida; huic calycis foliola ovata, illi lanceolata, utrique apice callosa.*

**Pes caprae.** 6. *OXALIS* scapo umbellifero, foliis ternatis subbipartitis apice subtus callosis. *Mill. ic.* 195. f. 1.

*Oxalis bulbosa pentaphylla & hexaphylla*, floribus magnis luteis & copiosis. *Burm. afr.* 80. t. 2. f. 28. f. 3.

*Habitat in Æthiopia.* 24

*Foliola obcordata, biloba f. semibifida, obtusa, glabra, tenera. Flos flavissimus. Radix granulata.*

**sensitiva.** 7. *OXALIS* scapo umbellifero, foliis pinnatis. *Fl. zeyl.* 180. \*

*Oxyoides malabarica sensitiva*, caule viridi glabro, flore luteo majore. *Garc. act. angl.* 1730. p. 379. t. 2.

*Herba Mimosa malabariensium.* *Zan. hist.* 199. t. 61.

*Herba fentiens.* *Rumph. amb.* 5. p. 301. t. 104. f. 2.

*Herba viva.* *Clus. exot.* 290.

*Herba viva foliis polypodii.* *Bauh. pin.* 259.

*Todda vaddi.* *Rheed. mal.* 9. p. 33. t. 19.

*Habitat in India.*

\* \* *Foliis caulinis alternis.*

**vericolor.** 8. *OXALIS* caule ramoso, pedunculis unifloris, foliis passim verticillatis linearibus emarginatis apice subtus callosis.

*Oxalis caule simpliciter, pedunculo unifloro abbreviato, foliis filiformibus subramosis.* *Sp. pl.* 434.

*Oxalis bulbosa trifolia*, foliis linearibus obtusis, flore externe rubro, intus albo. *Burm. afr.* 65. t. 27. f. 1.

*Oxys africana*, foliis tenuissimis in summitate caulis. *Raj. suppl.* 598.

β. *Oxys bulbosa africana angustifolia*, flore rubro obsoleto amplo. *Raj. suppl.* 548.

*Oxys africana*, foliis tenuissimis, flore amplo verticillato. *Pluk. amalth.* 169. t. 434. f. 5.

*Oxalis bulbosa angustifolia*, caule folioso, flore rubro. *Burm. afr.* 66. t. 27. f. 2.

*Habitat in Æthiopia.* 24

**incarnata.** 9. *OXALIS* caule subramoso bulbifero, pedunculis unifloris, foliis passim verticillatis foliolis obcordatis. *Oxalis pedunculis unifloris*, caule dichotomo. *Sp. pl.* 433.

*Oxalis caule bulbifero.* *Hort. cliff.* 175. *Roy. luc.* 458.

*Oxys*

*Oxys bulbosa æthiopica mindr, folio cordato, flore ex albedo purpurascēte. Comm. hort. 1. p. 43. t. 22.*

*Habitat in Æthiopia. 24*

*Folia respectīve ad lucem subtus incarnata. Bulbi ex alīs post florēscēntiam nascuntur.*

10. **OXALIS** caule ramofo, pedunculis unifloris; foliis hirta, sessilibus: foliolis lanceolatis.

*Oxalis caule ramofo, foliis lineari-lanceolatis sessilibus, floribus lateralibus solitariis. Roy. lugdb. 532.*

*Oxalis radice longa fibrosa, caulibus ramosis, foliis ternis angustis, florum petiolis longissimis. Burm. afr. 71. t. 28. f. 2.*

• *Oxys bulbosa africana, flore purpureo, foliis pilosis, pedunculis privis. Seb. mus. 1. p. 33. t. 22. f. 11.*

*Oxys africana, hirsuta latioribus foliis, flore magno purpureo. Pluk. amalth. 164.*

- β. *Oxalis bulbosa, foliis angustis ternis hirtis, flore purpureo. Burm. afr. 70. t. 28. f. 1.*

*Oxys africana bulbosa, flore purpureo, caule folioso. Raj. suppl. 599.*

*Oxys africana bulbosa, dorychnii monspeliensium parvis foliis, floribus purpureis amplis. Pluk. amalth. 164. t. 434. f. 7.*

*Habitat in Æthiopia.*

*Caulis solidior. Folia alterna, sessilia, obtusiuscula, subtus hirta. Pedunculi uniflori, hirti, axillares, longi uti Calyces. Radix bulbosa.*

11. **OXALIS** caule ramofo diffuso, pedunculis umbelliferis. *Hort. cliff. 175. Hort. upl. 116. Roy. lugdb. 158. Sauv. monsp. 173. Gort. gelr. 91.*

*Oxys flavo flore. Clus. hist. 2. p. 249.*

*Trifolium acetosum corniculatum. Bauh. pin. 330. luteum minus repens, etiam procumbens. Moris. hist. 2. p. 183. f. 2. t. 17. f. 2.*

*Habitat in Italia, Sicilia. ○*

*Confer. α. Oxalis lutea annua, floribus dentatis. Few. per. 3. p. 49. t. 24.*

- β. *Oxys lutea americana humilior annua. Dill. elth. 299. t. 221. f. 288.*

*Oxys zeylanica. Burm. zeyl. 179. differt tantum caule repente & quod minor.*

- fricta.** 12. *OXALIS* caule ramoso erecto, pedunculis umbelliferis. *Gron. virg.* 161. *Brown. jam.* 231.  
*Oxys americana* erectior. *Tournef. inst.* 80.  
*Trifolium acetosum corniculatum luteum majus rectum indicum* f. *virgineum*. *Morif. hist.* 2. p. 184. f. 2. t. 17. f. 3.  
*Habitat in Virginia.*  
*Affinis nimium precedenti, ut fere varietas.*
- frutescens.** 13. *OXALIS* caule fruticoso erecto, pedunculis umbelliferis, foliis ternatis ovatis: Intermedio petiolato. *Oxalis* caule erecto fruticoso, foliis ternatis impari maximo. *Mill. dict.* 7.  
*Oxys lutea frutescens*, trifolii bituminosi facie. *Plum. spec.* 2. ic. 213. f. 1.  
*Habitat in America meridionali.* ‡
- Barrelieri.** 14. *OXALIS* caule ramoso erecto, pedunculis bifidis racemiferis. †  
*Trifolium acetosum americanum*, rubro flore. *Barr. rar.* 64. t. 1139. *Bocc. mus.* 2. p. 63. t. 51.  
*Habitat in America.*

## S U R I A N A.

- maritima.** 1. *SURIANA.* *Hort. cliff.* 492.  
*Suriana* foliis portulacæ angustis. *Plum. gen.* 37. ic. 249. f. 1.  
*Suriana maritima*, foliis lanceolatis, floribus singularibus. *Brown. jam.* 190.  
*Arbor americana*, falicis folio, frondosa bermudensis. *Pluk. alm.* 44. t. 241. f. 5.  
*Habitat in Americæ calidioris maritimis.* ‡  
*Stamina huic 10 adscripta.* *D. Jacquin.*

## A G R O S T E M M A.

- Githago.** 1. *AGROSTEMMA* hirsuta, calycibus corollam æquantibus petalis integris nudis.  
*Agrostemma.* *Hort. cliff.* 175. *Pl. fvec.* 383.; 47. *Roy. lugd.* 449.  
*Lychnis segetum major.* *Bauh. pin.* 204.  
*Lolium.* *Fuchf. hist.* 127.  
*Habitat inter Europæe jegetes.* ☉
- Cœlirofa.** 2. *AGROSTEMMA* glabra, foliis lineari lanceolatis petalis emarginatis coronatis. *Hort. ups.* 115.  
*Lychnis*

*Lychnis segetum*, *Nigellastrum minus glabrum dicta*,  
flore eleganter rubello. *Morif. hist. 2. p. 543. f. 5.*  
*t. 22. f. 32.*

*Lychnis foliis glabris*, calyce duriore. *Bocc. sic. 27.*  
*Pseudo-Melanthium glabrum ficulum. Raj. hist. 999.*

*Habitat in Sicilia. Oriente. ☉*

3. AGROSTEMMA tomentosa, foliis ovato-lanceola- Coronaria  
tis, petalis integris coronatis. *Hort. upf. 115.*

*Coronaria. Hort. cliff. 174. Roy. lugdb. 449.*

*Lychnis coronaria dioscoridis sativa. Bauh. pin. 203.*

*Lychnis. Cam. epit. 569.*

*Habitat in Italia. ♂*

*Naturalis planta corolla alba medio rubra est.*

4. AGROSTEMMA tomentosa, petalis emarginatis. Flos Jovis.

*Lychnis coronaria sylvestris. Bauh. pin. 204. Morif.*  
*hist. 2. p. 540. f. 5. t. 36. f. 2.*

*Lychnis umbellifera montana helvetica. Zan. hist. 128.*

*t. 51. Raj. hist. 993.*

*Habitat in Helvetia.*

## LYCHNIS.

1. LYCHNIS floribus fasciculatis fastigiatis. *Hort. cliff. chalcidoni-*  
*174. Hort. upf. 115. Roy. lugdb. 449. ca.*

*Lychnis hirsuta, flore coccineo, major. Bauh. pin. 203.*

*Flos constantinopolitanus. Dod. pempt. 178.*

*Habitat in Tataria. 24*

2. LYCHNIS petalis quadrifidis, fructu subrotundo. Flos cuculi.  
*Hort. cliff. 174. Fl. svec. 384; 408. Roy. lugdb. 139.*

*Caryophyllus pratensis, flore laciniato simplici. Flos*  
*cuculi. Bauh. pin. 210.*

- β. *Caryophyllus pratensis, flore pleno. Bauh. pin. 210.*

*Mill. ic. 169. f. 2.*

*Odontites plinii. Clus. hist. 1. p. 292; 293.*

*Habitat in pratis Europæ humidiusculis. 24*

3. LYCHNIS petalis integris.

*Silene floribus pentagynis, capsulis quinquelocularibus.*

*Hort. cliff. 172. Fl. svec. 364; 409. Roy. lugdb. 445.*

*Lychnis sylvestris viscosa rubra angustifolia. Bauh.*  
*pin. 205.*

*Lychnis sylvestris IV. Clus. hist. 1. p. 289.*

*Habitat in Europæ septentrionalis pratis lucis. 24*

*Linæi Spec. plant. pars. I.*

R r

4. LYCH

Viscaria,

alpina.

4. **LYCHNIS** petalis bifidis, floribus corymbosis. *Fl. fœc. 2. n. 410. \**

*Silene* floribus corymbosis, caule erecto, foliis lanceolato-linearibus. *Roy. lugdb. 447.*

- Silene lapponica alpina*, facie viscaria. *Fl. lapp. 185. \**

*Silene* floribus in capitulum congestis. *Hall. helv. 376. t. 7.*

*Habitat in Alpibus Lapponicis, Helveticis, Sibiricis, Pyrenaicis. 2. ♂*

sibirica.

5. **LYCHNIS** petalis bifidis, caule dichotomo, foliis subhirtis.

*Habitat in Sibiria. Gmelin. 2.*

*Statura Silenes rupestris, sed Radix perennis, Folia minus glabra, & Styli 5 distinguunt plantam.*

dioica.

6. **LYCHNIS** floribus dioicis. *Hort. cliff. 171. Fl. fœc. 361; 411. Roy. lugdb. 448.*

*Cucubalus* caule composito, calycibus oblongo-ovatis. *Fl. lapp. 182.*

*Lychnis sylvestris* f. *aquatica purpurea simplex. Bauh. pin. 204.*

*Lychnis sylvestris noctiflora alba simplex*, calyce amplissimo. *Till. pis. 105. t. 41. f. 1.*

*Melandrium plinii genuinum. Clus. hist. 1. p. 394 mas.*

- β. *Cucubalus* floribus hermaphroditis pentagynis, capsulis unilocularibus. *Hort. cliff. 170.*

*Lychnis sylvestris alba simplex. Bauh. pin. 204.*

- γ. *Lychnis alba multiplex. Bauh. pin. 204.*

*Habitat in Europæ frigida pratis succulentis. 2*

*Variat floribus hermaphroditis, licet varius.*

apetala.

7. **LYCHNIS** calyce inflato, corolla calyce brevior. caule subuniflora, flore hermaphrodito.

*Cucubalus* caule simplicissimo uniflora, petalis calyce brevioribus. *Fl. fœc. 363; 412.*

*Cucubalus* caule simplicissimo uniflora, corolla inclusâ. *Fl. lapp. 181. t. 12. f. 1.*

*Habitat in Alpibus Lapponicis, Sibiricis. 2*

*In Sibiria planta caule multiflora lecta est. Hæc hermaphrodita.*

## CERASTIUM.

\* *Capsulis oblongis.*

1. CERASTIUM foliis connatis. *Hort. cliff.* 173. *Hort. perfoliatum.*  
*upf.* 115. *Roy. lugdb.* 450.

*Myosotis orientalis perfoliata, lychnidis folio. Tournesf.*

*cor.* 18. *Dill. elh.* 295. t. 217. f. 284.

*Habitat in Græcia.* ☉

2. CERASTIUM foliis ovatis, petalis calyci æquali-<sup>1</sup> vulgatum.  
 bus, caulibus diffusis. *Fl. svec.* 2. 415.

*Cerastium floribus decandris, petalis subtridentatis calyce majoribus. Læfl. it.* 143. \*

*Myosotis arvensis hirsuta, parvo flore albo. T. Vaill.*  
*parif.* 142. t. 30. f. 1.

*Alfene hirsuta, magno flore. Bauh. pin.* 251.

*Auricula muris quorundam, flore parvo, vasculo tenui longo. Bauh. hist.* 3. p. 359.

*Habitat in Scanlæ & Europæ australioris pratis, dreis.*

*Simile C. viscoso, sed denso cæspite crescens.*

3. CERASTIUM erectum villosa-viscosum. *Hort. cliff.* viscosum.  
 174. *Fl. svec.* 379; 414. *Roy. lugdb.* 450.

*Cerastium corolla calyci æquali. Fl. lapp.* 193.

*Myosotis hirsuta altera viscosa. Vaill. parif.* 142. t. 30.  
 f. 1. 3.

*Alfene hirsuta altera viscosa. Bauh. pin.* 251.

*Habitat in Europæ pratis macilentis.* ☉

4. CERASTIUM floribus pentandris, petalis emarginatis. *Hort. cliff.* 173. *Fl. svec.* 382; 416. \* *It. æl.* 17. <sup>semidecandrum.</sup>  
*Roy. lugdb.* 450. *Gron. virg.* 51. *Gort. gelr.* 90.

*Cerastium corolla calyce brevior. Fl. lapp.* 193.

*Cerastium hirsutum minus, parvo flore. Raj. angl.* 3.  
 p. 348. t. 15. f. 1.

*Myosotis arvensis hirsuta minor. Vaill. parif.* 142. t. 30. f. 2.

*Habitat in campis apricis sterilissimis Europæ borealis.* ☉

*Filamenta huic 3 sterilia, 3 antherifera.*

5. CERASTIUM floribus pentandris, petalis integris. <sup>pentadrum.</sup>  
*Læfl. it.* 142. \*

*Habitat in Hispania.*



*Planta minima, similis C. semidecandro, a quo differt: colore viridi nec incano. Staminibus 5 fertilibus; nec 5 aliis anthera destitutis. Petalis calyce longe brevioribus lanceolatis acutis; nec acute emarginatis.*

arvense.

6. **CERASTIUM** foliis lineari-lanceolatis obtusis glabris, corollis calyce majoribus. *Fl. suec.* 381; 417.  
*Cerastium foliis calycibusque hirsutis. Hort. cliff.* 174.  
*Roy. lugdb.* 450.  
*Myosotis arvensis hirsuta, flore majore. Vaill. parij.* 141. t. 30. f. 4.  
*Caryophyllus arvensis hirsutus, flore majore. Bauh. pin.* 210.  
*Habitat in Scania & australiori Europa. 24*

dichotomum.

7. **CERASTIUM** foliis lanceolatis, caule dichotomo ramosissimo, capsulis erectis. *Roy. lugdb.* 450.  
*Lychnis segetum minor. Bauh. pin.* 204.  
*Alfne corniculata. Clus. hist. 2. p. 184.*  
*Habitat inter segetes Hispaniæ. ☉*

alpinum.

8. **CERASTIUM** foliis ovato-lanceolatis; caule diviso, capsulis oblongis.  
*Cerastium foliis ovato lanceolatis, corollis calyce majoribus. Fl. suec.* 380; 418.  
*Cerastium foliis lanceolato-ovatis, caule subunisfloro. Roy. lugdb.* 451.  
*Cerastium corolla calyce majore. Fl. lapp.* 192.  
*Caryophyllus holosteus alpinus latifolius. Oed. dan. t. 6.*  
*Alfne myosotis facie lychnis alpina, flore amplo niveo, repens. Raj. angl. 3. p. 349. t. 15. f. 2.*  
*Habitat in Alpibus Europæ.*

\* \* *Capsulis subrotundis.*

repens.

9. **CERASTIUM** foliis lanceolatis, pedunculis ramosis, capsulis subrotundis.  
*Cerastium caule perenni procumbente, foliis lanceolatis tomentosis. Roy. lugdb.* 450.  
*Cerastium perenne procumbens. Hort. cliff.* 174.  
*Myosotis arvensis, polygoni folio. Vaill. parij.* 141. t. 30. f. 5.  
*Ocymoides lychnitis reptante radice. Col. phytob.* 115. t. 31.  
*Lychnis incana repens. Bauh. pin.* 206.  
*Habitat in Gallia, Italia. 24*

10. **CERASTIUM** foliis linearibus acuminatis glabris, *strictum.*  
pedunculis unifloris subtomentosis, capsulis globosis.  
*Alpine alpina subhirsuta, linearæ folio. Scheuch. alp. 2. p. 130.*  
*Caryophyllus holosteus alpinus, foliis strictissimis glabris acutis, flore magno. Burf. XI. 119.*  
*Habitat in Alpibus Austriacis.*  
*Facies Auriculæ muris pulchro flore, folio tenuissimo 1. B. 3. p. 360, sed Flos Arenariæ montanæ. Folia linearia, acuminatissima, stricta, glabra. Caules glabri, procumbentes. Pedunculus uniflorus s. biflorus, in medio Bracteis duabus, brevibus, lanceolatis, latioribus; subtomentosus uti Calyx.*
11. **CERASTIUM** caule perenni procumbente, foliis *suffruticosum.*  
lineari-lanceolatis subhirsutis.  
*Myosotis tenuissimo folio rigido. Tournes. inst. 205.*  
*Habitat in Europa australi. 5*
12. **CERASTIUM** foliis lanceolatis scabris, petalis cre- *maximum.*  
natis, capsulis globosis.  
*Habitat in Sibiria. D. Gmelin.*
13. **CERASTIUM** foliis cordatis sessilibus, floribus so- *aquaticum.*  
litariis, fructibus pendulis. *Fl. suec. 2. n. 413. \**  
*Alpine major. Bauh. pin. 250.*  
*Alpine maxima folanifolia. Mentz. pug. 2. t. 1. f. 3.*  
*Habitat ad littora lacuum Europæ. 4*
14. **CERASTIUM** glabrum, caule stricto, foliis lan- *mantium.*  
ceolatis, pedunculis longissimis, capsulis globosis.  
*Amæn. acad. 4. p. 315. \**  
*Alpine caryophylloides glabra, florum pediculis longissimis. Segu. veron. 3. p. 178. t. 4. f. 2.*  
*Habitat Veronæ. ☉*
15. **CERASTIUM** foliis ovatis subtomentosis, ramis *latifolium.*  
subunifloris, capsulis globosis.  
*Caryophyllus holosteus alpinus latifolius. Bauh. pin. 210. prodr. 104.*  
*Habitat in Alpibus Helvetiæ. 4*
16. **CERASTIUM** foliis oblongis tomentosis, *tomentoso-*  
culis ramosis, capsulis globosis. *sum.*

*Cerastium* foliis lanceolato-linearibus subhirsutis, corolla calycem superante. *Sauv. monsp.* 142.

*Caryophyllus* holosteus tomentosus latifolius. *Bauh. pin.* 210. *prodr.* 104.

β. *Caryophyllus* holosteus tomentosus angustifolius. *Bauh. pin.* 210. *prodr.* 104.

*Habitat in Granada.* 24

## S P E R G U L A.

- arvensis. 1. **SPERGULA** foliis verticillatis, floribus decandris. *Hort. cliff.* 173. *Fl. suec.* 377. 419. *Roy. lugdb.* 451. *Spergula* fructu pendulo. *Fl. lapp.* 190. *Spergula.* *Dod. pempt.* 527. *Alfina* *Spergula* dicta major. *Bauh. pin.* 251. *Habitat in Europæ agris.* ☉
- pentandra. 2. **SPERGULA** foliis verticillatis, floribus pentandris. *Læf. it.* 143. *Spergula* foliis filiformibus verticillatis raris, seminibus nigris. *Sauv. monsp.* 167. *Alfina* *spergulæ* facie minima, seminibus emarginatis. *Tournesf. inst.* 244. *Vaill. parisi.* 8. *Alfina* *spergulæ* facie minima. *Magn. monsp.* 14. *Arenaria* teretifolia verna, flore albo, semine limbo foliaceo cincto. *Rupp. jen.* 191. *Spergula* annua, semine foliaceo nigro circulo membranaceo albo cincto - *Morif. hist.* 2. p. 551. *Blas.* 28. *Dill. giff.* 46. *E. N. C. cent.* 5. p. 275. t. 4. *Raj. angl.* 3. p. 351. *Habitat in Germania, Gallia, Anglia, Hispania.* ☉
- nodosa. 3. **SPERGULA** foliis oppositis subulatis lævibus, caulibus simplicibus. *Spergula* foliis oppositis, pedunculis simplicibus. *Hort. cliff.* 173. *Fl. suec.* 378; 420. *Roy. lugdb.* 451. *Arenaria* *Bauh. hist.* 3. p. 720. *Raj. hist.* 1033. *Alfina* nodosa germanica. *Bauh. pin.* 251. *Alfina* palustris, ericæ folio, polygonoides, articulis crebrioribus, flore albo pulchello. *Pluk. alm.* 23. t. 7. f. 4. *Polygonum*, foliis gramineis, alterum. *Læf. præs.* 204. t. 64. *Habitat in Europæ frigidioris campis subhumidis.* 24

4. SPERGULA foliis oppositis subulatis ciliatis fasci- laricina.  
culatis.  
*Habitat in Sibiria. D. Gmelin.*
5. SPERGULA foliis oppositis linearibus lævibus, pe- taginoides.  
dunculis solitariis longissimis, caule repente.  
Alline tenuifolia, pedunculis florum longissimis. *Vaill.*  
*parif. 8. Sauv. monsp. 142.*  
*Habitat in Gallia, Sibiria ex D. Gmelin.*

## DECAGYNIA.

## NEURADA.

1. NEURADA. *Hort. upf. 117.* procum-  
bens:  
*Chamædrifolia tomentosa maseariensis. Pluk. alm. 97.*  
*t. 275 f. 6., Shaw. afr. 133.*  
*Habitat in Aegypto, Arabia, Numidia. ☉*

## PHYTOLACCA.

1. PHYTOLACCA floribus octandris octagynis. octandra.  
*Phytolacca mexicana, baccis sessilibus. Dill. elth. 218.*  
*t. 239. 308.*  
*Jama gobo. Kæmph. amæn. 828. t. 829.*  
*Habitat in Mexico. 24*  
*Statura Ph. decandra. Folia albidiora. Pedunculus*  
*communis brevissimus. Pedicelli breves, ut fere*  
*nulli. Calyx f. Corolla alba, omnino plana, nec con-*  
*cava. Stamina 8 Styli octo.*
2. PHYTOLACCA floribus decandris decagynis. *Hort. decandra.*  
*cliff. 177. Hort. upf. 117. Gron. virg. 161. Roy.*  
*lugdb. 222.*  
*Phytolacca vulgaris. Dill. elth. 318. t. 239. f. 309.*  
*Solanum racemosum americanum. Pluk. alm. 353. t.*  
*225. f. 3.*  
*Habitat in Virginia. 24*
3. PHYTOLACCA floribus icofandris decagynis. icofandra.  
*Phytolacca spicis florum longissimis, radice annua.*  
*Mill. dict. t. 207.*  
*Habitat in Malabar. ☉. 15*  
*Cauli, tripedalis, herbaceus sed perennans, paucis ra-*  
*mis subdivisus. Folia striatiora, acutiora. Pedun-*  
*cu-*


culus inferne triqueter, at intra racemum multangularis, sed in hac scaber. Pedicelli inferiores sepe triflori. Flores albi. Stamina, certe in inferioribus floribus, viginti. Bacca saturate purpurea.

dioica.

4. PHYTOLACCA floribus dioicis.

Habitat in - - - - Alstræmer ex horto Madritensi. h

Caulis perfecte fruticosus. Folia exacte ovata, glabra, magnitudine Populi nigrae, longius petiolata. Flores ut in reliquis, sed dioici.



Clas.

## Classis XL

## DODECANDRIA

Stamina inter XI &amp; XIX inclusive.

## MONOGYNIA.

## A S A R U M.

1. ASARUM foliis reniformibus obtusis binis. europæum,  
*Asarum foliis subcordatis petiolatis. Hort. cliff. 178.*  
*Fl. suec. 392; 421. Mat. med. 219. Guett. stamp. 198.*  
*Roy. lugdb. 207.*  
*Asarum. Bauh. pin. 197. Cam. epit. 19.*  
*Habitat in Europæ nemoribus. 24*
2. ASARUM foliis reniformibus mucronatis. canadense.  
*Asarum foliis subcordatis petiolatis. Gron. virg. 52.*  
*Asarum canadense, mucronato folio. Moris. hist. 3. p.*  
*511. f. 13. t. 7. f. 4.*  
*Asarum canadense. Corn. canad. 24. t. 25.*  
*Habitat in Canada. 24*  
*An varietas præcedentis? sic suadent Folia bina; dis-*  
*suadent alia.*
3. ASARUM foliis cordatis obtusis glabris petiolatis. virginicum.  
*Gron. virg. 162.*  
*Asarum virginianum, pistolochiæ foliis subrotundis,*  
*cyclaminis more maculatis. Pluk. alm. 53 t. 78. f. 2.*  
*Moris. hist. 3. p. 511. f. 17. t. 7. f. 3.*  
*Habitat in Virginia, terra Mariana, Carolina. 24*
4. ASARUM foliis sessilibus imbricatis, floribus quadri- Hypocistis.  
*fidis. Mat. med. 220.*  
*Asarum aphyllum squamosum. Sauv. monsp. 4.*  
*Hypocistis. Bauh. pin. 465. Cam. epit. 96, 97. Clus.*  
*hist. 1. p. 68, 79. Duham. arb. 1. p. 170. t. 68.*  
*Habitat in Hispania, Lusitania parasitica cisti.*

## GETHYLLIS.

1. GETHYLLIS. afræ.  
*Hort. cliff. 493. \**  
*Habitat in Africa.*  
*Statura Croci.*

## BOCCONIA.

- frutescens. 1. **BOCCONIA**. *Hort. cliff.* 202. *Jacq. amer.* 22.  
*Bocconia ramosa*, sphondylii folio tomentoso. *Plur.*  
*gen.* 55. *Trew. chr. t.* 4.  
*Bocconia ramosa*, foliis majoribus sinuatis, racemis terminalibus *Brown. jam.* 244.  
*Chelidonium majus arboreum*, foliis quercinis. *Sloar.*  
*jam.* 82. *hist.* 1. p. 195. t. 125.  
*Cocoxihvitl*. *Hern. mex.* 158.  
*Habitat in Mexico, Jamaica.* ‡

## RHIZOPHORA.

- conjugata. 1. **RHIZOPHORA** foliis ovato-oblongis obtusiusculis integerrimis, calycibus sessilibus, fructibus cylindraceo subulatis. *Fl. zeyl.* 181. \*  
*Habitat in India.* ‡
- gymnophiza. 2. **RHIZOPHORA** foliis ovato-lanceolatis integerrimis, radice terræ superimposita.  
*Rhizophora calycum laciniis persistentibus patentibus versus fructum incurvatis.* *Wachend. ult.* 89.  
*Mangium celsum.* *Rumph. amb.* 3. p. 102. t. 68.  
*Candel.* *Rheed. mal.* 6. p. 57. t. 31. 32. *Raj. hist.* 1769.  
*Habitat in Indiæ ripis salis.* ‡
- Candel. 3. **RHIZOPHORA** foliis obtusis, pedunculis bigeminatis folio longioribus, fructibus subulatis.  
*Tsierrou-Kandel.* *Rheed. mal.* 6. p. 63. t. 35. *Raj. hist.* 1772.  
*Habitat in Indiæ salis.* ‡
- Mangla. 4. **RHIZOPHORA** foliis acutis, fructibus subulato-clavatis. *Jacq. hist.* t. 89.  
*Rhizophora segmentis calycum persistentibus reflexis, fructu acuminato.* *Wach. ultr.* 90.  
*Magles aquatica*, foliis subrotundis & punctatis. *Plur.*  
*gen.* 13.  
*Mangle pyri foliis*, cum siliquis longis, ficui indicæ affinis. *Bauh. hist.* 1. p. 415. *Sloar. jam.* 155. *hist.* 2. p. 63.  
*Mangium candelarium.* *Rumph. amb.* 3. p. 108. t. 71; 72.  
*Peekandel.* *Rheed. mal.* 6. p. 91. t. 34. *Raj. hist.* 1772.  
*Habitat in paludibus Caribæarum, Malabariz.* ‡

5. RHIZOPHORA fructibus cylindricis obtusis. *cylindrica.*  
 Kari-Kandel. *Rheed. mal.* 6. p. 59. t. 33. *Raj. hist.* 1770.  
 Mangium minus. *Rumph. amb.* 3. p. 106. t. 69.  
*Habitat in uliginosis Malabarix.* ‡
6. RHIZOPHORA foliis ovatis, floribus confertis, fru- *corniculata.*  
 ctibus arcuatis mucronatis.  
 Mangium fruticosum corniculatum. *Rumph. amb.* 3. p.  
 117. t. 77.  
*Habitat in Moluccis.* ‡
7. RHIZOPHORA foliis ovatis obtusis, floribus foli- *cafeolaris.*  
 tariis, fructibus orbiculatis depressis mucronatis.  
 Mangium cafeolare. *Rumph. amb.* 3. p. 111. t. 73; 74.  
*Habitat in Moluccis.* ‡

B L A K E A.

1. BLAKEA. *trinervis.*  
 Blakea foliis ellipticis trinerviis, nitidis, floribus la-  
 teralibus. *Brown. jam.* 323. t. 35.  
*Habitat in Jamaica.* ‡  
 Folia oblongo-ovata, petiolata, integerrima, coriacea,  
 opposita, trinervia: nervis subtus protuberantibus ni-  
 gricantibus. Flores oppositi, pedunculati, solitarii.

G A R C I N I A.

1. GARCINIA foliis ovatis, pedunculis unifloris. *Hort. Mangostana.*  
*cliff.* 182.  
 Mangostans. *Garc. aff. angl.* 431. t. 1. *Bont. jam.* 115.  
 Mangostana. *Rumph. amb.* 1. p. 132. t. 43.  
 Laurifolia javanensis. *Bauh. pin.* 461. *Raj. hist.* 1662.  
*Habitat in Java.* ‡
2. GARCINIA foliis lanceolatis, pedunculis trifloris. *celebica.*  
 Mangostana celebica. *Rumph. amb.* 1. p. 134. t. 44.  
*Habitat in India.* ‡

S T Y R A X.

1. STYRAX. *Hort. cliff.* 187. *Hort. upf.* 122. *Mat. officinale.*  
*med.* 227. *Roy. lagdb.* 265. *Sauv. monsp.* 306. *Gron.*  
*orient.* 149.  
 Styrax folio mali cotonei. *Bauh. pin.* 452.  
 Styrax. *Cam. epit.* 38.  
*Habitat in Syria, Judæa, Italia.* ‡

1. WIN-



## WINTERANIA.

Canella.

1. WINTERANIA. *Hort. cliff.* 488.Winteranus cortex. *Clus. exot.* 75.Laurus foliis enerviis obovatis obtusis. *Mat. med.* 196.*Sp. pl.* 1. p. 371.Canella foliis oblongis obtusis nitidis, racemis terminalibus. *Brown. jam.* 215. t. 27. f. 3.Cassia lighea jamaicensis, laureolæ foliis subcinereis, cortice piperis modo acri. *Pluk. alm.* 89. t. 81. f. 1.Cassia cinnamomea f. Cinnamomum sylvestre barbadensium. *Pluk. alm.* 89. t. 161. f. 7.Arbor baccifera laurifolia aromatica, fructu viridi calyculato racemoso. *Sloan. jam.* 165. *hist.* 2. p. 87. t.191. f. 2. *Catesb. carol.* 2. p. 50. t. 50.*Habitat in America.* h.

## HALESIA.

tetraptera.

## 1. HALESIA foliis lanceolato-ovatis: petiolis glandulosis.

Halesia fructibus membranaceo-quadrangulatis. *Ellis. act. angl.* v. 51. p. 931. t. 22. f. A.Frutex padi foliis ferratis, floribus monopetalis albis campaniformibus, fructu crasso tetragono. *Cateb. car.* 1. p. 64. t. 64.*Habitat in Carolina.* h.

diptera.

## 2. HALESIA foliis ovatis: petiolis lævibus.

Halesia fructibus alatis. *Ellis. act. angl.* v. 51. t. 931 f. B.*Habitat in Carolina.* D. Garden. h.Folia hujus sextuplo majora præcedentis, nec subtus subtomentosa. Fructus mucronatus alis duabus oppositis magnis; duabus minutis. Ordine naturali proxima *Styraci*.

## CRATEVA.

gynandra.

## 1. CRATEVA inermis, foliis integerrimis, floribus gynandris.

Crateva arborea triphylla, foliis ovatis glabris, racemis terminalibus. *Brown. jam.* 246.Arbor americana triphylla, numerosis staminibus pureis apicibus præditis, floris umbilicum occupantibus. *Pluk. phyt.* 147. f. 6.

H2.

*Habitat in Jamaica.* ‡

Differt a *C. Tapia* foliis tenuibus nec crassis. Petalis lanceolatis nec ovatis; imprimis vero quod columna longitudine calycis attollat stamina, ut in *Cleome gynandra*. Pistillum in meis non aderat.

2. *CRATEVA inermis*, foliis integerrimis: foliolis 12-16. *Tapia*.  
teratibus basi antica brevioribus.

*Crateva inermis*. *Fl. zeyl.* 211. *Sp. pl.* 1. p. 444. *Hort. cliff.* 484.

*Apioscorodon* f. *Arbor americana* triphyllis, allii odore, poma terens. *Puk. alm.* 34. t. 137. f. 7.

*Malus americana* trifolia, fructu pomi aurantii instar colorato. *Comm. hort.* 1. p. 129. t. 67.

*Anona* trifolia, flore stamineo, fructu sphaerico ferrugineo scabro minore, allii odore. *Sloan. jam.* 208. *hist.* 2. p. 169. *Raj. dendr.* 79.

*Nürvala*. *Rheed. mal.* 3. p. 49. t. 22.

*Habitat in Indiis.* ‡

3. *CRATEVA spinosa*, foliis serratis. *Fl. zeyl.* 212. *Marmelos*.

*Cucurbitifera* trifolia spinosa medica, fructu pulpa cydonii æmula. *Pluk. alm.* 125. t. 170. f. 5.

*Cydonia exotica*. *Bauh. pin.* 435.

*Bilanus*. *Rumph. amb.* 1. p. 197. t. 81.

*Covalam*. *Rheed. mal.* 3. p. 37. t. 37.

*Habitat in India.* ‡

## TRIUMFETTA.

- . *TRIUMFETTA* foliis basi emarginatis. *Hort. cliff.* *Lappula*.

210. \* *Hort. upf.* 148. *Roy. lugdb.* 476.

*Triumfetta* floribus calyculatis, foliis semitrilobis. *Jacq. amer.* 22.

*Triumfetta* fructu echinato racemoso. *Plum. gen.* 40. *ic.* 255.

*Triumfetta villosa*, foliis inferioribus angulato-ovatis serrato dentatis, floribus ternatis fasciculis geminatis. *Brown. jam.* 233.

*Lappula bermudensis* althæoides spicata, fructu orbiculari majore. *Puk. alm.* 206. t. 245. f. 7.

*Agrimonia lappacea* inodora, folio subrotundo dentato. *Sloan. jam.* 92. *hist.* 1. p. 211. *Raj. suppl.* 247.

*Habitat in Jamaica, Brasilia, Bermudis.* ‡

- Bartramia. 2. **TRIUMFETTA** foliis basi integris.  
*Triumphetta* floribus calyculatis, foliis rhombeo-ovatis. *Jacq. amer.* 22. *hist.* t. 90.  
*Bartramia*. *Fl. zeyl.* 174. *Sp. pl.* 1. p. 389.  
*Lappula benghalensis* tetralpermos, ribesii folio, echinis orbiculatis ad foliorum ortum plurimis sessilibus. *Pluk. alm.* 206. t. 41. f. 5.  
*Agrimonia maderaspatana*, folio rotundo singulari subtus incano. *Pet. gaz.* 31. t. 42. f. 10.  
*Lappago amboinica*. *Rumph. amb.* 6. p. 59. t. 25. f. 2.  
*Habitat in India.* ☉

### PEGANUM.

- Harmala. 1. **PEGANUM** foliis multifidis. *Hort. upf.* 144. *Gron. orient.* 165.  
*Peganum*. *Hort. cliff.* 206. *Hort. upf.* 144. *Roy. lugdb.* 461.  
*Ruta sylvestris*, flore magno albo. *Bauh. pin.* 336.  
*Harmala*. *Dod. pempt.* 121.  
*Habitat in arena Madriti, Alexandriæ, Cappadociæ, Galatiæ.* 24
- daurica. 2. **PEGANUM** foliis indivisis.  
*Harmala montana daurica* perennis multicaulis, polygalæ folio, flore albo. *Amm. ruth.* 91.  
 β. *Harmala montana*, polygalæ foliis, floribus luteis. *Amm. ruth.* 92.  
*Habitat in Sibiria.* 24

### NITRARIA.

- Schoberi. 1. **NITRARIA**.  
*Oxyris* foliis obtusis. *Hort. upf.* 295. *Gmel. fibir.* 2. p. 237. t. 98.  
*Cassia fructu nigro*. *Amm. ruth.* 256.  
*Habitat ad Salsas aquas Vrunscinensis Lacus, Astrachanensis, Volgæ finitimis.* 5

### PORTULACA.

- oleracea. 1. **PORTULACA** foliis cuneiformibus, floribus sessilibus. *Roy. lugdb.* 473. *Gron. virg.* 59. *Hort. upf.* 146. *Mat. med.* 260. *Dalib. parif.* 158. *Hall. her.* 392. *Sauv. monsp.* 168. *Brown. jam.* 233.  
*Portulaca* foliis cuneiformibus verticillatis sessilibus floribus sessilibus. *Hort. cliff.* 207.

Por-

- Portulaca angustifolia sylvestris. Bauh. pin. 288.*  
 ②. *Portulaca latifolia sativa. Bauh. pin. 288.*  
*Portulaca domestica. Lob. ic. 388.*  
*Habitat in Europa, India, Ins. Ascensionis, America. ☉*  
*Horologii floræ vigilias 11—12 habet.*
2. **PORTULACA** foliis subulatis alternis: axillis pilosis, pilosa.  
 floribus sessilibus terminalibus.  
*Portulaca foliis linearibus acutis, caule procumbente piloso. Roy. lugdb. 473.*  
*Portulaca curassavica, angusto longo lucidoque folio, procumbens. Comm. hort. 1. p. 9. t. 5.*  
*Portulaca corassavica lanuginosa procumbens. Herm. par. 215. t. 215.*  
*Portulaca lanuginosa procumbens, vermiculatae foliis, americana. Pluk. alm. 304. t. 247. f. 7. Volk. norib. 341. t. 341.*  
*Anacampseros supina minor, foliis linearibus turgidis, floribus summis ramulis confertis. Brown. jam. 234.*
- β. *Portulaca corassavica lanuginosa, psylli folio, erectior & elatior, flore dilutius rubente. Herm. par. 214. t. 214.*  
*Portulaca curassavica lanuginosa, kali folio, erectior & elatior. Pluk. alm. 303. t. 105. f. 4.*  
*Habitat in America meridionali. ☉*  
*Folia alterna, sed terminalia s. floralia octo in orbem digesta. Flores sessiles, terminales, calyce bifido, Staminiibus quindecim, Stylo quinquesido, Capsula circumscissa.*
- 3 **PORTULACA** foliis oblongis carnosiss, caule coralloides.  
 rymbofo, floribus sessilibus.  
*Portulaca erecta, sedi minoris facie, capitulo tomentoso. Sloan. jam. 88. hist. 1. p. 205. t. 129. f. 3.*  
*Halimus minimus, foliis oblongis succulentis tumetibus, summis ramulis densissime sitis. Brown. am. 206.*  
*Habitat in Jamaica. ☉*  
*Calyx diphyllus. Capsula circumscissa.*
4. **PORTULACA** foliis ovatis gibbis, pedunculo multifloro, caule fruticoso. Anacampseros.  
*Anacampseros foliis ovatis convexis acutiusculis. Hort. upf. 146.*

*Anacampseros foliis acuminatis. Hort. cliff. 207. Roy. lugdb. 412.*

*Telephiastrum folio globofo. Dill. elth. 375. t. 281. f. 363.*

*Portulaca africana sempervirens, flore rubicundo Comm. hort. 2. p. 177. t. 89. Raj. suppl. 507.*

*Oxalis affinis planta aphylla mammillaris. Burm. afr. 76. t. 30. f. 2. vitata.*

*Habitat ad Cap. b. Spei. §*

*Axille foliorum pilosæ sunt, & Folia succulenta conjungunt hanc cum Portulacis, quamvis stamina plurima & Capsula trivalvis. Calyx diphyllus est.*

racemosa.

5. **PORTULACA** foliis obovatis planiusculis, racemo simplici triquetro.

*Portulaca pedunculis multifloris triangularibus. Jacq. amer. 22.*

*Helianthemum frutescens, portulacæ folio. Plum. spec. 7. t. 150. f. 2.*

*Habitat in America ad maris littora. §*

*Calyx diphyllus. Capsula trilocularis.*

paniculata.

6. **PORTULACA** foliis ovatis planis, panicula racemosa.

*Portulaca floribus paniculatis. Jacq. amer. 22.*

*Portulaca americana latifolia erecta, floribus albis. Comm. hort. 1. p. 7. t. 4.*

*Portulacæ facie maritima fruticosa erecta americana Sloan. jam. 87. hist. 1. p. 205.*

*Anacampseros foliis radicalibus mollibus ovatis glabris, scapo assurgenti paniculato. Brown. jam. 234.*

*Habitat in Americæ maritimis. §*

*Calyx pentaphyllus. Capsula trivalvis.*

## LYTHRUM.

Salicaria.

1. **LYTHRUM** foliis oppositis cordato-lanceolatis, floribus spicatis dodecandris.

*Lythrum foliis oppositis. Hort. cliff. 178. Fl. suec. 393; 422 Roy. lugdb. 458. Guett. stamp. 2. p. 122.*

*Lythrum floribus verticillatis. Fl. lapp. 197.*

*Lyfimachia spicata purpurea. Bauh. pin. 246.*

*Lyfimachia purpurea communis major. Clus. hist. 1. p. 51.*

B. Blat-

- β. *Blattaria rubra spicata major*, folio subrotundo. *Marif. hisp.* 2. p. 490. f. 5. t. 10. f. 11.
- γ. *Lythmachia trifolia spicata purpurea*. *Bocc. mus.* 167.
- δ. *Lythmachia quadrifolia purpurea*, mollioribus & longioribus foliis. *Bocc. mus.* 167.
- Habitat in Europa ad ripas aquarum.* 24
2. **LYTHRUM** foliis oppositis subtus tomentosis, floribus decandris, corollis calyce calyceque genitalibus brevioribus. fruticosum.
- Habitat in China.* 5
- Frutex cortice lacero. Folia opposita, sessilia, lanceolata, integerrima, subtus subtomentosa.*
3. **LYTHRUM** foliis oppositis ovalibus, floribus sessilibus alternis hexandris, caule diffuso. *Parsonia.*
- Parsonia herbacea, foliis ovatis oppositis, floribus singularibus foliis ad alterum latus interpositis.* *Brown. jam.* 199. t. 21. f. 2.
- Habitat in Jamaica.* 24
- Caules prostrati. Folia subsessilia, nuda, integerrima. Racemi alterni. Flores solitarii, alterni, sessiles.*
4. **LYTHRUM** foliis oppositis subtus tomentosis subpetiolatis, floribus verticillatis lateralibus. verticillatum.
- Lythrum foliis oppositis, floribus verticillatis.* *Gron. virg.* 52.
- Habitat in Virginia.*
5. **LYTHRUM** foliis oppositis linearibus petiolatis, floribus dodecandris. petiolatum.
- Lythrum foliis petiolatis.* *Gron. virg.* 52.
- Habitat in Virginia.*
6. **LYTHRUM** foliis oppositis linearibus, floribus oppositis hexandris. lineare.
- Lythrum foliis linearibus, floribus hexandris solitariis.* *Gron. virg.* 162.
- Habitat in Virginia.*
- Caulis filiformis erectus angulatus, foliis oppositis linearibus, terminatus spica majore & duabus lateralibus minoribus compositis ex floribus oppositis remotis, sub singulis folio. Facies L. Hyssopifoliae.*
7. **LYTHRUM** foliis oppositis ovatis, floribus alternis subdecandris, caule prostrato. *Melanium.*
- Limæi Spec. plant. pars I. S 3 Me*

*Melanium herbaceum reclinatum*, foliis ovatis oppositis, floribus singularibus ad alas alternas. *Brewerj. jam.* 215.

*Habitat in Jamaica.* 24

Hyssopifolia

8. **LYTHRUM** foliis alternis linearibus, floribus hexandris. *Hort. ups.* 118.

*Lythrum foliis alternis.* *Hort. cliff.* 178. *Sauv. monsp.* 53. *Roy. lugdb.* 458. *Gort. gelr.* 271.

*Salicaria hyssopi folio latiore.* *Hall. jen.* 147. t. 6. f. 3.

*Hyssopifolia.* *Bauh. pin.* 218.

β. *Lyfimachia rubra non siliquosa.* *Bauh. pin.* 246.

γ. *Lyfimachia linifolia purpureo-cærulea.* *Bauh. pin.* 246.

*Habitat in Germaniæ, Helvetiæ, Angliæ, Galliæ inundatis.* ☉

Thymifolia. 9. **LYTHRUM** foliis alternis linearibus, floribus tetrapetalis.

*Lythrum foliis linearibus.* *Sauv. monsp.* 49.

*Salicaria minima tenuifolia.* *Tournef. inst.* 254.

*Salicaria hyssopi folio angustiore.* *Buxb. hal.* 290. \*

*Hyssopifolia minor* s. potius *thymifolia maritima.* *Barr. hist.* 3. p. 792.

*Polygonum aquaticum minus.* *Barr. ic.* 773. f. 2.

*Habitat in Italiæ & G. Narbonensis uliginosis.* ☉

*Petala 4, stamina 2, calyx adentatus. Folia linear-lanceolata, alterna, ima opposita. Flores solitarii axillares. Calyx st. pulis 2 lateralibus fulcitus. D. Sauvages.*

virgatum.

10. **LYTHRUM** foliis oppositis lanceolatis, panicula virgata, floribus dodecandris.

*Salicaria orientalis, salicis folio acutissimo & glabro.* *Tournef. cor.* 18.

*Salicaria glabra ramosissima, longissimis & angustissimis foliis.* *Amm. ruth.* 89.

*Salicaria glabra, longioribus & angustioribus foliis.* *Amm. ruth.* 88.

*Habitat in Sibiria, Tataria.*

*Caulis paniculatus, ramis alternis longis. Flores 2. f. 2. pedicellati, ex singula ala alterna racemi.*

## GINORA.

americana. 1. **GINORA.** *Jacq. amer.* 22.

Ha.

*Habitat in America. b*  
*Frutex pulcher habet Myrti.*

## DIGYNIA.

### HELIOCARPUS.

1. HELIOCARPUS. *Hort. cliff. 211. t. 16. \* Trew. americana;*  
*chret. t. 45.*

*Habitat in Vera Cruce. b*

### AGRIMONIA.

1. AGRIMONIA foliis caulinis pinnatis: impari petio- *Eupatoria.*  
 lato, fructibus hispidis. *Hort. cliff. 179. Fl. sec. 394;*  
*423. Hort. upf. 118. Mat. med. 221. Gron. virg. 53.*  
*Roy. lugdb. 240. Hall. helv. 407. Guett. stamp. 1. p.*  
*293. Dalib. parif. 139.*

*Eupatorium veterum f. Agrimonia. Bauh. pin. 321.*

*Eupatorium. Fuchf. hist. 244. Cam. epit. 756. Mill. ic. 15.*

*Habitat in Europæ pratis apricis argillaceis. 24*

2. AGRIMONIA foliis caulinis pinnatis: impari sessili, *repen.*  
 fructibus hispidis.

*Agrimonia orientalis humilis, radice crassissima repen-*  
*te, fructu in spicam brevem & densam congesto.*

*Tournef. cor. 21.*

*Habitat in Oriente.*

*Affinis nimium A. Eupatoriæ, sed Caules breviores,*  
*crassiores, hirsuti. Stipulæ longitudine interstitio-*  
*rum, reflexæ, fere totum obumbrant caulem. Fo-*  
*liorum impar non pedicellatum. Spica brevis, den-*  
*sa, sessilis, Bracteis longitudine florum.*

3. AGRIMONIA foliis caulinis ternatis, fructibus gla- *Agrimonioid-*  
 bris. *Hort. cliff. 179. Roy. lugdb. 241. des.*

*Agrimariæ similis. Bauh. pin. 321.*

*Agrimonioides. Col. ceph. t. p. 145. t. 144. Boerh.*  
*lugdb. 1. p. 179.*

*Habitat in Italiæ nemoribus umbrosis humentibus. 24*

*Stamina sæpe vix ultra 8.*

## TRIGYNIA.

### RESEDA.

1. RESEDA foliis lanceolatis integris, calycibus quadri- *lateola.*  
 fidis.



## WINTERANIA.

Canella.

1. WINTERANIA. *Hort. cliff.* 488.Winteranus cortex. *Clus. exot.* 75.Laurus foliis enerviis obovatis obtusis. *Mat. med.* 196.*Sp. pl.* 1. p. 371.Canella foliis oblongis obtusis nitidis, racemis terminalibus. *Brown. jam.* 215. t. 27. f. 3Cassia ligtea jamaicensis, laureolæ foliis subcinereis, cortice piperis modo acri. *Pluk. alm.* 89. t. 81. f. 1.Cassia cinnamomea f. Cinnamomum sylvestre barbadensium. *Pluk. alm.* 89. t. 161. f. 7.Arbor baccifera laurifolia aromatica, fructu viridi calyculato racemoso. *Sloan. jam.* 165. *hist.* 2. p. 87. t. 191. f. 2. *Catesb. carol.* 2. p. 50. t. 50.*Habitat in America.* ☿.

## HALESIA.

tetraptera.

## 1. HALESIA foliis lanceolato-ovatis: petiolis glandulosis.

Halesia fructibus membranaceo-quadrangulatis. *Ellis. act. angl.* v. 51. p. 931. t. 22. f. A.Frutex padi foliis ferratis, floribus monopetalis albis campaniformibus, fructu crasso tetragono. *Catesb. car.* 1. p. 64. t. 64.*Habitat in Carolina.* ☿.

diptera.

## 2. HALESIA foliis ovatis: petiolis lævibus.

Halesia fructibus alatis. *Ellis. act. angl.* v. 51. t. 931 f. B.*Habitat in Carolina.* D. Garden. ☿.Folia hujus sextuplo majora præcedentis, nec subtus subtomentosa. Fructus mucronatus alis duabus oppositis magnis; duabus minutis. Ordine naturali proxima *Styraci*.

## CRATEVA.

gynandra.

## 1. CRATEVA inermis, foliis integerrimis, floribus gynandris.

Crateva arborea triphylla, foliis ovatis glabris, racemis terminalibus. *Brown. jam.* 246.Arbor americana triphylla, numerosis staminibus purpureis apicibus præditis, floris umbilicum occupantibus. *Pluk. phyt.* 147. f. 6.*Ha-*

*Habitat in Jamaica.* †

Differt a *C. Tapia* foliis tenuibus nec crassis. Petalis lanceolatis nec ovatis; imprimis vero quod columna longitudine calycis attollat stamina, ut in *Cleome gynandra*. Pistillum in meis n.n. aderat.

2. CRATEVA inermis, foliis integerrimis: foliolis lateribus basi antica brevioribus. *Tapia.*

*Crateva inermis.* *Fl. zeyl.* 211. *Sp. pl.* 1. p. 444. *Hort. cliff.* 484.

*Apioscorodon* f. *Arbor americana* triphyllos, allii odore, poma ferens. *Pluk. alm.* 34. t. 137. f. 7.

*Malus americana* trifolia, fructu pomi aurantii instar colorato. *Comm. hort.* 1. p. 129. t. 67.

*Anona* trifolia, flore stamineo, fructu sphaerico ferrugineo scabro minore, allii odore. *Sloan. jam.* 208. *hist.* 2. p. 169. *Raj. dendr.* 79.

*Nucifera.* *Rheed. mal.* 3. p. 49. t. 22.

*Habitat in Indiis.* †

3. CRATEVA spinosa, foliis serratis. *Fl. zeyl.* 212. *Marmelos.*

*Cucurbitifera* trifolia spinosa medica, fructu pulpa cydonii æmula. *Pluk. alm.* 125. t. 170. f. 5.

*Cydonia* exotica. *Bauh. pin.* 435.

*Balanus.* *Rumph. amb.* 1. p. 197. t. 81.

*Covalam.* *Rheed. mal.* 3. p. 37. t. 37.

*Habitat in India.* †

## TRIUMFETTA.

1. TRIUMFETTA foliis basi emarginatis. *Hort. cliff.* *Lappula.*

210. \* *Hort. upf.* 148. *Roy. lugdb.* 476.

*Triumfetta* floribus calyculatis, foliis semitrilobis. *Jacq. amer.* 22.

*Triumfetta* fructu echinato racemoso. *Plum. gen.* 40. ic. 255.

*Triumfetta* villosa, foliis inferioribus angulato-ovatis serrato dentatis, floribus ternatis: fasciculis geminatis. *Brown. jam.* 233.

*Lappula bermudensis* althæoides spicata, fructu orbiculari majore. *Pluk. alm.* 206. t. 245. f. 7.

*Agrimonia lappacea* inodora, folio subrotundo dentato. *Sloan. jam.* 92. *hist.* 1. p. 211. *Raj. juppl.* 247.

*Habitat in Jamaica, Brasilia, Bermudis.* †

- Bartramia.** 2. **TRIUMFETTA** foliis basi integris.  
*Triumphetta* floribus calyculatis, foliis rhombeo-ovatis. *Jacq. amer.* 22. *hist.* t. 90.  
**Bartramia.** *Fl. zeyl.* 174. *Sp. pl.* 1. p. 389.  
*Lappula benghalensis tetralpermos*, ribesii folio, echinis orbiculatis ad foliorum ortum plurimis sessilibus.  
*Pluk. alm.* 206. t. 41. f. 5.  
*Agrimonia maderaspatana*, folio rotundo singulari subtus incano. *Pet. gaz.* 31. t. 42. f. 10.  
*Lappago amboinica.* *Rumph. amb.* 6. p. 59. t. 25. f. 1.  
*Habitat in India.* ☉

## P E G A N U M.

- Harmala.** 1. **PEGANUM** foliis multifidis. *Hort. upf.* 144. *Gron. orient.* 165.  
*Peganum.* *Hort. cliff.* 206. *Hort. upf.* 144. *Roy. lugdb.* 461.  
*Ruta sylvestris*, flore magno albo. *Bauh. pin.* 336.  
**Harmala.** *Dod. pempt.* 121.  
*Habitat in arena Madriti, Alexandria, Cappadocia, Galatia.* 24
- daurica.** 2. **PEGANUM** foliis indivisis.  
*Harmala montana daurica perennis multicaulis*, polygalæ folio, flore albo. *Amm. ruth.* 91.  
 β. *Harmala montana*, polygalæ foliis, floribus luteis.  
*Amm. ruth.* 92.  
*Habitat in Sibiria.* 24

## N I T R A R I A.

- Schoberi.** 1. **NITRARIA.**  
*Osyris foliis obtusis.* *Hort. upf.* 295. *Gmel. fibir.* 2. f. 237. t. 98.  
*Casia fructu nigro.* *Amm. ruth.* 256.  
*Habitat ad Salfas aquas Vrunscinensis Lacus, Astrachanensis, Volgæ finitimis.* 5

## P O R T U L A C A.

- oleracea.** 1. **PORTULACA** foliis cuneiformibus, floribus sessilibus. *Roy. lugdb.* 473. *Gron. virg.* 59. *Hort. upf.* 146. *Mat. med.* 260. *Dalib. parif.* 158. *Hall. herb.* 392. *Sauv. monsp.* 168. *Brown. jam.* 233.  
*Portulaca foliis cuneiformibus verticillatis sessilibus floribus sessilibus.* *Hort. cliff.* 207.

Por-

*Portulaca angustifolia sylvestris. Bauh. pin. 288.*

4. *Portulaca latifolia sativa. Bauh. pin. 288.*

*Portulaca domestica. Lob. ic. 388.*

*Habitat in Europa, India, Ins. Ascensionis, America. ☉*

*Horologii floræ vigilias 11—12 habet.*

2. **PORTULACA** foliis subulatis alternis: axillis pilosis, pilosa. floribus sessilibus terminalibus.

*Portulaca foliis linearibus acutis, caule procumbente piloso. Roy. lugdb. 473.*

*Portulaca curassavica, angusto longo lucidoque folio, procumbens. Comm. hort. 1. p. 9. t. 5.*

*Portulaca corassavica lanuginosa procumbens. Herm. par. 215. t. 215.*

*Portulaca lanuginosa procumbens, vermiculata foliis, americana. Pluk. alm. 304. t. 247. f. 7. Volk. norib. 341. t. 341.*

*Anacampseros supina minor, foliis linearibus turgidis, floribus summis ramulis confertis. Brown. jam. 234.*

β. *Portulaca corassavica lanuginosa, psyllii folio, erectior & elatior, flore dilutius rubente. Herm. par. 214. t. 214.*

*Portulaca curassavica lanuginosa, kali folio, erectior & elatior. Pluk. alm. 303. t. 105. f. 4.*

*Habitat in America meridionali. ☉*

*Folia alterna, sed terminalia f. floralia octo in orbem digesta. Flores sessiles, terminales, calyce bifido, Staminiibus quindecim, Stylo quinquefido, Capsula circumscissa.*

3 **PORTULACA** foliis oblongis carnosiss, caule co- Halimoides. rymbofo, floribus sessilibus.

*Portulaca erecta, sedi minoris facie, capitulo tomentoso. Sloan. jam. 88. hist. 1. p. 205. t. 129. f. 3.*

*Halimus minimus, foliis oblongis succulentis tumetibus, summis ramulis densissime sitis. Brown. am. 206.*

*Habitat in Jamaica. ☉*

*Calyx diphyllus. Capsula circumscissa.*

4. **PORTULACA** foliis ovatis gibbis, pedunculo multifloro, caule fruticoso. Anacampseros.

*Anacampseros foliis ovatis convexis acutiusculis. Hort. upf. 146.*

*Anacampseros foliis acuminatis. Hort. cliff. 207. Roy. lugdb. 412.*

*Telephiastrum folio globofo. Dill. elth. 375. t. 281. f. 363.*

*Portulaca africana sempervirens, flore rubicundo Comm. hort. 2. p. 177. t. 89. Raj. suppl. 507.*

*Oxalis affinis planta aphylla mammillaris. Burm. afr. 76. t. 30. f. 2. vitiata.*

*Habitat ad Cap. b. Spei. §*

*Axillæ foliorum pilosæ sunt, & Folia succulenta conjungunt hanc cum Portulacis, quamvis stamina plurima & Capsula trivalvis. Calyx diphyllus est.*

racemosa.

5. **PORTULACA** foliis obovatis planiusculis, racemo simplici triquetro.

*Portulaca pedunculis multifloris triangularibus. Jacq. amer. 22.*

*Helianthemum frutescens, portulacæ folio. Plum. spec. 7. t. 150. f. 2.*

*Habitat in America ad maris littora. §*

*Calyx diphyllus. Capsula trilocularis.*

paniculata.

6. **PORTULACA** foliis ovatis planis, panicula racemosa.

*Portulaca floribus paniculatis. Jacq. amer. 22.*

*Portulaca americana latifolia erecta, floribus albis. Comm. hort. 1. p. 7. t. 4.*

*Portulacæ facie maritima fruticosa erecta americana. Sloan. jam. 87. hist. 1. p. 205.*

*Anacampseros foliis radicalibus mollibus ovatis glabris, scapo assurgenti paniculato. Brown. jam. 234.*

*Habitat in Americæ maritimis. §*

*Calyx pentaphyllus. Capsula trivalvis.*

## LYTHRUM.

Salicaria.

1. **LYTHRUM** foliis oppositis cordato-lanceolatis, floribus spicatis dodecandris.

*Lythrum foliis oppositis. Hort. cliff. 178. Fl. suec. 393; 422 Roy. lugdb. 458. Guett. stamp. 2. p. 122.*

*Lythrum floribus verticillatis. Fl. lapp. 197.*

*Lyfimachia spicata purpurea. Bauh. pin. 246.*

*Lyfimachia purpurea communis major. Cus. hist. 2 p. 51.*

3. Blat-

β. *Blattaria rubra spicata* major, folio subrotundo. *Marif.*

*hist. 2. p. 490. f. 5. t. 10. f. 11.*

γ. *Lythimachia trifolia spicata* purpurea. *Bocc. mus. 167.*

δ. *Lythimachia quadrifolia* purpurea, mollioribus & longioribus foliis. *Bocc. mus. 167.*

*Habitat in Europa ad ripas aquarum. 24.*

2. **LYTHRUM** foliis oppositis subtus tomentosis, floribus decandris, corollis calyce calyceque genitalibus brevioribus. fruticosum.

*Habitat in China. 5*

*Frutex cortice lacero. Folia opposita, sessilia, lanceolata, integerrima, subtus subtomentosa.*

3. **LYTHRUM** foliis oppositis ovalibus, floribus sessilibus alternis hexandris, caule diffuso. *Parsonia.*

*Parsonia herbacea, foliis ovatis oppositis, floribus singularibus foliis ad alterum latus interpolitis. Brown. jam. 199. t. 21. f. 2.*

*Habitat in Jamaica. 24.*

*Caules prostrati. Folia subsessilia, nuda, integerrima.*

*Racemi alterni. Flores solitarii, alterni, sessiles.*

4. **LYTHRUM** foliis oppositis subtus tomentosis subpetiolatis, floribus verticillatis lateralibus. verticillatum.

*Lythrum foliis oppositis, floribus verticillatis. Gron.*

*virg. 52.*

*Habitat in Virginia.*

5. **LYTHRUM** foliis oppositis linearibus petiolatis, floribus dodecandris. petiolatum.

*Lythrum foliis petiolatis. Gron. virg. 52.*

*Habitat in Virginia.*

6. **LYTHRUM** foliis oppositis linearibus, floribus oppositis hexandris. lineare.

*Lythrum foliis linearibus, floribus hexandris solitariis.*

*Gron. virg. 162.*

*Habitat in Virginia.*

*Caulis filiformis erectus angulatus, foliis oppositis linearibus, terminatus spica majore & duabus lateralibus minoribus compositis ex floribus oppositis remotis, sub singulis folio. Facies L. Hyssopifolia.*

7. **LYTHRUM** foliis oppositis ovatis, floribus alternis subdecandris, caule prostrato. *Melanium.*

*Linnaei Spec. plant. pars 1.*

*S 3*

*Me-*

*Melanium herbaceum reclinatum, foliis ovatis oppositis, floribus singularibus ad alas alternas. Brow. jam. 215.*

*Habitat in Jamaica. 24*

Hyssopifolia

8. **LYTHRUM** foliis alternis linearibus, floribus hexandris. *Hort. upf. 118.*

*Lythrum foliis alternis. Hort. cliff. 178. Sauv. monsp. 53. Roy. lugub. 458. Gort. gelr. 271.*

*Salicaria hyssopi folio latiore. Hall. jen. 147. t. 6. f. 3.*

*Hyssopifolia. Bauh. pin. 218.*

β. *Lythrum rubra non filiquosa. Bauh. pin. 246.*

γ. *Lythrum linifolia purpureo-cerulea. Bauh. pin. 246.*

*Habitat in Germaniæ, Helvetiæ, Angliæ, Galliæ inundatis. ☉*

Thymifolia. 9. **LYTHRUM** foliis alternis linearibus, floribus tetrapetalis.

*Lythrum foliis linearibus. Sauv. monsp. 49.*

*Salicaria minima tenuifolia. Tournef. inst. 254.*

*Salicaria hyssopi folio angustiore. Buxb. hal. 290. \**

*Hyssopifolia minor s. potius thymifolia maritima. Buxb. hist. 3. p. 792.*

*Polygonum aquaticum minus. Barr. ic. 773. f. 2.*

*Habitat in Italiæ & G. Narbonensis uliginosis. ☉*

*Petala 4, stamina 2, calyx adentatus. Folia linearilanceolata, alterna, ima opposita. Flores solitarii axillares. Calyx stipulis 2 lateralibus falcatus. D. Sauvages.*

virgatum.

10. **LYTHRUM** foliis oppositis lanceolatis, panicula virgata, floribus dodecandris.

*Salicaria orientalis, salicis folio acutissimo & glabro.*

*Tournef. cor. 18.*

*Salicaria glabra ramosissima, longissimis & angustissimis foliis. Amm. ruth. 89.*

*Salicaria glabra, longioribus & angustioribus foliis. Amm. ruth. 88.*

*Habitat in Sibiria, Tataria.*

*Caulis paniculatus, ramis alternis longis. Flores 2. f. 3. pedicellati, ex singula ala alterna racemi.*

## GINORA.

Americana. 1. **GINORA.** Jacq. amer. 22.

Ha.

*Habitat in America. ♀*  
*Frutex pulcher halitu Myrti.*

## DIGYNIA.

### HELIOCARPUS.

1. *HELIOCARPUS. Hort. cliff. 211. t. 16. \* Trew. americana; chret. t. 43.*

*Habitat in Vera Cruce. ♀*

### AGRIMONIA.

1. *AGRIMONIA foliis caulinis pinnatis: impari petio- Eupatoriā:  
 lato, fructibus hispidis. Hort. cliff 179. Fl. svec. 394;  
 423. Hort. upf. 118. Mat. med. 221. Gron. virg. 53.  
 Roy. lugdb. 240. Hall. helv. 407. Guett. stamp. 1: p.  
 293. Dalib. parif. 139.*

*Eupatorium veterum f. Agrimonia. Bauh. pin. 321.*

*Eupatorium. Fuchf. hist. 244. Cam. epit. 756. Mill. ic. 15.*

*Habitat in Europæ pratis apricis argillaceis. 4*

2. *AGRIMONIA foliis caulinis pinnatis: impari sessili, repens;  
 fructibus hispidis.*

*Agrimonia orientalis humilis, radice crassissima repen-  
 te, fructu in spicam brevem & densam congesto.*

*Tournef. cor. 21.*

*Habitat in Oriente.*

*Affinis nimium A. Eupatoriæ, sed Caules breviores,  
 crassiores, hirsuti. Stipulæ longitudine interstitio-  
 rum, reflexæ, fere totum obumbrant caulem. Fo-  
 liorum impar non pedicellatum. Spica brevis, den-  
 sa, sessilis, Bracteis longitudine florum.*

3. *AGRIMONIA foliis caulinis ternatis, fructibus gla- Agrimonoid-  
 bris. Hort. cliff. 179. Roy. lugdb. 241. des.*

*Agrimariæ similis. Bauh. pin. 321.*

*Agrimonoides. Col. ceph. 1. p. 145. t. 144. Boerh.  
 lugdb. 1. p. 179.*

*Habitat in Italiæ nemoribus umbrosis humentibus. 4*

*Stamina sæpe vix ultra 8.*

## TRIGYNIA.

### RESEDA.

1. *RESEDA foliis lanceolatis integris, calycibus quadri- Luteola:  
 fidis.*



*Reseda foliis simplicibus lanceolatis integris. Hort. cliff.*

212. *Vir. cliff.* 49. *Fl. suec.* 439; 424. *Roy. lugd.*

483. *Dalib. parif.* 160.

*Luteola herba salicis folio. Bauh. pin.* 100.

*Lutum herba. Dod. pempt.* 80.

*Habitat in Europa ad vias & pagos. ☉*

*Folia ad basin utrinque denticulum habent callosum.*

canescens.

2. RESEDA foliis subulatis sparsis. *Sauv. monsp.* 48.

*Reseda foliis simplicibus oblongis integris. Guett. stamp.*

1. p. 225.

*Reseda alba minor. Bauh. pin.* 100.

*Sesamoides salmanticum parvum. 2. Clus. hist.* 1. p.

296. t. 295.

*Habitat in Salmantica.*

glauca.

3. RESEDA foliis linearibus basi dentatis, floribus tetragynis.

*Reseda linariæ foliis. Bauh. pin.* 100. *prodr.* 41. *Burj.* IV. 88.

*Reseda, linariæ folio, glaucum pyrenaicum, flore stamineo. Morif. hist.* 3. p. 601. f. 15. t. 1. f. 4.

*Reseda, linariæ folio, pyrenaica, glauco folio. Fluk. alm.* 317. t. 107. f. 2. *Raj. suppl.* 511.

*Luteola pumila pyrenaica, linariæ folio. Tournesf. inf.* 424.

*Habitat in Pyrenæis.*

purpurascens.

4. RESEDA foliis linearibus obtusis, floribus pentagynis.

*Reseda alba minor. 1. Bauh. pin.* 100. *Raj. hist.* 1054.

*Sesamoides salmanticum parvum. 1. Clus. hist.* 1. p. 296.

*Sesamoides foliis crassius, floribus ex herbaceo purpurascentibus. Tournesf. inst.* 424.

*Habitat in collibus Salmanticis, Montspelli.*

Sesamoides.

5. RESEDA foliis lanceolatis integris, fructibus stellatis,

*Sesamoides fructu stellato. Tournesf. inst.* 424.

*Habitat Montspelli in Horto Dei.*

undata.

6. RESEDA floribus trigynis tetragynisque, calycibus quinquepartitis, foliis pinnatis undulatis.

*Reseda minor, foliis incis. Barr. rar.* 78. t. 588.

*Habitat in Hispania. 2.*

*Caulis pedalis, striato-angulatus, strictus. Radix radicalis & caulina pinnata: foliolis lanceolatis, de-*

*currentibus, acutis, undatis: alternis minoribus, erectioribus. - Superiora folia excludunt foliola alterna minora. Racemi terminales. Calyx 5partitus. Petala 5, trifida, alba. Antheræ luteæ, 10. Pistilla in aliis flosculis 3, in aliis 5. Capsulæ hujus generis maximæ.*

7. RESEDA foliis pinnatis apice recurvis, floribus suffruticoso-  
tragynis, calycibus quinquepartitis patentibus, loba-  
caule basi suffruticoso.

*Habitat in Hispania. 24*

*Radix f. caulis supra terram perennans, crassitie sæpe  
dupla pollicis, brevissimus, e cujus apice Caules  
plures, adscendentes, læves striis elevatis. Folia pin-  
nata: pinnis 5 f. 7, lanceolatis, decurrentibus, se-  
re undatis, lævibus, apice recurvis. Spicæ plures,  
alternæ, versus caulis apicem primores. Petala 6 f. 5,  
alba, trifida, calyce majora. Stamina 17, quasi me-  
dia inter R. undatam & albam.*

8. RESEDA foliis pinnatis, floribus tetrágynis, calyci- alba.  
cibus sexpartitis, *Hort. upf. 149.*

*Reseda foliis pinnatis integerrimis. Hort. cliff. 212.*

*Reseda maxima. Bauh. pin. 100. Lob. hist. 110.*

*Habitat Monspeli, in Hispania. ☉*

9. RESEDA foliis omnibus trifidis: inferioribus pinna- lutea.  
tis, *Hort. cliff. 212. Hort. upf. 149. Roy. lugdb. 483.*

*Hall. herb. 315. Dalib. paris. 159. Gort. gelr. 310.*

*Reseda vulgaris. Bauh. pin. 100. Raj. hist. 1053.*

*Reseda lutea. Bauh. hist. 3. p. 467.*

- β. Reseda gallica crispata. *Bocc. sic. 77. t. 41. f. 3.*

*Reseda massiliensis, foliis latioribus crispis. Pluk. alm.*

*317. t. 55. f. 4*

*Habitat in Europæ australioris montibus cretaceis. ☉*

10. RESEDA foliis integris trilobisque, calycibus sex- Phyteuma.  
partitis maximis, *Hort. cliff. 412. Dalib. paris. 159.*

*Reseda floribus trigynis, calycibus maximis hexaphyl-  
lis. Hort. upf. 150.*

*Reseda minor vulgaris. Tournef. inst. 423.*

*Reseda affinis Phyteuma. Bauh. pin. 100. prodr. 42.*

*t. 42.*

*Erucago apula trifida quinquefolia. Col. ceph. 267. t. 269.*

β. *Reseda minor vulgaris, folio minus inciso. Tournef. inst. 423.*

γ. *Reseda minor vulgaris, foliis integris. Tournef. inst. 423.*

*Phyteuma. Bauh. pin. 3. p. 386. fig.*

*Habitat in Gallia, Italia, Oriente. ☉*

odorata.

II. *RESEDA foliis integris trilobisque, calycibus florem æquantibus.*

*Reseda foliis integris trilobisque, floribus tetragynis. Mill. dict. 1. 217.*

*Reseda foliis integris, floribus odoratis. Hall. gen. 95. Zinn. gatt. 123.*

*Habitat in Ægypto. ♂*

*Simillima R. Phyteumæ, sed Folia magis undulata.*

*Calyces corolla non longiores. Rami declinati, setis vagis adpersi. Flores tristes ambrosiaci, sæpius tetragyni. Petalorum lacinie laterales in hac, intermedia vero in præcedenti, breviores.*

## EUPHORBIA,

\* *Fruticosa aculeata.*

antiquorum. I. *EUPHORBIA aculeata subnuda triangularis articulata: ramis patentibus. Hort. cliff. 196. \* Hort. upf. 138. Fl. zeyl. 199. Mat. med. 154. Roy, lugdb. 194. Amæn. acad. 3. p. 106. \**

*Euphorbium antiquorum verum. Comm. hort. 1. p. 23. t. 12.*

*Schadidacalli. Rheed. mal. 2. p. 81. t. 42.*

β. *Euphorbia aculeata nuda triangularis articulata, ramis erectis. Mill. dict.*

*Euphorbium trigonum & tetragonum spinosum, ramis compressis. Jsn. afl. 1720.*

*Tithymalus aizoides triangularis & quadrangularis articulatus & spinosus, ramis compressis. Comm. præf. 55. t. 5.*

*Habitat in India. ♀*

canariensis. α. *EUPHORBIA aculeata nuda subquadrangularis, aculeis geminatis. Hort. cliff. 196. \* Hort. upf. 138. Roy, lugdb. 194. Amæn. acad. 3. p. 107. \**

*Tithymalus aizoides fruticosus canariensis aphyllus*

qua-

quadrangularis & quinquangularis, spinis geminis aduncis atronitentibus armatus. *Comm. hort.* 2. p. 207. t. 104.

*Tithymalus aizoides lactifluus* f. *Euphorbia canariensis* quadrilatera & quinquelatera, cerei effigie. *Pluk. alm.* 370. t. 320. f. 2.

*Habitat in Canariis.* §

3. *EUPHORBIA* aculeata nuda septemangularis: spinis heptagona. solitariis subulatis floriferis. *Hort. cliff.* 196. \* *Roy. lugdb.* 194. *Amæn. acad.* 3. p. 109. \*

*Euphorbium heptagonum*, spinis longissimis in apice frugiferis. *Bærh. lugdb.* 1. p. 258. t. 258.

*Euphorbium capense*, spinis longis simplicibus. *Bradl. fvec.* 2. p. 4. t. 13.

*Habitat in Æthiopia.* §

4. *EUPHORBIA* aculeata nuda: angulis tuberosis spinis interstinctis. *Amæn. acad.* 3. p. 108. mamilla-  
ris.

*Euphorbium polygonum*, aculeis longioribus ex tuberculorum internodiis prodeuntibus. *Isn. aët.* 1720. p. 386.

*Tithymalus aizoides africanus*, validissimis spinis ex tuberculorum internodiis provenientibus. *Comm. præl.* 59. t. 9.

*Habitat in Æthiopia.* §

*Differt ab Heptagona angulis duplicatis torulosis; spinæ solitariae inter singula tubercula longitudinaliter digesta. Flores non vidi; an ex spinarum apice, ut heptagonæ?*

5. *EUPHORBIA* aculeata nuda multangularis, spinis cereiformis. solitariis subulatis. *Roy. lugdb.* 195. *Amæn. acad.* 3. p. 108.

*Euphorbium cerei effigie*, caulibus gracilioribus. *Boerh. lugdb.* 1. p. 258.

*Euphorbium aphyllum angulosum*, florum coma densissima. *Burm. afr.* 19. t. 9. f. 3.

*Tithymalus africanus spinosus*, cerei effigie. *Morif. hist.* 3. p. 245. *Pluk. alm.* 370. t. 231. f. 1.

*Habitat in Æthiopia.* §

6. *EUPHORBIA* aculeata nuda multangularis: aculeis officinarum. geminatis. *Hort. cliff.* 196. \* *Hort. upf.* 138. *Roy. lugdb.* 195. *Amæn. acad.* 3. p. 107. \*

*Euphorbium polygonum spinosum, cerei effigie. Isnard. aët. 1720. p. 500 t. 10*

*Euphorbium cerei effigie, caulibus crassioribus spinis validioribus armatum. Comm. hort. 1. p. 21. t. 11. Seb. thes. 1. p. 29. t. 19. f. 2.*

*Euphorbium. Bauh. pin. 387.*

*Habitat in Æthiopia & Africa calidiore. §*

*neriifolia.*

7. *EUPHORBIA aculeata seminuda: angulis oblique tuberculatis. Hort. cliff. 196. \* Hort. upf. 139. Fl. zeyl. 200. Roy. lugdb. 194. Amæn. acad. 3. p. 109. \**

*Euphorbium afrum spinosum, foliis latoribus non spinosis. Seb. thes. 1. p. 18. t. 9. f. 1.*

*Tithymalus aizoides arborescens spinosus, caudice angulari, nerii folio. Comm. præl. 22. 56. t. 6. Bradd. succ. 3. p. 10. t. 28.*

*Tithymalus indicus spinosus, nerii folio. Comm. hort. 1. p. 25. t. 13.*

*Ligularia Rumph. amb. 4. p. 88. t. 40.*

*Ela Calli. Rheed. mal. 2. p. 83. t. 43.*

*Habitat in India. §*

*\* \* Fruticosa inermes.*

*(Caulis nec dichotomus, nec umbelliferus.)*

*Caput me-  
dusa.*

8. *EUPHORBIA inermis imbricata tuberculis foliolo lineari instructis. Hort. cliff. 197. Hort. upf. 139. Roy. lugdb. 195. Amæn. acad. 3. p. 110.*

*Planta lactaria africana. Comm. hort. 1. p. 33. t. 17.*

*Tithymalus aizoides africanus, simplici squamato caule, chamænerii folio. Comm. præl. 58. t. 8. & p. 23. 57. t. 7.*

β. *Euphorbium anacantum squamosum, lobis florum tridentatis. Isnard. aët. 1720. p. 502. t. 11. Breyn. prodr. 3. p. 29. t. 19.*

γ. *Euphorbium erectum aphyllum, ramis rotundis, tuberculis tetragonis. Burm. afr. 16. t. 7. f. 2.*

δ. *Euphorbium humile procumbens, ramis simplicibus copiosis, caule crassissimo tuberoso. Burm. afr. 20. t. 10. f. 1.*

ε. *Euphorbium procumbens, ramis plurimis simplicibus squamosis, foliis deciduis. Burm. afr. 17. t. 8.*

ζ. *Euphorbium procumbens, ramis geminatis, caule glabro oblongo cinereo. Burm. afr. 18. t. 9. f. 1.*

*Habitat in Æthiopia. ‡*  
*Distinctissima planta petalis palmatis.*

9. EUPHORBIA inermis nuda fruticosa filiformis vominalis, lubilis: cicatricibus oppositis. *Hort. cliff. 197. \* Hort. upf. 140. Amæn. acad. 3. p. 110. \**  
 Felfel Tavi. *Alp. ægypt. 190. t. 190. Dill. elth. 368.*  
*Habitat in Africæ maritimis. ‡*
10. EUPHORBIA inermis seminuda fruticosa filiformis flaccida, foliis alternis. *Hort. cliff. 197. Hort. upf. 140. Roy. lugdb. 195. Amæn. acad. 3. p. 111. \* Gron. orient. 160.*  
*Tithymalus aphyllus mauritanicæ. Dill. elth. 384. t. 289. f. 373.*  
*Habitat in Africæ maritimis. ‡*
11. EUPHORBIA inermis seminuda fruticosa filiformis erecta, ramis patulis determinate confertis. *Hort. cliff. 197. Hort. upf. 139. Fl. zeyl. 197. Roy. lugdb. 195. Amæn. acad. 3. p. 111. \**  
*Tithymalus indicus frutescens. Comm. hort. 1. p. 27. t. 14.*  
*Tithymalus arborecens, caule aphylo. Pluk. phyt. 319. f. 9.*  
*Ossifraga lactea. Rumph. amb. 7. p. 62. t. 29.*  
*Tiru Calli. Rheed, mal. 8. p. --- t. 44.*  
*Habitat in India. ‡*
12. EUPHORBIA inermis fruticosa, foliis distiche alternis ovatis. *Hort. cliff. 198. \* Roy. lugdb. 195. des. Amæn. acad. 3. p. 111. \**  
*Tithymalus curassavicus myrtifolius, flore coccineo myrtifolia mellifluo. Herm. par. 234. t. 234.*  
*Tithymalus curassavicus myrtifolius, flore papilionaceo coccineo parvo. Comm. hort. 1. p. 31. t. 16. Pluk. alm. 369. t. 230. f. 2.*
- β. Tithymaloides laurocerasi folio non serrato. *Dill. elth. padifolia. 383. t. 288 f. 374.*  
*Habitat in Curassao, at β. forte in India. ‡*
13. EUPHORBIA inermis, foliis serratis petiolatis heterophyllis difformibus ovatis lanceolatis panduriformibus. *Amæn. acad. 3. p. 112.*  
*Tithymalus heterophyllus. Plum. ic. 250. t. 251. f. 3.*

*Tithymalus curassavicus*, salicis & atriplicis foliis variis, caulibus viridantibus. *Pluk. alm.* 369. t. 12. f. 6.  
*Tithymalus curassavicus*, salicis & atriplicis foliis hirsutis, caulibus subrubentibus. *Morif. hist.* 3. p. 336.  
*Habitat in America calidiore.*

cotinaifolia. 14. *EUPHORBIA* foliis oppositis subcordatis petiolatis, emarginatis integerrimis, caule fruticoso. *Amæn. acad.* 3. p. 112.

*Euphorbia inermis*, caule fruticoso, foliis oppositis subcordatis emarginatis, petiolis folio longioribus. *Hort. cliff.* 198. *Roy. lugdb.* 196.

*Tithymalus curassavicus*, folio cotini triphyllis, petalis florum serratis. *Pluk. alm.* 369. t. 230. f. 3.

*Tithymalus arboreus curassavicus*, cotini folio. *Seb. thes.* 1. p. 75. t. 46. f. 4.

*Tithymalus arboreus americanus*, cotini folio. *Comm. hort.* 1. p. 29. t. 15.

*Habitat in Curassao.* ♀

oeymoides. 15. *EUPHORBIA* inermis herbacea ramosa, foliis subcordatis integerrimis petiolo brevioribus, floribus solitariis. *Amæn. acad.* 3. p. 112.

*Euphorbia inermis*, foliis subcordatis obtusis integerrimis petiolatis, caule ramoso erecto. *Roy. lugdb.* 199.  
*Habitat in Campechia. Houst.*

\* \* \* *Dichotomæ* (umbella bifida aut nulla)

origanoides. 16. *EUPHORBIA* dichotoma, foliis serrulatis ovatis obtusis trinerviis, panicula terminali, caulibus simplicibus. *Amæn. acad.* 2. p. 114. \*

*Habitat in Insula Ascensionis.* 2

*Planta refert ita Origanum, ut primo intuitu nequeat non pro eo agnosc.*

hypericifolia. 17. *EUPHORBIA* dichotoma, foliis serratis ovali-oblongis glabris, corymbis terminalibus, ramis divaricatis. *Amæn. acad.* 3. p. 113. \*

*Euphorbia inermis*, foliis ovalibus oppositis hinc serratis uniformibus, ramis alternis, caule erectiusculo. *Hort. cliff.* 198. *Hort. ups.* 143. \* *Roy. lugdb.* 196.

*Euphorbia minima reclinata*, foliolis ovatis denticulatis, floribus subumbellatis terminalibus lateralibusque. *Brown. jam.* 235.

*Tithy-*

*Tithymalus americanus*, flosculis albis. *Comm. prel.*  
60. t. 50.

*Tithymalus erectus* acris, parietariae foliis glabris, floribus ad caulium nodos conglomeratis. *Sloan. jam.*  
82. *hist.* 1. p. 197. t. 126.

*Habitat in India.* ☉

18. *EUPHORBIA* *dichotoma*, foliis ferrulatis ovatis acuminatis, pedunculis capitatis axillaribus, caulibus pilosis *Amæn. acad.* 3. p. 114. \* hist.

*Euphorbia inermis*, foliis oppositis ovalibus serratis uniformibus, pedunculis capitatis axillaribus. *Fl. zeyl.*  
197. \*

*Euphorbia reclinata* minor subhirsuta, foliis serratis. *Brown. jam.* 234.

*Tithymalus botryoides zeylanicus*, cauliculis villosis. *Burm. zeyl.* 223. t. 104.

*Efula esculenta*. *Rumph. amb.* 6. p. 54. t. 23. f. 2.

*Habitat in India.* ☉

19. *EUPHORBIA* *dichotoma*, foliis serratis ovali-oblongis, pedunculis bicapitatis axillaribus, caule erecto. *Amæn. acad.* 3. p. 115. pilulifera.

*Tithymalus botryoides erectus*, florum capitulis conjugatis & longiori pediculo insidentibus. *Burm. zeyl.*  
224. t. 105. f. 1. *Pet. gal.* t. 80. f. 14.

*Habitat in India.* ☉

20. *EUPHORBIA* *dichotoma*, foliis subcrenatis linearibus, floribus fasciculatis terminalibus, caule erecto. hylophifolia.

*Euphorbia dichotoma erecta* tenuis, foliis linearibus, floribus quasi umbellatis terminalibus. *Brown. jam.*  
235.

*Habitat in America.*

21. *EUPHORBIA* *dichotoma*, foliis serratis ovali-oblongis, capitulis axillaribus glomeratis subsessilibus, caulibus procumbentibus. *Amæn. acad.* 3. p. 115. thymifolia.

*Tithymalus* foliis oppositis oblique cordatis obtusis serratis, pedunculis multifloris. *Fl. zeyl.* 198.

*Tithymalus indicus* annuus dulcis, floribus albis, caulibus viridantibus & rubentibus. *Pluk. alm.* 372. t. 113. f. 2.

β. *Tithymalus humilis* ramosissimus hirsutus, foliis thymiferratis. *Burm. zeyl.* 225. t. 105. f. 3.

*Habitat in India.* ☉

22. EU-



- maculata*. 22. **EUPHORBIA** *dichotoma*, foliis ferratis oblongis pilosis, floribus axillaribus solitariis, ramis patulis. *Euphorbia minima* supina rufescens. *Brown. jam.* 236. *Tithymalus* f. *Chamæfyce* altera virginiana, foliis crenatis & macula fusca eleganter notatis. *Pluk. alm.* 372. t. 65. f. 8.  
*Chamæfyce. Sloan. jam.* 83. hist. 1. p. 198.  
*Habitat in America septentrionali.* ☉  
 Caules dichotomi: Ramis alternis, patentibus, supra purpurascens. Folia ovali-oblonga, trinervia, subpilosa, serrata, altero latere maxima parte integerrima, tenera adhuc planta notata macula fusca. Flores axillares, solitarii, parvi, calyce rufo. Fructus laevis.
- parviflora*. 23. **EUPHORBIA** *subdichotoma*, foliis ferratis oblongis glabris, floribus solitariis, caule erectiusculo alternis ramofo.  
*Tithymalus erectus*, floribus rarioribus, foliis oblongis glabris integris. *Burm. zeyl.* 224. t. 105. f. 2.  
*Habitat in India.* ☉
- canescens*. 24. **EUPHORBIA** *dichotoma* foliis integris subrotundis pilosis, floribus solitariis axillaribus, caulibus procumbentibus.  
*Tithymalus exiguus villosus nummulariae folio.*  
*Tournef. inst.* 87.  
*Habitat in Hispania.* ☉  
*Affinis nimis E. Chamæfyces, sed tota undique albo-villosa. Folia obsolete emarginata, obsolete crenata.*
- Chamæfyce*. 25. **EUPHORBIA** *dichotoma*, foliis crenulatis subrotundis glabris, floribus solitariis axillaribus, caulibus procumbentibus. *Amæn. acad.* 3. p. 115. \*  
*Euphorbia inermis*, foliis oblique cordatis serrulatis uniformibus, ramis alternis, floribus solitariis. *Hort. cliff.* 198. \* *Roy. lugdb.* 196. *Gmel. fib.* 2. p. 237. *Groen. orient.* 160.  
*Chamæfyce. Bauh. pin.* 293. *Clus. hist.* 2. p. 187.  
*Habitat in Europæ australis, Sibiriae, Mesopotamiae aridis.* ☉  
*Fructus subhirsutus.*
- Pepilis*. 26. **EUPHORBIA** *dichotoma*, foliis integerrimis semicordatis, floribus solitariis axillaribus, caulibus procumb-

cumbentibus. *Amæn. acad. 3. p. 115. Gron. orient. 163.*

*Euphorbia foliis subrotundis obtusis, floribus solitariis in foliorum alis. Guett. stamp. 2. p. 420.*

*Peplis maritima, folio obtuso. Bauh. pin. 293.*

*Peplis. Clus. hist. 2. p. 187. Cam. epit. 970.*

*Peplion. Dalech. hist. 970.*

*Habitas in Narbonæ, Hispaniæ maritimis. ☉*

27. **EUPHORBIA** foliis oppositis integerrimis lanceo- polygoni-  
latis obtusis, floribus solitariis axillaribus, caulibus tia.  
procumbentibus. *Amæn. acad. 3. p. 116. \**

*Euphorbia procumbens, ramulis alternis, foliis lanceo-  
lato-linearibus, floribus solitariis. Gron. virg. 58.*

*Euphorbia minima ramosissima angustifolia. Raj. suppl. 431.*

*Habitat in Canada, Virginia. ☉*

28. **EUPHORBIA** dichotoma, foliis integerrimis lan- Ipecacuan-  
ceolatis, pedunculis axillaribus unifloris folia æquan- ha,  
tibus, caule erecto. *Amæn. acad. 3. p. 117. \**

*Tithymalus, flore exiguo viridi apicibus flavis, ante-  
quam folia emittit florens. Gron. virg. 58.*

*Habitat in Virginia, Canada. 24*

29. **EUPHORBIA** dichotoma, foliis integerrimis ova- portulacol-  
libus retusis, pedunculis axillaribus unifloris folia des.  
æquantibus, caule erecto. *Amæn. acad. 3. p. 117. \**

*Tithymalus perennis, portulacæ folio. Few. peruv. 707.  
t. 2?*

*Habitat in Philadelphia. Kalm. 24*

30. **EUPHORBIA** dichotoma, foliis integerrimis sub- myrsifolia.  
rotundis emarginatis subtus incanis, floribus solita-  
riis, caule erecto.

*Euphorbia erecta, foliolis ovatis oppositis, ramulis te-  
nuibus alaribus. Brown. jam. 235.*

*Habitat in Jamaica. ☉*

\* \* \* \* *Umbella trifida.*

31. **EUPHORBIA** umbella trifida : dichotoma, invo- Peplus.  
lucellis ovatis, foliis integerrimis obovatis petiolatis.  
*Amæn. acad. 3. p. 117. \**

*Euphorbia foliis obverse ovatis integerrimis, umbella  
universalis trifida triphylla, partialibus dichotomis di-  
phyl.*

phyllis. *Hort. cliff.* 199. *Fl. fvec.* 437 ; 426. \* *Roy. lugdb.* 197. *Dalib. parif.* 156.

Peplus f. *Esula* rotunda. *Bauh. pin.* 292.

Peplus. *Fuchf. hist.* 603. *Dod. pempt.* 375.

*Habitat in Europæ cultis oleraceis.* ☉

falcata.

32. EUPHORBIA umbella trifida: dichotoma, involucellis cordato-subfalcatis acutis, foliis lanceolatis obtusiusculis. *Amén. acad.* 3. p. 118. \*

*Habitat in Europa australi.*

terraccina.

33. EUPHORBIA umbella quadrifida dichotoma, foliis alternis lanceolatis retusis mucronatis. *Tithymalus marinus*, folio retuso, terracinus. *Barr. ic.* 823.

*Habitat in Hispania. Alstræmer.* ☉

Caulis herbaceus, ultra semipedalis, teres. Folia alternata, lanceolata, lævia, obliqua, margine scaberrimula, apice retusa & quasi truncata mucrone reflexo. Involucrum subquadriphyllum: foliis oblongo-ovatis, obtusis, folio latioribus, vix serratis. Involucella ovata, basi truncata. Petala flavescentia, b. f. tridentata. Fructus glabri. Ex inferioribus alis foliorum ramuli steriles. Figura Foliorum ad *Euph. exigua* retusam prope accedit, sed in illa lineam lata, huic digiti magnitudinis.

exigua.

34. EUPHORBIA umbella trifida: dichotoma, involucellis lanceolatis, foliis linearibus. *Amén. acad.* 3. p. 118. \*

acuta.

- α. Euphorbia inermis, foliis alternis linearibus acutis, umbella universali trifida, partialibus dichotomis diphyllis. *Hort. cliff.* 199. *Hort. upf.* 143 \* *Roy. lugdb.* 197. *Dalib. parif.* 157. *Gort. gelr.* 304.

*Tithymalus* f. *Esula* exigua. *Bauh. pin.* 291.

*Esula* minima tragi. *Dalech. hist.* 1656.

retusa.

- β. *Tithymalus* f. *Esula* exigua, foliis obtusis. *Bauh. pin.* 291. *prodr.* 132.

- γ. *Tithymalus* exiguus saxatilis. *Bauh. pin.* 291. *prodr.* 132. *Magn. monsp.* 259. t. 258. *Burf. XVI.* 33.

*Habitat in Lusatia, Gallia, Helvetia, Hispania inter segetes; β. Montpelii in saxosis. γ. Paduæ & Massiliæ: α. Acuta in collibus, Retusa in pratis.* ☉

suberosa.

35. EUPHORBIA umbella trifida, involuero tetraphyllo

lo, caule nudo, foliis oblongis emarginatis. *Amæn. acad. 3. p. 117.*

*Euphorbia inermis*, foliis oblongis obtusis emarginatis. *Roy. lugdb. 199.*

*Tithymalus humilis*, folio lapathi. *Ruxb. cent. 2. p. 27. t. 23.*

*Tithymalus tuberosus acaulos*, foliis oblongis cucullatis & planis. *Burm. afr. 9. t. 4.*

*Tithymalus africanus humilis*, foliis latioribus oblongis tuberosa radice. *Raj. suppl. 933.*

*Habitat in Aegypto & Aethiopia. †*

\* \* \* \* \* *Umbella quadrifida.*

36. *EUPHORBIA umbella quadrifida*: dichotoma, foliis oppositis integerrimis. *Amæn. acad. 3. p. 119. \* Lathyris.*

*Euphorbia inermis*, foliis oppositis lanceolatis integerrimis, umbella universali quadrifida tetraphylla, ulterioribus dichotomis. *Hort. upf. 140. Mat. med. 256.*

*Euphorbia inermis*, foliis oppositis lanceolatis, umbella universali trifida polyphylla, partialibus triphyllis, reliquis diphyllis. *Hort. cliff. 198. Roy. lugdb. 196. Dalib. parif. 155.*

*Lathyris major. Bauh. pin. 293.*

*Lathyris. Cam. epit. 968. Fuchf. hist. 454.*

*Habitat in Gallia, Italia ad agrorum margines. †*

\* \* \* \* \* *Umbella quinquefida.*

37. *EUPHORBIA umbella subquinquefida simplicis, spinosa.* involucrellis ovatis: primariis triphyllis, foliis oblongis integerrimis, caule fruticoso. *Amæn. acad. 3. p. 120. \**

*Euphorbia inermis fruticosa*, foliis lanceolatis integerrimis, floribus solitariis terminalibus, involucrellis triphyllis. *Hort. cliff. 201. Roy. lugdb. 196.*

*Euphorbia inermis fruticosa*, ramis siccis pungens, foliis ad umbellas ternis sæpius ovatis, floribus solitariis. *Sauv. monsp. 51.*

*Tithymalus ragulinus*, flore luteo pentapetalo. *Herm. lugdb. 600. t. 601.*

*Tithymalus maritimus spinosus. Bauh. pin. 291.*

*Habitat in Cretæ & Galloprovinciæ montibus maritimis. †*

*Rami senescentes fœci & aridi persflunt, unde quasi spinosa apparet.*

- Apios.** 38. EUPHORBIA umbella quinquefida: bifida, involucellis obcordatis. *Amæn. acad.* 3. p. 120.  
*Tithymalus tuberosa* pyriformi radice. *Bauh. pin.* 292.  
*Tithymalus tuberosa* radice. *Clus. hist.* 2. p. 290.  
*Habitat in Creta.*
- epithymio-**  
**des.** 39. EUPHORBIA umbella quinquefida: bifida, involucellis ovatis, foliis lanceolatis obtusis subtus villosis.  
*Tithymalus epithymi* fructu. *Col. ecphr.* 1. p. 52. t. 51.  
*Peplios* altera species. *Bauh. pin.* 292.  
*Habitat in Italia.* *D Jacquin.* 24.  
*Accedit ad E. dulcem, sed Folia magis margine scabra*  
*Et subtus villosa. Fructus hispidi, sevis subulnis*  
*purpurascentibus adpersi.*
- dulcis.** 40. EUPHORBIA umbella quinquefida: bifida, involucellis subovatis, foliis lanceolatis obtusis integerrimis. *Amæn. acad.* 3. p. 122.  
*Tithymalus montanus* non acris. *Bauh. pin.* 292.  
*Pithyusa* f. *Efula* minor altera, floribus rubris. *Lch.*  
*ic.* 358.  
*Habitat in Galliæ, Italiæ, Germaniæ umbrosis.*  
*Exsiccatione nigra evadit.*
- Pithyusa.** 41. EUPHORBIA umbella quinquefida: bifida, involucellis ovatis mucronatis, foliis lanceolatis: infimis involutis retrorsum imbricatis. *Amæn. acad.* 3. p. 122.  
*Tithymalus* foliis brevibus aculeatis. *Bauh. pin.* 292.  
*Pithyusa.* *Dalech. hist.* 1652.  
*Tithymalus maritimus, juniperi folio.* *Bocc. sic.* 9. t. 5.  
*Morif. hist.* 3. p. 337. f. 10. t. 1. f. 25.  
*Habitat in arenosis Belgii, Hispaniæ, Italiæ, Ma-*  
*liæ.* 24.
- perlandica.** 42. EUPHORBIA umbella quinquefida: dichotoma, involucellis subcordatis concavis, foliis lineari lanceolatis acutis glabris patentibus. *Huds. angl.* 152.  
*Tithymalus montanus, esulæ folio, minor italicæ.*  
*Barr. ic.* 822.  
*Tithymalus maritimus minor.* *Raj. syn.* 3. p. 313. t. 24. f. 6.  
*Habitat in Angliæ Devonshire.* 5.  
*Caules Fruticulosi, palmares, glabri, teretes, rubicundi.*  
*Folia alterna, subseffilia, lineari-lanceolata, mucro-*  
*na-*

nata, glabra, patentia, basi subtus rubra. Ramuli laterales: ex alis inferioribus steriles, demum ex-  
crescentes & caulem inumbrantes. Umbellæ termi-  
nales, quinquefidæ, dichotomæ, patentes. Involucrum  
foliis simile. Involucella subcordata, concava, mu-  
cronata. Flores subsessiles, lutei: Primi & secun-  
di masculi: Petalis muticis, obtusissimis; Reliqui  
hermaphroditi: Petalis bicornibus. Fructus glabri  
angulis muricatis.

43. **EUPHORBIA** umbella subquinquefida: bifida, in- Paralias.  
volucellis cordato-reniformibus, foliis sursum im-  
bricatis. *Amæn. acad.* 3. p. 129. *Huds. angl.* 183.

*Euphorbia inermis*, foliis setaceo-linearibus confertis,  
umbella universal multifida, partialibus ramosè bi-  
fidis. *Hort. cliff.* 200. *Roy. lugdb.* 193. *Gort. gelr.*  
306. *Gron. orient.* 162.

*Tithymalus maritimus*. *Bauh. pin.* 291. *Dod. pempt.*  
370. f. 1. 2.

*Tithymalus paralias*. *Cam. epit.* 962.

*Habitat in Europæ arena maritima.* 24

44. **EUPHORBIA** umbella quinquefida: dichotoma, aleppica.  
involucellis ovato-lanceolatis mucronatis, foliis in-  
ferioribus setaceis. *Amæn. acad.* 3. p. 122.

*Tithymalus* foliis inferioribus capillaceis: superioribus  
myrtosimilibus. *Morif. hist.* 3. p. 338.

*Tithymalus cyparissius*. *Alph. exot.* 65. t. 64.

*Habitat in Creta, Aleppo.* 24

45. **EUPHORBIA** umbella quinquefida: dichotoma, segetalis.  
involucellis cordatis acutis, foliis linearis lanceolatis,  
ramis floriferis. *Amæn. acad.* 3. p. 121. \* *Huds.*  
*angl.* 182.

*Euphorbia inermis*, foliis alternis linearibus acutis, par-  
tialibus umbellæ ovato-rhombeis, petalis bicornibus.  
*Hort. upf.* 142. \*

*Euphorbia inermis*, foliis linearibus acutis ad umbel-  
lam quinquefoliis, ad umbellulas ter dichotomas  
ovato-trigonis *Sauv. monsp.* 46.

*Tithymalus annuus*, lunato flore, linearis folio lon-  
giore. *Morif. hist.* 3. p. 339. f. 10. t. 2. f. 3.

*Tithymalus maritimo affinis*, linearis folio. *Bauh. pin.*  
291?

*Tithymalus segetum longifolius. Raj. angl. 3. p. 311.  
Habitat in Mauritania. ☉*

*helioscopia. 46. EUPHORBIA umbella quinquefida : trifida : dichotoma, involucellis obovatis, foliis cuneiformibus serratis. Amæn. acad. 3. p. 124. \**

*Euphorbia foliis crenatis, umbella universali quinquefida pentaphylla, partialibus trifidis, propriis triphyllis. Hort. cliff. 198. Fl. suec. 436. 425. Roy. lugdb. 197. Dalib. parif. 156.*

*Euphorbia inermis, foliis subrotundis crenatis. Fl. lapp. 220.*

*Tithymalus helioscopius. Bauh. pin. 291. Fuchf. hist. 811.*

*Habitat in Europæ cultis. ☉*

*Umbellæ radii bis trifidi, dein dichotomi ☿ Involucris foliis similia.*

*ferrata.*

*47. EUPHORBIA umbella quinquefida : trifida : dichotoma, involucellis diphyllis reniformibus, foliis amplexicaulibus cordatis ferratis. Amæn. acad. 3. p. 125. \**

*Euphorbia inermis, foliis denticulatis caulinis lanceolatis, umbellarum cordatis. Hort. cliff. 200. Hort. ups. 141. \* Roy. lugdb. 197.*

*Tithymalus characias, folio serrato. Bauh. pin. 290.*

*Tithymalus myrtites valentinus. Clus. hist. 2. p. 189.*

*Habitat in Narbona, Hispania, Italia, Oriente. ☉*

*verrucosa.*

*48. EUPHORBIA umbella quinquefida : subtrifida : trifida, involucellis ovatis, foliis lanceolatis ferrulatis villosis, capsulis verrucosis. Amæn. acad. 3. p. 120. \* Hudf. angl. 183.*

*Euphorbia inermis, foliis linearibus, umbella universali trifida triphylla, partialibus trifidis, propriis dichotomis diphyllis, foliolis subrotundis. Guett. fl. 269. Dalib. parif. 154.*

*Tithymalus myrsinites, fructu verrucæ simili. Bauh. pin. 291.*

*Tithymalus foliis subhirsutis ad caulem ellipticis, sub floribus binis subrotundis. Hall. hel. 191. \**

*Habitat in Gallia, Helvetia, Italia, Oriente. ♂*

*Fructus verrucosus simulque villosus.*

*corollata.*

*49. EUPHORBIA umbella quinquefida : trifida : dichotoma.*

chotoma, involucrellis foliisque oblongis obtusis, petalis membranaceis. *Amæn. acad.* 3. p. 122.

*Euphorbia inermis*, foliis lanceolatis obtusis alternis, ramis floriferis dichotomis, petalis maximis subrotundis. *Gron. Virg.* 58.

*Tithymalus marianus*, angustis rigidis foliis, summo caule ramosus, foliis atque ramulis ad divaricationem alternis. *Pluk. mant.* 182. t. 446. f. 3.

*Habitat in Virginia, Canada.*

10. *EUPHORBIA umbella quinquefida*: trifida: dichotoma, involucrellis ovatis, foliis lanceolatis, capsulis lanatis. *Amæn. acad.* 3. p. 123. \*

*Euphorbia inermis fruticosa*, foliis lanceolatis, involucro universaliter quinquefido, partialibus trifidis, reliquis bifidis. *Hort. ups.* 142. \* *Roy. lugdb.* 198.

*Tithymalus arboreus*, caule corallino, folio hyperici, pericarpio barbato. *Bærh. lugdb.* 1. p. 256.

*Habitat in Sicilia, Mauritania, Oriente.* 24

1. *EUPHORBIA umbella quinquefida*: trifida: bifida, involucris ovatis, petalis integris, foliis lanceolatis subpilosis apice serrulatis. pilosa.

*Euphorbia foliis alternis ex ovali-lanceolatis, umbellis diphyllis subtrifloris, capsulis erectis muricatis, caule simplici.* *Gmel. fib.* 2. p. 226. t. 93.

*Tithymalus palustris villosus mollior erectus.* *Barr. rar.* 41. t. 885.

*Tithymalus incanus hirsutus.* *Bauh. pin.* 292 ? prodr. 135?

*Tithymalus Characias pratensis incanus.* *Magn. monsp.* 255.

*Habitat in Sibiria.* 24

*Habitus exacte E. palustris, ut facile pro eadem sumeretur, eodemque tempore floret, paulo tamen major. Folia lato-lanceolata, alterna, utrinque vix manifeste pilosa, apice ita tenuissime serrata, ut vix observentur serraturæ. Umbellæ cum umbellulis lateralibus ita coacervatæ, ut primaria difficiliter eruatur, luteæ petalis 3 involucris. Flores primarii masculi pentapetali; reliqui hermaphroditi tetrapetali; petalis transverse ovalibus. Fructus verrucosi 3 pilis albis subtilissimis adpersi. Rami steriles ex alis foliorum inferiorum, ut ex summis alis pedunculi umbelluliferi.*



orientalis.

52. **EUPHORBIA umbella quinquefida** : quadrifida : dictotoma, involucrellis subrotundis acutis, foliis lanceolatis. *Amæn. acad.* 3. p. 123.

*Euphorbia inermis*, foliis lanceolatis, involucro universalis quinquefido lanceolato, partiali tetraphyllo subrotundo, propriis diphyllis. *Roy. lugdh.* 198.

*Tithymalus orientalis*, salicis folio, caule purpureo, flore magno. *Tournef. cor.* 2.

*Habitat in Oriente.*

- platyphylla. 53. **EUPHORBIA umbella quinquefida** : trifida : dictotoma, involucrellis carina pilosis, foliis serratis lanceolatis, capsulis verrucosis. *Syst. nat.* 10. p. 1049. 42. 47.

*Euphorbia umbella quinquefida* : subquadrifida : dictotoma, involucrellis primariis tetraphyllis, foliis serratis lanceolatis sessilibus. *Amæn. acad.* 3. p. 124.

*Euphorbia inermis*, foliis alternis lanceolatis amplius caulibus subserratis, umbella universalis quinquefida pentaphylla, partialibus bifidis. *Hort. ups.* 141.

*Euphorbia inermis*, foliis lanceolatis, umbella universalis partialiumque prima quinquefida : secunda trifida : reliquis bifidis. *Roy. lugdh.* 198. *Dalib. par.* 198.

*Tithymalus arvensis latifolius germanicus.* *Bak.* 291.

*Tithymalus platyphyllos.* *Fuchs. hist.* 813.

*Habitat in agris Gallie, Anglie, Germanie.* O.

*Caulis erectus, pedalis, glaber. Folia alterna, remote patentia, lanceolata, glabra, serrulata, carina tenuis rarissimis : inferiora subpetiolata, versus apicem latiora ; superiora basi emarginata & supra basin latera deflexa. Involucrum lanceolatum. Involucella partialia ovato-oblonga ; reliqua ovato-cordata, omnia serrulata. Petala suborbiculata, lutea. Germina obscure verrucosa. Ex alis foliorum bellula trifida. Planta superior viridi-flavesceus.*

\* \* \* \* \* *Umbella multifida.*

Esula.

54. **EUPHORBIA umbella multifida** : bifida : involucrellis subcordatis, petalis subbicornibus, ramis brevibus foliis uniformibus. *Amæn. acad.* 3. p. 127.

*Euphorbia inermis*, foliis lanceolato-linearibus involucri universalis foliis quinque ovato-acutis, partialibus semiorbiculatis. *Hort. ups.* 141. *Sauv. mons.* 47.

Es.

*Euphorbia inermis*, foliis ligulatis patulis ad umbellam  
quinis ovato-oblongis acutis, bracteis trigonis. *Sauv.*  
*monsp.* 51?

*Tithymalus lithospermi majoris* folio. *Magn. monsp.*  
304?

*Psula minor*. *Dalech. hist.* 1653. *Dod. pempt.* 374.

*Habitat in Germania, Belgio, Gallia.* 24

*Flores omnes fertiles* : petala *flavescentia*, 4, *obsoleta*  
*bicornia*? *Fructus glaber*. *Umbellæ caulis laterales*  
*dichotomæ*.

5. *EUPHORBIA umbella multifida* : *dichotoma*, in- *Cyparissias.*  
*volucellis subcordatis*, ramis sterilibus foliis setaceis,  
caulinis lanceolatis. *Amæn. acad.* 3. p. 127.

*Euphorbia inermis*, foliis confertis linearibus, umbella  
universali multifida, partialibus dichotomis : foliolis  
subrotundis. *Hort. cliff.* 199. \* *Hort. upf.* 142. \*  
*Roy. lugdb.* 197. *Dalib. paris.* 155. *Gort. geli.* 105.

*Tithymalus cyparissius*. *Bauh. pin.* 291. *Dalech. hist.*  
1644.

*Tithymalus cypressinus*. *Tabern. hist.* 990.

*Habitat in Minia, Bohemia, Helvetia, G. Narbo-*  
*nenfis collibus, viis, ficeis.*

6. *EUPHORBIA umbella suboctifida* : bifida, involu- *Myrsinites.*  
cellis subovatis, foliis spatulatis patentibus carnosus  
mucronatis margine scabris. *Amæn. acad.* 3. p. 128. \*

*Euphorbia inermis*, foliis superioribus reflexis latiori-  
bus lanceolatis, umbella universali trifida, partiali-  
bus bifidis. *Hort. cliff.* 199. *Hort. upf.* 141.

*Euphorbia inermis*, foliis ligulatis spinula terminatis, ad  
umbellam duodenis, bracteis trigonis spinula termi-  
natis. *Sauv. monsp.* 51.

*Tithymalus myrsinites latifolius*. *Bauh. pin.* 290.

*Tithymalus myrsinites legitimus*. *Clus. hist.* 2. p. 189.

*Habitat in Calabria, Monspeili.* 24

*Caules multi, pedales, deflexi, virides, inferne a ca-*  
*su foliorum cicatrifati.* *Folia alterna, spatulata,*  
*coriacea, concava, glauco-viridia, patentia, mucro-*  
*nata, margine subscabra, superiora reflexa.* *Umbel-*  
*la 7 - 9 fida, radiis semel bifidis.* *Involucrum u-*  
*niversale foliolis 7 - 9, ovatis, tenuioribus, acutis;*  
*Involucella biphylla, subcordata, latiora, concava,*  
*acuta, margine subscabra.* *Flores intra involucella*  
*primaria & secunda masculi, reliqui hermaphro.*

*ditii. Calyces ore serrati. Petala 4, flava, bicoloria apicibus teretibus, nitentia. Capsulæ glabra.*

- palustris.** 57. *EUPHORBIA umbella multifida: subtrifida: bifida, involuclis ovatis, foliis lanceolatis, ramis sterilibus.*  
*Amæn. acad. 3. p. 126. \**  
*Euphorbia foliis lanceolatis, umbella universali multifida polyphylla: partialibus trifidis triphyllis: prope dichotomis. Hort. cliff. 200. Fl. suec. 438; 427. Mat. med. 255. Roy. lugdb. 198. Dalib. parif. 15. Gort. gelr. 307.*  
*Tithymalus palustris fruticosus. Bauh. pin. 292.*  
*Efula major. Dalech. hist. 1653.*  
*Habitat in Suecia australi, Germania, Belgio. 2.*
- hiberna.** 58. *EUPHORBIA umbella sexfida: dichotoma, involuclis ovalibus, foliis integerrimis, ramis nullis, capsulis verrucosis. Amæn. acad. 3. p. 128. Hall. angl. 185.*  
*Tithymalus latifolius hispanicus. Bauh. pin. 291. Taber. hist. 987.*  
*Tithymalus hibernicus, vasculis muricatis erectis. D. elth. 387. t. 290. f. 374.*  
*Habitat in Hibernia, Sibiria, Austria, Pyrenæis. 2.*
- dendroides.** 59. *EUPHORBIA umbella multifida: dichotoma, involuclis subcordatis: primariis triphyllis, caule boreo. Amæn. acad. 3. p. 128. \**  
*Tithymalus myrtifolius arboreus. Bauh. pin. 290.*  
*Tithymalus dendroides. Cam. epit. 965.*  
*Habitat in Italia, Creta, Insulis Stœchadum. 6.*
- amygdaloides.** 60. *EUPHORBIA umbella multifida: dichotoma, involuclis perfoliatis orbiculatis, foliis obtusis. Amæn. acad. 3. p. 126.*  
*Tithymalus characias amygdaloides. Bauh. pin. 290.*  
*Habitat in Gallia, Germania.*
- Characias.** 61. *EUPHORBIA umbella multifida: bifida, involuclis perfoliatis emarginatis, foliis lanceolatis integerrimis, caule perenni. Amæn. acad. 3. p. 126. \**  
*Euphorbia inermis, foliis lanceolatis, umbella universali multifida, partialibus dichotomis, involuclis bifidis perfoliatis. Hort. cliff. 199. Hort. upf. 142. Roy. lugdb. 197.*

*Tithymalus characias rubens peregrinus. Bauh. pin. 290.*

*Tithymalus characias 1. Clus. hist. 2. p. 188.*

*Habitat in Gallia, Hispania, Italia, Germania. §*

*Caulis quadripedales, fruticosi, secundum annos ramosi, crassitie digiti. Rami annotini superne crassiores, subvillosi, infra folia adpersi cicatricibus transversis. Ramus florifer primo vere ex apice ramorum excrescit, fruticat, perit, cum ad latera baseos novi rami prodeunt. Folia lanceolata, coriacea, submeniosa, viridia, reflexa nervo utrinque elevato, versus flores latiora obtusiora. Umbella parva, conferta, terminalis, sessilis: Radiis numerosis, semel bifidis. Involucrum universale polyphyllum, sublan- ceolatum, parvum, reflexum; Involucella pefoliata, semibijida. Flores intra involucella primaria masculi, reliqui hermaphroditi; omnes Petalis 4, purpureis, humidis. Germina villosa.*

62. *EUPHORBIA umbella quinquefida: bifida, involu- sylvatica.*  
cellis perfoliatis subcordatis acutiusculis, foliis lanceolatis integerrimis.

*Tithymalus lunato flore. Col. ecphr. 2. p. 56. t. 57.*

*Tithymalus sylvaticus, lunato flore. Bauh. pin. 290.*

*Morif. hist. 3. p. 335. f. 10. t. 1. f. 3.*

*Habitat in Europa australiore. §*

*Affinis nimium E. Characie, unde etiam conjungitur ab Hallero, Allionio; Involucelli foliola unita quidem sunt, sed non emarginata.*

## PENTAGYNIA.

### GLINUS.

1. *GLINUS. Lxfl. it. 145. \* Gron. orient. 143.*

*Alfene lotoides sicula, Bocc. sic. 21. t. 11. Morif. hist. lotoides.*  
2. p. 552.

*Alfene heliotropio cognata, Cup. cath.*

*Portulaca batica, luteo flore, spuria aquatica. Barr. rar.*  
478. t. 336.

*Habitat ab Hispania in Asiam, ad fossas, locis in-  
datis. ☉*

*Tithymalus segetum longifolius. Raj. angl. 3. p. 312.  
Habitat in Mauritania. ☉*

*helioscopia. 46. EUPHORBIA umbella quinquefida : trifida : dichotoma, involucellis obovatis, foliis cuneiformibus serratis. Amæn. acad. 3. p. 124. \**

*Euphorbia foliis crenatis, umbella universali quinquefida pentaphylla, partialibus trifidis, propriis triphyllis. Hort. cliff. 198. Fl. Svec. 436. 425. Roy. lugdb. 197. Dalib. parif. 156.*

*Euphorbia inermis, foliis subrotundis crenatis. Fl. lapp. 220.*

*Tithymalus helioscopius. Bauh. pin. 291. Fuchf. hist. 811.*

*Habitat in Europæ cultis. ☉*

*Umbellæ radii bis trifidi, dein dichotomi ☿ Involucra foliis similia.*

*ferrata. 47. EUPHORBIA umbella quinquefida : trifida : dichotoma, involucellis diphyllis reniformibus, foliis amplexicaulibus cordatis ferratis. Amæn. acad. 3. p. 125. \**

*Euphorbia inermis, foliis denticulatis caulinis lanceolatis, umbellarum cordatis. Hort. cliff. 200. Her. upf. 141. \* Roy. lugdb. 197.*

*Tithymalus characias, folio serrato. Bauh. pin. 290.*

*Tithymalus myrtites valentinus. Clus. hist. 2. p. 189.*

*Habitat in Narbona, Hispania, Italia, Oriente. ☉*

*verrucosa. 48. EUPHORBIA umbella quinquefida : subtrifida : bifida, involucellis ovatis, foliis lanceolatis ferrulatis villosis, capsulis verrucosis. Amæn. acad. 3. p. 120. \* Hudf. angl. 183.*

*Euphorbia inermis, foliis linearibus, umbella universali trifida triphylla, partialibus trifidis, propriis dichotomis diphyllis, foliolis subrotundis. Guett. flamp. 269. Dalib. parif. 154.*

*Tithymalus myrsinites, fructu verrucæ simili. Bauh. pin. 291.*

*Tithymalus foliis subhirsutis ad caulem ellipticis, sub floribus binis subrotundis. Hall. hel. 191. \**

*Habitat in Gallia, Helvetia, Italia, Oriente. ☿*

*Fructus verrucosus simulque villosus.*

*corollata. 49. EUPHORBIA umbella quinquefida : trifida : dichotoma, involucellis ovatis, foliis lanceolatis serratis, capsulis villosis. Amæn. acad. 3. p. 120. \**

chotoma, involucellis foliisque oblongis obtusis, petalis membranaceis. *Amæn. acad.* 3. p. 122.

*Euphorbia inermis*, foliis lanceolatis obtusis alternis, ramis floriferis dichotomis, petalis maximis subrotundis. *Gron. Virg.* 58.

*Tithymalus marianus*, angustis rigidis foliis, summo caule ramosus, foliis atque ramulis ad divaricationes ternis. *Pluk. mant.* 182. t. 446. f. 3.

*Habitat in Virginia, Canada.*

50. **EUPHORBIA** umbella quinquefida: trifida: dichotoma, involucellis ovatis, foliis lanceolatis, capsulis lanatis. *Amæn. acad.* 3. p. 123. \*

*Euphorbia inermis fruticosa*, foliis lanceolatis, involucro universali quinquefido, partialibus trifidis, reliquis bifidis. *Hort. upf.* 142. \* *Roy. lugdb.* 198.

*Tithymalus arboreus*, caule corallino, folio hyperici, pericarpio barbato. *Bærh. lugdb.* 1. p. 256.

*Habitat in Sicilia, Mauritania, Oriente.* 24

51. **EUPHORBIA** umbella quinquefida: trifida: bifida, involucris ovatis, petalis integris, foliis lanceolatis subpilosis apice serrulatis. pilosa.

*Euphorbia foliis alternis ex ovali-lanceolatis, umbellis diphyllis subtrifloris, capsulis erectis muricatis, caule simplici.* *Gmel. fib.* 2. p. 226. t. 93.

*Tithymalus palustris villosus mollior erectus.* *Barr. rar.* 41. t. 885.

*Tithymalus incanus hirsutus.* *Bauh. pin.* 292 ? prodr. 135?

*Tithymalus Characias pratensis incanus.* *Magn. monsp.* 255.

*Habitat in Sibiria.* 24

*Habitus ex æste E. palustris, ut facile pro eadem sumeretur, eodemque tempore floret, paulo tamen major. Folia lato-lanceolata, alterna, utrinque vix manifeste pilosa, apice ita tenuissime serrata, ut vix observentur serraturæ. Umbellæ cum umbellulis lateralibus ita coacervatæ, ut primaria difficiliter eruatur, luteæ petalis & involucris. Flores primarii masculi pentapetali; reliqui hermaphroditi tetrapetali; petalis transverse ovalibus. Fructus verrucosi & pilis albis subtilissimis adspersi. Rami steriles ex alis foliorum inferiorum, ut ex summis alis pedunculi umbelluliferi.*

- orientalis. 52. **EUPHORBIA** umbella quinquefida : quadrifida : dichotoma, involucellis subrotundis acutis, foliis lanceolatis. *Amæn. acad. 3. p. 123.*  
*Euphorbia inermis*, foliis lanceolatis, involucro universalis quinquefido lanceolato, partiali tetraphyllo subrotundo, propriis diphyllis. *Roy. lugdb. 198.*  
*Tithymalus orientalis*, salicis folio, caule purpureo, flore magno. *Tournef. cor. 2.*  
*Habitat in Oriente.*
- platyphylla. 53. **EUPHORBIA** umbella quinquefida : trifida : dichotoma, involucris carina pilosis, foliis ferratis lanceolatis, capsulis verrucosis. *Syst. nat. 10. p. 1049. 42. 47.*  
*Euphorbia umbella quinquefida : subquadrifida : dichotoma*, involucellis primariis tetraphyllis, foliis ferratis lanceolatis sessilibus. *Amæn. acad. 3. p. 124.*  
*Euphorbia inermis*, foliis alternis lanceolatis amplexicaulibus subferratis, umbella universalis quinquefida pentaphylla, partialibus bifidis. *Hort. upf. 141.*  
*Euphorbia inermis*, foliis lanceolatis, umbella universalis partialiumque prima quinquefida : secunda trifida : reliquis bifidis. *Roy. lugdb. 198. Dalib. parif. 156.*  
*Tithymalus arvensis latifolius germanicus.* *Bauh. 7. 291.*  
*Tithymalus platyphyllos.* *Fuchf. hist. 813.*  
*Habitat in agris Galliæ, Angliæ, Germaniæ. ☉.*  
*Caulis erectus, pedalis, glaber. Folia alterna, ramentosa, patentia, lanceolata, glabra, serrulata, carina vilius rarissimis : inferiora subpetiolata, versus apicem litoria; superiora basi emarginata & supra basin et latera deflexa. Involucrum lanceolatum. Involucella partialia ovato-oblonga; reliqua ovato-cordata, omnia serrulata. Petala suborbiculata, lutea. Germina obscure verrucosa. Ex alis foliorum umbellula trifida. Planta superior viridi-flavescent.*  

\* \* \* \* \*

*Umbella multifida.*
- Esula. 54. **EUPHORBIA** umbella multifida : bifida : involucellis subcordatis, petalis subbicornibus, ramis ferrilibus foliis uniformibus. *Amæn. acad. 3. p. 127.*  
*Euphorbia inermis*, foliis lanceolato-linearibus involucri universalis foliis quinque ovato-acutis, partialibus semiobscure verrucosis. *Hort. upf. 141. Sauv. monsp. 47.*  

Es-

*Euphorbia inermis*, foliis ligulatis patulis ad umbellam  
quinis ovato-oblongis acutis, bracteis trigonis. *Sauv.*  
*monsp.* 51?

*Tithymalus lithospermi majoris folio.* *Magn. monsp.*  
304?

*Ffula minor.* *Dalech. hist.* 1653. *Dod. pempt.* 374.

*Habitat in Germania, Belgio, Gallia.* 24

*Flores omnes fertiles: petala flavescentia*, 4, *obsolete*  
*bicornia?* *Fructus glaber.* *Umbellæ caulis laterales*  
*dichotomæ.*

5. *EUPHORBIA umbella multifida: dichotoma*, involu- *Cyparissias.*  
cellis subcordatis, ramis sterilibus foliis setaceis,  
caulinis lanceolatis. *Amæn. acad.* 3. p. 127.

*Euphorbia inermis*, foliis confertis linearibus, umbella  
universali multifida, partialibus dichotomis: foliolis  
subrotundis. *Hort. cliff.* 199. \* *Hort. upf.* 142. \*  
*Roy. lugdb.* 197. *Dalib. parif.* 155. *Gort. gelr.* 105.

*Tithymalus cyparissius.* *Bauh. pin.* 291. *Dalech. hist.*  
1644.

*Tithymalus cypressinus.* *Tabern. hist.* 990.

*Habitat in Misiæ, Bohemiæ, Helvetiæ, G. Narbo-*  
*nenfis collibus, viis, ficcis.*

6. *EUPHORBIA umbella suboctifida: bifida*, involu- *Myrsinites.*  
cellis subovatis, foliis spatulatis patentibus carnosus  
mucronatis margine scabris. *Amæn. acad.* 3. p. 128. \*

*Euphorbia inermis*, foliis superioribus reflexis latiori-  
bus lanceolatis, umbella universali trifida, partiali-  
bus bifidis. *Hort. cliff.* 199. *Hort. upf.* 141.

*Euphorbia inermis*, foliis ligulatis spinula terminatis, ad  
umbellam duodenis, bracteis trigonis spinula termi-  
natis. *Sauv. monsp.* 51.

*Tithymalus myrsinites latifolius.* *Bauh. pin.* 290.

*Tithymalus myrsinites legitimus.* *Clus. hist.* 2. p. 189.

*Habitat in Calabria, Monspelli.* 24

*Caules multi, pedales, deflexi, virides, inferne a ca-*  
*su foliorum cicatrificati.* *Folia alterna, spatulata,*  
*coriacea, concava, glauco-viridia, patentia, mucro-*  
*nata, margine subscabra, superiora reflexa.* *Umbel-*  
*la 7 - 9 fida, raliis semel bifidis.* *Involucrum u-*  
*niversale foliolis 7 - 9, ovatis, tenuioribus, acutis;*  
*involucella biphylla, subcordata, latiora, concava,*  
*acuta, margine subscabra.* *Flores intra involucella*  
*primaria ♂ secundaria masculi, reliqui hermaphro.*



*diti. Calyces ore serrati. Petala 4, flava, bic-*  
*nia apicibus teretibus, nitentia. Capsulæ glabra.*

palustris.

57. *EUPHORBIA umbella multifida: subtrifida: bifida, involuclis ovatis, foliis lanceolatis, ramis sterilibus*  
*Amzn. acad. 3. p. 126. \**

*Euphorbia foliis lanceolatis, umbella universali multifida polyphylla: partialibus trifidis triphyllis: prope dichotomis. Hort. cliff. 200. Fl. suec. 438; 457. Mat. med. 255. Roy. lugdb. 198. Dalib. parif. 153. Gort. gelr. 307.*

*Tithymalus palustris fruticosus. Bauh. pin. 292.*

*Efula major. Dalech. hist. 1653.*

*Habitat in Suecia australi, Germania, Belgio. 4*

hiberna.

58. *EUPHORBIA umbella sexfida: dichotoma, involuclis ovalibus, foliis integerrimis, ramis nullis, capsulis verrucosis. Amzn. acad. 3. p. 128. H. angl. 185.*

*Tithymalus latifolius hispanicus. Bauh. pin. 291. Tabern. hist. 987.*

*Tithymalus hibernicus, vasculis muricatis erectis. P. elh. 387. t. 290. f. 374.*

*Habitat in Hibernia, Sibiria, Austria, Pyrenæis. 4*

dendroides.

59. *EUPHORBIA umbella multifida: dichotoma, involuclis subcordatis: primariis triphyllis, caule boreo. Amzn. acad. 3. p. 128. \**

*Tithymalus myrtifolius arboreus. Bauh. pin. 290.*

*Tithymalus dendroides. Cam. epit. 965.*

*Habitat in Italia, Creta, Insulis Stœchadum. 4*

amygdaloides.

60. *EUPHORBIA umbella multifida: dichotoma, involuclis perfoliatis orbiculatis, foliis obtusis. Amzn. acad. 3. p. 126.*

*Tithymalus characias amygdaloides. Bauh. pin. 290.*

*Habitat in Gallia, Germania.*

Characias.

61. *EUPHORBIA umbella multifida: bifida, involuclis perfoliatis emarginatis, foliis lanceolatis integerrimis, caule perenni. Amzn. acad. 3. p. 126. \**

*Euphorbia inermis, foliis lanceolatis, umbella universali multifida, partialibus dichotomis, involuclis bifidis perfoliatis. Hort. cliff. 199. Hort. ap. 147. Roy. lugdb. 197.*

**Tithymalus characias rubens peregrinus.** *Bauh. pin.* 290.

**Tithymalus characias** 1. *Clus. hist.* 2. p. 188.

*Habitat in Gallia, Hispania, Italia, Germania.* **h**

**Caulos quadripedales, fruticosi, secundum annos ramosi, crassitie digiti.** Ramis annotini superne crassiores, subvillosi, infra folia adpersi cicatricibus transversis. Ramus florifer primo vere ex apice ramorum excrescit, fruticat, perit, cum ad latera baseos novi rami prodeunt. Folia lanceolata, coriacea, submembrana, viridia, reflexa nervo utrinque elevato, versus flores latiora obtusiora. Umbella parva, conferta, terminalis, sessilis: Radiis numerosis, semel bifidis. Involucrum universale polyphyllum, sublan- ceolatum, parvum, reflexum; Involucella perfoliata, semibifida. Flores intra involucella primaria masculi, reliqui hermaphroditi; omnes Petalis 4, purpureis, humidis. Germina villosa.

**62. EUPHORBIA umbella quinquefida: bifida, involu-** *sylvatica.*  
**cellis perfoliatis subcordatis acutiusculis, foliis lan-**  
**ceolatis integerrimis.**

**Tithymalus lunato flore.** *Col. ecphr.* 2. p. 56. t. 57.

**Tithymalus sylvaticus, lunato flore.** *Bauh. pin.* 290.

*Morif. hist.* 3. p. 335. f. 10. t. 1. f. 3.

*Habitat in Europa australiore.* **h**

*Affinis nimium E. Characie, unde etiam conjungitur ab Hallero, Allionio; Involucelli foliola unita quidem sunt, sed non emarginata.*

## PENTAGYNIA.

### GLINUS.

**1. GLINUS.** *Læff. it.* 145. \* *Gron. orient.* 143.

**Albine lotoides sicula.** *Bocc. sic.* 21. t. 11. *Morif. hist.* *lotoides.*  
2. p. 552.

**Albine heliotropio cognata.** *Cup. cath.*

**Portulaca bætica, luteo flore, spuria aquatica.** *Barr. rar.*  
478. t. 336.

*Habitat ab Hispania in Asiam, ad fossas, locis in-  
datis.* ☉

## OCTAGYNIA.

## ILLICIAM.

ifatum.

## 1. ILLICIAM. †

Somo vulgo Skimmi. *Kämpf. amœn.* 880. t. 881.*Habitat in Japonia, China.* ‡*Planta a me non visa, fide Kämpferi recepta, forte**Anisum stellatum officinarum, quod adjectum Te-*  
*traodonti ocellari ejus auget venenum.*

## DODECAGYNIA.

## SEMPERVIVUM.

arborescens.

## 1. SEMPERVIVUM caule arborescente lævi ramofo.

*Sempervivum caule inferne nudo lævi ramofo. Hort. cliff.* 179. *Hort. upf.* 118. *Roy. lugdb.* 456.*Sedum majus arborescens, flosculis candidis. Bæd. pin.* 282. *Bradt. succ.* 4. p. 1. t. 31.*Sedum majus legitimum. Clus. hist.* 2. p. 58.*Habitat in Lusitania, Creta, Corcyra, Zacintho.* ‡

canariense.

## 2. SEMPERVIVUM caule foliorum ruderibus lacero, foliis retusis.

*Sempervivum caule infra folio lacero, foliis cuneiformibus: summis subrotundis. Hort. cliff.* 179. *Roy. lugdb.* 457.*Sedum canarium, foliis omnium maximis. Comm. hort.* 2. p. 189. t. 95.*Sedum majus canarium acaulon, pilis ad oras foliorum hispidis. Pluk. alm.* 340. t. 314. f. 1. *Raj. suppl.* 362.*Habitat in Canariis.* ‡

terrestrium.

## 3. SEMPERVIVUM foliis ciliatis, propaginibus patentibus.

*Sempervivum foliis radicalibus carnosissimis ciliatis, caulibus imbricatis membranaceis, corymbo racemoso reflexo. Hort. cliff.* 179. *Fl. succ.* 395; 428. *Mart. med.* 222. *Roy. lugdb.* 457.*Sempervivum majus alterum f. Barba jovis. Dod. pempt.* 127.*Sedum majus vulgare. Bæd. pin.* 283.*Habitat in Europæ tectis & collibus.* 2

4. SEM-

4. **SEMPERVIVUM** foliis ciliatis , propaginibus glo- globiferum.  
bolis.

*Sempervivum* foliis radicalibus in globum congestis  
ciliatis, propaginibus globosis. *Hort. cliff.* 180. *Roy.*  
*lugdb.* 457.

*Sedum* majus vulgari simile , globulis decidentibus.  
*Morif. hist.* 3. p. 472. f. 12. t. 7. f. 18.

*Sedum* vulgari magno simile. *Bauh. hist.* 3. p. 688.

*Habitat* in Rutheno. *D. Gmelin.* 24

*Flores* flavi, nec rubri.

5. **SEMPERVIVUM** foliis caule petalorumque apici- hirtum.  
bus hirtis. *Amæn. acad.* 4. p. 273.\*

*Sedum* majus montanum , foliis dentatis. *Bauh. pin.*  
283. *Raj. hist.* 1. p. 688.

*Cotyledon* altera montana. *Clus. hist.* 2. p. 63.

*Habitat* in Taurero rastiadiensi. 24

*Flores* albidus sunt , ceterum nimis affinis *S. montano*  
Et globifero.

6. **SEMPERVIVUM** foliis ciliatis, propaginibus patulis. montanum.

*Sedum* rosulis liberis, foliis lævibus. *Hall. helv.* 393.

*Sedum* alpinum, rubro magno flore. *Bauh. pin.* 284.

*Sedum* majus montanum , foliis non dentatis , floribus  
rubentibus, *Bauh. pin.* 283.

*Sedum* minus , flore rubente. *Refl. cyst. vern.* 6. t. 8.  
f. 2.

*Habitat* in rupibus Helvetiæ. 24

7. **SEMPERVIVUM** foliis pilis intertextis , propagini- arachnoide-  
bus globosis. um.

*Sempervivum* foliis radicalibus in globum congestis :  
villis reticulatim connexis. *Hort. cliff.* 180. \* *Roy.*  
*lugdb.* 457.

*Sempervivum* rubrum montanum gnaphaloides. *Col.*  
*ecphr.* 1. p. 292. t. 291.

*Sedum* montanum tomentosum. *Bauh. pin.* 284.

*Habitat* in Alpibus Italiæ, Helvetiæ, Pyrenæis. 24

## Classis XII

# ICOSANDRIA

## MONOGYNIA.

## CACTUS.

## \* Echino-Melocacti subrotundi.

mammilla-  
ris.1. CACTUS subrotundus testus tuberculis ovatis barbatus. *Hort. cliff.* 181. *Hort. upf.* 119. *Roy. lugdb.* 278.EchinoMelocactus minor lactescens, tuberculis f. mammillis majoribus. *Herm. par.* 136. s. 136.Ficoides f. Melocactus mammillaris glabra sulcis carens fructum suum undique fundens. *Pluk. alm.* 148. t. 29. f. 1.Ficoides f. Ficus americana sphaerica tuberculata lactescens, flore albo. *Comm. hort.* 1. p. 105. t. 55. *Bradl. succ.* 3. p. 11 t. 19.

Habitat in Americæ calidioris rupibus. §

Conf. Melocactus minimus lanuginosus & tuberosus. *Plum. spec.* 19. ic. 201. f. 1.Ficoides f. Melocactus minima lanuginosa, spinis minoribus, fructu sparsim egrediente. *Pluk. alm.* 148. t. 29. f. 2.Melocactus. 2. CACTUS subrotundus quatuordecim-angularis. *Hort. cliff.* 181. *Hort. upf.* 119. *Roy. lugdb.* 279.Cactus humilis subrotundus sulcatus & coronatus spinis confertis. *Brown. jam.* 238.Melocactus indiæ occidentalis. *Bauh. pin.* 384.EchinoMelocactus. *Clus. exot.* 92. t. 92. *Bradl. succ.* 4. p. 9. f. 32.

Habitat in Jamaica, America calidiore. §

## \* \* Cerei angulati erecti.

heptagonus. 3. CACTUS erectus oblongus septemangularis. *Hort. cliff.* 181. \* *Roy. lugdb.* 279.

Habitat in America. §

Historia Cactorum angulatorum 3 — 14, quorum plurimas species Plumier in iconibus sistit, etiamnum ob

*obscura permansit , quam itaque attenti per Indias peregrinatores graphice describant.*

4. CACTUS quadrangularis longus erectus : angulis tetragonus. compressis. *Hort. cliff.* 181. *Hort. upf.* 119. *Roy. lugdb.* 280.

*Cereus erectus minor , fructu spinoso , costarum numero varians. Herm. par.* 117.

*Habitat in Curassao , America calidiorē. ‡*

5. CACTUS erectus sexangularis longus. *Hort. cliff.* hexagonus 181. *Hort. upf.* 119. *Roy. lugdb.* 279.

*Cereus surinamensis. F. ph. N. C. 3. p. 349. t. 7, 8.*

*Cereus erectus altissimus surinamensis. Herm. par.* 116. *Raj. dendr.* 23.

*Cereus erectus maximus americanus hexangularis , flore alboradiato. Bradl. succ. 1. p. 1. t. 1.*

*Melocactus monoclonos , flore albo , fructu atropurpureo. Plum. spec. 19. ic. 191?*

*Habitat Surinami. ‡*

6. CACTUS subquingularis erectus longus artic. pentagonus. latus. *Hort. cliff.* 182. \* *Roy. lugdb.* 280.

*Habitat in America.*

7. CACTUS erectus longus octangularis : angulis compressis undatis ; spinis lana longioribus. *Hort. cliff.* 182. *Roy. lugdb.* 279. repandus.

*Cactus erectus cylindraceus sulcatus tenuior summi, tate attenuatus , aculeis confertis. Brown. jam.* 238.

*Cereus altissimus gracilior , fructu extus luteo , intus niveo seminibus nigris pleno. Sloan. jam.* 197. *hist.* 2. p. 158. *Raj. dendr.* 22. *Trew. chret.* 1. 14.

*Habitat in America calidiorē. ‡*

8. CACTUS erectus longus subnovemangularis : angulis obsoletis , spinis lana brevioribus. *Hort. cliff.* 182. *Roy. lugdb.* 279. lanuginosus

*Cereus curassavicus erectus maximus , fructu rubro non spinoso , lanugine flavescēte. Herm. par.* 115. t. 115.

*Habitat in Curassao. ‡*

9. CACTUS erectus longus suboctangularis : angulis obtusis. *Hort. cliff.* 181. *Hort. upf.* 120. *Roy. lugdb.* 279. peruvianus.

*Cactus cylindraceus erectus sulcatus major summitate obtusus, aculeis confertis. Brown, jam. 238.*

*Cereus erectus, fructu rubro non spinoso. Herm. par. 114.*

*Cereus peruvianus spinosus, fructu rubro nucis magnitudine. Bauh. pin. 458.*

*Euphorbii arbor cerei effigie. Lob. ic. 2. p. 25.*

*Habitat in Jamaica, Perupriceis aridis maritimis. h*

- Royenl. 10. **CACTUS** erectus articulatus subdecangularis: articulis subovatis, spinis lanam æquantibus. *Roy. lugdb. 279.*

*Habitat in America. h*

\* \* \* *Cerei repentes radiculis lateralibus,*

- grandiflorus 11. **CACTUS** repens subquinquangularis.

*Cactus scandens, angulis quinque pluribusque obtusis. Hort. cliff. 182. Hort. upf. 121. \* Roy. lugdb. 280.*

*Cereus scandens minor polygonus articulatus. Mill. icon. t. 90.*

*Cereus gracilis scandens ramosus plerumque sexangularis, flore ingenti atque fragranti. Trew. chret. t. 31, 32.*

*E. N. C. 1752. v. 9. app. 184. t. 11, 12, 13.*

*Cereus americanus major articulatus, flore maximo nocte se aperiente s. suavissimum odorem spirante.*

*Velk. hejp. 1. p. 133. t. 134.*

*Habitat in Jamaica, Vera Cruce. h*

- flagelliflor. 12. **CACTUS** repens decemangularis.

n. s.

*Cactus scandens, angulis decem obsoletis. Hort. upf. 121.*

*Cereus minor scandens polygonus spinosissimus, flore purpureo. Ehret. sel. 2. f. 2. Trew. chret. t. 30.*

*Ficoides americanum s. Cereus minima serpens americana. Pluk. alm. 148. t. 158. f. 6. Sloan. jam. 197. hist. 2. p. 158.*

*Habitat in America calidiora. h*

- parafiticus. 13. **CACTUS** repens teres striatus muticus.

*Cactus parafiticus inermis aphyllus ramosus propendens, ramulis gracilibus teretibus striatis. Brown jam. 238.*

*Opuntia minima flagelliformis. Plum. spec. 6. ic. 197. f. 2.*

*Habitat in America.*

14. CACTUS repens triangularis. triangularis.  
 Cactus triangularis scandens articulatus. *Hort. cliff.* 182.  
*Hort. upf.* 121. *Roy. lugdb.* 280.  
 Cactus debilis brachiatus æqualis triquetrus scandens f.  
 repens, spinis brevissimis confertis. *Brown. jam.* 468.  
 Cereus americanus triangularis radicosus. *Bradd. succ.*  
*1. p. 4. t. 3. E. N. C. 1752. v. 9. app. 199. t. 10. f.*  
*14. E. N. C. 1754. v. 9. app. 349. t. 3.*  
 Ficoides americanum, f. Cereus erectus cristatus, fo-  
 liis triangularibus protunde canaliculatis. *Pluk. alm.*  
*147. t. 29. f. 2.*  
*Habitat in Brasilia, Jamaica.*  
 \* \* \* \* Opuntia compressa articulis proliferis.
15. CACTUS articulato-prolifer, articulis globosis spi- monilifor-  
 nosis glomeratis. mis.  
 Melocactus, ex plurimis globulis opuntia modo na-  
 scentibus, spinosissimus. *Plum. spec.* 20. *ic.* 198.  
*Habitat in America calidiore. h*
16. CACTUS articulato-prolifer laxus, articulis ovatis, Opuntia.  
 spinis setaceis.  
 Cactus compressus articulatus ramosissimus, articulis  
 ovatis: spinis setaceis. *Hort. cliff.* 183. *Hort. upf.* 120.  
*Gron. virg.* 54. *Roy. lugdb.* 280.  
 Opuntia vulgo herbariorum. *Bauh. hist.* 1. *p.* 154. *Mill.*  
*ic.* 191.  
 Ficus indica, folio spinoso, fructu majore. *Bauh.*  
*pin.* 458.  
*Habitat in America, Peru, Virginia, nunc in Hispa-*  
*nia, Italia, Lusitania. h. 2*
17. CACTUS articulato-prolifer, articulis ovato-oblon- Ficus indi-  
 gis spinis setaceis. ca.  
 Cactus compressus articulatus ramosus, articulis ova-  
 to-oblongis, spinis setaceis. *Hort. cliff.* 183. *Hort.*  
*upf.* 120. *Roy. lugdb.* 280.  
*Habitat in America calidiore. h*
18. CACTUS articulato-prolifer, articulis ovato-ob- Tuna.  
 longis, spinis subulatis.  
 Cactus compressus articulatus ramosus, articulis ova-  
 to-oblongis, spinis subulatis. *Hort. cliff.* 183. *Hort.*  
*upf.* 120. *Roy. lugdb.* 280.



- Cactus brachiatus articulatus*, articulis oblongo-ovatis compressis, caudice tereti erecto serpissimo, aculeis brachiorum brevibus confertis. *Brown. jam.* 237.  
*Tuna major*, spinis validis flavicantibus, flore gilvo. *Dill. elth.* 396. t. 295. f. 238.  
*Opuntia major*, folio oblongo rotundo : spinis longis & validissimis, flore luteo. *Sloan. jam.* 103. *hist.* 2. p. 149. t. 224. f. 1.  
*Habitat in Jamaica* & *America calidiore.* §
- cochinillifer* 16. *CACTUS* articulato-prolifer, articulis ovato-oblongis subinermibus.  
*Cactus compressus articulatus ramosus*, articulis ovato-oblongis subinermibus. *Hort. upf.* 121.  
*Tuna mitior*, flore sanguineo, cochinillifera. *Dill. elth.* 399. t. 297. f. 383.  
*Ficus indica major lævis* f. spinosa vermiculos proferens. *Pluk. alm.* 146. t. 281. f. 383.  
*Opuntia maxima*, folio oblongo rotundo majore spinulis nonnullis & innocentibus obsito. *Sloan. jam.* 194. *hist.* 2. p. 152. t. 8. f. 1. 2.  
*Habitat in Jamaica* & *America calidiore.* §
- curassavicus.* 20. *CACTUS* articulato-prolifer, articulis cylindrico-ventricos compressis.  
*Cactus tereti-compressus articulatus ramosus.* *Hort. cliff.* 182. *Hort. upf.* 120. *Roy. lugdb.* 280.  
*Opuntia minima americana spinosissima.* *Bradl. succ.* 1. p. 5. t. 4.  
*Ficus indica* f. *Opuntia curassavica minima.* *Cogn. hort.* 1. p. 107. t. 56.  
*Ficus indica* f. *Opuntia minor caulescens arbuscula* in modum, ramis cineritiis, spinosissima. *Pluk. alm.* 147. t. 281. f. 3.  
*Habitat in Curassao.* §
- Phyllanthus* 21. *CACTUS* prolifer ensiformi-compressus serrato repandus.  
*Cactus foliis ensiformibus obtuse serratis.* *Hort. cliff.* 183. *Roy. lugdb.* 281.  
*Cactus mitis minor*, firmento flexili rotundo, frondibus longis compressis crenatis, ad crenas floridis. *Brown. jam.* 237.  
*Cereus scolopendri folio brachiato.* *Dill. elth.* 73. t. 64. f. 74.

Opun-

*Opuntia non spinosa minima caulescens, foliis pilosis strictissimis e crenis foliorum florem proferens. Sloan. jam. 216. hist. 2. p. 159.*

*Phyllanthos americana, sinuosis foliis longis. Pluk. alm. 296. t. 247. f. 5.*

*Ficus l. Opuntia non spinosa, scolopendriaefolio sinuato. Raj. dendr. 21.*

*Habitat in Brasilia, Surinamo, America meridionali. §*

22. CACTUS caule tereti arboreo : aculeis geminis recurvis, foliis lanceolato-ovatis. *Hort. upf. 122.* Pereskia.

*Cactus tamentosus foliatus spinosus, spinis geminis recurvis foliis mollibus ovatis. Brown. jam. 237.*

*Pereskia. Hort. cliff. 122. Roy. lugdb. 281.*

*Pereskia aculeata, flore albo, fructu flavesciente. Plum. gen. 37. Dill. elth. 305. t. 227. f. 294.*

*Malus americana spinosa, portulacæ folio, fructu folioso, semine reniformi splendente. Comm. hort. 1. p. 135. t. 70.*

*Portulaca americana latifolia ad foliorum ortum lanugine obducta, longioribus aculeis horrida. Pluk. alm. 135. t. 215. f. 6.*

*Grossulariæ fructu majore arbor spinosa, fructu folioso viridi-albicante. Sloan. jam. 165. hist. 2. p. 86. Raj. dendr. 27.*

*Habitat in America calidiore, Jamaica, Margaretha. §*

23. CACTUS caule tereti arboreo spinoso, foliis cuneiformibus retusis. portulacifolius.

*Opuntia arbor spinosissima, foliis portulacæ cordatis. Plum. spec. 6. ic. 197. f. 1.*

*Habitat in America calidiore. §*

*Fructus hujus minime foliis adpersus, uti precedentis.*

## PHILADELPHUS.

1. PHILADELPHUS foliis subdentatis.

coronarius.

*Philadelphus. Hort. cliff. 188. Hort. upf. 122.*

*Syringa alba f. Philadelphus athenæi. Bauh. pin. 399.*

*Frutex coronarius. Clus. hist. 1. p. 55.*

*Habitat Veronæ? §*

*Flos primarius quinquesidus calyce, corolla, pistillo, capsula. Phil. bot. 178. Petala æstivania obvoluta cum sole. Gemmis caret.*

2. PHI-

inodorus.

1. PHILADELPHUS foliis integerrimis. *Mill. dict.* 2.  
 Philadelphus flore albo majore inodoro. *Catesb. car.*  
 2. p. 84. t. 84.  
*Habitat in Carolina.* ♀

## PSIDIUM.

pyriferum.

1. PSIDIUM foliis lineatis obtusiusculis, pedunculis unin-  
 floris.  
*Psidium caule quadrangulo. Hort. cliff.* 148. *Fl. zeyl.*  
 192.  
*Guajava foliorum angulis quadrangulis, fructu oblon-*  
*go. Trew. chr.* t. 43.  
*Guajava alba dulcis. Comm. hort.* 1. p. 121. t. 63. *Mer-*  
*surin.* 19. t. 19.  
*Guajavos fructu pallido dulci. Burm. zeyl.* 112.  
*Cujavus domestica. Rumph. amb.* 1. p. 140. t. 47.  
*Pela. Rheed. mal.* 3. p. 31. t. 34.  
*Habitat in Indiis.* ♀

pomiferum.

2. PSIDIUM foliis lineatis acuminatis, pedunculis tri-  
 floris.  
*Guajava alba acida, fructu rotundiore. Pluk. alm.* 181.  
 t. 193. f. 4.  
*Guajabo pomifera indica: pomis rotundis. Bauh. pin.*  
 437.  
*Cujavus agrestis. Rumph. amb.* 1. p. 142. t. 4. *Mer-*  
*surin.* 57. t. 57.  
*Malacca pela. Rheed. mal.* 3. p. 33. t. 35.  
*Habitat in Indiis.* ♀

## EUGENIA.

malaccensis.

1. EUGENIA foliis integerrimis, pedunculis ramosis  
 lateralibus. *Fl. zeyl.* 187.  
*Perfici officulo fructus malaccensis rubens. Bauh. pin.*  
 441.  
*Jambosa domestica. Rumph. amb.* 1. p. 121. t. 37. 68.  
*Nati-schambu. Rheed. mal.* 1. p. 29. t. 18. *Roy. hij.*  
 1748.  
*Habitat in India.* ♀

Jambora.

2. EUGENIA foliis integerrimis, pedunculis ramosis  
 terminalibus. *Fl. zeyl.* 188.\*  
*Perfici officulo fructus malaccensis ex candido rube-*  
*scens. Bauh. pin.* 441.

*Jambosa sylvestris alba*. *Rumph. amb.* 1. p. 127. t. 39.  
*Malacca ichambu*. *Rheed. mal.* 1. p. 27. t. 17. *Raj. hist.*  
 1478.

*Habitat in India.* ‡

3. *EUGENIA* foliis integerrimis, pedunculis unifloris uniflora.  
 lateralibus. *Fl. zeyl.* 189.

*Eugenia*. *Roy. lugdb.* 265

*Eugenia indica*, myrti folio deciduo, flore albo, fructu suave rubenti molli leviter sulcato & odoro.

*Mich. gen.* 226. t. 108.

*Myrtus indica*, foliis rigescentibus latis ac recurvis parum odoratis. *Till. pif.* 117. t. 44.

*Habitat in India.* ‡

4. *EUGENIA* foliis crenatis, pedunculis terminalibus, acutangula.  
 pomis oblongis acutangulis. *Fl. zeyl.* 190. \*

*Butonica terrestris rubra*. *Rumph. amb.* 3. p. 181. t.

115.

*Tsieria-samitravadi*. *Rheed. mal.* 4. p. 51. t. 7. *Raj. hist.* 1480.

*Habitat in India.* ‡

5. *EUGENIA* foliis crenatis, racemis longissimis, po- racemosa.  
 mis ovatis. *Fl. zeyl.* 191. \*

*Butonica sylvestris alba*. *Rumph. amb.* 3. p. 181. t. 116.

*Samitravadi* s. *Caibatsumbu*. *Rheed. mal.* 4. p. 11. t. 16.

*Raj. hist.* 1479.

*Habitat in India.* ‡

## M Y R T U S.

1. *MYRTUS* floribus solitariis: involucro diphylo. communis.  
*Hort. cliff.* 42. *Hort. upf.* 123. *Mat. med.* 224. *Roy. lugdb.* 269.

2. *Myrtus* foliis ovatis, pedunculis longioribus. *Mill. dict. romana.*  
 t. 184. f. 1.

*Myrtus latifolia romana*. *Bauh. pin.* 468.

3. *Myrtus* foliis ovatis, baccis rotundioribus. *Mill. tarentina.*  
*dict.*

*Myrtus minor vulgaris*. *Bauh. pin.* 469.

4. *Myrtus* foliis ovato lanceolatis acutis, ramis rectio- italica.  
 ribus. *Mill. dict.*

*Myrtus communis italica*. *Bauh. pin.* 469.

5. *Myrtus* foliis ovato-lanceolatis confertis. *Mill. dict. baetica.*  
*Linnaei Spec. plant. pars. I.* U u *Myr-*

- Myrtus latifolia* boetica 2. *Bauh. pin.* 459. *Clus. hist.* 1. p. 65.
- lusitanica.* 1. *Myrtus* foliis lanceolato ovatis acutis. *Mill. dist.*  
*Myrtus sylvestris*, foliis acutissimis. *Bauh. pin.* 459.  
*Clus. hist.* 1. p. 66. f. 1.
- belgica.* 2. *Myrtus* foliis lanceolatis acuminatis. *Mill. dist.*  
*Myrtus latifolia belgica.* *Bauh. pin.* 459.
- mucronata.* 7. *Myrtus* foliis lineari lanceolatis acuminatis. *Mill. dist.*  
*Myrtus* foliis minimis & mucronatis. *Bauh. pin.* 459.  
*Clus. hist.* 1. p. 67.  
*Habitat in Europa australi, Asia, Africa.* 5
- brasiliensis.* 2. *MYRTUS* floribus solitariis, pedunculis nudis, petalis subciliatis.  
*Myrtus pomilera* latissimis foliis. *Plum. spec.* 18. 207. f. 1. (ob petala ciliata).  
*Philadelphus aborelensis*, foliis myrtinis nitidis oppositis, ramulis gracilibus, pedunculis bipartitis elatibus. *Brown jam.* 240.  
*Myrti folio* arbor, cortice argenteo, foliis oblongis ad basin latoribus inodoris. *Stean. jam.* 162. *hist.* 2. p. 78. t. 187. f. 3. *Kaj. dendr.* 35.  
*Arbor brasiliensis*, myrti laureæ foliis inodoris. *Comm. hort.* 1. p. 173. t. 8.  
*Habitat in Brasilia.* D. Wachend. 5  
*Folia ovata, obtusa, petiolata, glabra, opposita. Flores solitarii, axillares, pedunculati, pedunculo oppositi. Corollæ petala 4. f. 5, lacero-ciliata, cuneiforme ovata, concava, reflexa. Stamina cincta 40.*
- biflora.* 3. *MYRTUS* pedunculis bifloris, foliis lanceolatis. *Amæn. acad.* 5 p. 398. \*  
*Caryophyllus fruticulosus*, foliis lanceolatis oppositis floribus geminatis alaribus. *Brown. jam.* 248. t. 25. f. 3.  
*Habitat in Jamaica.* 5
- lucida.* 4. *MYRTUS* pedunculis subtrifloris, foliis subsessilibus lanceolatis attenuatis.  
*Habitat Surinamii. Kclander.* 5  
*Folia singulari structura, ex ovato admodum attenuata in apicem lanceolatum. Flores pentapetali.*
- cuminifolia.* 5. *MYRTUS* pedunculis multifloris, foliis lanceolatis ovatis. *Fl. Zeyl.* 185. \*

*Jambosa ceramica. Rumph. amb. 1. p. 130. t. 41.*

*Habitat in Zeylona. h*

6. MYRTUS pedunculis trichotomo-paniculatis, foliis oblongis floribus dioicis.

*Habitat in America. Miller. h*

*Folia opposita, lanceolato-ovata, crassa. Pedunculi axillares & terminales, brachiato-paniculati, longitudine foliorum. Petala pauca.*

7. MYRTUS pedunculis dichotomis paniculatis tomentosis, foliis geminis subovatis terminalibus. *Amzn. lia. acad. 5. p. 398. \**

*Chytraculia arborea, foliis ovatis glabris oppositis, racemis terminalibus. Brown. jam. 239 t. 37. f. 2.*

*Pluk. phyt. 274. f. 2.*

*Habitat in Jamaica. h*

8. MYRTUS pedunculis multifloris, foliis geminis subovatis terminalibus, ramis dichotomis. *Amzn. acad. 5. p. 398 \**

*Zuzygium fruticosum, foliis ovatis nitidis & ramulis ubique jugatis. Brown. jam. 240. t. 7. f. 2.*

*Habitat in Jamaica. h*

9. MYRTUS pedunculis multifloris, foliis ovatis subpetiolatis. *zeylanica.*

*Myrtus foliis ovatis acuminatis obtusiusculis. Fl. zeyl. 182. \**

*Myrtus zeylanica odoratissima, baccis niveis monococcis. Herm. agdb. 434. t. 435.*

*Myrtoides foliis ovatis. Hort. cliff. 489. Roy. lugdb. 535.*

*Habitat in Zeylona. h*

10. MYRTUS pedunculis trifido-multifloris, foliis subovalibus sessilibus. *Fl. zeyl. 184. \** *androsemaides*

*Habitat in Zeylona. h*

11. MYRTUS pedunculis trifido-multifloris, foliis obovatis. *Fl. zeyl. 183. Mat. med. 225.* *caryophyllata.*

*Caryophyllus aromaticus ind. occid. foliis & fructu rotundis. Pluk. alm. 188. t. 155. f. 3.*

*Habitat in Zeylona. h*

- Pimenta.** 12. *MYRTUS* foliis alternis. *Fl. Zeyl.* 186. *Mat. med.* 225.  
*Myrtus calycibus absque appendiculis. Hort. cliff. 31.*  
*Myrtus foliis oblongo-ovatis glabris alternis, racemis*  
*terminalibus & lateralibus. Brown, jam. 247.*  
*Myrtus arborea aromatica foliis laurinis. Stom. jor.*  
*161. hist. 2. 1. 76 t. 191 f. 1.*  
*Caryophyllus foliis oblongo-ovatis alternis, racemis*  
*terminalibus & lateralibus Brown, jam. 247*  
*Caryophyllus aromaticus ameri anus, lauri acuminatis*  
*foliis, fructu orbiculari. Pluk. alm. 88. t. 155 f. 4*  
*Habitat in India. b*
- Leucadendra.** 13. *MYRTUS* baccis sessilibus urceolatis, foliis lanceolatis. †  
*Arbor alba. Rumph. amb. 2. p. 72. t. 16, 17.*  
*Habitat in Ambo na. b*

## P U N I C A.

- Granatum.** 1. *PUNICA* foliis lanceolatis. caule arboreo. *Hort. cliff. 134. Hort. upf. 122. Mat. med. 223. Roy. lugd. 267 Gron. orient 144.*  
*Malus punica sylvestris. Bauh. pin. 438.*  
*Malus punica fativa. Bauh. pin. 438.*  
*Malus punica. Cam. epit. 130, 131.*  
 β. *Balaustia flore pleno majore & minore. Bauh. pin. 438.*  
*Habitat in Hispania, Italia, Mauritania, solo cretaceo.*
- nana.** 2. *PUNICA* foliis linearibus, caule fruticoso. *dict. †*  
*Punica americana nana s. humillima. Tournesf. 636.*  
*Habitat in Antiliis. b*

## A M Y G D A L U S.

- Perfica.** 1. *AMYGDALUS* foliorum serraturis omnibus tectis, floribus sessilibus solitariis. *Hort. cliff. 189. Hort. upf. 123. Mat. med. 230. Roy. lugd. 267.*  
*Perfica molli carne & vulgaris. Bauh. pin. 440.*  
*Perfica rubra. Cam. epit. 145.*  
 β. *Nuciperfica, quæ nucum juglandium faciem representat. Bauh. pin. 440. Roy. hist. 1516.*  
*Habitat - - - b*

*Flores sessiles, gemmales, solitarii. Folia e gemma terminali, nec infra flores.*

2. **AMYGDALUS** foliis serraturis infimis glandulosis, communis. floribus sessilibus geminis. *Hort. cliff.* 186 *Hort. upf.* 123. *Mat. med.* 429. *Roy. lugdb.* 267. *Gron. orient.* 146.

*Amygdalus sylvestris.* *Bauh. pin.* 442.

3. *Amygdalus sativa.* *Bauh. pin.* 441.

*Amygdalus.* *Dod. pempt.* 798.

- γ. *A. vgdalus amara.* *Tournef. infl.* 627.

*Habitat in Mauritaniae sepibus.* §

*Flores gemmales, sessilis s. subsessiles, saepius bini. Folia e gemmi infra & supra flores locatis.*

3. **AMYGDALUS** foliis basi attenuatis. *Hort. cliff.* 186. *Hort. upf.* 124. *Roy. lugdb.* 267. *nana.*

*Amygdalus indica nana.* *Fluk. alm.* 28. t. 11. f. 3. *Mill. dict.* t. 28. f. 2.

*Armeniaca persicae foliis, fructu exsucco.* *Ann. ruth.* 273. t. 30.

*Habitat in Asia septentrionali.*

## PRUNUS.

1. **PRUNUS** floribus racemosis, foliis deciduis basi sub. *Padus* tus biglandulosis

*Padus glandulis duabus basi foliorum subjectis.* *Hort. cliff.* 185. *Fl. sive.* 396 ; 431. *Roy. lugdb.* 269. *Hall. herb.* 327.

*Padus foliis annuis.* *Fl. lapp.* 198.

*Cerasus racemosa sylvestris, fructu non eduli.* *Bauh. pin.* 451.

*Habitat in Europa.* §

*Racemi foliacei infra folia. Petala serrata ut in sequenti, nec emarginata ut in Cerao.*

1. **PRUNUS** floribus racemosis, foliis deciduis basi an- *virginiana.* tice glandulosis.

*Cerasus sylv. stris, fructu nigricante in racemis longis pendulis phytolaccae instar congestis.* *Gron. virg.* 54. *Roy. lugdb.* 337.

*Cerasus latiore folio, fructu racemoso purpureo majore.* *Catesb. car.* 2. p. 94. t. 94

*Habitat in Virginia, Carolina.* §



Padi progenies, sed verrucæ in ramis duplo majores.  
 Petioli antice ad apicem binis paribus glandularum.  
 Folia minus subtus reticulata & fere glabra. Pe-  
 tala orbiculata, nec obovata. Fructus quadruplo ma-  
 jores rubri.

- canadensis. 3. PRUNUS floribus racemosis, foliis lato-lanceolatis  
 rugosis utrinque pubescentibus.  
 Cerasus pumila canadensis, oblongo angusto folio.  
 fructu parvo. *Duham. arb. 1. p. 149.*  
 Cerasus racemosa, foliis amygdalinis, americana. *Flak-  
 alm 97. t. 158. f. 4.*  
*Habitat in America septentrionali. ♀*  
 Rami læves. Folia lato-lanceolata absque glandulis.  
 venoso-rugosa, minusque rigida quam in reliquis.  
 acute serrata, utrinque viridia, ad tactum mar-  
 gesse villosa, & basi in petiolum desinentia.
- lusitanica. 4. PRUNUS floribus racemosis, foliis sempervirenti-  
 bus eglandulosis. *Mill. ic. 131. t. 196. f. 1.*  
 Padus folii glandula destitutis. *Vir. cliff. 42. Hort.  
 upf. 126. Roy. lugdb. 269.*  
 Padus foliis sempervirentibus ovatis. *Hort. cliff. 185.*  
 Lauro Cerasus lusitanica minor. *Dill. elh. 193. t. 155.  
 f. 193.*  
*Habitat in Lusitania, Pennsylvania. ♀*
- Lauro-Cera- 5. PRUNUS floribus racemosis, foliis sempervirenti-  
 sus, bus dorso biglandulosis.  
 Padus glandulis duabus dorso foliorum innatis. *V.  
 cliff. 42. Hort. upf. 126. Roy. lugdb. 269.*  
 Padus foliis sempervirentibus lanceolato-ovatis. *Hort.  
 cliff. 185.*  
 Cerasus folio laurino. *Bauh. pin. 450.*  
 Laurocerasus. *Clus. hist. 1. p. 4. Cam. hort. 1. 23.*  
*Habitat in Trapezunte, unde in Europam venit. 1576. ♀*
- Mahaleb. 6. PRUNUS floribus corymbosis, foliis ovatis.  
 Cerasus foliis ovatis. *Vir. cliff. 43. Hort. upf. 126.  
 Hall. helv. 360. Roy. lugdb. 268. crenatis. Hort.  
 487.*  
 Ceraso affinis. *Bauh. pin. 451.*  
 Mahaleb. *Cam. epit. 91.*  
*Habitat in Helvetia. ♀*

*Folia conduplicata, ovata, obtusa, crenata* Stipulæ  
*cor. n. æ dentibus fuscis. Corymbis convexus. bractææ brevissimæ, obtusissimæ, serratæ.*

7. PRUNUS floribus sessilibus, foliis subcordatis. Armeniaca.

*Prunus foliis ovato-cordatis. Hort. cliff. 186. Hort. upf. 124. Roy. lugdb. 268.*

*Mala Armeniaca majora. Bauh. pin. 442.*

*Mala Armeniaca majora, nucleo dulci. Bauh. pin. 442.*

β. *Malus Armeniaca minor. Bauh. pin. 442.*

*Habitat - - -*

*Folia veratitia convoluta.*

8. PRUNUS floribus sessilibus, foliis ovato-oblongis. sibirica.

*Armeniaca betulæ folio & facie, fructu exiucco. Amm. ruth 272, t. 29.*

*Habitat in Sibiria.*

*An hæc satis distincta a præcedenti?*

9. PRUNUS umbellis subsessilibus, foliis ovato lanceolatis conduplicatis glabris. Cerasus.

*Cerasus foliis ovato lanceolatis. Vir. cliff. 43. Hort. upf. 125. Mat. med. 233. Roy. lugdb. 267. Hall. helv. 356.*

α. *Cerasus sativa rotunda rubra & acida. Bauh. pin. capromiana. 449.*

β. *Cerasus hortensis, flore roseo. Bauh. pin. 450.*

γ. *Cerasus hortensis, flore pleno. Bauh. pin. 450.*

δ. *Cerasus alba dulcis. Bauh. pin. 450.*

ε. *Cerasus carne tenera & aquosa. Bauh. pin. 450.*

η. *Cerasus acidissima sanguineo succo. Bauh. pin. 450.*

θ. *Cerasus pumila. Bauh. pin. 450.*

ι. *Cerasus racemosa hortensis. Bauh. pin. 450.*

κ. *Cerasus sativa majora. Bauh. pin. 450.*

λ. *Cerasus crassa carne dura. Bauh. pin. 450.*

μ. *Cerasus hortensis, flore pleno. Bauh. pin. 450. Mill. dict. t. 89. f. 1.*

*Habitat in Europa. h*

*Folia conduplicata. Stipulæ dentatæ. Bractææ trifidæ. serratæ: intermedia foliacea. Gemmæ terminales foliosæ, laterales floriferæ.*

*rosea.  
 plena.  
 dulcis.  
 juliana.  
 austera.  
 pumila.  
 avium.  
 Bigar-lla.  
 Duracina,*

arvum.

10. PRUNUS umbellis sessilibus, foliis ovato-lanceolatis conduplicatis subtus pubescentibus. *Fl. svec. 1. p. 474. Hudf. angl. 187.*

Cerasus sylvestris, fructu nigro & rubro. *Bauh. hist. 1. p. Cerasus major s. sylvestris, fructu subdulci nigro colore inficiente. Bauh. pin. 1539.*

Cerasus sylvestris septentrionalis anglica, fructu rubro parvo serotino.

*Habitat in Europa borealiore. ‡*

*Nostra fructu rubro, nec nigro; horæo nec oportino est; magnitudine certat cum Quercu.*

domestica.

11. PRUNUS pedunculis subsolitariis, foliis lanceolato-ovatis convolutis, ramis muticis.

Prunus inermis, foliis lanceolato-ovatis. *Hort. cliff. 186. Hort. upf. 124. Mat. med. 232. Roy. lugdb. 258.*

Prunus sylvestris fructu majore. *Vaill. parif. 140.*

Prunus. *Bauh. pin. 443.*

Damascena.

- β. Pruna majora dulcia & parva atro-cærulea. *Bauh. pin. 443. n. 23.*

hungarica.

- γ. Pruna magna crassa subacida. *Bauh. pin. 443.*

juliana.

- δ. Pruna oblonga cærulea. *Bauh. pin. 443.*

perligona.

- ε. Pruna nigra, carne dura. *Bauh. pin. 443.*

cerea.

- ζ. Pruna coloris ceræ ex candido in luteum pallescente. *Bauh. pin. 443.*

acinaria.

- η. Pruna magna rubra rotunda. *Bauh. pin. 443.*

mali-jormis.

- θ. Pruna rotunda flava dulcia mali amplitudine. *Bauh. pin. 443.*

angustana.

- ι. Pruna angusto-maturefcentia minora & angustiora. *Bauh. pin. 443.*

præcox.

- κ. Pruna parva præcoccia. *Bauh. pin. 443.*

cærcola.

- λ. Pruna parva ex viridi flavescentia. *Bauh. pin. 443.*

amygdalina.

- μ. Pruna amygdalina. *Bauh. pin. 443.*

galatenfis.

- ν. Pruneoli albi oblongiusculi acidi. *Bauh. pin. 443.*

Brignola.

- ο. Pruna ex flavo rufescentia mixti saporis gratissima. *Bauh. pin. 443.*

myrobalan.

- ξ. Prunus fructu rotundo nigro-purpureo dulci. *Bauh. pin. 444.*

*Habitat in Europæ australioris l. cis. elevatis. ‡*

*Folia vernantia convoluta.*

infestitia.

12. PRUNUS pedunculis geminis, foliis ovatis subtus villosis convolutis, ramis spinefcentibus. *Amz. acad. 4. p. 273. \* Hudf. angl. 186.*

Pru-

*Prunus sylvestris major. Raj. hist. 1528.*  
*Pruna sylvestria præcocia. Bauh. pin. 444.*  
*Habitat in Germania, Anglia. §*

13. *PRUNUS pedunculis solitariis, foliis lanceolatis spinosa.*  
*glabris, ramis spinosis. Hort. cliff. 186. Fl. suec. 397;*  
*432. Mat. med. 231. Roy. lugdb. 268. Hall. helv. 355.*  
*Prunus sylvestris. Bauh. pin. 444. Tabern. ic. 992.*  
*Habitat in Europæ collibus apricis. §*

## CHRYSOBALANUS.

1. *CHRYSOBALANUS. Jacq. amer. 23. hist. t. 94.* Icaco.  
 2. *Icaco fructu ex albo rufescente. Plum. gen. 44. ic.*  
*158.*  
 3. *Icaco fructu nigro. Plum. gen. 44.*  
 4. *Icaco fructu purpureo. Plum. gen. 44.*  
 5. *Chrysobalanus fruticosus, foliis orbiculatis alternis, flo-*  
*ribus laxè racemosis. Brown. jæn. 250. t. 17. f. 5*  
*Habitat in America meridionali. §*  
*Species an una vel plures vix dixerò, quum tantum*  
*unica mihi visa δ, quæ caule scabro non est, sed*  
*Folia alterna, subrotunda, integerrima, subemargina-*  
*ta. Flores paniculati: Pedunculis exterius rotun-*  
*datis, interius planis.*

## DIGYNIA.

### CRATÆGUS.

1. *CRATÆGUS foliis ovatis inæqualiter serratis sub-* Aria.  
*tus tomentosus. Hort. cliff. 187. Fl. suec. 398; 433.*  
*Vir. cliff. 43. Mat. med. 234. Roy. lugdb. 271. Sauv.*  
*monsp. 306.*  
*Alni effigie, lanato folio major. Bauh. pin. 452.*  
*Aria. Dalch. hist. 202.*  
*Sorbus alpina. Bauh. hist. 1. p. 65.*  
 3. *Cratægus inermis, foliis ellipticis serratis transversaliter suecica,*  
*sinuatis subtus villosis. Fl. lapp. 199.*  
*Sorbus sylvestris anglica. Raj. hist. 1459.*  
*Habitat in Europæ, Helvetiæ frigidis; β. in Svecia,*  
*Anglia. §*  
 2. *CRATÆGUS foliis cordatis spetangulis: lobis infi-* terminall-  
*mis divaricatis.*

*Cratægus foliis cordatis acutis: lacinulis acutis serratis.*

*Hort. cliff. 187. Roy. lugdb. 271. Sauv. monsp. 286.*

*Sorbus torminalis & Cratægus theophrasti. Bauh. h.f. 1. p. 63.*

*Sorbus torminalis. Cam. epit. 162. Edw. av. 212.*

*Mespilus, apii folio, sy. vestris non spinosa, seu Sorbus torminalis. Bauh. pin. 454.*

*Habitat in Anglia, Germana, Helvetia, Burgundia. h*

coccinea.

3. *CRATÆGUS foliis ovatis repando-angulatis serratis grabris. Hort. cliff. 187. Hort. upj. 126. Gron. vi.g. 54. Roy. lugdb. 272.*

*Mespilus foliis cordatis ovatis acuminatis, marginibus acute serratis, ramis spinosis. Mill. d.f. t. 179.*

*Mespilus, apii folio virginiana spinis horrida fructu amplo coccineo. Pluk. alm. 249. t. 46. f. 4.*

*Mespilus virginiana, colore rutilo. Bauh. pin. 453.*

*Mespilus spinosa f. Oxyacantha virginiana maxima.*

*Argl. herb. 49. t. 13. f. 1.*

*Oxyacantha spina sancta dicta. Raj. hist. 1799.*

*Habitat in Virginia, Canada. h*

*Variat cum validis spinis lateralibus & absque spinis.*

*Stipulae lineares. Folia basi sulciformia: gemma obovata, uti stipula & petioli glanduloso-ciliosi.*

Cnus galli.

4. *CRATÆGUS foliis lanceolato-ovatis serratis grabris, ramis spinosis. Kalm. it. 1. p. 244.*

*Mespilus foliis lanceolatis serratis, spinis robustioribus, floribus corymbosis. Mill. d.f. t. 178. f. 2.*

*Mespilus aculeata pyrifolia denticulata splendens, fructu insigni rutilo, virginienfis, Pluk. alm. 249. t. 46. f. 1.*

*Mespilus pruni foliis, spinis longissimis fortibus, fructu rubro magno. Clayt. virg. 55.*

*Habitat in Virginia. h*

tomentosa.

5. *CRAIÆGUS foliis cuneiformi-ovatis serratis subangulatis subtus villosis; ramis spinosis.*

*Mespilus inermis, foliis ovato-oblongis serratis subtus tomentosis. Gron. vi.g. 55.*

*Mespilus caroliniana, apii folio, vulgari similis major, nuctu luteo. Trew. chret. t. 17.*

*Mespilus virginiana, grossulariæ foliis. Pluk. phyt. 100. f. 1.*

*Habitat in Virginia. h*

6. CRA-

6. CRATÆGUS foliis lanceolato-ovatis subtrilobis viridis.  
ferratis glabris, caule inermi.  
Mespilus inermis foliis oblongis integris acuminatis  
ferratis parvis utrinque viridibus. *Gron. virg.* 163.  
*Habitat in Virginia.*  $\S$   
*Stipula semicordata.*
7. CRATÆGUS foliis lanceolatis ferratis, caule iner- *indica,*  
mi corymbis squamosis.  
*Habitat in India.*  $\S$   
*Frutex ramis inermibus. Folia lato-lanceolata, obtu-*  
*se ferrata, crassa, uscula, petiolata. Corymbi termi-*  
*nales, pedunculis squamosis, bracteis subulatis.*
8. CRATÆGUS foliis obtusis subtrifidis ferratis. *Hort. Oxyacan-*  
*th.* 188. *Fl. svec.* 399; 434. *Roy. lugdb.* 272.  
Mespilus, apii folio, sylvestris spinosa f. *Oxyacantha.*  
*Bauh. pin.* 454.  
*Oxyacantha f. Spina acuta. Dod. pempt.* 751.  
*Habitat in Europæ pratis apricis duris.*  $\S$
9. CRATÆGUS foliis obtusis subtrifidis subdentatis. *Azarolus.*  
Mespilus apii folio laciniato. *Bauh. pin.* 453.  
Mespilus Aronia veterum. *Bauh. hist.* 1 p. 67.
10. Mespilus orientalis; apii folio, subtus hirsuto. *Poch. Aronia.*  
*orient.* 189. t. 85.  
*Habitat Florentiæ, Monspeli.*  $\S$   
*Differre videtur ab Oxyacantha sola magnitudine, ut*  
*Sorbus domestica a sylvestri; in hac quadruplo majo-*  
*ra omnia; spinæ in culta pauciores & sæpe nullæ.*

## TRIGYNIA.

### SORBUS.

1. SORBUS foliis pinnatis utrinque glabris. *Hall. herb. aucuparia.*  
250.  
Sorbus foliis pinnatis. *Hort. cliff.* 188. *Fl. svec.* 400;  
435. *Mat. med.* 235. *Roy. lugdb.* 274. *Gron. orient.*  
151.  
Sorbus sylvestris, foliis domesticæ similis. *Bauh. pin.*  
415.  
Sorbus sylvestris. *Cam. epit.* 161.  
*Habitat in Europæ frigidioribus, Libano.*  $\S$

2. SOR-

- hybrida.** 2. **SORBUS** foliis semipinnatis subtus tomentosis.  
*Crataegus feminea. Fl. rec. 2. n. 433.*  
*Habitat in Gotlandia. h*  
*Nova arbor e Cratægo Aria 3. & Sorbo Aucuparia,*  
*jed speciem distinctam propagans. Folia pinnata,*  
*at extrema foliola conflunt in unum, textura Arie.*  
*Sapor medius ex dulci acefçens.*
- domestica.** 3. **SORBUS** foliis pinnatis subtus villosis. *Hall. herb.*  
 351.  
*Sorbus sativa. Bauh. pin. 415. Edw. av. t. 211.*  
*Sorbus legitima. Clus. hist. 1. p. 19.*  
*Habitat in Europæ calidioribus. h*  
*Fructus magnitudine minoris pomi Mali.*

## SESUVIUM.

- Portulaca-**  
**strum.** 1. **SESUVIUM.** *Jacq. hist. 1. p. 95.*  
*Portulaca foliis lanceolatis convexis, pedunculis uni-*  
*floris. Sp. pl. 1. p. 446. Læfl. it. 191.*  
*Portulaca curassavica, a. gusto longo lucidoque folio,*  
*procumbens, floribus rufis. Pluk. alm. 303. t. 216,*  
*f. 1. Hrn. par. 212. t. 212.*  
*Portulaca marina latifolia, flore suaverubente. Plum.*  
*sp. c. 6. ic. 223 f. 2.*  
*Portulaca aizoides maritima procumbens, flore purpu-*  
*reo. Sloan. jam. 88. hist. 1. p. 204.*  
*Halimus indicus. Rumph. amb. 6 p. 163. t. 72. f. 1.*  
*Habitat in ludæ maritimis. ☉*

## PENTAGYNIA.

## MESPILUS.

- germanica.** 1. **MESPILUS** inermis, foliis lanceolatis subtus tomentosis, floribus sessilibus solitariis. *Hort. cliff. 189.*  
*Hort. ups. 129. Mat. med. 236. Roy. lugdb. 270.*  
*Bachm. hist. 174.*  
*Mespilus germanica, folio laurino non ferrato. Bauh.*  
*p. n. 453.*  
*Mespilus. Dod. pempt. 801.*  
 2. **Mespilus**, folio laurino, major. *Bauh. pin. 453.*  
*Habitat in Europa australi. h*  
*Rami villosi. Folia ovali-lanceolata; ad apicem ser-*  
*rata, subvillosa, Petiolis brevissimis canaliculatis. Ca-*  
*ly.*

lyces terminales, hirsuti, bractea longitudine corollæ.

2. **MESPILUS** spinosa, foliis lanceolato-ovatis crenatis, calycibus fructus obtusis. *Hort. cliff.* 189. *Vir. cliff.* 44. *Roy. lugdb.* 271. *Pyracantha.*

*Oxyacantha dioscoridis* f. *Spina acuta*, pyri folio. *Bauh. pin.* 454.

*Uva ursi.* *Dalech. hist.* 164.

*Habitat in Galloprovinciæ, Italiæ sepibus.* §

3. **MESPILUS** inermis, foliis lanceolatis crenatis subarbutifolia. *Hort. cliff.* 189. *Roy. lugdb.* 271. *Cratægus virginiana*, foliis arbuti. *T. Mill. dict.* 1, 109. *Sorbus virginiana*, folio arbuti. *Herm. lugdb.* 578. t. 669.

*Sorbus aucuparia virginiana*, foliis arbuti. *Breyn. prodr.* 1. p. 15.

*Habitat in Virginia.* §

4. **MESPILUS** inermis, foliis ovalibus ferratis, caulis hirsutis. *Mill. dict.* 178. f. 1. *Amelanchier.*

*Mespilus inermis*, foliis ovalibus ferratis acutis. *Hort. cliff.* 189. *Roy. lugdb.* 271.

*Alni effigie*, lanato folio, minor. *Bauh. pin.* 452.

*Vitis idæa* III. *Clus. hist.* 1. p. 75.

*Habitat in Helvetia, Austria, Galloprovincia.* §

*Tenella* adhuc herba ramis percolis p. dunculis foliis subtus villosa tomentosa est, adulta fructifera glabra lanam exuit.

5. **MESPILUS** inermis, foliis ovato-oblongis glabris ferratis. *canadensis.*

*Mespilus inermis*, foliis subtus glabris obverse-ovatis.

*Gron. vurg.* 54.

*Habitat in Virginia, Canada.* §

6. **MESPILUS** inermis, foliis ovalibus ferratis glabris, floribus capitatis, bracteis diducis linearibus. *Chama Mespilus.*

*Cratægus* foliis ovalibus utrinque glabris plicatis indivisis ferratis. *Hort. cliff.* 497. *Vir. cliff.* 43. *Roy. lugdb.* 272. *Hall. helv.* 353.

*Cotoneaster* folio oblongo serrato. *Bauh. pin.* 452.

*Cotoneaster* torte generi. *Clus. hist.* 1. p. 63.

*Habitat in Alpibus Austriacis, Pyrenaicis.* §



*Folia pruni argute serrata: serraturis versus apicem inflexis tenuissimis, utrinque nuda. Sripulæ lineares, petiolis longiores, deciduæ. Corymbus ex alis pedunculatus, conglobatus in capitulum, pedunculo subvillosa, foliis dimidio brevior. Bractææ inter flores, lineares, flore longiores, coloratæ, deciduæ.*

**Cotoneaster.** 7. **MESPILUS** inermis, foliis ovatis integerrimis. *Hort. cliff.* 189. *Fl. suec.* 403; 438. \* *Roy. lugdb.* 271. *Hall. helv.* 352.

Cotoneaster folio rotundo non serrato. *Bauh. pin.* 452.

Chamæ Mespilus gesneri. *Clus. hist.* 1. p. 60.

Chamæ Mespilus cordi. *Bauh. pin.* 452.

*Habitat in Europæ frigidioris collibus apricis, inque Pyrenæis, Ararat. h*

## P Y R U S.

**communis.** 1. **PYRUS** foliis serratis, pedunculis corymbosis.  
*Pyrus foliis serratis, pomis basi productis Hort. cliff.*  
 190. *Hort. ups.* 130. *Fl. suec.* 401; 436. *Roy. lugdb.*  
 266. *Hall. helv.* 351.

*Pyrastr.* *Pyrus sylvestris. Bauh. pin.* 439. *Dod. pempt.* 351.

*falerna.* 2. *Pyrus Bergamotta gallis. Bauh. hist.* 1. p. 44.

*pompejana.* 3. *Pyrus boni christiani. Bauh. hist.* 1. p. 44.

*favoria.* 4. *Pyrus jesu s. moschatellina rubra. Bauh. hist.* 1. p. 44.

*Volema.* 5. *Pyra dorfalia eademque liberalia dicta. Bauh. hist.* 1.  
 p. 53.

*Habitat in Europa. h*

*Varietates hujus 80 videndæ in Milleri dictionario Hortulan.*

**Malus.** 2. **PYRUS** foliis serratis, umbellis sessilibus.  
*Pyrus foliis serratis, pomis basi concavis. Hort. cliff.*  
 189. *Hort. ups.* 130. *Fl. suec.* 402; 437. *Nat. med.*  
 237. *Roy. lugdb.* 266. *Hall. helv.* 351.

*sylvestris.* *Malus sylvestris Bauh. pin.* 433. *Dod. pempt.* 790.

*paradisæa.* 3. *Malus pumila, quæ potius frutex quam arbor. Bauh.*  
*pin.* 433.

*prasomila.* 4. *Malus prasomila. Bauh. pin.* 433.

*rubelliana.* 5. *Malus sativa, fructu sanguinei coloris ex austero sub-*  
*dulci. Tournes. inst.* 635.

*cestiana.* 6. *Mala curtispindula dicta. Bauh. hist.* 1. p. 21.

*cavillea.* 7. *Mala sativa, fructu magno intense rubente, violæ*  
*odore. Tournes. inst.* 635.

r. Po-

## ICOSANDRIA PENTAGYNIA.

687

1. *Poma orbiculata. Ruell. stirp. Epirotica.*  
*Habitat in Europa. h*
3. **PYRUS** foliis ferrato-angulatis, umbellis. peduncu- coronaria.  
 latis.  
*Pyrus foliis ferrato-angulosis. Sp. pl. 1. p. 480. Kalm.*  
*it. 3. p. 10.*  
*Malus sylvestris, floribus odoratis. Gron. virg. 55.*  
*Habitat in Virginia. h*  
*Umbella pedunculata. Pedunculi glabri. Calyces gla-*  
*bri, interne tomentosi. Folia Mali, sed magis glabra*  
*& argutius serrata.*
4. **PYRUS** foliis integerrimis, floribus solitariis. *Hort. Cydonia.*  
*cliff 160. Mat. med. 238. Roy. lugdb. 267. Kalm.*  
*it. 3. p. 107.*  
*Malus cotonea sylvestris. Bauh. pin. 435.*  
*β. Malus cotonea major. Bauh. pin. 434.*  
*γ. Malus Cotonea minor. Bauh. pin. 434.*  
*Cotonea & Cydonia. Lob hist. 580.*  
*Habitat ad ripas petrosas Danubii. h*  
*Calyx patens, serratus, longitudine corollæ.*

## TETRAGONIA.

1. **TETRAGONIA** foliis linearibus. *Roy. lugdb. 250. fruticosa*  
*Tetragonia. Hort. cliff. 188. Hort. upf. 126.*  
*Tetragonocarpus africana, ruti ans, foliis longis & an-*  
*gustis. Comm. hort. 2. p. 205. t. 103.*  
*Evonymo affinis africana, portulacæ folio. Seb. thes. 2.*  
*p. 13. t. 11. f. 8.*  
*Habitat in Æthiopia. 2.*  
*Pedunculi terni, unilori, ex alis inter caulem & ra-*  
*mulum.*
2. **TETRAGONIA** foliis ovatis. *Roy. lugdb. 450. herbacea.*  
*Tetragonocarpus africana, radice magna crassa & car-*  
*nosa. Comm. hort. 2. p. 203. t. 102.*  
*Habitat in Æthiopia. h*

## MESEMBRYANTHEMUM.

\* *Albis corollis.*

1. **MESEMBRYANTHEMUM** foliis alternis tereti- nodiflorum.  
 uiculis obtusis basi ciliatis. *Hort. upf. 129*  
 Kali

*Kali crassulæ minoris foliis. Bauh. pin. 289. Mori. hist. 2. p. 610. f. 5. t. 33. f. 7.*

*Kali neapolitanum aizoides repens. Col. ecphr. 2. f. 72. t. 73.*

*Kali II. Alp. ægypt. 125. t. 127.*

*Habitat in Ægypto, Neapoli, locis arenosis editis, quæ ab æstu maris adsparguntur. ☉*

crystallinum.

2. MESEMBRYANTHEMUM foliis alternis ovatis papulosis undulatis. *Hort. cliff. 216. n. 1. Hort. 427. 127. Roy. lugdb. 281.*

*Meembryanthemum crystallinum, plantaginis folio undulato. Dill. elth. 231. t. 180. f. 221.*

*Ficoides africana, folio plantaginis undulato micis argenteis adsperso T. Bradl. succ. 5. t. 15. f. 48.*

*Habitat in Africa? ☉*

*Caulis primum erectus, digitalis foliis oppositis: Rami horizontales, longissimi, oppositi, prostrati, alterne ramosi, Foliis alternis.*

copticum.

3. MESEMBRYANTHEMUM foliis semiteretibus papulosis distinctis, floribus sessilibus axillaribus, calycibus quinquefidis.

*Kali ægyptiacum, foliis valde longis hirsutis. Bauh. pin. 286.*

*Kali III. ægypt. 129. t. 128.*

*Habitat in Ægypto. ☉*

*Statura & magnitudo M. nodiflori. Caulis trichotomus, herbaceus. Folia opposita, semiteretia, papulosa, ramulis terminantia, nulla in internodiis, longitudine internodiorum. Flores sessiles, solitarii, in divaricatura ramorum. Calyx quinquefidus, corolla longior. Capsula turbinata, retusa, quinqueangularis angulis dilatatis, compressis. Semina subrotunda, subcochleata, striata. Kali 3. Alp. videtur potius ad hanc speciem pertinere, quam ad Plantaginem indicam, quamvis figura vitiosa & planta minime hirsuta.*

geniculiflorum.

4. MESEMBRYANTHEMUM foliis semiteretibus papulosis distinctis, floribus sessilibus axillaribus, calycibus quadrifidis.

*Meisenbr. ramis undique papillofis folio crassioribus. Hort. cliff. 218. n. 22. Roy. lugdb. 285.*

Me-

**Mesemb. capense** geniculiflorum neapolitanum creditum. *Dill. elth.* 271. t. 205. f. 261.

**Ficoides neapolitana**; flore candido. *H. Bradl. succ.* 5: t. 17. f. 34.

**Ficoides capense**; folio tereti; flore albidō. *Pet. gaz.* t. 78. f. 8.

*Habitat ad Cap. b. Spei.* b

*Calyx quadrifidus. Styl. 4. Fructus quadrilocularis.*

5. **MESEMBRYANTHEMUM** foliis semicylindricis noctiflorum; impunctatis distinctis; floribus pedunculatis; calycibus quadrifidis.

**Mesembr. foliis subulatis semicylindricis lævibus vix connatis apice recurvis.** *Hort. cliff.* 220. n. 28. *Roy. Lugdb.* 286.

**Mesembr. noctiflorum**, flore intus candido extus phoeniceo odoratissimo. *Dill. elth.* 273. t. 206. f. 262.

- β. **Mesembr. noctiflorum**, flore intus candido extus stramineo odoratissimo. *Dill. elth.* 274. t. 206. f. 263.

*Habitat ad Cap. b. Spei.* b

*Calyx huic quadrifidus. Styl. 4. Fructus quadrilocularis.*

6. **MESEMBRYANTHEMUM** foliis semiteretibus impunctatis recurvis distinctis congestis; calycibus terminalibus digitiformibus. *splendens.*

**Mesembr. foliis confertis splendentibus**, flore pallido: *Dill. elth.* 270. t. 204. f. 260.

**Ficoides capensis frutescens**; foliis teretibus confertis; glaucis, flore albo: *Bradl. succ.* 1. p. 7. t. 6.

*Habitat ad Cap. b. Spei.*

**Caulis punctis albidis in ramis non ultimis; Folia triquetra-teretiungula, apice patula acuta: axillaria vero incurva. Corolla flavescenti-albida. Calyces erecti, digitiformes, mutici.**

7. **MESEMBRYANTHEMUM** foliis subulatis scabrido-punctatis connatis apice patulo; caule erecto, corymbo trichotomo. *umbellatum*

**Mesembr. frutescens**, floribus albis umbellatis: *Dill. elth.* 277. t. 208. f. 266.

**Ficoides africana erecta teretifolia**; floribus albis umbellatis: *Herm. par.* 166. t. 166. *Bradl. succ.* 4. p. 12. t. 34. *Pluk. phyt.* 117. f. 1.

*Habitat ad Caput b. Spei.* b

calamiforme 8. **MESEMBRYANTHEMUM** acaule, foliis subteretibus adscendentibus impunctatis connatis, floribus octagynis. *Mill. dict.* 176. f. 1.

Mesembr. calamiforme. *Dill. elth.* 239. t. 186. f. 228.

Ficoides capensis humilis, cepææ folio, flore stramineo. *Bradl. succ.* 2. p. 10. f. 19.

*Habitat ad Cap. b. Spei.* 24

*Folia quamvis basi connata recta, tamen sæpe horizontaliter patentia, apice recurva, superficie punctato-scabrida.*

Tripolium. 9. **MESEMBRYANTHEMUM** foliis alternis lanceolatis planis impunctatis, caulibus laxis simplicibus, calycibus pentagonis. *Hort. cliff.* 217. n. 11. *Hort. upf.* 128. *Roy. lugdb.* 283.

Mesembr. tripolii folio, flore argenteo. *Dill. elth.* 230. t. 179. f. 220.

Ficoides africanum procumbens, tripolii succulento folio, flore candido. *Bradl. succ.* 5. p. 14. t. 47.

Ficoides africanum, Mesembryanthemum s. Ficus aizoides major procumbens, tripolii folio magis succulento denticulis fimbriato. *Pluk. mant.* 77. t. 329. f. 4.

*Habitat ad Cap. b. Spei.* ♂. 24

\* \* *Rubicundis corollis.*

bellidiflorum. 10. **MESEMBRYANTHEMUM** acaule, foliis triquetris linearibus impunctatis apice trifariam dentatis. *Hort. cliff.* 218. n. 13. *Roy. lugdb.* 283.

Mesembr. bellidiflorum. *Dill. elth.* 244. t. 189. f. 233.

*Habitat ad Cap. b. Spei.* 24

deltoides. 11. **MESEMBRYANTHEMUM** foliis deltoidibus triquetris dentatis impunctatis distinctis. *Hort. cliff.* 218. n. 14. \* *Roy. lugdb.* 284.

Mesembr. deltoides & dorso & lateribus muricatis. *Dill. elth.* 255. t. 195. f. 246.

Ficus aizoides africana erecta, foliis triangulari breviusculo fimbriato, floribus roseis odoratis. *Volk. h. b.* 223. t. 224. f. 5.

Mesembr. deltoides, dorso & lateribus muricatis, majus. *Dill. elth.* 254. t. 196. f. 247.

Mesembr. deltoides, non dorso sed lateribus muricatis. *Dill. elth.* 253. t. 195. f. 243. 244.

*Ha.*

*Habitat ad Cap. b. Spei. h*

*Pedunculi uniflori, diphylli. Corollæ dilute purpureæ, odoratæ, non calyce longiores. Stamina erecta in conum, alba. Antheris flavis.*

12. **MESEMBRYANTHEMUM** foliis subovatis *par-* *barbatiss.*  
pulosis distinctis apice barbatis. *Hort. cliff.* 216. n.  
5. \* *Hort. ups.* 127. *Roy. lugdb.* 282. *Mill. dict.* t.  
176. f. 3.

*Mesembr. radiatum, ramulis prolixis recumbentibus.*  
*Dill. elth.* 245. t. 190. f. 234.

*Ficoides capensis, tereti folio: apicibus hirtis. Pet. gaz.*  
*t.* 77. f. 6.

*Ficoides capensis frutescens, folio tumido extremitate*  
*stellata, flore purpureo: Bradl. succ.* 1. p. 6. t. 5.  
3 15.

*Ficus aizoides, folio tereti in villos radiatos abeunte,*  
*flore rubro. Volk. hesp.* 222. t. 124. f. 6.

13. *Mesembr. radiatum humile, foliis minoribus. Dill.*  
*elth.* 246. t. 190. f. 235.

γ. *Mesembr. radiatum humile, foliis majoribus. Dill.*  
*elth.* 248. t. 190. f. 236.

*Habitat ad Cap. b. Spei. h*

*Calycis foliola apice barbata, æque ac folia plantæ,*  
*demonstrant Perianthii ortum. Petala minima interio-*  
*ra, quæ cingunt stamina, alba. Varietas 3, hybrida*  
*forte, differt ab 2. Calyce hexaphyllo, nec 5 phyllo:*  
*Stylis Loculisque 6, nec 5. f. 8. Pedunculis hispidis*  
*pilosis, nec lævibus. Foliorum oppositionibus imbrica-*  
*tis, nec remotis.*

13. **MESEMBRYANTHEMUM** foliis cylindricis *par-* *hispidiss.*  
pulosis distinctis, caule hispido: *Mill. dict.* t. 176.  
f. 3.

*Mesembr. caule hispido. Hort. cliff.* 217. n. 7. \* *Roy.*  
*lugdb.* 282.

*Mesembr. pilosum micans, flore saturate purpureo:*  
*Dill. elth.* 249. t. 214. f. 278.

β. *Mesembr. pilosum micans, flore purpureo pallidiore.*  
*Dill. elth.* 290. t. 214. f. 280, 279.

γ. *Mesembr. pilosum micans, flore purpureo striato:*  
*Dill. elth.* 291. t. 215. f. 281.

*Ficoides s. Ficus aizoides africana, folio longo tenui,*  
*flore aurantio. Bradl. succ.* 4. p. 13. t. 35.

*Ficus aizoides teretifolia*, foliis crystallino rore elegantèr conspersis, floribus dilute roseis. *Volk. hesp.* 221. t. 224. f. 2.

*Habitat ad Cap. b. Spei.* §

*Pedunculus valde deorsum scaber magis visu quam tactu.* Calyx muticus. Flos odoratus, simillimus *M. barbati* (a quo forte olim ortum), densa barba foliorum, sed cauli inspersa.

- villosum. 14. **MESEMBRYANTHEMUM** foliis pubescentibus connatis impunctatis, caule piloso. *Hort. cliff.* 217. n. 6. \* *Roy. lugdb.* 282.

*Habitat ad Cap. b. Spei.* §

Folia linearia, semiteretia, canaliculato-plana, opaca, margine, versus basin imprimis, villis adperso. Caulis pubescens villis vix exstantibus. Flores non vidi.

- scabrum. 15. **MESEMBRYANTHEMUM** foliis subulatis distinctis subtus undique muricatis, calycibus muticis. *Hort. cliff.* 219. n. 20. \* *Hort. ups.* 129.

*Mesembr. purpureum scabrum*, staminibus collectis. *Dill. elth.* 260. t. 197. f. 251.

*Habitat ad Cap. b. Spei.* §

- emarginatum. 16. **MESEMBRYANTHEMUM** foliis subulatis congestis subscabris, calycibus spinosis, petalis emarginatis. *Mesembryanthemum purpureum scabrum*, staminibus expansis. *Dill. elth.* 259. t. 197. f. 250.

*Ficoides capensis*, triangulari folio acuto, flore purpureo. *Pet. gaz.* t. 77. f. 3.

*Habitat ad Cap. b. Spei.* §

- uncinatum. 17. **MESEMBRYANTHEMUM** articulis caulinis terminatis in folia connata acuminata subtus dentata. *Hort. cliff.* 218. n. 16. *Roy. lugdb.* 284.

*Mesembr. perfoliatum*, foliis minoribus diacanthis. *Dill. elth.* 250. t. 193. f. 239.

*Ficoides asra*, folio triangulari glauco perfoliato brevissimo apice spinoso. *Bradl. succ.* 3. p. 8. t. 27. 26. *Burm. afr.* t. 26. f. 3.

- β. *Mesembr. perfoliatum*, foliis majoribus triacanthis. *Dill. elth.* 251. t. 193. f. 240.

*Ficoides africana* frutescens perfoliata, folio triangulari glauco punctato. *Bradl. succ.* 5. p. 13. t. 46.

*Habitat ad Cap. b. Spei.* §

18. ME-

18. **MESEMBRYANTHEMUM** foliis tereti-triquetris punctatis distinctis, spinis ramosis. *Hort. cliff.* 216. n. 3. \* *Roy. lugdb.* 281. spinosum.  
*Mesembr. frutescens, ramis triacanthis. Dill. elth.* 276. t. 208. f. 265.  
*Ficoides africana, aculeis longissimis & foliis nascentibus ex foliorum alis. Tournef. afl.* 1705. p. 316. *Bradt. succ.* 41. t. 39.  
*Habitat ad Cap. b. Spei.* 5
19. **MESEMBRYANTHEMUM** foliis subulatis papulosis distinctis apice patulis, radice capitata. *Hort. cliff.* 216. n. 4. \* *Roy. lugdb.* 282. tuberosum.  
*Mesembr. frutescens, radice ingenti tuberosa. Dill. elth.* 275. t. 207. f. 264.  
*Habitat ad Cap. b. Spei.* 5
20. **MESEMBRYANTHEMUM** foliis semiteretibus subulatis glabris distinctis internodio longioribus, *Hort. cliff.* 220. n. 26. *Hort. upf.* 128. *Roy. lugdb.* 286. tenuifolium.  
*Mesembr. tenuifolium procumbens, folio tenuiore viridi, flore coccineo. Dill. elth.* 264. t. 201. f. 256.  
*Ficoides capensis humilis teretifolia, flore coccineo. Bradt. succ.* 1. p. 13. t. 9.  
*Ficoides africana minor procumbens, folio tenuiore viridi, flore coccineo. Moris. hist.* 3. p. 507. f. 12. t. 8. f. 6.  
*Habitat ad Cap. b. Spei.* 5
21. **MESEMBRYANTHEMUM** foliis subtriquetris compressis incurvatis punctatis distinctis congestis basi marginatis. *Hort. cliff.* 220. n. 29. *Roy. lugdb.* 287. stipulaceum.  
*Mesembr. frutescens, flore purpureo rariore. Dill. elth.* 279. t. 209. f. 267; 268.  
*Habitat ad Cap. b. Spei.* 5  
*Caulis florens rigidus, subgriseus. Folia linearia, carnosa, lateribus valde compressa & punctis pellucidis adpersa, incurva, supra planiuscula, mucronata. Flores subcorymbosi, purpurei, Staminibus luteis.*
22. **MESEMBRYANTHEMUM** foliis semicylindricis impunctatis connatis apice triquetris, caule repente semicylindrico. *Hort. cliff.* 217. n. 9. \* *Roy. lugdb.* 283. crassifolium.  
*Mesembr. crassifolium repens, flore purpureo. Dill. elth.* 266. t. 201. f. 257.



*Ficoides africana* reptans, folio triangulari viridi, flore saturate purpureo. *Bradl. succ.* 4. p. 16. t. 38.  
*Habitat ad Cap. b. Spei.* 24

falcatum. 23. **MESEMBRYANTHEMUM** foliis subacinaciformibus incurvis punctatis distinctis, ramis teretibus.  
*Hort. cliff.* 219. n. 19. \* *Roy. lugdb.* 285.

*Mesembr. falcatum* minimum, flore purpureo parvo.  
*Dill. elth.* 288. t. 213. f. 276.

*Ficoides* asra, folio triangulari ensiformi brevissimo, flore dilute purpurascente filamentoso. *Bradl. succ.* 5. p. 9. t. 42.

*Habitat ad Cap. b. Spei.* 25

*Corolla purpurea, parva, petalis distocatis.*

glomeratum. 24. **MESEMBRYANTHEMUM** foliis teretiusculis compressis punctatis, caule paniculato multifloro.  
*Mesembryanthemum falcatum* minus, flore carneo minore. *Dill. elth.* 281. t. 213. f. 274.

*Habitat ad Cap. b. Spei.* 25

*Caulis diffusus, paniculatus. Folia angulis obtusis trigona, glauca punctis diaphanis. Ramulorum rudimenta ex alis, foliis confertissimis incurvis. Calyx longius pedunculatus; Germine subverrucoso punctis eminentibus obsoletis; Laciniis calycinis apice membranaceis; sub apice spina patula. Corolla purpurea M. deltoideis.*

lanceolatum. 25. **MESEMBRYANTHEMUM** foliis semicylindricis recurvis congestis basi interiore gibbis connatis, caule pendulo.

*Mesembryanthemum loreum.* *Dill. elth.* 264. t. 200. f. 255.

*Habitat ad Cap. b. Spei.* 24

*Flores rarissimi prodire apud Engelb. Gother, unde patuit distinctum a M. corniculato plantam esse. Flos ex alis foliorum infimis rami, pedunculo brevi; Calyce aequali, mediocri; Corolla purpurea petalis lanceolato-linearibus; Stamina albidis.*

filamentosum.

26. **MESEMBRYANTHEMUM** foliis æquilateri-tri-  
quetris acutis subpunctatis subconnatis angulis scabris, ramis hexagonis.

*Mesembr. falcatum* majus, flore purpureo mediocri.  
*Dill. elth.* 285. t. 211. f. 273.

*Habitat ad Cap. b. Spei. 24*

Folia punctata, vix connata. Flores sessiles, pentagyni. Corollæ misere, parvæ, obsolete violaceæ, petalis subulatis nec lanceolatis.

27. MESEMBRYANTHEMUM foliis acinaciformibus impunctatis connatis : angulo carinali scabris, petalis lanceolatis. *Hort. cliff.* 219. n. 18. \* *Roy. lugdb.* 284. acinaciforme.

Mesembr. acinaciforme, flore amplissimo purpureo.

*Dill. elth.* 282. t. 211. f. 270. & t. 212. f. 271.

*Habitat ad Cap. b. Spei. 24*

28. MESEMBRYANTHEMUM foliis acinaciformibus obtusis impunctatis connatis apice spinosis, caule ancipiti. forficatum.

*Habitat ad Cap. b. Spei. 24*

Caulis debilis. Flores non vidi, affine *M. acinaciformi*.

\* \* \* Luteis corollis.

29. MESEMBRYANTHEMUM foliis æquilateri-tri- edule.  
quetris acutis strictis impunctatis connatis carina  
subferratis, caule ancipiti.

Mesembryanthemum falcatum majus, flore amplo luteo. *Dill. elth.* 283. t. 212. f. 272.

Ficoides f. Ficus aizoides africana major procumbens, triangulari folio, fructu maximo eduli. *Herm. lugdb.* 244. t. 245. *Morif. hist.* 3. p. 506. f. 12. t. 7. f. 1.

*Habitat ad Cap. b. Spei. 24. 6*

Folia crassitie digiti, nitida, recta, triquetra, æquilatera, longa, basi connata, carina cartilaginea scabra.

30. MESEMBRYANTHEMUM foliis subulatis lævi- bicolorum.  
bus punctatis distinctis, caule frutescente, corollis  
bicoloribus. *Mill. dict.* t. 177. f. 1.

Mesembr. tenuifolium frutescens, flore croceo. *Dill. elth.* 267. t. 202. f. 258.

Ficoides f. Ficus aizoides africana minor erecta, triangulari folio viridi, flore intus aureo; foris purpureo. *Herm. lugdb.* 250. t. 249. *Morif. hist.* 3. p. 507. f. 12. t. 6. f. 14.

Ficoides capensis frutescens, folio tereti punctato, petalis luteis. *Bradl. succ.* 1. p. 8. t. 7.

*Habitat ad Cap. b. Spei. §*

*Corollæ extus purpureæ, intus flavæ, apice extus purpureæ.*

- serratum. 31. **MESEMBRYANTHEMUM** foliis subulatis triquetris punctatis distinctis angulo carinali retrorsum ferratis. *Hort. cliff.* 218. n. 15. *Roy. lugdb.* 284.

Mesembr. serratum flore acetabuliformi luteo. *Dill. elth.* 249. t. 192. f. 238.

*Habitat ad Cap. b. Spei. §*

- micans. 32. **MESEMBRYANTHEMUM** foliis subcylindricis papulosis distinctis, caule scabro. *Hort. cliff.* 222. n. 30. \* *Roy. lugdb.* 287.

Mesembr. micans, flore phoeniceo : filamentis atris. *Dill. elth.* 292. t. 215. f. 282.

Ficoides capensis, tereti folio, flore croceo. *Pet. gar.* t. 7. f. 9.

Ficoides capensis, folio tereti argenteo, petalis plurimis aurantiacis. *Bradl. succ.* 1. p. 9. t. 8.

*Habitat ad Cap. b. Spei. §*

- rostratum. 33. **MESEMBRYANTHEMUM** acaule, foliis semicylindricis connatis externe tuberculatis. *Hort. cliff.* 219. n. 23. *Roy. lugdb.* 285.

Mesembr. rostrum ardeæ referens. *Dill. elth.* 240. t. 186. f. 229.

*Habitat ad Cap. b. Spei. §*

- veruculatum. 34. **MESEMBRYANTHEMUM** foliis triquetro-cylindricis acutis connatis arcuatis impunctatis distinctis. *Hort. cliff.* 220. n. 24. *Hort. ups.* 128. *Roy. lugdb.* 285.

Mesembr. foliis veruculiformibus, floribus mellinis umbellatis. *Dill. elth.* 268. t. 203. f. 259.

*Habitat in Africa. §*

Folia subtriquetro-cylindrica, pellucida, apice purpurascentia. Flores odoratissimi, conferti, parvi, flavi. Corolla vix calyce longior, petalis apice subdentatis. Stamina parva : Filamentis basi villosis.

- glaucum. 35. **MESEMBRYANTHEMUM** foliis triquetris acutis punctatis distinctis, calycinis foliolis ovato-cordatis. *Hort. cliff.* 220. n. 27. *Roy. lugdb.* 283.

Mesembr. scabrum, flore sulphureo convexo. *Dill. elth.* 256. t. 196. f. 248.

*Ficoides africana frutescens*, folio triangulari breviorre glauco. *Morif. hist.* 3. p. 507. f. 12. t. 6. f. 3.

*Ficoides afra*, caule lignoso, erecta, folio triangulari ensiformi scabro, flore luteo magno. *Bradl. succ.* 4. p. 15. t. 37.

*Ficoides* f. *Ficus aizoides africana minor erecta*, folio triangulari glauco, flore luteo. *Herm. lugdb.* 247. t. 248.

*Habitat ad Cap. b. Spei.* 5

Folia mucronata, rore glauca, punctis sparsis non scabra, nisi carina.

36. MESEMBRYANTHEMUM foliis triquetro-semicylindricis scabrido-punctatis, supra basin lineae elevatis connatis. corniculatum.

Mesembr. foliis corniculatis longioribus. *Dill. elth.* 262. t. 199. f. 253, 254.

*Ficoides capensis*, folio triangulari, flore luteo intus pallido. *Pet. gar.* t. 77. f. 10.

3. Mesembr. foliis corniculatis brevioribus. *Dill. elth.* 261. t. 198. f. 252.

*Ficoides africana reptans*, folio triangulari crasso longissimo, flore intus luteo extus aurantiaco. *Bradl. succ.* 4. p. 18. t. 40.

*Habitat in Africa.* 24

Flores decagyni.

37. MESEMBRYANTHEMUM foliis planiusculis lanceolatis impunctatis patentibus distinctis oppositis alternisque remotis. expansum.

Mesembr. tortuosum, foliis sempervivi expansis. *Dill. elth.* 234. t. 182. f. 223.

*Ficoides capensis*, folio lato acuto, flore albo intus luteo. *Pet. gar.* t. 78. f. 10.

*Ficoides africana procumbens*, foliis planis conjugatis lucidis. *Bradl. succ.* 3. p. 7. t. 16.

*Habitat ad Cap. b. Spei.* 24

Flores pentagyni.

38. MESEMBRYANTHEMUM foliis planiusculis oblongo-ovatis subpapillofis confertis connatis, calycibus triphyllis bicornibus. tortuosum.

Mesembr. foliis planis oppositis ovatis acuminatis connatis integerrimis. *Hort. cliff.* 217. n. 10.

Mesembr. tortuosum, foliis sempervivi congestis. *Dill. elth.* 233. t. 181. f. 222.

Ficoides capensis procumbens, oleæ folio, flore albo medio croceo. *Bradl. succ.* 2. p. 7. t. 16.

*Habitat ad Cap. b. Spei.* 24

*Calycis foliola 3 expansa, 2 cornuta. Corolla calyce brevior. Pistillum tetragynam.*

pomeridianum.

39. MESEMBRYANTHEMUM foliis planiusculis lato-lanceolatis lævibus subciliatis distinctis, caule pedunculis germinibusque hirtis.

*Habitat ad Cap. b. Spei.* ☉

*Nova, e seminibus enata primo anno 1762, adolevit & floruit. Caulis herbaeus, palmaris, teres: pilis patulis, confertis, albis. Folia spatulata s. lanceolata petiolo basi dilatato, distincta, lævia, subcarnea, margine inprimis petiolorum subciliata, internodiis longiora, patentia. Pedunculus terminalis, caule longior, hirtus uti caulis. Floris Calyx reliquis major, subquinquepartitus, basi albo-villosus, laciniis lanceolato-foliaceis, corolla longioribus, patulis, basin versus interius fusco-membranaceis. Corolla sulphurea, multiplex petalis linearibus, acutis, numerosissimis: intus setaceis, umbilicum floris obtegentibus instar cancelli. Stamina lutea. Styli 12, totidemque fructus loculamenta. Rami e singulis alii, primo similes, unde dichotoma planta, sed detilis. Floret horis pomeridianis ab 1 — 6. nisi pluvia moneat corollam citius claudi, aut flos etiam provectior diutius expandatur.*

- ringens. 40. MESEMBRYANTHEMUM subacaule, foliis ciliato-dentatis punctatis. *Mill. dist.* 1. 177. f. 2. *Hort. cliff.* 218. n. 12. \* *Roy. lugdb.* 283.

caninum.

Mesembr. riëtum caninum referens. *Dill. elth.* 241. t. 188. f. 231.

Ficoides capensis humilis, folio triangulari prope summitatem dentato, flore luteo. *Bradl. succ.* 2. p. 8. t. 17.

felinum. β. Mesembr. riëtum felinum representans. *Dill. elth.* 240. t. 187. f. 230.

Ficoides afra, folio triangulari ensiformi crasso brevi ad

ad margines laterales multis majoribus spinis aculeato. *Mart. cent.* 30. t. 30.

*Habitat ad Cap. b. Spei.* 5

41. MESEMBRYANTHEMUM foliis dolabriformibus punctatis. *Hort. cliff.* 219. n. 19. *Roy. lugdb.* 284. *Mesembr. folio dolabriformi. Dill. elth.* 248. t. 191. f. 237. *Ficoides capensis humilis, foliis cornua cervi referentibus, petalis luteis, noctiflora. Bradl. succ. r. p.* 11. t. 10.

dolabriforme.

*Habitat ad Cap. b. Spei.* 4

42. MESEMBRYANTHEMUM foliis difformibus punctatis connatis. *Roy. lugdb.* 287. *Mill. dict.* t. 176. f. 2.

difforma.

*Mesembryanth. foliis difformibus, flore luteo. Dill. elth.* 252. t. 194. f. 241, 242.

*Ficoides Mesembryanthemum crassissimo & valde succulento folio nigris punctis notato. Pluk. alm.* 148. t. 325. f. 4.

*Habitat ad Cap. b. Spei.* 5

43. MESEMBRYANTHEMUM acaule, foliis triquetris integerrimis.

albidum.

*Mesembryanthemum foliis robustis albicantibus, Dill. elth.* 243. t. 189. f. 232.

*Ficoides africana, folio triangulari ensiformi glauco crasso, flore amplo aureo. Bradl. succ.* 5. p. 20. t. 42.

*Habitat in Æthiopia,* 4

44. MESEMBRYANTHEMUM acaule, foliis linguiformibus altero margine crassioribus impunctatis.

linguiforme.

*Hort. cliff.* 219. n. 8. *Hort. upf.* 128. *Roy. lugdb.* 282.

*Mesembryanth. folio scalprato. Dill. elth.* 235. t. 183. f. 224.

- β. *Mesembr. folio linguiformi latiore. Dill. elth.* 236. t. 184. f. 224.

- γ. *Mesembr. folio linguiformi angustiore. Dill. elth.* 238. t. 185. f. 226.

- δ. *Mesembr. folio linguiformi longiore. Dill. elth.* 238. t. 185. f. 227.

*Habitat ad Cap. b. Spei.* 4

*Flores Decagyni Calyce quadripartito,*

45. MESEMBRYANTHEMUM foliis alternis subuliginiformibus.

la-me.

latis triquetris longissimis impunctatis. *Hort. cliff.*  
 216. n. 2. \* *Hort. upf.* 129. *Roy. lugdb.* 281.  
 Mesembr. folio pugioniformi, flore aureo stramineo.  
*Dill. elth.* 280. t. 210. f. 269.  
 Ficus capensis, caryophylli folio, flore aureo specioso.  
*Bradd. succ.* 2. p. 5. t. 14.  
*Habitat ad Cap. b. Spei.* ‡  
*Flores decagyni.* (\*)

## AIZOON.

- canariense. 1. AIZOON foliis cuneiformio-vatis, floribus sessilibus  
*Hort. upf.* 127.  
 Aizoon foliis obverse ovatis. *Hort. cliff.* 215. *Roy.*  
*lugdb.* 221.  
 Ficoides procumbens, portulacæ folio. *Niss. aet.* 1711.  
 p. 422. t. 13. f. 1.  
 Kali aizoides canariensis procumbens. *Pluk. alm.* 202.  
 t. 303. f. 4. *Volk. norib.* 236. t. 236.  
*Habitat in Canariis.* ☉  
 hispanicum. 2. AIZOON foliis lanceolatis, floribus sessilibus.  
 Ficodea hispanica annua, folio longiore. *Dill. elth.*  
 143. t. 117. f. 143.  
*Habitat in Hispania, Africa.* ☉  
 Varietas capensis caule foliisque hirsutis, at pagina su-  
 periore minus.  
 panicula- 3. AIZOON foliis lanceolatis, floribus paniculatis. †  
 rum. Aizoon foliis lanceolatis subtus hirsutis. *Roy. lugdb.* 221  
*Habitat in Africa.*

## SPIRÆA.

\* Frutices.

- fallicifolia. 1. SPIRÆA foliis lanceolatis obtusis serratis nudis, flo-  
 ri-

(\*) Mesembryanthemi Species possunt variis modis dispeſci.

Annua: nodiflorum, crystallinum, copticum.

Acaulla; ringens, bellidiflorum, linguiforme, albidum; rostratum, ce-  
 lamiforme, difforme, dolabriforme.Laxa caule pendulo: Tripolium, tortuosum, expansum, lorum, cor-  
 niculatum, acinaciforme, edule, forficatum, filamentosum, crassifol-  
 ium, tenuifolium, pomeridianum.

Fruticosa caule lignoso duro: reliqua: 4, 5, 6, 7, 11, 12, 13, 14,

15, 16, 17, 18, 19, 21, 23, 24, 30, 31, 32, 34, 35, 45.

Alternifolia: nodiflorum, crystallinum, Tripolium, pugioniforme.

Latifolia: crystallinum, Tripolium, expansum, tortuosum, pomeridianum.

Tetragyna: gemiculiflorum, nodiflorum, tortuosum.

Decagyna: linguiforme, pomeridianum, pugioniforme.

ribus duplicato racemosis. *Hort. cliff.* 191. *Hort. upf.* 131. *Roy. lugdb.* 277.

*Spiræa theophrasti* forte. *Clus. hist.* 1. p. 80.

*Frutex spicatus*, foliis serratis salignis. *Bauh. pin.*

475-

*Spiræa salicis* folio longiore serrato, floribus rubris.

*Amm. ruth.* 265.

*Habitat in Sibiria, Tataria.* §

2. *SPIRÆA* foliis lanceolatis inæqualiter serratis subtus tomentosa, tomentosis, floribus duplicato-racemosis.

*Ulmaria pentacarpus*, integris serratis foliis parvis sub-

tus incanis, virginiana. *Pluk. alm.* 393. t. 321. f. 5.

*Raj. suppl.* 330. *Morif. hist.* 3. p. 323.

*Habitat in Philadelphia.* *Kalm.* §

*Differt a præcedenti Foliis magis profunde & inæqualiter serratis; subtus valde tomentosis adeoque crassis; Racemo terminali composito densiori longiori. Floribus demum magis rubris, & Corollis oblongo-minoribus.*

3. *SPIRÆA* foliis ovatis integerrimis, umbellis sessilibus. *Hort. upf.* 131. *hypericifolia.*

*Spiræa* foliis integerrimis, pedunculis simplicibus. *Hort. cliff.* 190. *Roy. lugdb.* 277.

*Prunus sylvestris* affinis canadensis. *Bauh. pin.* 517. *Pluk. alm.* 308. t. 218. f. 5.

*Habitat in Canada.* §

4. *SPIRÆA* foliis ovatis inciso-serratis glabris, umbellis pedunculatis. *chamædryosifolia.*

*Spiræa chamædryos* foliis. *Amm. ruth.* 269.

*Habitat in Sibiria.* *D. Gmelin.* §

5. *SPIRÆA* foliis oblongiusculis apice serratis, corymbis lateralibus. *crenata.*

*Spiræa hispanica*, hyperici folio crenato. *Barr. rar.* 1376. t. 564.

*Spiræa hyperici* folio ampliore, modo integro modo dentato. *Amm. ruth.* 270.

*Habitat in Sibiria, Hispania.* §

*Folia ovato-oblonga, alia apice crenata s. serrata, alia integerrima. Corymbi terminales, perfectissimi, albi, totum fruticem sua copia nivem reddentes.*



- opulifolia.** 6. *SPIRÆA* foliis lobatis serratis, corymbis terminalibus.  
*Spiræa* foliis incisis angulatis, floribus corymbosis.  
*Hort. cliff.* 190. *Gron. virg.* 55. *Roy. lugdb.* 277.  
*Evonymus virginiana*, ribesii folio, capsulis eleganter  
bullatis. *Comm. hort.* 1. p. 169. t. 87.  
*Habitat in Virginia, Canada.* ‡
- serbifolia.** 7. *SPIRÆA* foliis pinnatis: foliolis uniformibus serratis.  
caule fruticoso, floribus paniculatis.  
*Spiræa* foliis pinnatis. *Amén. acad.* 2. p. 352.\*  
*Spiræa* serbi folio tenuiter crenato, floribus in thyrsis  
albis. *Amm. ruth.* 186.  
*Habitat in Sibiria uliginosis.* ‡  
\* \* *Herba.*
- Aruncus.** 8. *SPIRÆA* foliis supradecompositis, spicis paniculatis,  
floribus dioicis.  
*Aruncus.* *Hort. cliff.* 463. *Roy. lugdb.* 278. *Gron. virg.*  
121.  
Barba capræ floribus oblongis. *Bauh. pin.* 163.  
Barba capræ. *Cam. hort.* 26. t. 9.  
*Habitat in Austria, Albornia montanis.* ‡
- Filipendula.** 9. *SPIRÆA* foliis pinnatis: foliolis uniformibus serratis,  
caule herbaceo, floribus cymosis.  
*Filipendula* foliis pinnatis: foliolis uniformibus. *Hort.*  
*cliff.* 191. *Fl. suec.* 404, 439. *Mat. med.* 239. *Roy. lugdb.*  
277.  
*Filipendula vulgaris.* *Bauh. pin.* 163.  
*Filipendula.* *Cam. epit.* 608.  
*Habitat in Europa pascuis.* ‡
- Ulmaria.** 10. *SPIRÆA* foliis pinnatis: impari majore lobato, floribus  
cymosis. *Fl. lapp.* 201. *Hort. cliff.* 191. *Fl. jic.*  
405, 440. *Mat. med.* 241. *Roy. lugdb.* 277.  
Barba capræ floribus compactis. *Bauh. pin.* 164.  
Regina prati. *Dod. pempt.* 57.  
*Habitat in Europa pratis uliginosis, umbrosis.* ‡
- trifoliata.** 11. *SPIRÆA* foliis ternatis serratis subæqualibus, floribus  
subpaniculatis.  
*Filipendula* foliis ternatis. *Hort. cliff.* 191. *Hort. virg.*  
131. *Roy. lugdb.* 277. *Gron. virg.* 55.  
*Ulmaria major trifolia*, flore amplo pentapetalo.  
*virginiana.* *Fluk. alm.* 293. t. 286. f. 5. *Raj. suppl.* 3.  
U

*Ulmaria virginiana trifolia*, floribus candidis amplis  
longis & acutis. *Morif. hist.* 3. p. 323.  
*Habitat in Virginia, Canada.*

# POLYGYNIA.

## ROSA.

1. *ROSA* germinibus ovatis, pedunculis subhispidis, alpina.  
caule inermi.  
*Rosa campestris*, spinis carens, biflora *Bauh. pin.* 484.  
*Rosa rubello flore simplici non spinosa. Bauh. hist.* 2.  
p. 39.  
*Rosa non spinosa. Hall. opusc.* 218.  
*Habitat in alpinis Helvetiæ.* ♀  
*Folia glabra. Calyces simplices. Petala obcordata,*  
*biloba.*
2. *ROSA* germinibus globosis pedunculisque glabris, pimpinelli-  
caule aculeis sparsis rectis, petiolis scabris, foliolis folia.  
obtusis.  
*Habitat forte in Europa.* ♀
3. *ROSA* germinibus globosis pedunculisque glabris, eglanteria.  
caule aculeis sparsis rectis, petiolis scabris, foliolis  
acutis.  
*Rosa aculeata, foliis odoratis subtus rubiginosis. Hall.*  
*helv.* 350. *Fl. svec.* 2. n. 443.  
*Rosa sylvestris, foliis odoratis. Bauh. pin.* 483.  
*Rosa eglanteria. Tabern. ic.* 1087.  
*Habitat in Helvetia, Anglia.* ♀  
*Folia odorata. Flores lutei.*
4. *ROSA* germinibus globosis pedunculisque glabris, cinnamo-  
caule aculeis stipularibus, petiolis subinermibus. mea.  
*Rosa, odore cinnamomi, simplex. Bauh. pin.* 483.  
*Rosa cinnamomea, floribus subrubentibus, spinosa.*  
*Bauh. hist.* 2. p. 39.  
*Habitat in Europa australi.* ♀
5. *ROSA* germinibus globosis hispidis, pedunculis sub- carolina.  
hispidis, caule aculeis stipularibus, petiolis aculeatis.  
*Rosa foliis serratis medio tenus integerrimis. Sp. pl.*  
492.  
*Rosa carolina fragrans, foliis medio tenus serratis. Dill.*  
*elth.* 225. t. 245. f. 316

Ha-

*Habitat in America septentrionali.* †

*Caulis lævis. Aculei stipulares, duo. Folia foliolis 7, oblongo-ovatis, glabris, ferratis, subtus pallidioribus. Petiolis aculeatis. Pedunculi plures, ramosi, inermes, adspersi pilis glandulosis. Calycis foliola indivisa subtus hispida. Petala obcordata, tubicunda. Flos serotinus.*

villosa. 6. *ROSA* germinibus globosis aculeatis; pedunculis hispidis, caule aculeis sparsis, petiolis aculeatis, foliis tomentosis.

*Rosa foliis utrinque villosis; fructu spinoso. Hall. herb. 350. Fl. suec. 2. p. 1295.*

*Rosa sylvestris pomifera major. Bauh. pin. 484.*

*Habitat in Europa.* †

centifolia. 7. *ROSA* germinibus ovatis pedunculisque hispidis, caule hispido aculeato, petiolis inermibus.

*Rosa caule aculeato pedunculis hispidis, calycibus semipinnatis glabris. Sp. pl. 1. p. 491.*

*Rosa multiplex media. Bauh. pin. 482.*

*Rosa centifolia batavica 11. Cus. hist. 1. p. 114.*

*Habitat - - - †*

*Conf. Rosa rubra plena spinosissima; pedunculo mucoso. Mill. dict. 1. 221. f. 1.*

gallica. 8. *ROSA* germinibus ovatis pedunculisque hispidis, caule petiolisque hispido-aculeatis.

*Rosa rubra multiplex. Bauh. pin. 481.*

*Rosa rubra, flore semipleno. Bauh. hist. 2. p. 34.*

versicolor. 9. *Rosa prænestina variegata plena. Mill. dict. 1. 221. f. 2.*

*Rosa versicolor. Bauh. pin. 481.*

*Rosa prænestina alba & versicolor. Bauh. hist. 2. p. 2.*

*Habitat in Europa.* †

sempervirens. 9. *ROSA* germinibus ovatis pedunculisque hispidis, caule petiolisque aculeatis.

*Rosa caule aculeato, foliis quinque glabris perennantibus. Sp. pl. 1. p. 492.*

*Rosa moschata sempervirens Bauh. pin. 482.*

*Rosa sempervirens jungermanni. Cus. hist. 2. app. 2. Dill. elth. 326. t. 246. f. 318.*

*Habitat in Germania.* †

canina. 10. *ROSA* germinibus ovatis pedunculisque glabris, caule petiolisque aculeatis.

Ro-

Rosa caule aculeato, petiollis inermibus, calycibus semipinnatis. *Fl. suec.* 406; 441.

Rosa sylvestris vulgaris, flore odorato incarnato. *Bauh. pin.* 483.

Rosa canina vulgo dicta. *Dod. pempt.* 187.

*Habitat in Europa.* ♀

11. ROSA germinibus ovatis pedunculisque glabris, indica. caule subinermi, petiollis aculeatis.

Rosa subinermis, foliolis quinis subtus tomentosis: impari majori, stipulis obsoletis. *Sp. pl.* 1. p. 492.

Rosa cheufan glabra, juniperi fructu. *Pet. gaz.* 56. t. 25. f. 11.

*Habitat in China.* ♀

Rami inermes; varius armati una alterave spina tenuissima versus folia vel in petiollis. Folia pinnata: foliolis quinis, subtus tomentosis, supra glabris, serratis: extrema duplo majore. Pedunculi longi, nudi, simplices. Calyx incisus, lavis. Fructus magnitudine Sorbi aucupariae.

12. ROSA germinibus ovatis glabris, pedunculis caule. pendula. que hispidis, petiollis inermibus, fructibus pendulis.

Rosa sanguisorbæ majoris folio, fructu longo pendulo.

*Dill. elth.* 325. t. 245. f. 317.

*Habitat in Europa.* ♀

13. ROSA germinibus ovatis glabris, pedunculis hispidis, caule petiolisque aculeatis.

Rosa caule aculeato, pedunculis hispidis, calycibus semipinnatis glabris. *Sp. pl.* 1. p. 492.

Rosa alba vulgaris major. *Bauh. pin.* 482.

3. Rosa alba, flore pleno. *Best. syst. vern.* 6. t. 3. f. 1.

*Habitat in Europa.* ♀

14. ROSA germinibus ovatis glabris, pedunculis caule spinosissima. petiolisque aculeatissimis.

Rosa caule petiolisque aculeatis calycis foliolis indivisis. *Fl. suec.* 407; 442.

Rosa campestris spinosissima, flore albo odorato.

*Bauh. pin.* 483.

*Habitat in Europa.* ♀

Species Rosarum difficillime limitibus circumscribuntur ☹  
forte natura vix eos posuit.

## RUBUS.

\* *Frutescentes.*

idæus.

1. RUBUS foliis quinato-pinnatis ternatisque, caule aculeato, petioliis canaliculatis. *Fl. suec.* 408; 446. *Mat. med.* 242.

Rubus caule erecto hispido, foliis ternatis. *Fl. lap.* 204. *Hort. cliff.* 192. *Hort. ups.* 133. *Roy. lugdb.* 273.

Rubus idæus spinosus. *Bauh. pin.* 479.

- β. Rubus idæus fructu albo. *Bauh. pin.* 479.

- γ. Rubus idæus lævis. *Bauh. pin.* 479.

*Habitat in Europæ lapidosis.* 24. ♂

occidentalis.

2. RUBUS foliis ternatis subtus tomentosis, caule aculeato, petioliis teretibus. *Kalm. it.* 2. p. 283.

Rubus foliis longioribus subtus molli lanugine obductis & incanis, flore & fructu minoribus. *Siccan.* 173. *hist.* 2. p. 109. t. 213. f. 1. *Raj. dendr.* 76.

Rubus idæus, fructu nigro, virginianus. *Dill. elth.* 327. t. 287. f. 319.

Rubus americanus magis erectus, spinis rarioribus, stipite cæruleo. *Pluk. alm.* 325.

*Habitat in Canada.* *Kalm.* 24.

Caulis teret, lævis, aculeis recurvis, nebula cærulea obductus. Petioli teretes aculeis recurvis. Folia ternata intermedio pedicellato, ovata, inciso-serrata, subtus albo-tomentosa. Racemus terminalis. Fructus niger; variat etiam fructu rubro, magis aculeatus grato quam *R. idæi*. Differt imprimis a *R. idæi* cui simillimus: Foliis sæpius ternis, quamvis lateralia foliola sæpe duo evadant & tum ternatis, & pinnatis; Foliolis lateralibus sæpius angulatis; Aculeis alternis, recurvis, nec sparsis & confertis. Petioli etiam aculeati, sed primo intuitu nebula cærulea caulis dignoscitur.

hispidus.

3. RUBUS foliis ternatis nudis, caulibus petiolisque hispidis.

*Habitat in Canada.* *Kalm.* 25.

Caules viticulis longis, procumbentibus lignosis, cum petioliis adpersi setis rigidis. Folia ternata, ovata, inciso-serrata: intermedio pedicellato; Pediculi etiam hispidi.

caesus.

4. RUBUS foliis ternatis subnudis: lateralibus bilobis.

ca.

caule tereti aculeato. *Hort. cliff.* 192. *Fl. suec.* 410;

445. \* *Roy. lugdb.* 273. *Hall. helv.* 444.

*Rubus repens*, fructu cæsius. *Bauh. pin.* 479.

*Rubus minor*. *Dod. pempt.* 742.

*Habitat in Europæ dumetis.* §

*Folia subtus non tomentosa, quamvis sæpe pubescentia.*

5. *RUBUS* foliis quinato-digitatis ternatisque, caule fruticosus. petiolisque aculeatis. *Fl. suec.* 409; 444. \*

*Rubus* caule aculeato, foliis ternatis ac quinatis. *Hort.*

*cliff.* 192. *Gron. virg.* 163. *Roy. lugdb.* 273.

*Rubus vulgaris* f. *Rubus* fructu nigro. *Bauh. pin.* 479.

*Rubus.* *Cam. epit.* 751.

6. *Rubus* vulgaris major, fructu albo. *Raj. angl.* 3. p. 467.

7. *Rubus* flore albo pleno. *Magn. hort.* 175.

*Habitat in sepibus præsertim maritimis Europæ.* §

*Caulis subangulatus, longissimus. Folia subtus subvillosa.*

6. *RUBUS* foliis digitatis denis quinis ternatisque, canadensis. caule inermi. *Mill. icon.* t. 223.

*Habitat in Canada.* *Kalm.*

*Caulis subpurpureus. Foliola lanceolata, utrinque nuda, tenuissima, argute serrata. Bractæ lanceolatae. Stipulæ lineares, acutæ.*

7. *RUBUS* foliis ternatis subtus tomentosis, caule parvifolius. petiolisque aculeis recurvis.

*Rubus moluccanus parvifolius.* *Rumph. amb.* 5. p. 88.

t. 47. f. 1.

*Habitat in India.* *Osbeck.* §

*Caules fruticosi, teretes, incani. Folia subtus albo-tomentosa, venosa; petiolis subtus recurvato-aculeatis.*

*Calyces tomentosi, in racemum digesti.*

8. *RUBUS* foliis simplicibus cordatis lobatis, caule a-moluccanus. culeato decumbente.

• *Rubus moluccanus latifolius.* *Rumph. amb.* 5. p. 88.

t. 47. f. 2.

*Habitat in Amboina.* §

9. *RUBUS* foliis simplicibus palmatis, caule inermi odoratus. multifolio multifloro. *Hort. cliff.* 192. \* *Hort. upf.*

133. *Roy. lugdb.* 274. *Mill. ic.* 223.

*Rubus odoratus.* *Corn. canad.* 149. t. 150.

*Habitat in Canada.* 24. §

\* \* *Herbacei.*

- saxatilis.* 10. RUBUS foliis ternatis flagellis nudis reptantibus herbaceis. *Fl. suec.* 411; 447.  
 Rubus caule repente annuo, foliis ternatis. *Fl. lapp.* 206. *Roy. lugdb.* 274. *Hall. helv.* 344. *Roy. lugdb.* 274  
*Chamærubus saxatilis.*  
*Habitat in Europæ collibus lapidosis.*
- arcticus.* 11. RUBUS foliis ternatis, caule inermi unifloro. *Fl. suec.* 412; 448. \* *Mat. med.* 244.  
 Rubus caule unifloro, foliis ternatis. *Fl. lapp.* 207. *t. 5. f. 2. Hort. cliff.* 292. *Roy. lugdb.* 274.  
 Rubus humilis, flore purpureo. *Buxb. cens.* 5. p. 12. *t. 26. mala.*  
 Rubus trifolius humilis non spinosus, sapore & odore fragariæ, fructu rubro polycocco. *Amm. ruh.* 183.  
*Habitat in Bothnia Sveciæ, Sibiria, Canada.* 24
- Chamæmoros.* 12. Rubus foliis simplicibus lobatis, caule inermi unifloro. *Fl. suec.* 413; 449. \* *Mat. med.* 244.  
 Rubus caule bifolio unifloro, foliis simplicibus. *Fl. lapp.* 208. *t. 5. f. 1. Hort. cliff.* 192. *Roy. lugdb.* 279.  
*Chamærubus foliis ribes.* *Bauh. pin.* 480.  
*Morus norvegica.* *Till. aboenf.* 47. *t. 150. Oed. dan.* 1. 1.  
*Habitat in Sveciæ paludibus uliginosis, turfosis, frutuosus.* 24  
*Planta dioica est.*
- Dalibarda.* 13. RUBUS foliis simplicibus cordatis indivisis crenatis, scapo aphylo unifloro.  
*Dalibarda.* *Sp. pl.* 1. p. 491.  
*Habitat in Canada.* *Kalm.* 24  
*Radix repens. Stolones repentes, herbacei. Folia (viola) cordata, obtusa, crenata, venosa, pilis raris adspersa. Petioli pilosi, longitudine foliorum. Scapi nudi s. aphylli, pilosi, longitudine foliorum, uniflori. Calyx 5partitus. Corolla alba. Semina 5, ovata, nuda. Stamina longitudine corollæ. Anthe Flores masculi & feminei in eadem planta? Habitus junctus conjunctionem cum Rubo, imprimis Chamæmoro, utraque distinguenda a Rubis.*

## FRAGARIA.

- vesca.* 1. FRAGARIA flagellis reptans. *Hort. cliff.* 192. *Hort.* 25.

*upf.* 133. *Fl. suec.* 414; 450. *Mat. med.* 245. *Roy. lugdb.* 274. *Hall. helv.* 243.

*Fragaria vulgaris.* *Bauh. pin.* 326. *Fl. lapp.* 209. *Gron. sylvestris. virg.* 56.

*Fragaria fructu albo.* *Bauh. pin.* 326.

6. *Fragaria fructu parvi pruni magnitudine.* *Bauh. pin.* 327. *pratenfis.*

7. *Fragaria chilensis, fructu maximo, foliis carnosis chilensis. hirsutis.* *Dill. eth.* 145. *t.* 120. *f.* 146.

*Habitat in Europæ borealis sterilibus, duris, apricis.* 24

*Varietas β pratenfis, quæ in hortis sativa audit, distinguitur. Baccæ a calyce non sponte secedente, & ejusdem scrobiculis pro singulis seminibus.*

2. **FRAGARIA** caule erecto suffruticoso, foliis hirsutis. †

*Fragaria arborea, flore herbaceo.* *Zan. hist.*

*Fragaria major vesca, flore herbaceo.* *Morif. hist.* 2. p. 186.

*Habitat in* - - - †

*Folia inter flores distinguunt hanc a prima.*

3. **FRAGARIA** caule decumbente repente. *Roy. lugdb.* sterilis. 274. *Dalib. parif.* 14. *Sauv. monsp.* 177. *Huds. angl.* 195.

*Fragaria sterilis.* *Bauh. pin.* 327.

*Fragaria sterilis f. minime vesca hirsuta minime incana.* *Morif. hist.* 2. p. 186. *f.* 2. *t.* 19. *f.* 5.

*Fragaria sylvestris minime vesca f. sterilis.* *Lob. ic.* 698.

*Habitat in Anglia, Helvetia.* 24

*Caules procumbentes; nullus rectus; Pedunculi axillares, uniflori; Foliola ovalia; Baccæ exsuccæ.*

## POTENTILLA.

\* *Foliis pinnatis.*

1. **POTENTILLA** foliis pinnatis, caule fruticoso. *Hort. fruticosa. cliff.* 193. *Fl. suec.* 416; 451. *Roy. lugdb.* 275. *Huds. angl.* 195. *Kalm. it.* 2. p. 263.

*Pentaphylloides rectum fruticosum eboracense.* *Morif. hist.* 2. p. 193. *f.* 2. *t.* 23. *f.* 5. *Hortul. angl.* 54. *t.* 54.

*Pentaphylloides rectum frutescens.* *Walth. hort.* 95. *t.* 17.

*Pentaphylloides fruticosa elatior.* *Ann. ruth.* 88. *t.* 17.

89. *t.* 18. *f.* 1.

*Habitat in Eboraco Angliæ, Oelandia australi, Sibiria.* †



- Anserina. 2. **POTENTILLA** foliis pinnatis ferratis, caule repente. *Fl. lapp.* 210. *Fl. svec.* 415; 452. *Hort. cliff.* 193. *Mat. med.* 246. *Roy. lugdb.* 275.  
*Potentilla. Bauh. pin.* 321. *Cam. epit.* 758.  
*Habitat in Europæ pascuis; in argillofis argentea.* 24
- sericea. 3. **POTENTILLA** foliis bipinnatis utrinque tomentosis: segmenti parallelis approximatis, caulibus decumbentibus.  
*Potentilla foliis pinnatis hirsutis, pinnis tredecim ovatis crenatis decurrentibus. Hall. gætt. 2. p. 106.*  
*Potentilla foliis duplicato-pinnatis: pinnulis linearibus integerrimis brevibus. Gmel. mss.*  
*Habitat in Sibiria.* 24  
*Habitus foliorum P. anserinae argentea, quamvis minima, cæulis vero & fructificationes P. verna. Radix squamis foliorum rudibus nigris vestita. - Folia pinnata ex 13 circiter foliolis ovatis, tomentosis, sericeis, prætinato-pinnatifidis, & parallelis. Pedunculi pilosi. Caples foliis duplo s. triplo longiores decumbentes absque fermenis, nec repentes, in medio folio unico minori magisque divaricato instructi, teretes, apice Floribus aliquot luteis, alternis, pedunculatis.*
- multifida. 4. **POTENTILLA** foliis bipinnatis: segmentis integerrimis distantibus subtus tomentosis, caule decumbente  
*Pentaphylloides supinum minus, folio glabro non ferrato. Amm. ruth.* 116.  
*Pentaphylloides repens, foliis pinnatis. Buxb. cent.* 1 p. 30. t. 49. f. 1.  
 β. *Pentaphylloides supina minor, foliis alatis hirsutis varie dissectis. Amm. ruth.* 117.  
*Habitat in Sibiria, Tataria, Cappadocia.* 24  
*Habitus multa indicat affinia cum P. argentea. Folia ex septem foliolis: segmentis linearibus s. lanceolatis, quinque s. summum septenis, alternis, at in impari oppositis, a se invicem hiantibus sive apice remotis, subtus tomentosis.*
- fragarioides. 5. **POTENTILLA** foliis pinnatis ternatisque: extimis majoribus, flagellis reptantibus.  
*Potentilla foliis ternatis hirsutis utrinque viridibus lobatis accessoriis. Gmel. mss.*

**Potentilla foliis novenis palmatis apice serratis.** *Hall.*  
*gatt. 2. p. 106.*

*Habitat in Sibiria. 24*

**Bacies Fragariæ sterilis.** *Radix subtuberosa. Folia ferrica petiolis pilosis, pinnata ex 3. 5. vel 7. Foliolis ovatis, oppositis, serratis, basi quo propioribus co minoribus. Stolones decumbentes.*

6. **POTENTILLA foliolis pinnatis alternis: foliolis** *rupestris.*  
*quinis ovatis crenatis, caule erecto. Hort. cliff.*  
*193. It. w. goth. 63; 76. Fl. suec. 2. p. 453.*

*Pentaphylloides erectum. Bauh. hist. 2. p. 398. Moris.*  
*hist. 2. p. 192. f. 2. t. 20. f. 1.*

*Quinquefolium fragiferum. Bauh. pin. 326. Clus. hist. 2.*  
*p. 107. n. 5.*

*Habitat ad latera montium Westrogothiæ, Sibiria,*  
*Germaniæ. 24*

7. **POTENTILLA foliis pinnatis subæqualibus: folio-** *bifurc.*  
*lis oblongis subbifidis: extimis confluentibus.*

*Pentaphylloides orientale supinum, folis alatis bifidis*  
*vel trifidis. Tournesf. cor. 21.*

*Habitat in Sibiria. 24*

8. **POTENTILLA foliis pinnatis, foliolis subrotun-** *pimpinell.*  
*dis dentatis æqualibus, caule erecto. loides.*

*Pentaphylloides orientale erectum, folio & facie pim-*  
*pinellæ. Tournesf. cor. 21. Buxb. cent. 1, p. 30. t. 48.*

*Habitat in Armeniæ saxosis. 24*

9. **POTENTILLA foliis pinnatis, caule dichotomo** *supina.*  
*decumbente. Hort. cliff. 193. Hort. ups. 133. Roy.*  
*lugdb. 275.*

*Pentaphyllum alpinum minus supinum. Pluk. alm. 285.*  
*t. 106. f. 7.*

*Pentaphyllum supinum quorundam, potentillæ facie.*  
*Clus. hist. 2. p. 107.*

*Pentaphylli species flore luteo enneaphyllos hirsuta*  
*Amrosf. phyt. t. 414.*

*Quinquefolio fragifero affinis. Bauh. pin. 326.*

*Habitat ad Moguntiam & in Sibiria. ☉*

\* \* *Foliis digitatis.*

10. **POTENTILLA foliis septenatis lanceolatis serratis recta.**  
*utrinque subpilosis, caule erecto.*

Y y 4

Po-

- Anserina 2. **POTENTILLA** foliis pinnatis serratis, caule repente. *Fl. lapp.* 210. *Fl. suec.* 415; 452. *Hort. cliff.* 193. *Mat. med.* 246. *Roy. lugdb.* 275.  
*Potentilla.* *Bauh. pin.* 321. *Cam. epit.* 758.  
*Habitat in Europæ pascuis; in argillofis argentea.* 24
- sericea. 3. **POTENTILLA** foliis bipinnatis utrinque tomentosis: segmenti parallelis approximatis, caulibus decumbentibus.  
*Potentilla foliis pinnatis hirsutis, pinnis tredecim ovatis crenatis decurrentibus.* *Hall. gætt.* 2. p. 106.  
*Potentilla foliis duplicato-pinnatis: pinnulis linearibus integerrimis brevibus.* *Gmel. mss.*  
*Habitat in Sibiria.* 24  
*Habitus foliorum P. anserinae argentea, quamvis minima, Caulis vero & fructificationes P. verna. Radix squamis foliorum rudibus nigris vestita. Folia pinnata ex 13 circiter foliolis ovatis, tomentosis, sericeis, prætinato-pinnatifidis, & parallelis. Pedunculi pilosi. Caples foliis duplo s. triplo longiores, decumbentes absque sarmentis, nec repentes, in medio folio unico minori magisque divaricato instructi, teretes, apice floribus aliquot luteis, alternis, pedunculatis.*
- multifida. 4. **POTENTILLA** foliis bipinnatis: segmentis integerrimis distantibus subtus tomentosis, caule decumbente.  
*Pentaphylloides supinum minus, folio glabro non serrato.* *Amm. ruth.* 116.  
*Pentaphylloides repens, foliis pinnatis.* *Buxb. cent.* 1. p. 30. t. 49. f. 1.  
 β. *Pentaphylloides supina minor, foliis alatis hirsutis varie dissectis.* *Amm. ruth.* 117.  
*Habitat in Sibiria, Tataria, Cappadocia.* 24  
*Habitus multa indicat affinia cum P. argentea. Folia ex septem foliolis: segmentis linearibus s. lanceolatis, quinque s. summum septenis, alternis, at in impari oppositis, a se invicem hiantibus sive apice remotis, subtus tomentosis.*
- fragarioides. 5. **POTENTILLA** foliis pinnatis ternatisque: extimis majoribus, flagellis reptantibus.  
*Potentilla foliis ternatis hirsutis utrinque viridibus: lobatis accessoris.* *Gmel. mss.*

**Potentilla** foliis novenis palmatis apice serratis. *Hall. Gart. 2. p. 106.*

*Habitat in Sibiria. 2.*

*Bacis Fragariæ sterilis. Radix subtuberosa. Folia ferrica petiolis pilosis, pinnata ex 3. 5. vel 7. Foliolis ovatis, oppositis, serratis, basi quo propioribus eo minoribus. Stolones decumbentes.*

6. **POTENTILLA** foliis pinnatis alternis: foliolis *rupestris.* quinis ovatis crenatis, caule erecto. *Hort. cliff. 193. It. w. goth. 63; 76. Fl. svec. 2. p. 453.*

*Pentaphylloides erectum. Bauh. hist. 2. p. 398. Moris. hist. 2. p. 192. f. 2. t. 20. f. 1.*

*Quinquefolium fragiferum. Bauh. pin. 326. Clus. hist. 2. p. 107. n. 5.*

*Habitat ad latera montium Westrogothiæ, Sibiria, Germaniæ. 2.*

7. **POTENTILLA** foliis pinnatis subæqualibus: foliolis *bifurca.* oblongis subbifidis: extimis confluentibus.

*Pentaphylloides orientale supinum, foliis alatis bifidis vel trifidis. Tournes. cor. 21.*

*Habitat in Sibiria. 2.*

8. **POTENTILLA** foliis pinnatis, foliolis subrotundis *pimpinell.* dentatis æqualibus, caule erecto. *loides.*

*Pentaphylloides orientale erectum, folio & facie pimpinellæ. Tournes. cor. 21. Buxb. cent. 1, p. 30. t. 48.*

*Habitat in Armeniæ saxosis. 2.*

9. **POTENTILLA** foliis pinnatis, caule dichotomo *supina.* decumbente. *Hort. cliff. 193. Hort. ups. 133. Roy. lugdb. 275.*

*Pentaphyllum alpinum minus supinum. Pluk. alm. 285. t. 106. f. 7.*

*Pentaphyllum supinum quorundam, potentillæ facie. Clus. hist. 2. p. 107.*

*Pentaphylli species flore luteo enneaphyllos hirsuta. Amros. phyt. 1. 414.*

*Quinquefolio fragifero affinis. Bauh. pin. 326.*

*Habitat ad Moguntiam & in Sibiria. ☉*

\* \* *Foliis digitatis.*

10. **POTENTILLA** foliis septenatis lanceolatis serratis recta. utrinque subpilosis. caule erecto.

Y y 4

Po-

- Potentilla foliis digitatis, caule erecto corymboso.*  
*Hort. ups. 133. \* Roy. lugdb. 275.*  
*Quinquefolium rectum luteum. Bauh. pin. 325.*  
*Quinquefolium alterum vulgare. Dod. pempt. 116.*  
*Habitat in Italia, Narbona, ad agrorum margines. 2.*  
*Folia utrinque viridia, at adulta fere nuda.*
- argentea. 11. **POTENTILLA** foliis quinatis cuneiformibus incis-  
 sis subtus tomentosis, caule erecto.  
*Potentilla foliis digitatis, caule erecto corymboso. Fl.*  
*svec. 417; 454.*  
*Potentilla foliis digitatis inciso-ferratis, caule erecto.*  
*Hort. cliff. 193.*  
*Quinquefolium folio argenteo. Bauh. pin. 325.*  
*Pentaphyllum minus. Cam. epit. 760.*  
*Habitat in Europæ ruderatis. 2.*
- hirta, 12. **POTENTILLA** foliis septenatis quinatisque cunei-  
 formibus incisfis pilosis, caule adscendente.  
*Quinquefolium montanum rectum hirsutum. Bauh.*  
*pin. 325. prodr. 139.*  
*Habitat Montpelii. D. Sauvages; inque Pyrenæis. 2.*  
*Differt a præcedente Foliis radicalibus septematis, & sul-*  
*tus longis distinctis pilis hirsutis, nec tomentosis, api-*  
*ce argutius furcatis. Pedunculi uniflori, longi.*
- alpularis. 13. **POTENTILLA** foliis septenatis sessilibus stipulæ  
 dilatatæ insidentibus.  
*Potentilla foliis novenis stipulis insidentibus. Gmel. mss.*  
*Habitat in Sibiria. 2.*  
*Caulis adscendens, filiformis, levis, apice divisus. Sti-*  
*pulæ amplexicaules, solitariae, emarginatæ. Foliola*  
*8 ad 9 lineari-lanceolata, subserrata, glabra, ex*  
*emarginatura stipulæ pendentia absque petiolo.*
- aurea. 14. **POTENTILLA** foliis radicalibus quinatis serratis  
 acuminatis; caulinis ternatis, caule declinato. *Amx.*  
*acad. 4. p. 316. \**  
*Potentilla foliis quinatis acute serratis ora sericea, pe-*  
*talis maculatis. Hall. helv. 339. t. 6. f. 4.*  
*Quinquefolium minus repens alpinum aureum. Bauh.*  
*pin. 325.*  
*Habitat in Alpibus Helvetiæ. 2.*
- verna. 15. **POTENTILLA** foliis radicalibus quinatis acute  
 ser-

*ferratis retusis; caulinis ternatis, caule declinato.*

*Fl. svec. 419. 455. \**

*Potentilla foliis quinatis incisif, caule affurgente. Fl. lapp. 212. Hort. cliff. 194. Roy. lugdb. 276.*

*Quinquefolium minus repens luteum. Bauh. pin. 325.*

*Pentaphyllum f. Quinquefolium minus. Tabern. ic. 123.*

*Habitat in Europæ pascuis fœcis, frigidioribus. 24*

16. *POTENTILLA* foliis radicalibus quinatis cuneiformibus ferratis, caulinis suboppositis, ramis filiformibus decumbentibus. *Amæm. acad. 4. p. 274. \* Hudf. angl. 196.*

*Quinquefolium minus repens lanuginosum luteum. Bauh. pin. 325.*

*Quinquefolium IV, flavo flore 2. species. Clus. hist. 2. p. 106. f. 106.*

*Quinquefolio similis enneaphyllos hirsuta. Bauh. pin. 325. prodr. 139.*

*Habitat in Austria, Helvetia, Baldo. 24*

17. *POTENTILLA* foliis quinatis villosis, caule ascendente hirsuto. *canadensis.*

*Quinquefolium canadense humilius. Morif. hist. 2. p. 188.*

*Habitat in Canada. Kalm.*

*Habitus Fragariæ. Folia radicalia quinata: foliolis ovatis, acute serratis, supra pubescentibus, subtus villosis albo-sericeis, petiolis valde albo-hirsutis. Caulis digitalis, ramosus, rarioribus foliis. Corollæ luteæ. Similis P. vernæ, a qua differt Foliis valde pilosis, argute serratis, subrotundis & Caule valde hirsuto.*

18. *POTENTILLA* foliis quinatis apice conniventiferratis, caulibus filiformibus procumbentibus, receptaculis hirsutis. *Hort. cliff. 194. Roy. lugdb. 276.*

*Quinquefolium album. Hall. it. helv. 70.*

*Quinquefolium album majus alterum. Bauh. pin. 325.*

*Quinquefolium tragi. Trag. hist. 507.*

*Pentaphyllum majus, flore albo. Clus. hist. 2. p. 105.*

*Habitat in alpibus Stiriz, Austriæ, Pannoniæ. 24*

19. *POTENTILLA* foliis quinatis apice conniventicaulescens.

Y y 5

fer-

U. B. GENT

ferratis, caulibus multifloris erectis, receptaculis hirsutis. *Amen. acad.* 4. p. 317. \*

Quinefolium album majus. *Bauh. prodr.* 139. caulescens. *Bauh. pin.* 325. *Hall. opusc.* 263. n. 50. 47.

Quinefolium album minus alterum. *Bauh. pin.* 325.

Quinefolium 2. minus, albo flore. *Clus. hist.* 2. p. 105.

*Habitat in Alpibus Helvetiae, Austriae, Stiriae, Horto Dei.* 24.

valderia.

30. **POTENTILLA** foliis septenatis obovatis ferratis tomentosis, caule erecto, petalis calyce brevioribus, receptaculis lanatis.

*Habitat in Alpibus Valderianis & Vinadenfibus.* 24.

Folia radicalia septena: foliolis obovatis, conniventiferratis, subtus valde tomentosis. Caulis erectus paucis foliis ternatis. Flores dichotomi, corymbosi, conferti. Petala alba, calyce breviora.

alpida.

31. **POTENTILLA** foliis quinatis tomentosis conniventitridentatis, caulibus unifloris, receptaculis lanatis. *Amen. acad.* 4. p. 316. \*

Heptaphyllum argenteum alpinum trifoliatum sabaudum. *Bocc. mus.* 2. p. 20. t. 9.

Trifolium alpinum argenteum, perfici flore. *Bauh. pin.* 328. *Pon. bald.* 222. *Raj. hist.* 615.

Comaroides alpina argentea sericea, perfici flore. *Segg. ver.* 1. p. 497.

*Habitat in Baldo. Seguier.* 24.

septans.

32. **POTENTILLA** foliis digitatis, caule repente, pedunculis unifloris. *Fl. suec.* 418; 456. *Mat. med.* 247.

Potentilla foliis digitatis longitudinaliter patentiferratis, caule repente. *Hort. cliff.* 194. *Gron. virg.* 163.

Quinefolium majus repens. *Bauh. pin.* 325.

Quinefolium majus luteum. *Fuchs. hist.* 624.

*Habitat in Europae apricis argilleosis.* 24.

\* \* \* *Foliis ternatis.*

monspellen-  
sis.

23. **POTENTILLA** foliis ternatis, caule ramoso erecto, pedunculis supra genicula enatis. *Hort. upf.* 134.

Pentaphylloides majus erectum, flore luteo. *Morij. hist.* 2. p. 192. f. 2. t. 20. f. 2.

Pen-

*Pentaphylloides fragariæ folio.* *Magn. Monsp.* 304.  
*Fragaria sterilis alpina caulescens.* *Boerh. lugdb.* 1.  
 p. 42.

*Habitat Monspelli.* ☉

*Foliola ovata, obtuse & æqualiter serrata.*

24. **POTENTILLA** foliis ternatis, caule dichotomo, *Norvegia.*  
*pedunculis axillaribus.*

*Potentilla foliis ternatis, caule corymboso erecto.* *Fl. fvec.* 420; 457. \*

*Potentilla foliis ternatis incisif, caule diffuso.* *Fl. lapp.* 211. *Hort. cliff.* 193. *Roy. lugdb.* 275.

*Quinquefolium hirsutum luteum paucioribus laciniis.*  
*Læsc. pruss.* 218. t. 70.

*Habitat in Norvegiæ, Sueciæ, Borussia, Canadæ*  
*agris.* ☉

*Foliola lanceolata, serrata, dentata.*

25. **POTENTILLA** foliis ternatis incisif subtus tomentosis, caule ascendente. *Fl. fvec.* 2. n. 458. \*

*Potentilla foliis ampliter crenatis subtus tomentosis.*  
*Hall. gatt.* 2. p. 108.

*Fragaria sterilis procumbens, foliis betonicæ instar*  
*ferratis.* *Amm. ruth.* 109. t. 14. f. 2.

*Habitat in Alpihus Lapponiæ, Sibiria.*

*Structura P. vernæ, seu folia P. argentiæ, at terna &*  
*latiora, magisque hirsuta, subtus tomentosa nivea. Co-*  
*rolla lutea.*

26. **POTENTILLA** foliis ternatis dentatis utrinque grandiflora.  
*subpilosis, caule decumbente foliis longiore.*

*Potentilla foliis ternatis, petalis cordatis luteis.* *Hall.*  
*helv.* 342. t. 6. f. 2.

*Fragaria sterilis, amplissimo folio & flore, petalis*  
*cordatis.* *Vaill. paris.* 55. t. 10. f. 1.

*Habitat in Helvetia, Sibiria, Pyrenæis.* ☉

*Folia utraque pagina concoloria, nec inferiori argentea.*  
*Caules pedales, ascendentes, rufescentes.*

27. **POTENTILLA** foliis ternatis dentatis utrinque tomentosis, scapo decumbente. *subcaulis.*

*Fragaria sterilis sylvestris sericea f. incana.* *Morif. hist.*  
 2. p. 187.

*Fragariæ affinis sericea incana.* *Bauh. pin.* 327. prodr.  
 139.

*Ha-*



*Habitat in Sibiria, Gallia australi, Granada, Gallo-provinciæ rupe victoriæ. 24*

*Folia ternata, rugosa, dense tomentosa, pilosa, coriacea, cuneiformia, obtusa, dentata. Stipulæ latæ, acutæ, pilosæ. Caulis s. Schæpus decumbens, foliis sæpe longior, biflorus s. uniflorus. Corolla maxima, lutea: petalis emarginatis; basi flavis. Stamina lutea.*

### TORMENTILLA.

*erecta.*

1. **TORMENTILLA** caule erectiusculo, foliis sessilibus  
*Tormentilla. Cam. epit. 685. Fl. lapp. 213. Fl. suec. 421; 459. Mat. med. 249. Hort. cliff. 194. Roy. lugdb. 276.*

*Tormentilla sylvestris. Bauh. pin. 326.*

*Habitat in Europæ pascuis siccis. 24*

*reptans.*

2. **TORMENTILLA** caule repente, foliis petiolatis  
*Huds. angl. 197.*

*Pentaphyllum minus viride, flore aureo tetrapetalo, radicalas in terram e geniculis dimittens. Moris. hist. 2. p. 190.*

*Pentaphyllum reptans alatum, foliis profundius serratis. Plot. oxf. 6. §. 7. t. 9. f. 5.*

*Habitat in Anglia. 24*

### GEUM.

*virginiana.*  
*nana.*

1. **GEUM** floribus erectis, fructu globoso: aristis uncinatis nudis, foliis ternatis. *Hort. cliff. 195. Gron. virg. 56.*

*Caryophyllata virginiana, albo flore minore, radice inodora. Herm. par. III. t. III.*

*Habitat in Virginia, Sibiria. 24*

*Differt a G. urbano simillimo: Petalis albis calyce brevioribus; Floribus florescentibus mutantibus; Pedunculis versus florem incrassatis; Seminibus glabris; Foliis radicalibus subbipinnatis.*

*urbanum.*

2. **GEUM** floribus erectis, fructu globoso villosa: aristis uncinatis nudis, foliis lyratis. *Hort. cliff. 195. Fl. suec. 423; 460. Mat. med. 249. Roy. lugdb. 276.*

*Caryophyllata vulgaris. Bauh. pin. 321.*

*Caryophyllata. Dod. pempt. 137.*

*Habitat in Europæ umbrosis. 24*

*Folia radicalia lyrata in hac, & stipulæ dentatæ.*

3. **GE-**

3. GEUM floribus nutantibus, fructu oblongo: aristis rivale.  
plumosis. *Hort. cliff.* 195. *Fl. suec.* 424, 461. *Roy.*  
*lugdb.* 276.  
Geum rivale. *Fl. lapp.* 216.  
Caryophyllata aquatica, nutante flore. *Bauh. pin.* 321.  
Caryophyllata septentrionalium. *Lob. ic.* 694.  
β. Caryophyllata aquatica altera. *Bauh. pin.* 322.  
*Habitat in Europæ pratis subhumidis.* 24  
*Seminum aristæ apex in hac pilosus, in antecessente*  
*nudus.*
4. GEUM flore inclinato solitario, fructu oblongo: a- montanum.  
ristis plumosis rectis.  
Caryophyllata pinnis confertioribus extrema subre-  
tunda, tubis rectis. *Hall. helv.* 336. \*  
Caryophyllata alpina lutea. *Bauh. pin.* 322.  
Caryophyllata alpina. *Pou. bald.* 342.  
Caryophyllata montana. *Cam. epit.* 727.  
β. Caryophyllata alpina minor. *Bauh. pin.* 322. *prodr.*  
139.  
Caryophyllata alpina minima, flore aureo. *Barr. rar.*  
588. t. 399.  
*Habitat in Alpibus Helvetiæ, Austriæ, Delphinatus.* 24  
*Folia hirsuta. Flos patens. Caulis uniflorus. Petala*  
*lutea.*
5. GEUM foliolis uniformibus incis: alternis mino- reptans.  
ribus, flagellis reptantibus.  
Caryophyllata flagellis foliosis. *Hall. enum.* 19.  
Caryophyllata alpina, apii folio. *Bauh. pin.* 322.  
Caryophyllata alpina tenuifolia incana, flore luteo lon-  
gius radicata. *Barr. rar.* 589. t. 400. *Bocc. mus.* 160.  
t. 128.  
*Habitat in Helvetia & valle Barfiliionensi.*

# D R Y A S.

1. DRYAS floribus pentapetalis, foliis pinnatis. *Amæn. pentapetalis*  
*acad.* 2. p. 353.  
Caryophyllata pentaphyllea. *Bauh. hist.* 2. p. 398.  
*quoad solam figuram.*  
*Habitat in Camtschatka.* 24
2. DRYAS floribus octopetalis, foliis simplicibus. octopetals.  
*Dryas. Fl. lapp.* 215. *Fl. suec.* 426; 462. *Hort. cliff.*  
195. *Roy. lugdb.* 279. *Hall. helv.* 335. *Segu. ver.* 512.  
Char

*Chamædrys* 3 f. *montana*. *Clus. hist. 2. p. 351.*

*Leucas*; *Chamædrys alpina*. *Oed. dan. 1. 31.*

*Habitat in Alpibus Lapponicis, Helveticis, Austriacis, Sabaudicis, Hibernicis, Sibiricis. 24*

## COMARUM

palustre.

1. *COMARUM*. *Fl. lapp: 214. Fl. svec. 422; 463. Hort. cliff. 195. Roy. lugdb. 276. Hall. herb. 337.*

*Quinquefolium palustre rubrum. Bauh. pin. 325.*

*Pentaphyllum palustre. Cord. hist. 95.*

*Habitat in Europæ uliginosis. 24*

## CALYCANTHUS.

floridus.

1. *CALYCANTHUS* petalis interioribus longioribus.

*Butneria anemones flore. Duham. arb. 1. p. 114. t. 45.*

*Basteria foliis ovatis acuminatis, caule fruticoso. Mill. dict. 1. 60.*

*Beureria petalis coriaceis oblongis, calycis foliolis reflexis. Ehret. pict. 1. 13.*

*Frutex corni foliis, floribus instar anemones stellatæ, petalis crassis rigidis colore sordide rubente, cortice aromatico. Carexb. car. 1. p. 46. t. 46.*

*Habitat in Carolina. 5*

præcox.

2. *CALYCANTHUS* petalis interioribus minutis. †

*Abai f. Robai. Kæmph. aman, 878. t. 879.*

*Habitat in Japonia. 5*

*Ignota mihi, videtur præcedentis generis, sed diversæ speciei, quod floreat ante folia, quodque petala interiora lutea minora flori adscribantur.*

## Classis XIII

POLYANDRIA  
MONOGYNIA.

## MARCRAVIA.

1. MARCRAVIA. Jacq. *hif.* t. 96.

umbellata.

*Marcgravia scandens*, fructu radiatim pōsito. *Plum.*  
*gen.* 7. *ic.* 173.

*Marcgravia scandens*, foliis caulinis, subrotundis ad  
margines glandulatis, ramorum integris ovatis alter-  
nis distiche pōsitis, floribus umbellatis terminalibus.  
*Brown.* *jam.* 244. t. 26.

*Phyllitidi scandenti affinis major*, folio crasso subro-  
tundo. *Sloan.* *jam.* 15. *hif.* 1. p. 74 t. 28. f. 1. mala.

*Habitat in America calidiore.* b

*Folia alterna*, petiolata, ovato-lanceolata, integerrima.  
*Umbella simplex* pedunculis numerosis, in orbem po-  
sitis, equalibus, horizontaliter patentibus. Flores  
adscendentes, nudi, corolla calyceque caducis. Dis-  
cus umbellæ constans glandulis nectariferis, sessili-  
bus, sæpius quinis, erectis, forma corollæ ringentis  
galeæ s. supremi petali aconiti, ore ambitum respi-  
cientibus.

## R H E E D I A

## 1. RHEEDIA. †

lateriflora.

*van Rheedia* folio subrotundo, fructu luteo. *Plum.*  
*gen.* 45. t. 257.

*Habitat in America.*

*Plantam equidem numquam vidi; certiore locum indigi-  
tent autoptæ. Folia opposita, petiolata, ovata, in-  
tegrerrima. Pedunculi laterales s. axillares foliorum,  
simplices, conferti, uniflori. Pl.*

## M O R I S O N I A.

1. MORISONIA. Jacq. *hif.* t. 87.

americana.

*Morisona arbor* nucis juglandis folio. *Plum.* *gen.* 36.  
*ic.* 203.

Ha-

*Habitat in America calidiorē. †*

*Folia alterna, petiolata, ovata, obtusa.*

## C A P P A R I S.

- spinosa.* 1. CAPPARIS pedunculis solitariis unifloris, stipulis spinosis, foliis annuis, capsulis ovalibus.  
*Capparis aculeata. Hort. cliff. 203. Mat. med. 259. Roy. lugdb. 476. Gron. orient. 154.*  
*Capparis spinosa, fructu minore, folio rotundo. Bauh. pin. 480.*  
 β. *Capparis folio acuto. Bauh. pin. 480.*  
*Capparis sicula, duplicata spina, folio acuto. Bocc. sic. 79. t. 42. f. 3.*  
*Habitat in Europæ australis & Orientis ruderalis, muris. †*
- zeylanica.* 2. CAPPARIS pedunculis solitariis unifloris, stipulis spinosis, foliis ovatis utrinque acutis. *Fl. zeyl. 210.*  
*Capparis zeylanica, duplicatis spinis, folio acuto. Per. mus. 625.*  
*Capparis spinosa, foliis oblongis. Burm. zeyl. 53.*  
*Habitat in Zeylona. †*  
*Utrum hæc distincta species, aut etiam varietas præcedentis, dicant qui fructum obtinent.*
- sepiaria.* 3. CAPPARIS pedunculis umbellatis, stipulis spinosis, foliis annuis ovatis emarginatis.  
*Arbuscula baccifera scandens spinosa, foliis buxi palidioribus. Pluk. mant. 27. t. 338. f. 3.*  
*Habitat in India. †*  
*Rami teretes, lignosi. Spinæ stipulares, geminæ, recurvæ, breves, nigrae. Folia subpetiolata, ovata, nuda, integerrima, emarginata. Umbellæ terminales, simplices, multifloræ: Pedunculo communi, brevi.*
- Raducca:* 4. CAPPARIS pedunculis subfoliariis, foliis persistentibus ovato-oblongis nudis determinate confertis.  
*Capparis inermis, foliis ovato-oblongis per spatia confertis perennantibus. Hort. cliff. 204. \**  
*Capparis alia arborescens, lauri foliis, fructu oblongo ovato. Plum. spec. 7. ic. 73. f. 2.*  
*Breynia fruticosa, foliis singularibus oblongo-ovatis superne nitidis, siliquis minoribus teretibus æqualibus. Brown. jam. 246. t. 27. f. 2.*

*Baducca. Rheed. mal. 6. p. 105. t. 57. Rai. dendr. 1630.*

*Habitat in Indiis. h*

*Folia utrinque glabra, venis obsolete reticulata, imprimis subtus.*

5. CAPPARIS pedunculis multifloris terminalibus angulatis, foliis persistentibus ovalibus obtusis. Cynophal-  
lophora.

*Capparis inermis, foliis ovalibus alternis perennantibus.*

*Hort. cliff. 204. \**

*Capparis arborecens, lauri foliis, fructu longissimo.*

*Plum. spec. 7. ic. 73. f. 1.*

*Breynia fruticosa, foliis oblongis obtusis. Brown. jam.*

*246. t. 27. f. 1.*

*Cynophallophorus f. Penis caninus caribæorum arbor, foliis subrotundis clavos pulpiferos phalloides pro siliquis gerens. Pluk. alm. 126. t. 172. f. 4*

*Acaciæ affinis arbor siliquosa, folio subrotundo singulari, flore stamineo albido, siliqua tereti ventricosa interiore tunica mucosa miniata. Sloan. jam. 1538.*

*hist. 2. p. 59. Rai. dendr. 102.*

*Habitat in America meridionali. h*

6. CAPPARIS pedunculis unifloris compressis, foliis siliquosa, persistentibus lanceolato-oblongis acuminatis subtus punctatis.

*Breynia arborecens, foliis ovatis utrinque acuminatis, siliqua torosa longissima. Brown. jam. 247.*

*Planta forte. Pluk. phyt. 327. f. 6.*

*Habitat in Jamaica. h*

*Simillima præcedenti. Folia supra nitida, subtus pulvere ferrugineo adpersa, valde acuminata.*

7. CAPPARIS pedunculis racemosis, foliis persistentibus oblongis, pedunculis calycibusque tomentosis, floribus octandris. Breynia.

*Breynia foliis oblongo-ovalibus. Roy. lugdb. 476. Sp.*

*pl. 1. p. 503.*

*Breynia elæagni foliis. Plum. gen., 40. Breyn. ic. 13.*

*c. fig.*

*Breyniæ affinis arbor octandra. Læfl. it. 207. \**

*Habitat in America. h*

8. CAPPARIS pedunculis umbellatis, foliis persistentibus lanceolatis subtus tomentosis, floribus octandris. ferruginea.

*Aman. acad. 5. p. 398. \**

*Linnaei Spec. plant. pars. 1.*

*Z 2*

*Cra-*

*Cratæva fruticosa*, foliis singularibus oblongis utrinque acutis subtus quasi villosis, floribus octandris racemis comosis alaribus. *Brown. jam. 247. t. 28 f. 1.*

*Habitat in Jamaica. h*

*flexuosa.* 9. CAPPARIS pedunculis congestis terminalibus, foliis persistentibus oblongis obtusis glabris, ramis flexuosis.

*Morisona flexuosa. Amæn. acad. 5. p. 398. \**

*Habitat in Jamaica. h*

*hastata.* 10. CAPPARIS pedunculis multifloris, foliis hastatis lanceolatis nitidis. *Jacq. amer. 23. t. 174. f. 55.*

*Habitat in America. h*

### ACTÆA.

*spicata.* 1. ACTÆA racemo ovato, fructibusque baccatis. *Actæa. Hort. cliff. 209. Fl. suec. 431; 464 Hort. 138. Mat. med. 253. Roy. lugdb. 480. Dalib. parij. 132.*

*Actæa caule inermi. Fl. lapp. 217.*

*nigra.*

*Aconitum bacciferum. Bauh. pin. 183.*

*Christophoriana. Clus. hist. 2. p. 86.*

*alba.*

3. ACTÆA racemo ovato, fructibus niveis, radice tuberosa. *Mill. dist.*

*Aconitum baccis niveis. Corn. canad. 76. t. 77.*

*Christophoriana americana racemosa baccis, niveis rubris. Moris. hist. 2. p. 8. f. 1. t. 2. f. 7.*

*Habitat in nemoribus Europæ; 3. Americæ. 4*

*Petala rhombea, plano-membranacea.*

*racemosa.* 2. ACTÆA racemis longissimis, fructibus unicarpis.

*ribus.*

*Actæa racemis longissimis. Gron. virg. 59. Mat. 253.*

*Christophoriana americana procerior & longius spicata.*

*Dill. elth. 79. t. 67. f. 78.*

*Christophoriana facie herba spicata. Pluk. amstel. t. 383. f. 3.*

*Habitat in Florida, Virginia, Canada. 4*

*Petala crassa, gibba, minima, pedicellata, setaceo-acuminata, calycis foliolis plura.*

*cimicifuga*

3. ACTÆA racemis paniculatis, fructibus quad-

*laribus*

*Cimicifuga. Amæn. acad. 2. p. 54.*

*Thalictroides foetidissimum, christophorianæ facie.*

*Ann. ruth. 102.*

*Habitat in Sibiria. D. Kraschenninnikof. 24.*

*Petala 4. subrotunda, calyce majora, tenaxiora. Styli. 4.*

## SANGUINARIA.

1. SANGUINARIA. *Hort. cliff. 202. Gron. virg. 57. canadensis.*

*Sanguinaria minor, flore simplici. Dill. elth. 335. t.*

*252. f. 326.*

*Chelidonium majus canadense acaulon. Corn. canad.*

*212. Moris. hist. 2. p. 257. f. 3. t. 11. f. 1. Raj. hist.*

*1887.*

*Ranunculus virginienfis albus. Park. theatr. 327. Raj.*

*suppl. 314.*

2. Sanguinaria major, flore simplici. *Dill. elth. 335. t.*

*252. f. 325.*

3. Sanguinaria major, flore pleno. *Dill. elth. 335. t. 252.*

*f. 326.*

*Habitat in America septentrionali. 24.*

## PODOPHYLLUM.

1. PODOPHYLLUM foliis peltatis palmatis.

*peltatum.*

*Podophyllum. Hort. cliff. 202. Hort. ups. 137. Gron.*

*virg. 57. Roy. lugdb. 480. Trew chet. t. 29.*

*Anapodophyllum canadense. Cuesb. car. 1. p. 24. t. 24.*

*Aconitifolia humilis, flore albo unico campanulato, fructu cynosbati. Mentz. pug. t. 11.*

*Habitat in America septentrionali. 24.*

2. PODOPHYLLUM foliis binatis semicordatis.

*diphyllum.*

*Habitat in Virginia. Collinson. 24.*

*Folia radicalia, petiolata, binata ut in Hymenæa, gla-*

*bra, integerrima, semicordata, absque pedicellis.*

*Scapus radicalis, uniflorus, fructu antecedentis.*

*Flos mihi non visus.*

## CHELIDONIUM.

1. CHELIDONIUM pedunculis umbellatis.

*majus.*

*Chelidonium pedunculis multifloris. Hort. cliff. 201.*

*Fl. suec. 430; 465. Hort. ups. 137. Max. med. 252.*

*Roy. lugdb. 478. Dalib. parif. 152.*

*Chelidonium majus vulgare. Bauh. pin. 144.*

*Z z 2*

*Che-*



*Chelidonium majus.* *Fuchf. hist.* 865.

β. *Chelidonium majus*, foliis quernis. *Bauh. pin.* 144.

*Chelidonium majus*, folio laciniato. *Clus. hist.* 2. p. 203.

*Habitat in Europæ ruderatis.* 2.

*Glaucium.* 2. **CHELIDONIUM** pedunculis unifloris, foliis amplexicaulibus sinuatis, caule glabro.

*Chelidonium glabrum*, pedunculis unifloris. *Hort. upf.* 137. *Roy. lugdb.* 478. *Dalib.-parif.* 152.

*Chelidonium pedunculis unifloris.* *Hort. cliff.* 201. *Gron. virg.* 57.

*Papaver corniculatum luteum.* *Bauh. pin.* 171.

*Papaver corniculatum*, flavo flore. *Clus. hist.* 2. p. 91.

*Habitat in Angliæ, Helvetiæ, Galliæ, Italiæ, Virginiæ arenosis.* ☉

*Planta glauca.* *Caulis lævis.* *Folia scabriuscula: radicalibus pinnatifidis; caulinis solum lobatis.* *Flores flavi.* *Siliquæ scabræ, dissepimento in loculamenta, uti sequentes, distinctæ*

*corniculatum.* 3. **CHELIDONIUM** pedunculis unifloris, foliis sessilibus pinnatifidis, caule hispido.

*Chelidonium hispidum*, pedunculis unifloris. *Hort. upf.* 137. *Sauv. monsp.* 286.

*Glaucium hirsutum*, flore phœniceo. *Tournef. inst.* 254.

*Papaver corniculatum phœniceum hirsutum.* *Bauh. pin.* 171.

*Papaver cornutum*, phœniceo flore. *Clus. hist.* 2. p. 91.

β. *Papaver corniculatum phœniceum glabrum.* *Bauh. pin.* 171.

*Habitat in Hungaria, Bohemia, Monspelli.* ☉

*Planta viridis.* *Folia omnia pinnatifida.* *Flores rubro-lutei.* *Siliquæ hispide.*

*hybridum.* 4. **CHELIDONIUM** pedunculis unifloris, foliis pinnatifidis linearibus, caule lævi, siliquis trivalvibus.

*Papaver caule ramoso*, foliis linearibus pinnatifidis, capsulis sulcatis. *Hort. upf.* 136. *Sauv. monsp.* 252.

*Papaver corniculatum violaceum.* *Bauh. pin.* 172.

*Papaver corniculatum*, violaceo flore. *Clus. hist.* 2. p. 92.

*Habitat in Europa australiore.* ☉

*Stigmata.* 3.

P A P A V E R.

\* *Capsulis hispidis.*

1. PAPAVER capsulis subglobosis torosis hispidis, cau- hybridum.  
le folioso multifloro.

Papaver foliis ternato-pinnatifidis, capsulis sulcatis hispidis. *Hort. cliff.* 201.

Papaver caule ramoso, foliis lineari-pinnatifidis, capsulis sulcatis hispidis. *Hort. upf.* 136.

Argemone [capitulo brevior. *Bauh. pin.* 172.

Argemone capitulo hirsuto rotundo torulis canaliculato. *Morif. hist.* 2. p. 278. f. 3. t. 14. f. 9. *Lob. ic.* 276.

*Habitat in Europa australiore. ☉*

2. PAPAVER capsulis clavatis hispidis, caule folioso Argemone.  
multifloro.

Papaver capsulis hispidis oblongo-turbinatis sulcatis. *Pl. suec.* 429, 466. *Dalib. parif.* 152. *Gron. orient.* 158.

Papaver foliis ternato-pinnatis, fructu angulato. *Hort. cliff.* 201. *Roy. lugdb.* 419. *Sauv. monsp.* 252.

Argemone capitulo longiore. *Bauh. pin.* 172. *Lob. ic.* 276.

Argemone capitulo tenuiore longiore hirsuto. *Morif. hist.* 2. p. 278. f. 3. t. 14. f. 10.

*Habitat in Europæ campis arenosis. ☉*

Folia tripartita pinnatifida. Pedunculi scabri. Calycees capsulaeque pilosæ.

3. PAPAVER capsula hispida, scapo unifloro nudo hi- alpinum.  
spido, foliis bipinnatis.

Papaver perenne, pilis rigidissimis, foliis pinnatis, flore luteo hirsuto. *Hall. helv.* 304.

Papaver alpinum saxatile, coriandri folio. *Segu. ver.* 1. p. 416. t. 4. f. 4.

Argemone alpina, foliis scandicis, lutea. *Bauh. pin.* 172. *prodr.* 93. *Burf. IX.* 58.

Argemone alpina, coriandri folio. *Bauh. pin.* 172.

Argemone minor penz. *Dalech. hist.* 1714.

*Habitat in Helvetia, Schneeberg Austriae, Pyrenæis. 21*  
Flores plantæ Burserianæ apud C. Bauhinum albi erant,  
nec lutei, afferente Bursero.

4. PAPAVER capsulis hispidis, scapo unifloro nudo nudicute.  
Z z 3 hi-

hispido, foliis simplicibus pinnato-sinuatis. *Hort. upf.* 136.

*Papaver erraticum*, nudicaule flore flavo odorato. *Dill. elth.* 202. t. 224. f. 291. *Oed. dan.* t. 41.

*Papaver erraticum*, luteo flore, capite oblongo hispido. *Amm. ruth.* 61. n. 81.

*Habitat in Sibiria.* ♂

*Pedunculi* & *calyces* pilosi.

\* \* *Capsulis glabris.*

dubium.

6. **PAPAVER** capsulis oblongis glabris, caule multifloro setis adpressis, foliis pinnatifidis incisis. *Fl. suec.* 2. n. 467.

*Papaver erraticum*, capite longissimo glabro. *Tournef. hist.* 238. *Kupp. jen.* 3. p. 71.

*Argemone* capitulo longiore glabro. *Morif. blaf.* 233. *hist.* 2. p. 279. f. 3. t. 14. f. 11.

*Habitat inter segetes Europæ septentrionalioris.* ☉

Rhœas.

6. **PAPAVER** capsulis glabris globosis, caule piloso multifloro, foliis pinnatifidis incisis.

*Papaver* foliis pinnatifidis hispidis, fructu ovato. *Hort. cliff.* 201. *Fl. suec.* 428; 468. *Mat. med.* 251. *Roy. lugdb.* 478. *Dalib. parif.* 151. *Hall. helv.* 303.

*Papaver* caule multifloro folioso hispido, foliis pinnatifidis, capsulis lævibus. *Hort. upf.* 136. *Saurv. morif.* 252. *Gron. orient.* 157.

*Papaver erraticum majus.* *Bauh. pin.* 171.

*Papaver rhœas prius.* *Dalech. hist.* 439.

- β. *Papaver erraticum*, pleno flore. *Bauh. pin.* 171.

- γ. *Papaver erraticum minus.* *Bauh. pin.* 171.

*Habitat in Europæ arvis, agris.* ☉

*Fili in caule patentes sunt.*

- somniferum. 7. **PAPAVER** calycibus capsulisque glabris, foliis amplexicaulibus incisis.

*Papaver* caule multifloro, foliis simplicibus glabris incisis caulinis. *Hort. upf.* 135. *Mat. med.* 250.

*Papaver* foliis simplicibus glabris incisis. *Hort. cliff.* 200. *Hall. helv.* 303. *Roy. lugdb.* 879.

*Papaver hortense*, semine albo. *Bauh. pln.* 170.

- ε. *Papaver hortense*, semine nigro. *Bauh. pin.* 170.

- ζ. *Papaver cristatum*, floribus & semine album. *Bauh. pin.* 171.

δ. Pz

- δ. *Papaver cristatum*, floribus rubris, semine nigro. *Bauh. pin.* 171.
- ε. *Papaver flore pleno*, album. *Bauh. pin.* 171.  
*Habitat in Europæ australioris ruderalis.* ☉  
*Caulis, Folia, Calyces, Capsula glabra.* Stigmata 10.
8. PAPAVER capsulis glabris oblongis, caule multifloro lævi, foliis pinnatis incisis. *Huds. angl.* 203.  
*Papaver foliis pinnatis, fructu acuminato.* *Hort. cliff.* 201. *Roy. lugdb.* 479.  
*Papaver cambricum perenne, flore sulphureo.* *Dill. elth.* 300. t. 223. f. 290.  
*Argemone cambro-britannica lutea, capite longiore glabro.* *Morif. hist.* 2. p. 297. f. 3. t. 14. f. 12.  
*Papaver erraticum pyrenaicum, flore flavo.* *Bauh. pin.* 171. *prodr.* 92. *Burf. IX.* 45.  
*Habitat in Cambriæ septentrionalis nemorosis, Pyrenæis.* 24  
*Stigmata 5 f. 6, distincta. Capsula oblonga, inferne angustior, apice subrostrata.*
9. PAPAVER capsulis glabris, caulibus unifloris scarabris foliosis, foliis pinnatis ferratis. *Hort. upf.* 136.  
*Papaver foliis pinnatis, fructu globoso.* *Roy. lugdb.* 279.  
*Papaver orientale hirsutissimum, flore magno.* *Tournef. cor.* 17. *itiz.* 3. p. 127. t. 127. *Comm. rar.* 34. t. 34.  
*Habitat in Oriente.* ☉  
*Stigmata 16. Setæ in caule sparsæ, cauli adpressæ, basi prominula asperæ.*

## A R G E M O N E

1. ARGEMONE capsulis quinquevalvibus, foliis spinosis. *mexicana.*  
*Argemone.* *Hort. cliff.* 201. *Hort. upf.* 135. *Roy. lugdb.* 479. *Brown. jam.* 244. *Merian. surin.* 24. t. 24.  
*Papaver spinosum.* *Bauh. pin.* 171. *prodr.* 93. t. 92.  
*Clus. hist.* 2. p. 93.  
*Papaver spinosum luteum, foliis albis venis notatis.* *Morif. hist.* 2. p. 277. f. 3. t. 14. f. 5.  
*Habitat in Mexico, Jamaica, Caribæis, nunc in Europa australi.* ♂

2. ARGEMONE capsulis trivalvibus.

Z z 4

Pa-

*armeniaca.*

*Cratæva fruticosa*, foliis singularibus oblongis utrinque acutis subtus quasi villosis, floribus octandris, racemis comosis alaribus. *Brown. jam.* 247. t. 28. f. 1.

*Habitat in Jamaica.* b

*flexuosa.* 9. *CAPPARIS* pedunculis congestis terminalibus, foliis persistentibus oblongis obtusis glabris, ramis flexuosis.

*Morisona flexuosa.* *Amæn. acad.* 5. p. 398. \*

*Habitat in Jamaica.* b

*hastata.* 10. *CAPPARIS* pedunculis multifloris, foliis hastatolanceolatis nitidis. *Jacq. amer.* 23. t. 174. f. 56.

*Habitat in America.* b

### A C T Æ A.

*spicata.* 1. *ACTÆA* racemo ovato, fructibusque baccatis. *Actæa.* *Hort. cliff.* 209. *Fl. suec.* 431; 464. *Hort. uf.* 138. *Mat. med.* 253. *Roy. lugdb.* 480. *Dalib. parif.* 152.

*Actæa* caule inermi. *Fl. lapp.* 217.

*nigra.* *Aconitum bacciferum.* *Bauh. pin.* 183.

*Christophoriana.* *Clus. hist.* 2. p. 86.

*alba.* 3. *ACTÆA* racemo ovato, fructibus niveis, radice tuberosa. *Mill. dict.*

*Aconitum baccis niveis.* *Corn. canad.* 76. t. 77.

*Christophoriana americana* racemosa baccis, niveis & rubris. *Morif. hist.* 2. p. 8. f. 1. t. 2. f. 7.

*Habitat in nemoribus Europæ; 3. Americæ.* 2

*Petala rhombea, plano-membranacea.*

*racemosa.* 2. *ACTÆA* racemis longissimis, fructibus unicapsularibus.

*Actæa* racemis longissimis. *Gron. virg.* 59. *Mat. med.* 253.

*Christophoriana americana* procerior & longius spicata. *Dill. elth.* 79. t. 67. f. 78.

*Christophoriana* facie herba spicata. *Pluk. amalt.* 54. t. 383. f. 3.

*Habitat in Florida, Virginia, Canada.* 2

*Petala crassa, gibba, minima, pedicellata, acutiuscula, setaceo-acuminata, calycis foliolis plura.*

*cimicifuga.* 2. *ACTÆA* racemis paniculatis, fructibus quadrilobis

*Cimicifuga. Amæn. acad. 2. p. 54.*

*Thalictroides foetidissimum, christophorianæ facie.*

*Amn. ruth. 102.*

*Habitat in Sibiria D. Kraschenninnikof. 24*

*Petala 4. subrotunda, calyce majora, tenaciora. Styli. 4.*

## SANGUINARIA.

1. SANGUINARIA. *Hort. cliff. 202. Gron. virg. 57. canadensis.*

*Sanguinaria minor, flore simplici. Dill. elth. 335. t.*

*252. f. 326.*

*Chelidonium majus canadense acaulon. Corn. canad.*

*212. Moris. hist. 2. p. 257. f. 3. t. 11. f. 1. Raj. hist.*

*1887.*

*Ranunculus virginienfis albus. Park. theatr. 327. Raj.*

*suppl. 314.*

2. Sanguinaria major, flore simplici. *Dill. elth. 335. t.*

*252. f. 326.*

3. Sanguinaria major, flore pleno. *Dill. elth. 335. t. 252.*

*f. 326.*

*Habitat in America septentrionali. 24*

## PODOPHYLLUM.

1. PODOPHYLLUM foliis peltatis palmatis.

*peltatum.*

*Podophyllum. Hort. cliff. 202. Hort. ups. 137. Gron.*

*virg. 57. Roy. lugdb. 480. Trew chret. t. 29.*

*Anapodophyllum canadense. Catesb. car. 1. p. 24. t. 24.*

*Aconitifolia humilis, flore albo unico campanulato,*

*fructu cynosbati. Menz. pug. t. 111.*

*Habitat in America septentrionali. 24*

2. PODOPHYLLUM foliis binatis semicordatis.

*diphyllum.*

*Habitat in Virginia. Collinson. 24*

*Folia radicalia, petiolata, binata ut in Hymenæa, gla-*

*bra, integerrima, semicordata, absque pedicellis.*

*Scapus radicalis, uniflorus, fructu antecedentis.*

*Flos mihi non visus.*

## CHELIDONIUM.

1. CHELIDONIUM pedunculis umbellatis.

*majus.*

*Chelidonium pedunculis multifloris. Hort. cliff. 201.*

*Fl. fvec. 430: 465. Hort. ups. 137. Mat. med. 252.*

*Roy. lugdb. 478. Dalib. parif. 152.*

*Chelidonium majus vulgare. Bauh. pin. 144.*

*Z z 2*

*Che-*

- Chelidonium majus.* *Fuchf. hist.* 865.  
 β. *Chelidonium majus*, foliis quernis. *Bauh. pin.* 144.  
*Chelidonium majus*, folio laciniato. *Clus. hist.* 2. p. 203.  
*Habitat in Europæ ruderatis.* 24
- Glaucium. 2. **CHELIDONIUM** pedunculis unifloris, foliis amplexicaulibus sinuatis, caule glabro.  
*Chelidonium glabrum*, pedunculis unifloris. *Hort. upf.* 137. *Roy. lugdb.* 478. *Dalib. parif.* 152.  
*Chelidonium pedunculis unifloris.* *Hort. cliff.* 201. *Gron. virg.* 57.  
*Papaver corniculatum luteum.* *Bauh. pin.* 171.  
*Papaver corniculatum*, flavo flore. *Clus. hist.* 2. p. 91.  
*Habitat in Angliæ, Helvetiæ, Galliæ, Italiæ, Virginiæ arenosis.* ○  
*Planta glauca.* *Caulis levis.* *Folia scabriuscula: radicalibus pinnatifidis; caulinis solum lobatis.* *Flores flavi.* *Siliquæ scabræ, dissepimento in loculamenta, uti sequentes, distinctæ*
- corniculatum. 3. **CHELIDONIUM** pedunculis unifloris, foliis sessilibus pinnatifidis, caule hispidulo.  
*Chelidonium hispidum*, pedunculis unifloris. *Hort. upf.* 137. *Sauv. monsp.* 286.  
*Glaucium hirsutum*, flore phœniceo. *Tournef. inst.* 254.  
*Papaver corniculatum phœniceum hirsutum.* *Bauh. pin.* 171.  
*Papaver cornutum*, phœniceo flore. *Clus. hist.* 2. p. 91.  
 β. *Papaver corniculatum phœniceum glabrum.* *Bauh. pin.* 171.  
*Habitat in Hungaria, Bohemia, Monspelli.* ○  
*Planta viridis.* *Folia omnia pinnatifida.* *Flores rubro-lutei.* *Siliquæ hispide.*
- hybridum. 4. **CHELIDONIUM** pedunculis unifloris, foliis pinnatifidis linearibus, caule lævi, siliquis trivalvibus.  
*Papaver caule ramoso*, foliis linearibus pinnatifidis, capsulis sulcatis. *Hort. upf.* 136. *Sauv. monsp.* 252.  
*Papaver corniculatum violaceum.* *Bauh. pin.* 172.  
*Papaver corniculatum*, violaceo flore. *Clus. hist.* 2. p. 92.  
*Habitat in Europa australiore.* ○  
*Stigmata.* 3.

## P A P A V E R.

\* *Capsulis hispida.*

1. PAPAVER capsulis subglobosis torosis hispida, cau- hybridum.  
le folioso multifloro.

Papaver foliis ternato-pinnatifidis, capsulis sulcatis hispidis. *Hort. cliff.* 201.

Papaver caule ramoso, foliis lineari-pinnatifidis, capsulis sulcatis hispida. *Hort. ups.* 136.

Argemone [capitula breviora. *Bauh. pin.* 172.

Argemone capitula hirsuta rotunda torulis canaliculata. *Morif. hist.* 2. p. 278. f. 3. t. 14 f. 9. *Lob. ic.* 276.

*Habitat in Europa australiore.* ☉

2. PAPAVER capsulis clavatis hispida, caule folioso Argemone.  
multifloro.

Papaver capsulis hispida oblongo-turbinatis sulcatis. *Fl. suec.* 429, 466. *Dalib. parif.* 152. *Gron. orient.* 158.

Papaver foliis ternato-pinnatis, fructu angulato. *Hort. cliff.* 201. *Roy. lugdb.* 419. *Sauv. monsp.* 252.

Argemone capitula longiora. *Bauh. pin.* 172. *Lob. ic.* 276.

Argemone capitula tenuiora longiora hirsuta. *Morif. hist.* 2. p. 278. f. 3. t. 14. f. 10.

*Habitat in Europæ campis arenosis.* ☉

*Folia tripartita pinnatifida. Pedunculi scabri. Calyces capsulaeque pilosæ.*

3. PAPAVER capsula hispida, scapo unifloro nudo hi- alpinum.  
spido, foliis bipinnatis.

Papaver perenne, pilis rigidissimis, foliis pinnatis, flore luteo hirsuto. *Hall. helv.* 304.

Papaver alpinum saxatile, coriandri folio. *Segu. ver.* 1. p. 416. t. 4. f. 4.

Argemone alpina, foliis scandicis, lutea. *Bauh. pin.* 172. *prodr.* 93. *Burf. IX.* 58.

Argemone alpina, coriandri folio. *Bauh. pin.* 172.

Argemone minor penz. *Dalech. hist.* 1714.

*Habitat in Helvetia, Schneeberg Austria, Pyrenæis.* 2.

Flores plantæ Burserianæ apud C. Bauhinum albi erant, nec lutei, afferente Bursero.

4. PAPAVER capsulis hispida, scapo unifloro nudo nudicula.  
hi-



hispido, foliis simplicibus pinnato-sinuatis. *Hort. ups.* 136.

*Papaver erraticum*, nudicaule flore flavo odorato. *Dill. elr.* 202. t. 224. f. 291. *Oed. dan.* t. 41.

*Papaver erraticum*, luteo flore, capite oblongo hispido. *Amm. ruth.* 61. n. 81.

*Habitat in Sibiria.* ♂

*Pedunculi* ♂ *calyces pilosi.*

\* \* *Capsulis glabris.*

dubium.

6. **PAPAVER** capsulis oblongis glabris, caule multifloro setis adpressis, foliis pinnatifidis incis. *Fl. suec.* 2. n. 467.

*Papaver erraticum*, capite longissimo glabro. *Tournef. inst.* 238. *Kupp. jen.* 3. p. 71.

*Argemone* capitulo longiore glabro. *Morif. bles.* 233. *bist.* 2. p. 279. f. 3. t. 14. f. 11.

*Habitat inter segetes Europæ septentrionalioris.* ☉

Rheas.

6. **PAPAVER** capsulis glabris globosis, caule piloso multifloro, foliis pinnatifidis incis.

*Papaver* foliis pinnatifidis hispidis, fructu ovato. *Hort. cliff.* 201. *Fl. suec.* 428; 468. *Mat. med.* 251. *Roy. lugdb.* 478. *Dalib. paris.* 151. *Hall. helv.* 303.

*Papaver* caule multifloro folioso hispido, foliis pinnatifidis, capsulis lævibus. *Hort. ups.* 136. *Sauv. mont.* 252. *Gron. orient.* 157.

*Papaver erraticum* majus. *Bauh. pin.* 171.

*Papaver rheas* prius. *Dalech. hist.* 439.

β. *Papaver erraticum*, pleno flore. *Bauh. pin.* 171.

γ. *Papaver erraticum* minus. *Bauh. pin.* 171.

*Habitat in Europæ arvis, agris.* ☉

*Fili in caule patentes sunt.*

somniferum.

7. **PAPAVER** calycibus capsulisque glabris, foliis amplexicaulibus incis.

*Papaver* caule multifloro, foliis simplicibus glabris incis. caulinis. *Hort. ups.* 135. *Mat. med.* 250.

*Papaver* foliis simplicibus glabris incis. *Hort. cliff.* 200. *Hall. helv.* 303. *Roy. lugdb.* 879.

*Papaver hortense*, semine albo. *Bauh. pin.* 170.

β. *Papaver hortense*, semine nigro. *Bauh. pin.* 170.

γ. *Papaver cristatum*, floribus & semine album. *Bauh. pin.* 171.

δ. Pz

- δ. *Papaver cristatum*, floribus rubris, semine nigro. *Bauh. pin.* 171.
- ε. *Papaver* flore pleno, album. *Bauh. pin.* 171.  
*Habitat in Europæ australioris ruderalis.* ☉  
*Caulis, Folia, Calyces, Capsula glabra. Stigmata 10.*
8. PAPAVER capsulis glabris oblongis, caule multiflo- *cambricum.*  
 ro lævi, foliis pinnatis incis. *Huds. angl.* 203.  
*Papaver* foliis pinnatis, fructu acuminato. *Hort. cliff.*  
 201. *Roy. lugdb.* 479.  
*Papaver cambricum* perenne, flore sulphureo. *Dill.*  
*elh.* 300. t. 223. f. 290.  
*Argemone cambro-britannica* lutea, capite longiore  
 glabro. *Morif. hist.* 2. p. 297. f. 3. t. 14. f. 12.  
*Papaver erraticum pyrenaicum*, flore flavo. *Bauh. pin.*  
 171. *prodr.* 92. *Burf. IX.* 45.  
*Habitat in Cambriæ septentrionalis nemorosis, Pyre-*  
*næis.* 24.  
*Stigmata 5 f. 6, distincta. Capsula oblonga, inferne*  
*angustior, apice subrostrata.*
9. PAPAVER capsulis glabris, caulibus unifloris sca- *orientale.*  
 bris foliosis, foliis pinnatis serratis. *Hort. ups.* 136.  
*Papaver* foliis pinnatis, fructu globoso *Roy. lugdb.*  
 279.  
*Papaver orientale* hirsutissimum, flore magno. *Tournef.*  
*cor.* 17. *itiz.* 3. p. 127. t. 127. *Comm. rar.* 34. t. 34.  
*Habitat in Oriente.* ☉  
*Stigmata 16. Setæ in caule sparsæ, cauli adpressæ, basi*  
*prominula asperæ.*

## A R G E M O N E

1. ARGEMONE capsulis quinquevalvibus, foliis spi- *mexicana,*  
 nois.  
*Argemone. Hort. cliff.* 201. *Hort. ups.* 135. *Roy. lugdb.*  
 479. *Brown. jam.* 244. *Merian. surin.* 24. t. 24.  
*Papaver spinosum. Bauh. pin.* 171. *prodr.* 93. t. 92.  
*Clus. hist.* 2. p. 93.  
*Papaver spinosum luteum*, foliis albis venis notatis.  
*Morif. hist.* 2. p. 277. f. 3. t. 14. f. 5.  
*Habitat in Mexico, Jamaica, Caribæis, nunc in Eu-*  
*ropa australi.* ♂
2. ARGEMONE capsulis trivalvibus. *Pa-* *armeniaca.*  
 Z z 4

*Papaver orientale*, hypecoi follo, fructu minime.

*Tourn. f. cor.* 17.

*Habitat in Armenia.*

pyrenaica. 3. ARGEMONE capsulis quadrivalvibus, caule nudo.

*Habitat in Pyrenæis. Tourn. f.*

Hæc semiquadrivalvis, uti species secunda semitrivalvis, & prima semiquirquevalvis. Hæ species vel sub distincto genere simul tradendæ, vel etiam ad genus *papaverinum* amandandæ omnes. Hujus habitus plurimum accedit *Papaveri alpino*.

## CAMBOGIA.

Gutta.

1. CAMBOGIA. *Amæn. acad.* 1. p. 402. *Fl. syl.* 195.

*Nat. med.* 257.

Coddam-pull. *Rheed. mal.* 1. p. 41. t. 24.

*Habitat in India.* §

## MUNTINGIA.

Calabura.

1. MUNTINGIA. *Jacq. hist.* 1. 107.

*Muntingia pedunculis unifloris. Hort. cliff.* 203.

*Muntingia fruticosa villosa: foliis ferratis oblongis uno latere brevioribus. Brown. jam.* 245.

*Muntingia folio sericeo molli, fructu majori. Plun. gen.* 41. ic. 205.

*Calabura alba. Pluk. alm.* 75. *mant.* 34. t. 152. f. 4.

*Loti arboris folio angustiore, rubi flore, fructu polyspermo umbilicato. Sloan. jam.* 162. *hist.* 2. p. 80. t. 194. f. 1. *Raj. dendr.* 32.

*Habitat in Jamaica.* §

## SARRACENIA.

purpurea.

1. SARRACENIA foliis gibbis. *Hort. cliff.* 427. *Gron. virg.* 164.

*Sarracena foliis brevioribus latioribus. Catesb. car.* 2. p. 70. t. 70.

*Coilophyllum virginianum, brevioris folio, flore purpurascens. Moris. hist.* 3. p. 533.

*Bucanephyllum americanum, limonio congener dictum. Pluk. alm.* 71. *amst. h.* 46. t. 376. f. 6.

*Limonium peregrinum, foliis forma floris aristolochiæ. Bauh. pin.* 192.

*Habitat in Americæ septentrionalis udibus.* §

2. SAR-

## 2. SARRACENIA foliis strictis.

1278.

*Sarracena foliis rectis. Hort. cliff. 497. Gron. virg. 164.**Sarracena foliis longioribus & angustioribus. Careb. car. 2. p. 69 t. 69.**Coilophyllum virginianum, longiore folio erecto, flore luteo. Moris. hist. 3. p. 533.**Bucanephyllum elatius virginianum f. Limonio congeneris altera species elatior, foliis triplo longioribus.**Pluk. alm. 72. amalib. 46. t. 152. f. 3. (mala). t. 376. f. 5. bona.**Thuris limpidi folium. Bauh. hist. 1. p. 307. Dalech. hist. 1754. Lob. adv. 430.**Habitat in Americæ septentrionalis udis. 24*

## N Y M P H Æ A.

1. NYMPHÆA foliis cordatis integerrimis, calyce pe-<sup>luteo.</sup>*talis majore pentaphyllo. Fl. lapp. 218. Fl. fvec. 426;**469. Hort. cliff. 203. Roy. lugdb. 480. Dalib. parif. 150.**Nymphæa lutea major. Bauh. pin. 193.**Nymphæa lutea. Cam. epit. 635.**Habitat ad Europæ littora, sub dulci aqua. 24**In Canada eadem, floris magnitudine viz Calthæ.**Kalm.*2. NYMPHÆA foliis cordatis integerrimis, calyce qua-<sup>alba.</sup>  
drifido.*Nymphæa calyce tetraphyllo, corolla multiplici. Fl.**lapp. 219. Fl. fvec. 427. 470. Hort. cliff. 203. Mat.**med. 258. Roy. lugdb. 480. Dalib. parif. 150.**Nymphæa alba major. Bauh. pin. 193.**Nymphæa alba. Cam. epit. 634.**Habitat in Europæ & Americæ aquis dulcibus. 24*3 NYMPHÆA foliis cordatis dentatis. <sup>Lotus.</sup> *Fl. zeyl. 194.**Hasselt. it. 471.**Nymphæa foliis amplioribus profunde crenatis subtus areolatis. Brown. jam. 343.**Nymphæa indica, flore candido, folio in ambitu serrato. Sloan. jam. 120. hist. 1. p. 252. Raj. suppl. 630.**Lotus ægyptia. Alp. ægypt. 103. exot. 214. t. 213. 216. 218. 220. 222. 226.**Ambel. Rhæd. mal. 11. p. 51. t. 26.**Habitat in calidis Indiæ, Africa, Americæ. 24*

- Nelumbæ.* 4. *NYMPHÆA* foliis peltatis undique integris. *Hort. cliff.* 203. *Fl. zeyl.* 193.  
*Nymphæa* foliis orbiculatis peltatis subtus radiatis. *Brown. jam.* 343.  
*Nymphæa indica*, *Faba ægyptia dicta*, flore incarnato. *Herm. par.* 205. t. 205.  
*Nymphæa fabifera*, indiæ paludibus gaudens, foliis umbilicatis amplis, pediculis spinosis, flore roseo purpureo. *Pluk. alm.* 267. t. 322. f. 1. apt.  
*Taratti. Rumph. amb.* 6. p. 168. t. 73.  
*Tamara. Rheed. mal.* 11. p. 59. t. 30.  
*Bem - Tamara. Rheed. mal.* 11. p. 61. t. 31. *Raj. suppl.* 632.  
 β. *Nymphæa* affinis glandifera virginiana umbilicato folio, flore luteo pleno. *Morif. hist.* 3. p. 514. *Raj. suppl.* 632.  
*Nymphæa* glandifera virginiana lactescens, minori umbilicato folio, pediculo leui, flore amplo luteo pallescente. *Pluk. alm.* 267.  
*Habitat in Indiis.* 2.  
*Varietas β. differt seminibus duris, in reliquis convenit ut vix distingvenda videatur.*

## B F X A.

- Orellana.* 1. *BIXA. Hort. cliff.* 211. *Mat. med.* 266. *Roy. luxdb.* 477. *Brown. jam.* 254.  
*Orleana f. Orellana* folliculis lappaceis. *Pluk. alm.* 272. t. 209. f. 4. *Comm. hort.* 1. p. 65. f. 33.  
*Arbor mexiocana*, fructu castaneæ, coccifera. *Bauh. pin.* 419.  
*Urucu. Sloan. jam.* 150. *hist.* 2. p. 52. t. 181. f. 1.  
*Habitat in America calidiore, ad fossas & fontes.* 5

## S L O A N E A.

- dentata.* 1. *SLOANEA* foliis cordato-ovatis denticulatis, stipulis serratis. *Last. it.* 311?  
*Sloanea. Hort. cliff.* 210?  
*Sloanea* amplis castaneæ foliis, fructu echinato. *Plum. gen.* 49. ic. 244.  
*Castanea* foliis oblongo ovatis serratis, fructu rotundo maximo echinato. *Mill. dict.*  
*Apeiba. Marccr. bras.* 123. t. 123. *Raj. hist.* 1643.  
*Habitat in America meridionali.* 5  
*Sloaneæ genus etiamnum obscurum est, & certe ex Foliis, Pericarpii valvulis ac hispiditate nobis verosimil-*

*millima occurrit Milleri sententia, quæ Fagi speciem hanc facit, quam tamen non admittit Læflingii descriptio.*

8. SLOANEA foliis oblongis integerrimis emarginatis. *emarginata.*  
*Anona foliis laurinis in summitate incis, fructu com-*  
*presso scabro fusco in medio acumine longo. Catesb.*  
*car. 2. p. 87. t. 87.*  
*Habitat in Carolina* ♀

## M. A M M E A.

1. MAMMEA staminibus flore brevioribus. *americana.*  
*Mammea foliis ovalibus nitidis, fructu subrotundo*  
*scabro. Brown. jam. 248.*  
*Mammei magno fructu, persicæ sapore. Plum. gen.*  
*44. id. 170.*  
*Mamay. Bauh. hist. 1. p. 172. Dalech. hist. 1836. Laet.*  
*amer. 356.*  
*Malus persicæ maxima, foliis rotundis splendentibus*  
*glabris, fructu maximo scabro rugoso. Sloan. jam.*  
*179. hist. 2. p. 123. t. 217. f. 3.*  
*Arbor indica mamei dicta. Bauh. pin. 417. Raj. hist.*  
*1665.*  
*Habitat in Hispaniola, Jamaica.* ♀  
*Rami scabri; Folia oblonga, obtusa.*

2. MAMMEA staminibus flore longioribus. *Ojb. it. asiatica.*  
*278.*  
*Mammea maxima, foliis longioribus, cortice fulcato*  
*cinereo. Brown. jam. 248?*  
*Mali Persicæ Mameyæ folio longiore arbore maxima,*  
*cortice fulcato cinereo amaro. Sloan. jam. 180. hist.*  
*2. p. 124?*  
*Habitat in Java. Osbeck.* ♀  
*Arbor magna, sæpius procumbens. Folia pedalia, obo-*  
*vata s. oblonga, lævia. Pedunculi uniflori. Calyx*  
*ovatus, bipartitus. Petala 4, ovata, calyce triplo*  
*longiora, brevi decidua, cum staminibus corolla du-*  
*plo longioribus, numerosis, subulatis, erectis. An-*  
*theræ subrotundæ, minime. Fructus tetragonus.*

## O C H N A.

1. OCHNA racemis lateralibus. *Fl. zeyh. 209. \** *squarrosa.*  
*Japo.*

*Japotapita cinnamomi* folio, floribus spicatis. *Burm. zeyl.* 123. t. 56.

β. *Arbor africana*, subrotundo folio margine denticulis acutis aspero, floribus pentapetalis atropurpureis.

*Pluk. alm.* 41. t. 263. f. 1. 2.

*Habitat in India, β. in Africa.*

Racemi *perflunt squarrosi infra folia a casu pedicellorum & fructificationum.*

*Japotapita.*

2. *OCHNA* racemis terminalibus. †

*Japotapita pyramidata* flore luteo, fructu rubro. *Plum. amer.* 42. ic. 153.

*Japotapita. Marcgr. bras.* 101. *Raj. hist.* 1632.

*Habitat in America meridionali. ‡*

## CALOPHYLLUM.

*Inophyllum.*

1. *CALOPHYLLUM* foliis ovalibus. *Fl. zeyl.* 101. ‡

*Arbor indica*, mali medicæ amplioribus foliis. *Pluk.*

*alm.* 41. t. 147. f. 3.

*Bintangor maritima. Kumph. amb.* 2. p. 211. t. 71.

*Ponna f. Ponna-maram. Rheed. mal.* 4. p. 79. t. 38.

*Raj. hist.* 1525.

*Habitat in India. ‡*

*Calaba.*

2. *CALOPHYLLUM* foliis ovatis obtusis. *Fl. zeyl.*

202. \* *Jacqu. hist.* t. 165.

*Calophyllum* foliis ovatis obtusis : striis parallelis transversis. *Hort. cliff.* 206. *Roy. lugdb.* 476.

*Inophyllum* flore quadrifido. *Burm. zeyl.* t. 60.

*Tsierou-ponna. Rheed. mal.* 4. p. 81. t. 39.

*Habitat in Indiis. ‡*

*An Polygama?*

## GRIAS.

*cauliflora.*

1. *GRIAS.*

*Calophyllum* foliis tripedalibus obovatis, floribus per caulem & ramos sparsis. *Brown. jam.* 245.

*Palmis affinis Malus Persica maxima*, caudice non ramoso, foliis longissimis, flore tetrapetalo pallide luteo, fructu ex arboris trunco. *Sloan. jam.* 179. *hist.*

2. p. 122. t. 217. f. 1, 2.

*Habitat in Jamaica. ‡*

## T I L I A.

1. *TILIA* floribus nectario destitutis.

europea.

*Tilia. Hort. cliff. 204. Fl. fvec. 432, 471. Mat. med. 262. Phil. bot. 259. \* Roy. lugdb. 477. Dalib. parif. 153.*

*Tilia femina, folio majore. Bauh. pin. 426.*

*Tilia urbana. Gesn. tab. III.*

β. *Tilia montana, maximo folio. Bauh. pin. 426.*

γ. *Tilia femina, folio minore. Bauh. pin. 426.*

δ. *Tilia ulmi folio, semine hexagono. Raj. angl. 3. p. 473.*

ε. *Tilia foliis molliter hirsutis, viminibus rubris, fructu tetragono. Raj. angl. 3. p. 473.*

ζ. *Tilia bohemica, foliis minoribus glabris, fructu oblongo utrinque acuminato minimò costulato. Till. pis. 165. t. 49. f. 9.*

*Habitat in Europæ pratis. ‡*

americana.

2. *TILIA* floribus nectario instructis. *Kalm.*

*Tilia foliis majoribus mucronatis. Gron. virg. 58.*

*Tilia, anplissimis glabris foliis, nostrati similis. Pluk. mant. 181.*

*Habitat in Virginia, Canada. ‡*

## L A E T I A.

1. *LAETIA* floribus apetalis. *Jacq. amer. 24. hist. t. apetal. 108. Amæn. acad. 5. p. 413. \**

*Guidonia Laëtia. Læfl. it. 190.*

*Guidonia foliis ovatis utrinque porrectis alternis quandoque crenatis, racemis laxis alaribus. Brown. jam. 249.*

*Habitat in America. ‡*

2. *LAETIA* floribus petaloideis completis. *Jacq. completa. amer. 24.*

*Thamnia foliis ovatis levissime crenatis late virentibus nitidis alternis, petiolis brevibus, pedunculis geniculatis. Brown. jam. 245?*

*Habitat in America. ‡*

## M I C R O C O S.

1. *MICROCOS* panicula terminali. *Fl. zeyl. 207. \** paniculata.

*Microcos foliis alternis oblongis acuminatis. Burm.*

*zeyl. 159. t. 74.*

*Schageri-Cottam. Rheed. mal. 1. p. 105. t. 56.*

*Habitat in India. ‡*

2. MI-



- latiflora* 2. **MICROCOS** pedunculis axillaribus confertis dichotomis. *Fl. zeyl.* 208. \*  
Hamdamanius. *Herm. zeyl.* 3.  
*Habitat in Zeylona.* §

## ELÆOCARPUS.

- ferrata* 1. **ELÆOCARPUS**. *Fl. zeyl.* 206. \*  
*Elajocarpus foliis lauri ferratis, floribus spicatis.*  
*Burm. zeyl.* 39. t. 40.  
*Ganitrus. Rumph. amb.* 3. p. 160. t. 101.  
*Perin-Kara. Rheed. mal.* 4. p. 51. t. 24. *Raj. hist.* 1546.  
*dendr.* 42.  
*Habitat in India.* §

## LECYTHIS.

- ollaria* 1. **LECYTHIS**. *Læfl. ut.* 189. \* *Jacq. amer.* 5.  
*Habitat in America meridionali.* §  
*Folia alterna sessilia. Flores racemosi.*

## MESUA.

- ferrea* 1. **MESUA** foliis lanceolatis. *Fl. zeyl.* 203. \*  
*Naghas. Herm. zeyl.* 7.  
*Balutta-tsiampacam C. Castanea rosea indica. Rheed.*  
*mal.* 3. p. 63. t. 53. *Raj. hist.* 1680.  
*Habitat in India.* §

## VATERIA.

- indica* 1. **VATERIA**. *Fl. zeyl.* 204.  
*Pænoe. Rheed. mal.* 4. p. 33. t. 15.  
*Habitat in India.* §

## LAGERSTROEMIA.

- indica* 1. **LAGERSTROEMIA**.  
*Tsin Kin. Rumph. amb.* 7. p. 61. t. 28.  
*Habitat in China.* §  
*Arbor. Folia alterna, subsessilia, oblonga, integerrima, glabra; Florolia subrotunda. Thyrsus terminalis Læxus, pedicellis trifidis f. trifloris.*

## THEA.

- bohea* 1. **THEA** floribus hexapetalis. *Hort. cliff.* 204. *Alc.*  
*med.* 264. *Hill. exot.* t. 22.  
*Thee. Kämpf. jap.* 605, t. 606.

Thee.

*Thee frutex.* Bart. afl. 4. p. 1. t. 1. Bont. jav. 87.  
t. 88. Barr. rar. 128. t. 904.

*Thee sinensium.* Breyn. cent. 111. t. 112. ic. 17. t. 3.  
Bocc. mus. 114. t. 94.

*Chaa.* Bauh. pin. 147.

*Evonymo affinis arbor orientalis nucifera, flore roseo.*  
Pluk. alm. 139. t. 88. f. 6.

*Habitat in Japonia, China.* h

2. *THEA floribus enneapetalis.* Hill. exot. t. 22.

viridis.

*Habitat in China.* h

*Folia hujus longiora, illius breviora.*

## CARYOPHYLLUS.

1. *CARYOPHYLLUS.* Hort. cliff. 207. Mat. med. 26. aromaticus.

*Caryophyllus aromaticus fructu oblongo.* Bauh. pin.  
410.

*Caryophyllus aromaticus ind. orient. fructu clavato  
monopyreno.* Pluk. alm. 88. t. 155. f. 1.

*Caryophyllus.* Clus. exot. 16. Rumph. amb. 2. p. 1. t.  
1, 2, 3.

*Habitat in Moluccis, solo aridissimo.* h

## MENTZELIA.

1. *MENTZELIA.* Hort. cliff. 492. Amæn. acad. 5. p. aspera.  
398. \* Læst. it. 223.

*Mentzelia foliis & fructibus asperis.* Plum. gen. 42. ic.  
174. f. 1.

*Mentzelia fetis uncinatis munita, foliis lobatis, fructi-  
bus sessilibus singularibus ad divaricationes ramorum.*

*Brown. jam. 249.*

*Onagra americana, folio betonicæ, fructu hispido.*

*Tournef. inst. 302.*

*Habitat in America.*

## PLINIA.

1. *PLINIA.*

pinnata.

*Plinia fructu croceo odorato.* Plum. gen. 9. ic. 225.

*Habitat in America.* h

*Arbor. Folia pinnata, abrupta; foliolis circiter 12. ova-  
to-lanceolatis, integerrimis. Flores sessiles, sparsi per  
ramos nudos seniores. Fructus edules.*

DE-

## D E L I M A.

farmentosa. 1. DELIMA. *Fl. Zeyl.* 205. \*. *Amzn. acad.* 1. p. 402.  
*Habitat in Zeylona.* §

## C I S T U S.

\* *Exstipulata fruticosa.*

capensis. 1. CISTUS arborescens exstipulatus, foliis ovato-lanceolatis petiolatis trinerviis denticulatis utrinque nudis.

*Habitat ad Cap. b. Spei.* §

*Differt a sequentibus foliorum denticulis.*

pilosus. 2. CISTUS arborescens exstipulatus, foliis ovatis petiolatis hirtis.

Cistus mas, folio rotundo hirsutissimo. *Bauh. pin.* 464.

Cistus mas major, folio rotundiore. *I. B. Duham. arb.* 1. p. 167. t. 64.

Cistus mas matthioli. *Dalech. hist.* 222.

*Habitat in Italia, Hispania.* §

*Folia petiolata, ovata, subundata, magnitudine folii pyri, pilis albis raris dispersa. Flores purpurei.*

populifolius. 3. Cistus arborescens exstipulatus, foliis cordatis lobis acuminatis petiolatis. *Hort. cliff.* 205. *Roy. lugdb.* 474.

Cistus Ledon, foliis populi nigrae, major. *Bauh. pin.* 467.

*Ledum latifolium* 2. majus. *Clus. hist.* 1. p. 78.

β. Cistus Ledon, foliis populi nigrae, minor. *Bauh. pin.* 467.

*Ledum latifolium* 2. minus. *Clus. hist.* 1. p. 78.

*Habitat in Lusitania.* §

*Planta ramis & foliis pilosa, etiam pilosissima calycibus. Flores albi.*

laurifolius. 4. CISTUS arborescens exstipulatus, foliis oblongo-ovatis petiolatis trinerviis supra glabris, petiolis basi connatis.

Cistus Ledon, foliis laurinis. *Bauh. pin.* 467.

Cistus Ledon latifolium creticum. *Bauh. hist.* 2. p.

*Ledon* 1 latifolium. *Clus. hist.* 1. p. 77.

*Habitat in Hispania.* §

Folia ovato-oblonga exterioribus ovato-lanceolatis, trinerviis, nuda, subtus subtomentosa; petiolis basi connatis. Bractæ caducæ. Pedunculi longissimi, subumbellati. Flores albi ungue violaceo.

5. CISTUS arborescens exstipulatus, foliis lanceolatis ladaniferus. supra lævibus: petiolis basi coalitis vaginantibus.

Hort. cliff. 205. Sauv. monsp. 147, 150.

Cistus ladanifera hispanica incana. Bauh. pin. 467.

Cistus Ledon 1. angustifolius. Clus. hist. 1. p. 77.

- β. Cistus Ledon flore macula nigricante notato. I. B.

Comm. hort. 1. p. 39. t. 20.

Habitat in Hispaniæ, Lusitaniæ collibus. §

Flores umbellati. Folia lineari-lanceolata. Corollæ albæ ungue violaceo.

6. CISTUS arborescens exstipulatus, foliis lineari-lanceolatis sessilibus utrinque villosis trinerviis. Hort. cliff. 205. Hort. upf. 144. Roy. lugdb. 475. Sauv. monsp. 147.

Cistus ladanifera monspeliensium. Bauh. pin. 467.

Ledum. Dalech. hist. 230.

- β. Cistus Ledon foliis oleæ, sed angustioribus. Bauh. pin. 467.

Ledon 5. Clus. hist. 1. p. 79.

Habitat in G. Narbonensi & Regno Valentino. §

Folia suprema basi latiora, reliqua linearia, subtus reticulata trinervia. Racemus longe pedunculatus, sæpe bifidus. Corollæ albæ.

7. CISTUS arborescens exstipulatus, foliis spatulatis incanatomentosis rugosis: interioribus basi vaginantibus connatis. Hort. cliff. 205. Hort. upf. 143. Roy. lugdb. 475.

Cistus mas angustifolius. Bauh. pin. 464.

Cistus mas 2. Clus. hist. 1. p. 69.

Habitat in Hispania, G. Narbonensi. §

Rami villosi. Folia minime nervosa, in mucronem desinentia, disco parum flexuosa, desinunt basi in Petiolos coalitos vaginantes, vel potius obovato-spatulata, inferioribus magis connatis & fere vaginantibus. Calyces pilosi foliolis subcordatis. Petala rubro-purpurea, emarginata s. obcordata, integerrima, concava.

1. CISTUS arborescens exstipulatus, foliis ovato-lanceolatis. Linnæi Spec. plant. pars I. A 2 a ceo-

ceolatis tomentosis incanis sessilibus subtrinerviis.  
*Sauv. monsp.* 150.

*Cistus mas*, folio oblongo incano. *Bauh. pin.* 464.

*Cistus mas*. 1. *Clus. hist.* 1. p. 68.

*Cistus mas* 4. monspeliensis, folio oblongo albido.  
*Bauh. hist.* 2. p. 3.

*Habitat in G. Narbonensi, Hispania.* §

*Differt a C. incano*, cui *simillimus*, Ramis tomentosis nec pilosis; Foliis pallidioribus, omnino planis, basi sessilibus, minime petiolatis aut vaginantibus, lanceolatis, subtrinerviis. Calyces non hirti foliis exterioribus longioribus, lateribus reflexis. Petala rubro-purpurea, non emarginata, margine subcrenata, planiuscula.

creticus. 9. *CISTUS* arborescens exstipulatus foliis spatulato-ovatis petiolatis enerviis scabris, calycinis lanceolatis.

*Cistus ladanifera cretica.* *Tournef. cor.* 19. *Buxb. cer.* 3. p. 34. t. 64. f. 1.

*Cistus Ledon cretense.* *Bauh. pin.* 467. non synonyma.

*Ladanum creticum.* *Alp. exot.* 89. t. 88.

*Habitat in Creta, Syria.* §

Flores rubri.

salvifolius. 10. *CISTUS* arborescens exstipulatus, foliis ovatis petiolatis utrinque hirsutis. *Hort. cliff.* 205. *Roy. lugd.* 475. *Sauv. monsp.* 150.

*Cistus femina*, folio salvix. *Bauh. pin.* 464.

*Cistus femina.* *Clus. hist.* 1. p. 70.

*Habitat in Italia, Sicilia, Narbona.* §

*Pedunculi uniflori, diphylli. Corollæ albæ aut flavae. Folia ovata, scabra, rugosa: prima utrinque hirsuta: mollia.*

crispus. 11. *CISTUS* arborescens exstipulatus, foliis lanceolatis pubescentibus trinerviis undulatis. *Hort. cliff.* 206. *Roy. lugd.* 475. *Sauv. monsp.* 147.

*Cistus mas*, foliis chamædryos. *Bauh. pin.* 464.

*Cistus mas* 5. *Clus. hist.* 1. p. 69.

*Habitat in Lusitania.* §

*Ramuli, pedunculi & calyces tenui lana adpersi. Flores purpurei. Calyces lanceolati.*

halimifolius 12. *CISTUS* arborescens exstipulatus, foliolis duobus

lycinis linearibus. *Hort. cliff.* 205. *Roy. lugdb.* 475.  
*Sauv. monsp.* 147.

Cistus femina, portulacæ marinæ folio latiore obtuso.  
*Bauh. pin.* 465.

Cistus folio halimi 1. *Clus. hist.* 1. p. 71.

β. Cistus femina, portulacæ marinæ folio angustiore mucronato. *Bauh. pin.* 465.

Cistus halimi folio 2. *Clus. hist.* 1. p. 71.

*Habitat in Lulitaniæ maritimis.* §

Flores flavi.

13. CISTUS arborescens exstipulatus, foliis revolutis linearibus, floribus umbellatis. Libanotis.

Cistus angusto libanotidis folio, flore singulari. *Barb.*  
*ic.* 294.

Cistus Ledon angustis foliis. *Bauh. pin.* 467.

*Habitat in Hispania. Læsing.* §

Frutex habitu Rosmarini. Caulis nudus, purpurascens.

Folia opposita, exstipulata, sessilia, linearia, sed utrinque revoluta, lævia, marginibus basiæ hirsuta, internodiis longiora, ad apices ramorum approximata.

Folia floralia s. suprema breviora, ovata, plana.

Pedunculi terminales, foliis longiores: Floribus aliquot, albis. Calyces ovati, viscidi.

\* \* Exstipulati, suffruticosi.

4. CISTUS suffruticosus procumbens exstipulatus, foliis oppositis linearibus, floribus umbellatis. umbellatus.

Cistus foliis linearibus non stipulatis, caule florali umbellato lignoso. *Guet. stamp.* 2. p. 180. *Dalib. parif.*  
 158.

Cistus Ledon foliis thymi. *Bauh. pin.* 467.

Ledon X. *Clus. hist.* 1. p. 81.

*Habitat in Gallia, Hispania.* §

Umbellæ pedunculatæ sunt.

CISTUS suffruticosus ascendens exstipulatus, foliis alternis fasciculatis linearibus glabris, pedunculis racemosis. lævipes.

*Amæn. acad.* 4. p. 275. \*

Chamæ Cistus, ericæ folio, luteus elatior. *Bauh. pin.*  
 466.

Helianthemum tenuifolium glabrum erectum, luteo flore. *Bauh. hist.* 2. p. 18.

*Habitat Monspeli.* §

A 2 2 2

16. Cl-

Fumana.

16. **CISTUS** suffruticosus procumbens exstipulatus, foliis alternis linearibus margine scabris, pedunculis unifloris.

Cistus caule procumbente, foliis alternis. *Fl. succ.* 435  
474.

**Helianthemum** foliis linearibus sæpius alternis non stipulatis, pedunculis unifloris, caule lignoso. *Gaz. stamp.* 2. p. 18.

**Cistus** fruticosus procumbens, foliis aciformibus oppositis, floribus auriculatis. *Sauv. monsp.* 46.

**Chamæ Cistus**, ericæ folio, luteus humilior. *Bauh.* 466.

- **Helianthemum** tenuifolium glabrum, luteo flore, per humum sparsum. *Bauh. hist.* 2. p. 18.

*Habitat in Gallia, Gotlandia, Helvetia.* §

*Corolla calyce vix major est.*

- italicus. 17. **CISTUS** suffruticosus exstipulatus, foliis oppositis hispidis: inferioribus ovatis; superioribus lanceolatis ramis patentibus.

**Helianthemum**, serpilli folio villoso, flore pallido. *alicum. Barr. rar.* 510. t. 366.

*Habitat in Italia.* §

**Frutex** Caule erecto, spithameo: Ramis oppositis, brevioribus, patentibus, subdeflexis, rufis. Folia opposita, exstipulata, utrinque adpersa pilis strigosis, raris, subgeminis. *Et* subciliata: inferiora petiolata, ovata; superiora subsessilia. Racemus terminatus. Calyces hispidi. Corolla pallida petalis vix exsertatis.

- canus. 18. **CISTUS** suffruticosus exstipulatus procumbens, foliis oppositis obovatis villosis subtus tomentosis, floribus subumbellatis.

**Cistus** foliis ovatis subtus incanis, alis nudis, caule procumbente. *Sauv. monsp.* 148.

**Chamæ Cistus** foliis myrti minoris incanis. *Bauh.* 466.

**Chamæ Cistus** foliis myrti tarentinæ canis & cinereus. *Bauh. hist.* 2. p. 18.

**Chamæ Cistus** 3. *Clus. hist.* 1. p. 74.

- β. **Helianthemum** alpinum serpilli folio nigricante & suto. *Segu. v.r.* 3. p. 195. t. 6. f. 2.

**Helianthemum** serpilli folio villoso, flore pallido. *Barr. rar.* 50. t. 366.

*Habitat in G. Narbonensi, Hispania. §*

*Variat 3. Foliis ellipticis utrinque pilis raris albis adspersis, nec subtus tomentosis. Calyces semper hirsuti*  
*§ umbellæ terminales sessiles.*

19. *CISTUS* suffruticosus exstipulatus, foliis oppositis *marifolius*, oblongis petiolatis planis subtus incanis.

*Helianthemum alpinum, foliis pilosellæ minoris fuchsi. Bauh. hist. 2. p. 18. Segu. ver. 3. p. 196.*

*Helianthemum foliis ad terram congestis: superne pilosis, inferne tomentosis. Hall. helv. 359.*

*Helianthemum luteum, thymi durioris folio. Barr. var. 521. t. 441.*

*Habitat Massiliæ, Veronæ; in Helvetia. §*

*Variat Foliis lanceolatis supra glabris, § ovalibus supra subpilosis, semper vero Folia subtus dense tomentosa § incano-alba.*

20. *CISTUS* suffruticosus procumbens exstipulatus, foliis oppositis oblongis utrinque glabris: petiolis ciliatis, petalis emarginatis. *Oelandicus.*

*Cistus caule procumbente, foliis oblongis utrinque glabris, stipulis nullis. Fl. suec. 434. \* 473. \**

*Habitat in rupibus apricis Oelandiæ. §*

\* \* \* *Exstipulati, herbacei.*

21. *CISTUS* exstipulatus perennis, foliis radicalibus, *Tuberaria*, ovatis trinerviis tomentosis; caulinis glabris lanceolatis; summis alternis.

*Cistus perennis, foliis humi appressis villosis; summis glabris. Sauv. monsp. 149.*

*Cistus folio plantaginis. Bauh. pin. 465.*

*Helianthemum, plantaginis folio, perenne. Buxb. cent. 3. p. 33. t. 63.*

*Tuberaria nostras. Bauh. hist. 2. p. 12.*

*Tuberaria major. Bauh. hist. 2. p. 12.*

*Habitat in Galloprovincia, Hispania, Pisis. 21*

22. *CISTUS* herbaceus exstipulatus, foliis oppositis *guttatus*, lanceolatis trinerviis, racemis ebracteatis. *Huds. angl. 204.*

*Cistus annuus, foliis radicalibus oppositis lanceolatis trinerviis, ramiferis ternatis linearibus. Sauv. monsp. 148.*



Cistus foliis oblongo-acuminatis non stipulatis, caule florali racemoso. *Guett. stamp. 2. p. 181. Dalib. parif. 158.*

Cistus flore pallido, punicante macula insignito. *Bauh. pin. 465.*

Helianthemum flore maculoso. *Col. ecphr. 2. p. 78. t. 77. Habitat in G. Narbonensi, Italia, Anglia. ☉*

Racemi capillares omnino, absque ullis bracteis. Calycis foliola duo exteriora majora ciliata.

canadensis. 23. CISTUS herbaceus exstipulatus, foliis omnibus alternis lanceolatis caule adscendente.

*Habitat in Canada. Kalm. 24*

Facies C. Helianthemis, sed Folia alterna.

\* \* \* \* Stipulati, herbacei.

ledifolius. 24. CISTUS herbaceus erectus glaber stipulatus, floribus solitariis subsessilibus, folio ternato oppositis. *Hort. upf. 144. Sauv. monsp. 178.*

Cistus stipulis quaternis, foliis oblongis, caule erecto. *Hort. cliff. 475.*

Cistus ledi folio. *Bauh. pin. 465.*

Cistus annuus, foliis ledi. *Lob. ic. 2. p. 118.*

*Habitat Monpelii. ☉*

Calyx corolla longior: foliolis 3. erectis, 2. patulis.

salicifolius. 25. CISTUS herbaceus patulus villosus stipulatus, floribus racemosis erectis: pedicellis horizontalibus. *Hort. upf. 144.*

Cistus stipulis quaternis, foliis subovatis subtus tomentosis, caule procumbente. *Hort. cliff. 206.*

Cistus folio salicis. *Bauh. pin. 465.*

Cistus annuus 1. *Clus. hist. 1. p. 76.*

Helianthemum annuum humile, foliis ovatis, flore fugaci. *Segu. ver. 3. p. 197. t. 6. f. 3. opt.*

*Habitat in Lusitania, Hispania. ☉*

egyptiacus. 26. CISTUS herbaceus erectus stipulatus, foliis linearilanceolatis petiolatis, calycibus inflatis corolla majoribus.

*Habitat in Aegypto, B. Jusseus. ☉*

Conf. Cyticistus foliis inferioribus confertis, superioribus oppositis. *Burm. zeyl. 85. t. 36.*

\* \* \* \* \* Sti-

\* \* \* \* \* *Stipulati, suffruticosi.*

27. **CISTUS** suffruticosus stipulatus, foliis obteetis squamatis.  
squamis orbiculatis.

Cistus humilis, floribus compactis in verticillos, minoris  
halimi foliis. *Barr. ic.* 328. *Bocc. mus.* 2. p. 76.  
t. 64.

*Habitat in Hispania. Læst. it.* 67. †

*Suffrutex Ramis erectiusculis, inferne tetragonis. Folia ovali-lanceolata, crassiuscula, petiolata, opposita aut tria utrinque. Stipulae minutissima, racemosa, marcescentes, sessiles. Pedunculi teretes, conferti. Planta tota obteeta squamis orbiculatis, centro depressis.*

28. **CISTUS** suffruticosus procumbens stipulatus, foliis ovato-oblongis subpilosis, petalis lanceolatis. *Hudf. angl.* 205.

*Hellanthemum vulgare, petalis florum perangustis. Dill. elih.* 177. t. 145. f. 174.

*Habitat in Angliæ comitatu Surrejano prope Croydon.* †

29. **CISTUS** suffruticosus stipulatus, foliis inferioribus orbiculatis, superioribus ovatis. *nummularius.*

Cistus humilis f. *Chamaecistus nummulariae* folio.  
*Magn. monsp.* 293.

*Hellanthemum ad nummulariam accedens. Bauh. hist.* 2. p. 20.

*Habitat Monspelii.* †

30. **CISTUS** suffruticosus stipulatus, foliis oblongis, ca- lycopis laevibus. *Sauv. monsp.* 148. *serpyllifolius.*

*Chamaecistus repens serpyllifolia lutea. Bauh. pin.* 466.

*Chamaecistus serpylli folio nigricante & hirsuto, flore aureo odorato. Bauh. hist.* 2. p. 17.

*Chamaecistus* 2. *Clus. hist.* 1. p. 73.

*Habitat in Alpibus Austriæ, Stiria.* †

31. **CISTUS** suffruticulosus stipulatus procumbens, foliis linearibus oppositis brevissimis congestis. *Sauv. monsp.* 148. *thymifolius.*

*Chamaecistus luteus, thymi folio, oliganthes. Barr. ic.* 444.

*Habitat in G. Narbonensi, Hispania.* †

ceolatis tomentosis incanis sessilibus subtrinerviis.  
*Sauv. monsp.* 150.

Cistus mas, folio oblongo incano. *Bauh. pin.* 464.

Cistus mas. 1. *Clus. hist.* 1. p. 68.

Cistus mas 4. monspeliensis, folio oblongo albido.  
*Bauh. hist.* 2. p. 3.

*Habitat in G. Narbonensi, Hispania.* §

*Differt a C. incano, cui simillimus, Ramis tomentosis nec pilosis; Foliis pallidioribus, omnino planis, basi sessilibus, minime petiolatis aut vaginantibus, lanceolatis, subtrinerviis. Calyces non hirti foliis exterioribus longioribus, lateribus reflexis. Petala rubro-purpurea, non emarginata, margine subcrenata, planiuscula.*

creticus. 9. CISTUS arborescens exstipulatus foliis spatulato-ovatis petiolatis enerviis scabris, calycinis lanceolatis.

Cistus ladanifera cretica. *Tournef. cor.* 19. *Buxb. cer.* 3. p. 34. t. 64. f. 1.

Cistus Ledon cretense. *Bauh. pin.* 467. non synonyma.

Ladanum creticum. *Alp. exot.* 89. t. 88.

*Habitat in Creta, Syria.* §

Flores rubri.

salvifolius. 10. CISTUS arborescens exstipulatus, foliis ovatis petiolatis utrinque hirsutis. *Hort. cliff.* 205. *Roy. lugd.* 475. *Sauv. monsp.* 150.

Cistus femina, folio salvie. *Bauh. pin.* 464.

Cistus femina. *Clus. hist.* 1. p. 70.

*Habitat in Italia, Sicilia, Narbona.* §

Pedunculi uniflori, diphylli. Corollæ albæ aut flavæ. Folia ovata, scabra, rugosa: prima utrinque hirsuta, mollia.

crispus. 11. CISTUS arborescens exstipulatus, foliis lanceolatis pubescentibus trinerviis undulatis. *Hort. cliff.* 206. *Roy. lugd.* 475. *Sauv. monsp.* 147.

Cistus mas, foliis chamædryos. *Bauh. pin.* 464.

Cistus mas 5. *Clus. hist.* 1. p. 69.

*Habitat in Lusitania.* §

Ramuli, pedunculi & calyces tenui lana adpersi. Flores purpurei. Calyces lanceolati.

halimifolius 12. CISTUS arborescens exstipulatus, foliis duobus cal-

lycinis linearibus. *Hort. cliff.* 205. *Roy. lugdb.* 475.

*Sauv. monsp.* 147.

*Cistus femina*, portulacæ marinæ folio latiore obtuso.

*Bauh. pin.* 465.

*Cistus folio halimi* 1. *Clus. hist.* 1. p. 71.

β. *Cistus femina*, portulacæ marinæ folio angustiore mucronato. *Bauh. pin.* 465.

*Cistus halimi folio* 2. *Clus. hist.* 1. p. 71.

*Habitat in Lulitaniæ maritimis.* 5

*Flores flavi.*

13. *CISTUS* arborescens exstipulatus, foliis revolutis libanotis. linearibus, floribus umbellatis.

*Cistus angusto libanotidis folio*, flore singulari. *Barb. ic.* 294.

*Cistus Ledon angustis foliis.* *Bauh. pin.* 467.

*Habitat in Hispania.* *Læfing.* 5

*Frutex habitu Rosmarini. Caulis nudus, purpureascens.*

*Folia opposita, exstipulata, sessilia, linearia, sed utrin-*

*que revoluta, levia, marginibus bascos hirsuta, inter-*

*nodis longiora, ad apices ramorum approximata.*

*Folia floralia s. suprema breviora, ovata, plana.*

*Pedunculi terminales, foliis longiores: Floribus*

*aliquot, albis. Calyces ovati, viscidi.*

\* \* *Exstipulati, suffruticosi.*

4. *CISTUS* suffruticosus procumbens exstipulatus, foliis oppositis linearibus, floribus umbellatis.

*Cistus foliis linearibus non stipulatis, caule florali umbellato lignoso.* *Guett. stamp.* 2. p. 180. *Dalib. parif.* 158.

*Cistus Ledon foliis thymi.* *Bauh. pin.* 467.

*Ledon X.* *Clus. hist.* 1. p. 81.

*Habitat in Gallia, Hispania.* 5

*Umbellæ pedunculatæ sunt.*

5. *CISTUS* suffruticosus ascendens exstipulatus, foliis alternis fasciculatis linearibus glabris, pedunculis racemosis. *Amæn. acad.* 4. p. 275.

*Chamæ Cistus, ericæ folio, luteus elatior.* *Bauh. pin.* 466.

*Helianthemum tenuifolium glabrum erectum, luteo flore.* *Bauh. hist.* 2. p. 18.

*Habitat Monspeli.* 5

A 2 2 2

16. Cl-

Fumana.

16. CISTUS suffruticosus procumbens exstipulatus, foliis alternis linearibus margine scabris, pedunculis unifloris.

Cistus caule procumbente, foliis alternis. *Fl. succ.* 435  
474.

Helianthemum foliis linearibus sæpius alternis non stipulatis, pedunculis unifloris, caule lignoso. *Guss. stamp.* 2. p. 18.

Cistus fruticosus procumbens, foliis aciformibus nudis, floribus auriculatis. *Sauv. monsp.* 46.

Chamæ Cistus, ericæ folio, luteus humilior. *Bauh.* 466.

- Helianthemum tenuifolium glabrum, luteo flore, per humum sparsum. *Bauh. hist.* 2. p. 18.

Habitat in Gallia, Gotlandia, Helvetia. *§*

Corolla calyce vix major est.

- italicus. 17. CISTUS suffruticosus exstipulatus, foliis oppositis hispidis: inferioribus ovatis; superioribus lanceolatis ramis patentibus.

Helianthemum, serpilli folio villoso, flore pallido. *alicum. Barr. rar.* 510. t. 366.

Habitat in Italia. *§*

Frutex Caule erecto, spithameo: Ramis oppositis, majoribus, patentibus, subdeflexis, rufis. Folia sessile, exstipulata, utrinque adpersa pilis singulis, raris, subgeminis & subciliata: inferiora petiolata ovata; superiora subsessilia. Racemus terminalis. Calyces hispidi. Corolla pallida petalis vix ciliatis.

- canus. 18. CISTUS suffruticosus exstipulatus procumbens foliis oppositis obovatis villosis subtus tomentosis, floribus subumbellatis.

Cistus foliis ovatis subtus incanis, alis nudis, caule procumbente. *Sauv. monsp.* 148.

Chamæ Cistus foliis myrti minoris incanis. *Bauh.* 466.

Chamæ Cistus foliis myrti tarentinæ canis & cinereus. *Bauh. hist.* 2. p. 18.

Chamæ Cistus 3. *Clus. hist.* 1. p. 74.

- β. Helianthemum alpinum serpilli folio nigricante & bifido. *Segu. v.r.* 3. p. 195. t. 6. f. 2.

Helianthemum serpilli folio villoso, flore pallido. *Barr. rar.* 50. t. 366.

*Habitat in G. Narbonensi, Hispania, §*

*Variat 3. Foliis ellipticis utrinque pilis raris albis adspersis, nec subtus tomentosis. Calyces semper hirsuti*  
*§ umbellæ terminales sessiles.*

19. CISTUS suffruticosus exstipulatus, foliis oppositis, marifolius, oblongis petiolatis planis subtus incanis.

*Helianthemum alpinum, foliis pilosellæ minoris fuchsi. Bauh. hist. 2. p. 18. Segu. ver. 3. p. 196.*

*Helianthemum foliis ad terram congestis: superne pilosis, inferne tomentosis. Hall. helv. 359.*

*Helianthemum luteum, thymi durioris folio. Barr. var. 521. t. 441.*

*Habitat Massiliæ, Veronæ; in Helvetia. §*

*Variat Foliis lanceolatis supra glabris, § ovalibus supra subpilosis, semper vero Folia subtus dense tomentosa § incano-alba.*

20. CISTUS suffruticosus procumbens exstipulatus, foliis oppositis oblongis utrinque glabris: petiolis ciliatis, petalis emarginatis.

*Cistus caule procumbente, foliis oblongis utrinque glabris, stipulis nullis. Fl. suec. 434. \* 473. \**

*Habitat in rupibus apricis Oelandiæ. §*

\* \* \* *Exstipulati, herbacei.*

21. CISTUS exstipulatus perennis, foliis radicalibus, Tuberaria, ovatis trinerviis tomentosis; caulinis glabris lanceolatis; summis alternis.

*Cistus perennis, foliis humi appressis villosis; summis glabris. Sauv. monsp. 149.*

*Cistus folio plantaginis. Bauh. pin. 465.*

*Helianthemum, plantaginis folio, perenne. Buxb. cent. 3. p. 33. t. 63.*

*Tuberaria nostras. Bauh. hist. 2. p. 12.*

*Tuberaria major. Bauh. hist. 2. p. 12.*

*Habitat in Galloprovincia, Hispania, Pisis. 21*

22. CISTUS herbaceus exstipulatus, foliis oppositis guttatus, lanceolatis trinerviis, racemis ebracteatis. Hudf. angl. 204.

*Cistus annuus, foliis radicalibus oppositis lanceolatis trinerviis, ramiferis ternatis linearibus. Sauv. monsp. 148.*

Cistus foliis oblongo-acuminatis non stipulatis, caule florali racemoso. *Guett. stamp. 2. p. 181. Dalib. parif. 158.*

Cistus flore pallido, punicante macula insignito. *Bauh. pin. 465.*

Helianthemum flore maculoso. *Col. ecphr. 2. p. 78. t. 77. Habitat in G. Narbonensi, Italia, Anglia. ☉*

Racemi capillares omnino, absque ullis bracteis. Calycis foliola duo exteriora majora ciliata.

canadensis. 23. CISTUS herbaceus exstipulatus, foliis omnibus alternis lanceolatis caule adscendente.

*Habitat in Canada. Kalm. 24.*

Facies C. Helianthemis, sed Folia alterna.

\* \* \* \* Stipulati, herbacei.

ledifolius. 24. CISTUS herbaceus erectus glaber stipulatus, floribus solitariis subsessilibus, folio ternato oppositis. *Hort. upf. 144. Sauv. monsp. 178.*

Cistus stipulis quaternis, foliis oblongis, caule erecto. *Hort. cliff. 475.*

Cistus ledi folio. *Bauh. pin. 465.*

Cistus annuus, foliis ledi. *Lob. ic. 2. p. 118.*

*Habitat Monspeli. ☉*

Calyx corolla longior: foliolis 3. erectis, 2. patulis.

allicifolius. 25. CISTUS herbaceus patulus villosus stipulatus, floribus racemosis erectis: pedicellis horizontalibus. *Hort. upf. 144.*

Cistus stipulis quaternis, foliis subovatis subtus tomentosis, caule procumbente. *Hort. cliff. 206.*

Cistus folio salicis. *Bauh. pin. 465.*

Cistus annuus 1. *Clus. hist. 1. p. 76.*

Helianthemum annuum humile, foliis ovatis, flore fugaci. *Segu. ver. 3. p. 197. t. 6. f. 3. opt.*

*Habitat in Lusitania, Hispania. ☉*

egyptiacus. 26. CISTUS herbaceus erectus stipulatus, foliis linearilanceolatis petiolatis, calycibus inflatis corolla majoribus.

*Habitat in Aegypto. B. Jusseus. ☉*

Conf. Cyticistus foliis inferioribus confertis, superioribus oppositis. *Burm. zeyl. 85. t. 36.*

\* \* \* \* \* Sti-

\* \* \* \* \* *Stipulati, suffruticosi.*

27. *CISTUS* suffruticosus stipulatus, foliis obtectis squamatus, squamis orbiculatis.

*Cistus humilis*, floribus compactis in verticillos, minoris halimi foliis. *Barr. ic.* 328. *Bocc. mus.* 2. p. 76. t. 64.

*Habitat in Hispania. Læst. it.* 67. ‡

*Suffrutex Ramis erectiusculis, inferne tetragonis. Folia ovali-lanceolata, crassiuscula, petiolata, opposita aut tria utrinque. Stipulae minutissimæ, racemosæ, marcescentes, sessiles. Pedunculi teretes, conferti. Planta tota obtecta squamis orbiculatis, centro depressis.*

28. *CISTUS* suffruticosus procumbens stipulatus, foliis ovato-oblongis subpilosis, petalis lanceolatis. *Huds. angl.* 205. surrejanus.

*Helianthemum vulgare, petalis florum perangustis. Dill. elh.* 177. t. 145. f. 174.

*Habitat in Angliæ comitatu Surrejano prope Croydon. ‡*

29. *CISTUS* suffruticosus stipulatus, foliis inferioribus orbiculatis, superioribus ovatis. nummularius.

*Cistus humilis f. Chamæcistus nummulariæ folio. Magn. monsp.* 293.

*Helianthemum ad nummulariam accedens. Bauh. hist.* 2. p. 20.

*Habitat Monspeli. ‡*

30. *CISTUS* suffruticosus stipulatus, foliis oblongis, calycibus lævibus. *Sauv. monsp.* 148. serpillifolius.

*Chamæcistus repens serpillifolia lutea. Bauh. pin.* 466.

*Chamæcistus serpylli folio nigricante & hirsuto, flore aureo odorato. Bauh. hist.* 2. p. 17.

*Chamæcistus 2. Clus. hist.* 1. p. 73.

*Habitat in Alpibus Austriæ, Stiriæ. ‡*

31. *CISTUS* suffruticulosus stipulatus procumbens, foliis linearibus oppositis brevissimis congestis. *Sauv. monsp.* 148. thymifolius.

*Chamæcistus luteus, thymi folio, oliganthes. Barr. ic.* 444.

*Habitat in G. Narbonensi, Hispania. ‡*



pilosus.

32. **CISTUS** suffruticosus stipulatus, foliis linearibus subtus bifurcatis incanis, calycibus lævibus. *Sauv. monsp.* 147.

*Chamæcistus* foliis thymi incanis. *Bauh. pin.* 466.

*Chamæcistus* 4. *Clus. hist.* 1. p. 74.

- β. *Cistus* stipulis quaternis, foliis lineari-ovalibus incanis, calycibus tomentosis. *Sauv. monsp.* 148.

*Helianthemum* flore albo, folio angusto hirsuto. *Bauh. hist.* 2. p. 17.

- γ. *Cistus* foliis villosis lanceolatis, axillis foliosis, stipulis subulatis. *Sauv. monsp.* 148.

*Helianthemum* f. *Cistus* humilis, folio samplici, capitulis valde hirsutis. *Bauh. hist.* 2. p. 20.

*Habitat* Montpelii. §

*Hæc tres α. β. γ. vel varietates vel valde affines: certiora determinent autoptæ.*

*Helianthemum.*

33. **CISTUS** suffruticosus procumbens, stipulis lanceolatis, foliis oblongis revolutis subpilosis.

*Cistus* stipulis quaternis, foliis oblongis utrinque nudis, caule procumbente. *Hort. cliff.* 206. *Fl. fvec.* 433; 472.

*Chamæcistus* vulgaris, flore luteo. *Bauh. pin.* 465. *Læs. pruss.* 43. t. 8.

Flos folis, *Panax chironium*. *Cam. epit.* 501.

*Habitat* in Europæ pascuis siccis. §

*Petala* suborbiculata, integerrima, flava, annulo receptaculum ambiente sæpius fulvo. Racemus ante florescentiam nutans. Folia pilis raris adpersa.

hirtus.

34. **CISTUS** suffruticosus stipulatus, foliis ovatis, calycibus hispidis.

*Cistus* foliis lanceolato-linearibus integerrimis subtus incanis. *Grou. orient.* 169.

*Cistus* foliis oblongo-linearibus margine revolutis. *L. Monier. obs.* 229. *Sauv. monsp.* 148.

*Cistus* *Ledon rorismarini* foliis subtus incanis. *Bauh. pin.* 467.

*Helianthemum*, angusto serpilli folio, villosum, flore aureo, italicum. *Barr. rar.* 511. t. 488.

*Ledon* 8. *Clus. hist.* 1. p. 8.

*Habitat* in Hispania, Narbona. §

apenninus.

35. **CISTUS** fruticosus stipulatus patulus, foliis lanceolatis hirtis.

He-

*Helianthemum album germanicum. Tabern. icon. 1062.*

*Helianthemum saxatile, foliis & caulibus incanis oblongis, floribus albis. Mentz. pug. 8. f. 3. Dill. elth. 170.*

*Habitat in Apenninis, Italiæ montibus. §*

*Folia supra viridia hirta, subtus incana.*

36. *CISTUS* suffruticosus stipulatus procumbens, foliis pollifolius. oblongo-ovatis incanis, calycibus lævibus, petalis ferratis. *Huds. angl. 205.*

*Cistus humilis alpinus durior, polii nostratis folio candicante. Pluk. alm. 107. t. 23. f. 6.*

*Chamaëcistus montanus, polii folio. Raj. angl. 4. p. 274. t. 274. f. 2.*

*Helianthemum montanum, polii folio incano, flore candido. Dill. elth. 175. t. 145. f. 172.*

*Habitat in æstuario Sabrinæ Anglorum ad Brent-down. §*

37. *CISTUS* suffruticosus stipulatus, foliis alternis lanceolatis planis lævibus. *Amæn. acad. 4. p. 275.\**

*Habitat in Arabia. §*

*Cistorum historia maxime omnium obscura ob varietatum copiam, illustrabitur si determinent autoptæ in singulis.*

1. *Caudicem arborescentem, fruticulosum, annum f. perennem.*
2. *Caulum erectum f. decumbentem.*
3. *Folia opposita f. alterna: cum figura eorundem.*
4. *Stipulas binas f. nullas.*
5. *Pedunculum uniflorum f. multiflorum: nudum f. bracteatum.*
6. *Petalorum figuram.*
7. *Capsulas gloculares f. 3valves.*
8. *Calycem æqualem f. inæqualem.*

## PROCKIA.

1. *PROCKIA. Brown.*

crucis.

*Habitat in Inf. S. Crucis. §*

*Frutex Ramis teretibus. Folia alterna, petiolata, ovata, acuminata, serrata, glabra. Stipulæ lineares. Flores nonnulli, terminales, alterni, pedicellati: Bracteis linearibus.*

## CORCHORUS.

- aliquosus.** 1. **CORCHORUS** capsulis linearibus compressis, foliis lanceolatis æqualiter serratis.  
*Corchorus*, foliolumi, minor. *Plum. spec.* 7. ic. 103. f. 1.  
*Corchorus* Coreta. *Amæn. acad.* 5. p. 398. \*  
*Corchoro* affinis *chamædryos* folio, flore stamineo, seminibus atris quadrangulis duplici serie dispositis.  
*Sloan jam.* 50. *hist.* 1. p. 145. t. 94. f. 1. *Raj. suppl.* 516.  
*Coreta* foliis minoribus ovatis crenatis, floribus singularibus. *Brown. jam.* 147.  
*Habitat in America meridionali.* ‡  
*Caulis fere fruticosus, erectus, paniculatus; subpubescens. Folia lanceolata s. ovato-oblonga, serraturis æqualibus absque dente elongato ad basin. Pedunculi uniflori, lateriflori, absque bracteis. Germina hirta. Siliquæ bivalves, compressæ. Flores Vernales apetalæ, tetrandri. calyce aphylo. Autumnales quinquefidi, completi, polyandri.*
- olitorius.** 2. **CORCHORUS** capsulis oblongis ventricosis, foliorum serraturis infimis setaceis. *Fl. zeyl.* 213. *Hort. upf.* 147. *Gron. orient.* 170.  
*Corchorus* foliorum infimis serraturis maximis reflexis.  
*Hort. cliff.* 209. *Roy. lugdb.* 478.  
*Corchorus* plinii. *Bauh. pin.* 317.  
*Corchorus.* *Comm. hort.* 47. t. 12.  
*Habitat in Asia, Africa, America.* ☉
- astuans.** 3. **CORCHORUS** capsulis oblongis sexsulcatis sexcuspidatis, foliis cordatis infimis serraturis setaceis.  
*Læfl. it.* 224. n. 112. β.  
*Alcea* cibaria s. *Corchorus* americana, carpini foliis, sextuplici capsula longiore. *Pluk. phyt.* 127. f. 3.  
*Triumfetta* subvillosa, foliis rotundioribus undulatis atque dentatis, postremis in setas inermes. *Brown. jam.* 232. t. 25. f. 1.  
*Habitat in America calidiore.* ‡
- capsularis.** 4. **CORCHORUS** capsulis subrotundis depressis rugosis. *Fl. zeyl.* 214. \*  
*Corchorus* foliorum infimis serraturis minoribus. *Hort. cliff.* 210. *Roy. lugdb.* 478.  
*Alcea* olitoria s. *Corchorus* americanus, prælongis

foliis , capsula striata subrotunda brevi. *Pluk. alm.*

18. t. 255. f. 4.

*Habitat in India.* ☉

5. CORCHORUS capsulis oblongis cauleque pilosis, hirtus.  
foliis oblongis serraturis æqualibus.

Corchorus, folio ulmi, major *Plum. spec. 7. ic. 103. f. 2.*

*Habitat in America meridionali.*

6. CORCHORUS capsulis subrotundis lanatis , foliis hirsutis.  
ovatis obtusis tomentosis æqualiter ferratis.

Corchorus americanus lanuginosus, folio chamædryos,  
siliqua tricapsulari lanuginosa & subaspera. *Breyn.*  
*prodr. 3. p. 56.*

Guazuma frutex chamædryfolia , fructu lanuginoso,  
major & minor. *Plum. gen. 36. ic. 104.*

*Habitat in America meridionali.*

### SEGUIERIA.

1. SEGUIERIA. *Læfl. it. 191. \**

*americana.*

*Habitat in Americæ meridionalis umbrosis.* ‡

### SYMPLOCOS.

1. SYMPLOCOS. *Jacqu. amer. 24.*

*martinicensis.*

*Cof r. Læfl. it. 309.*

*Habitat in America.* ‡

## DIGYNIA.

### PÆONIA.

1. PÆONIA foliolis oblongis, *Hort. cliff. 212. Hort. officinalis.*  
*upf. 149. Mat. med. 267. Sauv. monsp. 307.*

- α. Pæonia foliis difformiter lobatis. *Hall. helv. 311.*

*feminea.*

Pæonia communis f. femina. *Bauh. pin. 323.*

Pæonia femina. *Fuchf. hist. 202. Lob. ic. 602.*

- β. Pæonia foliis lobatis ex ovato-lanceolatis. *Hall. helv. 311.*

Pæonia folio nigricante splendido , quæ mas. *Bauh.*  
*pin. 323.*

Pæonia mas. *Lob. ic. 684.*

*Habitat in Nemoribus montium Idæ, Helvetiæ. 24*

*Flores explicati regulares , sed æstivatione omnino irregulares sunt.*

2. PÆ-

tenuifolia.

2. *PÆONIA* foliolis linearibus multipartitis.*Pæonia* laciniis foliorum linearibus. *Zinn. gatt.* 127.*Habitat in Ucraina. Gorter. 2.**Radix repens. Folia Delphinii grandiflori f. Anthoræ, supra decomposita multipartita in lacinias numerosas, lineares, laeves. Flos terminalis P. officinalis, sed minor, pedunculatus, solitarius. Caulis neque ramosus. Verbo quasi Filia Pæoniæ ex Adonide apennina.*

## CURATELLA.

americana.

1. *CURATELLA. Læff. it.* 260. \**Habitat in America meridionali. ‡**Statura Coccolobæ. Folia alterna, subsessilia, oblonga.*

## CALLIGONUM.

Polygonoides.

1. *CALLIGONUM. Hort. cliff.* 212.*Polygonoides orientale, ephedræ facie. Tournef. cor.**47. itin. 3. p. 214. t. 214.**Habitat in monte Ararat. ‡*

## TRIGYNIA.

## DELPHINIUM.

\* *Unicapsularia.*

Consolida.

1. *DELPHINIUM* nectariis monophyllis, caule subdiviso. *Hort cliff.* 212. *Fl. svec.* 440; 476. *Mat. med.**268. Roy. lugdb.* 482. *Dalib. paris.* 158.*Consolida regalis arvensis. Bauh. pin.* 142.*Consolida regalis. Cam. epit.* 521.*Habitat in agris Europæ restibilibus. ☉*

Ajacis.

2. *DELPHINIUM* nectariis monophyllis, caule simplici. *Hort. cliff.* 213. *Hort. upf.* 150. *Roy. lugdb.**482.**Consolida regalis hortensis, flore majore & simplici.**Bauh. pin.* 142.*Consolida regalis hortensis, flore minore. Bauh. pin.**142.**Consolida regalis flore majore & multiplici. Bauh.**pin.* 142.*Flos regius. Dod. pempt.* 252.*Habitat - - - ☉.*\* \* *Tri-*

\* \* *Tricapularia.*

2. **DELPHINIUM** nectariis monophyllis, corollis hexa- ambiguum.  
petalis, capsulis ternis, foliis multipartitis.  
Delphinium elatius, simplici flore. *Clus. hist.* 2. p. 205.  
Consolida regalis, flore minore. *Bauh. pin.* 142.  
*Habitat in Mauritania.* ☉  
Statura *D. Ajacis*, sed magis canescens. Rami divaricati in caule simplici. Corollæ cæruleæ, extus virides. Nectarium monophyllum, bifidum lobis acutiujculis. Calcar sæpe apice bidentatus. Differt a *D. monogynis* numero capsularum; a *D. trigynis* Nectario interiore monophyllo s. calcare interiore non bipartito; adeoque *Staphisagriae* genere separari nequeunt. Pistilla variant in eadem planta 1. ☿ 3. ut vix constans numerus.
4. **DELPHINIUM** nectariis diphyllis, corollis enneaperegrinum.  
petalis, capsulis ternis, foliis multipartitis obtusis.  
*Hort. cliff.* 213.  
Consolida regalis latifolia, parvo flore. *Bauh. pin.* 142.  
*prodr.* 74. t. 74. *Morif. hist.* 3. p. 466. f. 12. t. 4. f. 3.  
*Habitat in Italia, Sicilia, Melita, Palæstina.* ☉  
Caulis simplex: Ramis virgatis, nudis, striculis. Folia *D. Ajacis* facie. Bractææ subulatæ. Corolla cærulea, præter Nectaria ☿ quatuor petala inferiora, habet præterea Petala 2. lateralia subrotunda, unguibus longis affixa, quæ defleſcuntur versus petala exteriora s. infima, ☿ iis incumbunt. Nectaria nulla habent labella deflexa. Capsulæ ternæ.
5. **DELPHINIUM** nectariis diphyllis labellis integris, grandiflorum.  
floribus subsolitariis, foliis compositis lineari-multipartitis. *Hort. upf.* 150. \*  
Delphinium lusitanicum glabrum, aconiti folio. *Rolof. hort.* 61. t. 3.  
Delphinium elatius subincanum perenne, floribus amplis azureis. *Amm. ruth.* 175. *Mill. dict.* 1. 119.  
*Habitat in Sibiria.* 24  
Capsule incanæ sunt.
6. **DELPHINIUM** nectariis diphyllis: labellis bifidis apice barbatis, foliis incisis, caulè recto. *Hort. upf.* 151. \* elatum.

Del-

*Delphinium nectariis diphyllis*, foliis peltatis multipartitis acutis. *Hort. cliff.* 213.

*Delphinium perenne*, aconiti folio ampliori, floribus cæruleis. *Ann. ruth.* 174. *Hall. gatt.* 93.

*Aconitum cæruleum hirsutum*, flore consolidæ regalis. *Bauh. pin.* 181.

*Aconitum lycoctonum*, flore delphinii filesiaci. *Clus. hist.* 2. p. 94.

*Habitat in Sibiria, Helvetia, Silesia.* 24

**Staphisagria 7. DELPHINIUM** nectariis diphyllis petalo brevioribus, foliis palmatis: lobis obtusis. *Hort. cliff.* 213. *Hort. upf.* 150. *Mat. med.* 269. *Roy. lugdb.* 482. *Sauv. monsp.* 214.

*Staphisagria.* *Bauh. pin.* 324. *Dod. pempt.* 366.

*Habitat in Istria, Dalmatia, Calabria, Apulia, Creta, Galloprovincia.*

*Foliorum lobi trifidi. Nectarii cornu brevissimum, obtusum.*

## ACONITUM.

**lycoctonum. 1. ACONITUM** foliis palmatis multifidis villosis.

*Aconitum foliis peltatis multifidis, petalo supremo cylindraceo.* *Fl. lapp.* 211. *Hort. cliff.* 213. *Fl. jvec.* 441; 476. *Roy. lugdb.* 482. *Hall. helv.* 312.

*Aconitum lycoctonum luteum.* *Bauh. pin.* 183.

*Aconitum lycoctonum*, flore luteo. *Besl. cyst. æst.* 25. f. 2.

*Aconitum II.* *Cam. epit.* 827.

*Habitat in Alpibus Lapponiæ, Helvetiæ, Austriæ, Italiæ.* 24

**uncinatum. 2. ACONITUM** foliis multilobis, corollarum galeis apice basi longius producto.

*Habitat in Philadelphia.* 24

*Folia triloba f. quinqueloba, angulato-dentata, nuda.*

*Flores cærulei, singulares, apice galeæ uncinato, recto ♂ longius, quam ipse unguis galeæ, extenso.*

**variegatum. 3. ACONITUM** foliis multifidis: laciniis semipartitis superne latis. *Hort. cliff.* 214. *Hort. upf.* 151. *Roy. lugdb.* 482.

*Aconitum cæruleum minus f. Napellus minor.* *Bauh. pin.* 183.

Aco-

*Aconitum cæruleum parvum.* *Dod. pempt.* 441. *Dalech. hist.* 1743.

*Habitat in Italiæ, Bohemiæ montibus.* 24

4. *ACONITUM* foliorum laciniis linearibus superne Napellus. latioribus linearibus exaratis. *Hort. cliff.* 214. *Fl. suec.* 442; 477. *Gron. virg.* 165. *Hall. helv.* 312. *Afl. stokh.* 1739. p. 41. t. 2.

*Aconitum cæruleum* f. *Napellus* 1. *Bauh. pin.* 183.

*Napellus verus.* *Lob. hist.* 387.

*Habitat in Helvetia, Bavaria, Gallia.* 24

5. *ACONITUM* foliis multipartitis: laciniis linearibus pyrenaicum incumbens: squarrosis. *Hort. upf.* 152. \*

*Aconitum pyrenaicum luteum*, foliorum segmentis fibi invicem incumbentibus. *Raj. europ.* 367.

*Habitat in Sibiria, Tataria, Pyrenæis.* 24

*Racemus ante anthesin nutat.*

6. *ACONITUM* floribus subpentagynis, foliorum laciniis cuneiformibus incisus acutis.

*Aconitum violaceum* f. *Napellus* 2. *Bauh. pin.* 183.

*Aconitum lycoctonum tauricum.* *Clus. hist.* 2. p. 95.

- β. *Aconitum purpureum* f. *Napellus* 3. *Bauh. pin.* 183.

*Aconitum lycoctonum neubergense.* *Clus. hist.* 2. p. 96.

- γ. *Aconitum cæruleo-purpureum*, flore maximo f. *Napellus* 4. *Bauh. pin.* 183.

*Aconitum lycoctonum judenbergense.* *Clus. hist.* 2.

p. 97.

*Habitat in Stiria, Taurero.* 24

7. *ACONITUM* floribus pentagynis, foliorum laciniis linearibus. *Anthora.*

*Aconitum foliorum laciniis linearibus distinctis ubique ejusdem latitudinis.* *Hort. cliff.* 214. *Hort. upf.* 152.

*Mat. med.* 270. *Hall. helv.* 313. *Roy. lugdb.* 483.

*Aconitum salutarium* f. *Anthora.* *Bauh. pin.* 184.

*Anthora* f. *Antithora.* *Cam. epit.* 837.

*Habitat in Alpibus Pyrenæis, Helveticis, Taurinis, Allobrogicis.* 24

## TETRAGYNIA

### TETRACERA.

1. *TETRACERA.* *Hort. cliff.* 214. \*

volubilis.

Pe-



*Petræa* floribus spicatis, scabris lauri foliis. *Amm. herb.* 581.

*Tetracera* foliis amplis serratis obovatis cum acumine, capsulis bigeminis. *Brown. jam.* 255.

*Fagus americanus*, ulmi amplissimis foliis, capsulis bigemellis. *Pluk. amalt.* 87.

*Arbor maxima* forte prunifera, cortice cannabino, folio longissimo latissimoque. *Sloan. jam.* 184. *hist.* 2. p. 130. *Raj. dendr.* 43.

*Arbor americana* convolvulacea platyphyllos barbadensibus dicta. foliis serratis. *Pluk. alm.* 48. t. 146. l. 1?

*Habitat in America meridionali.* 5

## PENTAGYNIA.

### AQUILEGIA.

vulgaris.

1. *AQUILEGIA nectariis incurvis.* *Hort. upf.* 153. *Mat. med.* 272. *Fl. svec.* 2. n. 478.

*Aquilegia.* *Hort. cliff.* 215. *Roy. lugdb.* 482. *Dalib. parif.* 160.

*Aquilegia sylvestris.* *Bauh. pin.* 144.

β. *Aquilegia hortensis simplex.* *Bauh. pin.* 144.

γ. *Aquilegia hortensis multiplex, flore magno.* *C. B.* 144.

δ. *Aquilegia hortensis multiplex, flore inverso.* *C. B.* 144.

ε. *Aquilegia flore roseo multiplici.* *C. B.* 144.

ζ. *Aquilegia degener virescens.* *Bauh. pin.* 144.

*Habitat in Europæ nemoribus saxosis.* 4

alpina.

2. *AQUILEGIA nectariis rectis petalo lanceolato brevioribus.*

*Aquilegia montana, magno flore.* *Bauh. pin.* 144. *prodr.*

75. *Burf. VII.* 108. *Bauh. hist.* 3. p. 484. *Raj. hist.* 707.

*Habitat in Helvetia.* ♂

*Folia duplicato-ternata, parva: foliolis multifidis: lobis sublinearibus, obtusis, facie 3 teneritudine sequentis. Nectarii cornicula erectiuscula, brevia, petalis dimidio breviora. Petala lanceolata, longissima.*

canadensis.

3. *AQUILEGIA nectariis rectis, staminibus corolla longioribus.* *Hort. upf.* 153.

*Aquilegia corolla simplici, nectariis fere rectis.* *Grew. virg.* 59.

*Aquilegia pumila præcox canadensis.* *Corn. canad.* 60. t. 60. *Lob. ic.* 26. *Mill. dict.* t. 47.

Aqui-

# POLYANDRIA PENTAGYNIA. 753

*Aquilegia præcox canadensis*, flore externe rubicundo: medio luteo. *Morif. hist. 3. p. 457. f. 12. t. 2. f. 4.*  
*Habitat in Virginia, Canada.* 24

## NIGELLA.

\* *Pentagynæ.*

1. **NIGELLA** floribus involucro folioso cincti. *Hort. damascena. cliff. 215. Hort. upf. 153. Roy. lugdb. 481. Mill. dict. t. 187. f. 2.*

*Nigella angustifolia*, flore majore simplici cæruleo.  
*Bauh. pin. 145.*

*Melanthium sylvestre. Matth. diosc. 529.*

- β. *Nigella* flore majore pleno cæruleo. *Bauh. pin. 145.*  
*Habitat inter segetes Europæ australis.* ☉

- γ. **NIGELLA** pistillis quinis, capsulis muricatis subrotundis, foliis subpilosis. *Hort. upf. 154. Mat. med. 271. Bæhm. lipf. 173. Mill. dict. t. 187. f. 1.*

*Nigella* flore minore simplici candido. *Bauh. pin. 145.*

*Melanthium sativum. Cam. epit. 551.*

*Nigella* flore minore pleno & albo. *Bauh. pin. 146.*

*Habitat in Germania, Ægypto, Creta.* ☉

3. **NIGELLA** pistillis quinis, petalis integris, capsulis arvensibus turbinatis.

*Nigella* flore foliis nudo, pistillis corollam æquantibus. *Hort. cliff. 215. Roy. lugdb. 481. Hall. helv. 316. Dalib. parif. 160. Bæhm. lipf. 174.*

*Nigella arvensis cornuta. Bauh. pin. 145.*

*Melanthium sylvestre alterum. Cam. epit. 553.*

*Habitat in Germaniæ, Galliæ, Italiæ agris.* ☉

*Hujus Capsula inferne angustior, præcedentis inferne ventricosior, superne utraque contra.*

\* \* *Decagynæ.*

4. **NIGELLA** pistillis denis corollam æquantibus. *Hort. hispanica. upf. 154. Sauv. monip. 253.*

*Nigella latifolia*, flore majore simplici cæruleo. *Bauh. pin. 145. prodr. 75. Morif. hist. 3. p. 516. f. 12. t. 18. f. 8.*

*Habitat in Hispania, Monspelii.* ☉

5. **NIGELLA** pistillis denis corolla longioribus. *Hort. orientalis. cliff. 215. \* Hort. upf. 153. Roy. lugdb. 481.*

*Linnei spec. plant. pars I.*

B b b

NI-

*Nigella chalepensis lutea*, corniculis longioribus. *Morif. hist.* 3. p. 516. f. 12. t. 18. f. 10. *Raj. app.* 525.  
*Habitat circum Alepum.* ☉

## REAUMURIA.

vermiculata

## 1. REAUMURIA.

*Sedum ficulum vermiculatum*, flore saxifragæ albæ, semine villoso. *Bocc. fice.* t. 6. f. 7. *Morif. hist.* 3. p. 481. f. 12. t. 9. f. 6.

*Sedum minus arborescens vermiculatum.* *Lob. ic.* 380.

*Sedum minus fruticosum.* *Bauh. pin.* 284.

*Kali vermiculatum*, albo & amplo sedî rosei flore. *Barr. ic.* 888.

*Kali arabicum primum genus.* *Rauv. it.* 37. t. 37. *Bauh. pin.* 289.

*Habitat in Ægypti, Syriæ, Siciliæ litoribus.* *D. Roque.* ☉

*Statua Salsolæ f. Tamarisci, foliis punctis roridis aspersis. Flores terminales, sessiles inter folia, versus apicem majora. Corollæ albæ: Lamina ciliata, lateribus petalorum utrinque adnexa, constituit nectarium. Flores inferi sunt.*

## HEXAGYNIA.

## STRATIOTES.

Aloldes,

## 1. STRATIOTES foliis ensiformi-triangularibus ciliato-aculeatis.

*Stratiotes.* *Fl. lapp.* 222. *Fl. svec.* 444; 479. *Hort. cliff.* 221. *Roy. lugdb.* 10. *Gort. petr.* 313.

*Stratiotes militaris aizoides.* *Lob. hist.* 904.

*Aloë palustris.* *Bauh. pin.* 286.

*Habitat in Europæ septentrionalis aquis pigris puris.* 24

Alismoides,

2. STRATIOTES foliis cordatis. *Fl. zeyl.* 223.

*Ottel-ambel.* *Rheed. mal.* 11. p. 95. t. 46.

*Habitat in India.* 24

## POLYGYNIA.

## DILLENIA.

*Indica.* 1. *DILLENIA.* *Hort. cliff.* 221.

*Songium.* *Rumph. amb.* 2. p. 14. t. 45.

Sym

*Syalita. Rheed. mal. 3. p. 39. t. 38. 39.*

*Habitat in Malubaria. §*

## LIRIODENDRON,

1. LIRIODENDRON foliis lobatis. *Hort. cliff. 223. Tulipifera.*  
*Hort. upf. 154. Gron. virg. 60. Roy. lugdb. 494. Kalm,*  
*it. 2. p. 322. Trew. chr. t. 10.*

*Tulipifera arbor virginiana. Herm. lugdb. 612. t. 613.*

*Tulipifera virginiana, tripartito aceris folio: media lacinia velut abscissa. Pluk. alm. 379. t. 117. f. 5. §*  
*t. 248. f. 7. Catesb. car. 1. p. 48. t. 48. Raj. hist.*  
*1798.*

2. *Tulipifera caroliniana, foliis productioribus magis angulosis. Pluk alm. 379. t. 68. f. 3.*

*Habitat in America septentrionali. §*

2. LILIODENDRON foliis lanceolatis.

*Liliifera.*

*Sampacca montana. Rumph. amb. 2. p. 204. t. 69.*

*Habitat in Amboina. §*

## MAGNOLIA.

1. MAGNOLIA foliis lanceolatis perennantibus.

*grandiflora.*

*Magnolia foliis lanceolatis persistentibus, caule erecto arboreo. Mill. dict. t. 172.*

*Magnolia altissima, flore ingenti candido. Catesb. car.*  
*2. p. 61. t. 61.*

*Magnolia maximo flore, foliis subtus ferrugineis.*  
*Trew. chr. t. 33.*

*Magnolia altissima, laurocerasi folio, flore ingenti candido. Anoen. angl. t. 14.*

*Habitat in Florida, Carolina. §*

- MAGNOLIA foliis ovato-oblongis subtus glaucis.

*glauc.*

*Magnolia foliis ovato-lanceolatis. Hort. cliff. 222.*  
*Gron. virg. 61. Kalm. it. 2. p. 324.*

*Magnolia laurifolia subtus albicante. Dill. elth. 207. t. 207. t. 168. f. 205. Catesb. car. 1. p. 39. t. 39. Trew.*  
*chr. t. 9.*

*Tulipifera virginiana, laurinis foliis averfa parte rore caeruleo tinctis, conibaccifera. Pluk. alm. 379. t. 68. f. 4.*

*Laurus tulipifera, baccis calyculatis. Raj. hist. 1690.*  
*§ 1798. n. 4.*

*Habitat in Virginia, Pennsylvania. §*

- acuminata. 3. *MAGNOLIA* foliis ovato-oblongis acuminatis.  
*Magnolia* foliis ovato-lanceolatis acuminatis annuis,  
 floribus obtusis. *Mill. dict.*  
*Magnolia* flore albo, folio majore acuminato haud  
 albicante. *Catesb. car. 3. p. 15. t. 15. Gron. virg. 61.*  
*Kalm. it. 2. p. 324.*  
*Habitat in Pennsylvania. ♀*

- tripetala 4. *MAGNOLIA* foliis lanceolatis, petalis exterioribus dependentibus.  
*Magnolia* foliis lanceolatis amplissimis annuis. *Mill. dict.*  
*Magnolia* amplissimo flore albo, fructu coccineo.  
*Catesb. car. 2. p. 80. t. 80.*  
*Habitat in Carolina, rarius in Virginia. ♀*

## MICHELIA.

- Champaca. 1. *MICHELIA*. *Fl. zeyl. 144. Osb. it. 93.*  
*Sampacca. Rumph. amb. 2. p. 199. t. 67, 68.*  
*Champacam. Rheed. mal. 1. p. 31. t. 19. Raj. h. 1641.*  
*Habitat in India. ♀*

## UVARIA.

- zeylanica. 1. *UVARIA* foliis integerrimis.  
*Uvaria. Fl. zeyl. 224. \**  
*Funis mufarius. Rumph. amb. 5. p. 78. t. 42.*  
*Narum-panel. Rheed. mal. 2. p. 11. t. 9. Raj. h. 1636.*  
*Habitat in India. ♀*  
*Conferantur Canangæ. Rumph. amb. 2. t. 65. 66. f.*  
 japonica. 2. *UVARIA* foliis serratis.  
*Frutex viscosus procumbens, folio telephii vulgaris æmulo, fructu racemoso. Kämpf. amæn. 476. t. 47.*  
*Habitat in Japonia. ♀*

## ANNONA.

- muricata. 1. *ANNONA* foliis ovali-lanceolatis glabris nitidis pinnatis, pomis muricatis. *Hort. cliff. 222. \* Gron. virg. 61. Roy. lugdb. 494.*  
*Annona* foliis oblongo-ovatis nitidis, fructibus spinosis mollibus tumentibus obatis. *Brown. jam. 254.*

*Anona indica* latifolia, fructu squamoso aspero. *Pluk. alm.* 31. t. 134. f. 2.

*Anona indica* fructu conoide viridi squamis veluti aculeato. *Pluk. alm.* 32. t. 135. f. 2.

*Anona maxima*, foliis latis splendentibus, fructu maximo viridi conoide: tuberculis f. spinulis innoceptibus aspero. *Sloan. jam.* 203. *hist.* 2. p. 166. t. 225.

*Anona. Comm. hort.* 1. p. 133. t. 69. *Mer. surin.* 14. t. 14.

*Guanabanus* fructu molliter aculeato. *Plum. gen.* 43. t. 143. f. 1.

*Habitat in America calidiore.* ♀

3. *ANNONA* foliis oblongis, fructibus obtuse sub-squamosis squamatis.

*Annona* foliis oblongo-ovatis undulatis venosis, floribus tripetalis, fructibus mammillatis. *Brown. jam.* 256.

*Anona tuberosa.* *Rumph. amb.* 1. p. 138. t. 46.

*Anona* foliis odoratis minoribus, fructu conoide squamoso parvo dulci. *Sloan. jam.* 205. *hist.* 2. p. 168. t. 227. *Raj. dendr.* 77.

*Atamaram.* *Rheed. mal.* 3. p. 21. t. 29.

*Habitat in America meridionali.* ♀

3. *ANNONA* foliis lanceolatis, fructibus ovatis reticulato-areolatis.

*Annona* foliis oblongis undulatis venosis, fructibus areolatis. *Brown. jam.* 256.

*Anona maxima* foliis oblongis angustis, fructu maximo luteo conoide: cortice glabro in areolas angulares distincto. *Sloan. jam.* 264. *hist.* 2. p. 167. t. 226.

*Catesb. car.* 2. p. 62. t. 62. *Raj. dendr.* 77.

*Anona.* *Rumph. amb.* 1. p. 136. t. 45.

*Habitat in America meridionali.* ♀

4. *ANNONA* foliis oblongis obtusiusculis glabris, palustris, fructibus areolatis.

*Annona uliginosa*, foliis nitidis ovatis, fructibus areolatis odoratis. *Brown. jam.* 256.

*Annona* foliis oblongis obtusis glabris, fructu rotundo cortice glabro. *Mill. dict.*

*Anona aquatica*, foliis laurinis atro-virentibus, fructu minore conoide luteo, cortice glabro in areolas distincto. *Sloan. jam.* 205. *hist.* 2. p. 169. t. 228. f. 1. *Raj. dendr.* 78.

*Anona americana* juxta fluviorum ripas innascens pyramidiformi fructu. *Pluk. alm.* 32. t. 240. f. 6.

*Habitat in America ad aquas.* §

glabra. 5. *ANNONA* foliis lanceolato-ovatis, fructibus conoidibus glabris.

*Anona maxima*, foliis latis, fructu maximo luteo conoide: cortice glabro. *Catesb. car.* 2. p. 64. t. 64

*Habitat in Carolina.* §

trilobâ. 6. *ANNONA* foliis lanceolatis, fructibus trifidis. *Mel. diff.* t. 35.

*Anona* fructu lutescente levi serotum arietis referente. *Catesb. car.* 2. p. 85. t. 85. *Trew. chr.* t. 5. *Duk. arb.* t. p. 56. t. 19.

*Habitat in Carolina.* §

asiatica. 7. *ANNONA* foliis lanceolatis glabris nitidis lineatis. *Hort. cliff.* 222. \* *Fl. zeyl.* 225. \* *Roy. lugdb.* 494

*Annona sylvestris.* *Burm. zeyl.* 21.

*Guanabanus* fructu purpureo. *Plum. spec.* 43. ic. 14. f. 2.

*Habitat in Zeylona.* §

africana. 8. *ANNONA* foliis lanceolatis pubescentibus. *Hort. cliff.* 222. *Roy. lugdb.* 494.

*Guanabanus* fructu subcæruleo. *Plum. gen.* 43.

*Guanabanus* persæ folio. *Trew. chr.* t. 49?

*Habitat in America.* §

## ANEMONE.

\* *Hepaticæ flore calycato.*

*Hepatica.* 1. *ANEMONE* foliis trilobis integerrimis.

*Hepatica.* *Hort. cliff.* 223. *Fl. suec.* 445; 480. *H. ups.* 155. *Mat. med.* 277. *Gron. virg.* 62. *Roy. lug.* 487.

*Trifolium hepaticum*, flore simplici & pleno. *Bapin.* 339.

*Habitat in Europæ nemoribus lapidosis.* §

palmata. 2. *ANEMONE* foliis reniformibus lobatis crenatis, involucri multifido, calyce hexaphyllo colorato.

*Pulsatilla* foliis palmatis. *Hort. cliff.* 224. *Roy. lug.* 487.

*Anemone, cyclaminis* f. *malvæ folio, lutea. Bauh. pin. 173. Moris. hist. 2. p. 425. f. 4. t. 25. f. 3.*

*Anemone latifolia flava. Bauh. pin. 176.*

*Anemone hortensis latifolia* 1. *Clus. hist. 1. p. 248.*

*Habitat in Lusitania ad Tagum. 2.*

*Calyx sexpartitus, integerrimus, villosus, coloratus, nec a flore remotus, conjungit Anemones T. & Hepaticas. D.*

\* \* *Pulsatillæ pedunculo involucreto, seminibus lanatis.*

3. ANEMONE pedunculo involucreto, foliis digitatis *patens. multifidis.*

*Pulsatilla folio anemones secundæ f. subrotundo. Bauh. pin. 94. prodr. 94.*

*Pulsatilla polyanthos violacea, anemones folio. Breyn. cent. 132. Helv. puls. 52. t. 2. 3.*

*Pulsatilla, anemones folio dissecto, lanuginosa, flore majore dilute luteo patente. Amm. ruth. 104.*

*Habitat in Tobolsko Sibiriz, Lusatia inferiore. 2.*

*Folia radicalia ternato-digitata: Foliolis sessilibus, laciniatis, acuminatis: Foliolo intermedio tripartito; lateralibus bipartitis. Scapus uniflorus, foliis brevior. Involucrum remotum, villosum. Corolla alba, subtus villosa. Stamina lutea.*

4. ANEMONE pedunculo involucreto, foliis pinnatis, vernalis. flore erecto.

*Pulsatilla foliis simpliciter pinnatis: foliolis lobatis, flore erecto. Fl. svec. 448; 483.*

*Pulsatilla involucreto piloso nitido. Fl. lapp. 223.*

*Pulsatilla, apii folio, vernalis, flore majore. Bauh. pin. 177. prodr. 94. Helv. puls. 63. t. 9.*

*Pulsatilla ex vallesia. Oed. dan. t. 29.*

*Pulsatilla lutea, apii hortensis folio. Bauh. pin. 177.*

β. *Pulsatilla, apii folio, vernalis, flore minore. Bauh. pin. 177. prodr. 94. Helv. puls. 64. t. 10.*

γ. *Pulsatilla lutea, apii hortensis folio. Bauh. pin. 177. Hall. helv. - t. 6.*

*Habitat in Sueciæ, Helvetiæ sylvis sterilissimis. 2.*

5. ANEMONE pedunculo involucreto, petalis rectis, *Pulsatilla, foliis bipinnatis.*

*Pulsatilla foliis decompositis pinnatis, flore nutante: limbo erecto. Hort. cliff. 223. Fl. svec. 446; 481.*



*Mat. med.* 276. *Roy. lugdb.* 487. *Dalib. parif.* 160  
*Sauv. monsp.* 254.

*Pulsatilla* folio crassiore & majore flore. *Bauh. pin.* 177. *Helw. pulf.* 62. t. 8.

*Pulsatilla.* *Cam. epit.* 298.

*Habitat in campis sylvestribus exaridis collibusque apricis Europæ borealis.* 2.

pratensis. 6. ANEMONE pedunculo involocrato, petalis apice reflexis, foliis bipinnatis.

*Pulsatilla* foliis decompositis pinnatis, flore pendulo: limbo reflexo. *Hort. cliff.* 223. \* *Fl. suec.* 447; 482.

*Roy. lugdb.* 487. *Sauv. monsp.* 254.

*Pulsatilla* flore minore nigricante, *Bauh. pin.* 177. *Helw. pulf.* 66. t. 12.

*Pulsatilla vulgaris*, saturatione flore. *Clus. hist.* 1. p. 246.

*Pulsatilla* flore clauso obsoleto; petalis reflexis. *Helw. pulf.* 65. t. 11.

*Habitat in Scaniz, Germaniz campis pratensibus apricis aridis.* 2.

\*\*\* *Anemones caule folioso, seminibus caudatis.*

alpina. 7. ANEMONE foliis caulinis ternis connatis supradecompositis multifidis, seminibus hirsutis caudatis.

• *Anemone* tubis caudatis, foliis omnibus duplicatopinnatis: pinnulis remotis. *Hall. herb.* 331.

*Anemone sylvestris.* 2. *Clus. hist.* 1. p. 245.

*Pulsatilla* flore albo. *Bauh. pin.* 177. *Lob. ic.* 282.

β. *Anemone alpina alba major.* *Bauh. pin.* 176. *prodr.* 94. *Burf. IX.* 80.

γ. *Anemone alpina alba minor.* *Bauh. pin.* 176. *prodr.* 194. *Burf. IX.* 81.

*Habitat in Alpibus Styriacis, Helveticis.* 2.

*Confer: Pulsatilla lutea, pastinacæ sylvestris folio C. B.* 177. *quæ vix videtur ab hac diversa.*

coronaria. 8. ANEMONE foliis radicalibus ternato-decompositis, involucri folioso. *Hasselt. it.* 477. *Hist. anat.* 12.

*Pulsatilla* foliis decompositis ternatis. *Hort. cliff.* 223.

*Hort. ups.* 155. *Roy. lugdb.* 487.

*Anemone tenuifolia*, simplici flore. *Bauh. pin.* 174. 2.

2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12.

β. A-

β. *Anemone tenuifolia multiplex rubra.* Bauh. pin. 176.  
*Anemone.* Cam. epit. 386.

γ. *Anemone angustifolia multiplex*, mutata florum facie quotannis nova. H. R. P. Mill. dist. t. 31.  
*Habitat in Oriente, Constantinopoli allata.* 2.

9. ANEMONE foliis digitatis.

hortensis.

*Pulsatilla foliis digitatis.* Hort. cliff. 224.

*Anemone hortensis latifolia.* Bauh. pin. 176.

*Anemone, geranii rotundo folio, purpurea* cens. Bauh. pin. 173.

*Anemone hortensis latifolia* 3. Clus. hist. 1. p. 249.

*Anemone* 1. Dod. pempt. 434.

*Habitat ad Rhenum* 3 in Italia. 2.

• • • • *Anemoneidæ flore nudo, seminibus ecaudatis.*

10. ANEMONE pedunculo nudo, seminibus subrotundis hirsutis. sylvestris.

*Anemone seminibus hirsutis.* It. oel. 112. Fl. suec. 449; 484.

*Anemone seminibus seta plumosa terminatis.* Hort. cliff. 224. Roy. lugdb. 224.

*Anemone sylvestris alba major.* Bauh. pin. 176.

*Anemone sylvestris.* 1. Clus. hist. 1. p. 244.

β. *Anemone sylvestris alba minor.* Bauh. pin. 176. prodr. 93. Burf. IX. 79.

*Habitat in Germania.* 2.

11. ANEMONE pedunculis alternis longissimis, fructibus cylindricis, seminibus hirsutis mucicis. virginiana.

*Anemone caule ramoso, petalis lanceolatis.* Roy. lugdb. 488. Gron. virg. 165.

*Anemone virginiana tertiæ similis, flore parvo.* Herm. lugdb. 645. parad. 18. t. 18.

*Habitat in Virginia.* 2.

Caulis nudus, terminatus, Foliis ternis, ternatis, medio trilobo, acuto, serrato, lateralibus bipartitis, lobato-serratis. Pedunculus exit e centro petiolorum, longus, nudus. Flos parvus petalis ovatis, viridibus, instar calycis cujusdam. Pistillorum capitulum elongatum, cylindricum, resolvitur in semina hirsuta. Juxta pedunculi basin exit caulis ulterius, terminatus pedunculo florifero 3 foliis duobus oppositis 3 sic ulterius.

- dichotoma. 12. **ANEMONE** caule dichotomo, foliis sessilibus oppositis amplexicaulibus trifidis incisis. *Amæn. acad.* 1. p. 155. \*  
*Habitat in Canada, Sibiria.* 24  
*Statura A. nemorosa, sed altior, radice repens. Folia opposita, sessilia, amplexicaulia, profunde trifida, incisa. Flores A. nemorosa, solitarii e dichotomia caulis, pedunculati.*
- trifolia. 13. **ANEMONE** foliis ternatis ovatis integris serratis, caule unifloro. *Hort. ups.* 155. *Hort. cliff.* 224. *Roy. lugdb.* 488. *Dalib. parif.* 161.  
*Anemone trifolia, flore albo. Bauh. hist.* 3. p. 412.  
*Anemone trifolia. Dod. pempt.* 436.  
*Habitat in Gallia.* 24
- quinquefo-  
lia. 14. **ANEMONE** foliis quinatis ovalibus serratis, caule unifloro.  
*Ranunculus nemorum, fragariae foliis, virginianus. Pluk. alm.* 310. t. 106. f. 3.  
*Habitat in Virginia, Canada. Kalm.* 24
- nemorosa. 15. **ANEMONE** feminibus acutis, foliolis incisis, caule unifloro. *Hort. cliff.* 224. *Fl. svec.* 450; 485. *Mat. med.* 275. *Dalib. parif.* 161.  
*Anemone nemorosa, flore majore. Bauh. pin.* 176.  
*Ranunculus sylvarum. Clus. hist.* 247.  
*Habitat in Europæ asperis, duris, nemoribus.* 24
- apennina. 16. **ANEMONE** feminibus acutis, foliolis incisis, petalis lanceolatis numerosis. *Huds. angl.* 209.  
*Ranunculus nemorosus, flore cæruleo, apennini montis. Mentz. pug.* 1. 8. f. 2.  
*Ranunculus nemorosus, flore purpureo-cæruleo. Park. theatr.* 325. *Raj. angl.* 3. p. 259.  
*Anemone, geranii robertiani folio, cærulea. Bauh. pin.* 174.  
*Anemone hortensis tenuifolia, simplici flore 1. Clus. hist.* 1. p. 254.  
*Habitat in Apenninis, Romæ, Anglia.*
- ranunculol-  
des. 17. **ANEMONE** feminibus acutis, foliolis incisis, petalis subrotundis, caule sub-bifloro. *Hort. cliff.* 125. *Fl. svec.* 451; 486. *Roy. lugdb.* 488. *Dalib. parif.* 161.

*Ranunculus nemorosus luteus. Bauh. pin. 178.*

*Ranunculus phragmites luteus nemorosus. Bauh. hist.*

*3.p. 413.*

*Habitat in Europæ borealis pratis nemorosis. 24*

18. ANEMONE caule unifloro, involucri folio ob-  
tusulo. sibirica.

*Habitat in Sibiria. D. Gmelin. 24*

*Folia radicalia digitata, multifida, marginibus præsertim hirsuta: petiolis pilosis basi vaginantibus. Scapus nudus, pilosus. Involucrum sub flore tripartitum, lobatum: laciniis lanceolatis. Flos patens: petalis 6, subrotundis, fulvis. Stamina flava. Pistillorum capitula subrotunda, muricata.*

19. ANEMONE floribus umbellatis, seminibus depresso-ovalibus nudis. narcissiflora.

*Anemone multiflora, semine lævi latissimo, petalis senis ex rotundo-acutis. Hall. herb. 330.*

*Ranunculus montanus hirsutus humilior, narcissi flore.*

*Bauh. pin. 182.*

*Ranunculus montani 2, species altera. Clus. hist. 1.p. 235.*

*Caryophyllata alpina quinquefolia. Bauh. pin. 322.*

*Habitat in Alpibus Austriæ, Helvetiæ, Sibiricæ. 24*

20. ANEMONE floribus umbellatis congestis, foliis multifidis. fasciculata.

*Ranunculus orientalis, aconiti lycoctoni folio, flore magno albo. Tournef. cor. 20. itin. 3.p. 92. t. 92.*

*Habitat in Oriente.*

*Obscure adhuc distincta a præcedente (cui simillima), quamdiu hujus semina non descripta sunt.*

21. ANEMONE foliis caulinis simplicibus verticillatis, radicalibus duplicato-ternatis. Thalictroides.

*Ranunculus nemorosus, aquilegiæ foliis, virginianus, asphodeli radice. Phl. alm. 310. t. 106. f. 4.*

- β. Thalictrum caule unifloro, ex eodem puncto foliis quatuor simplicibus instructo. *Gron. Virg. 62.*

*Habitat in Virginia, Canada.*

*Canadensis plantæ β, a D. Kalmio lectæ, Radix tuberosa. Folia radicalia petiolis longis, duplicato-ternata: foliolis subcordatis: angulatis. Scapus foliis paulo altior, terminatus foliis quatuor simplicibus, folio-*

*foliolis radicalibus similibus, petiolis longis insidentibus. Pedunculi terminales, conferti s. fasciculati 1, 2, 3, s. 5, simplices, uniflori, aequales. Corollæ albæ: petalis ovatis, obtusis, patentibus. Stamina flavescentia. Pistilli capitulum ovatum, muricatum.*

### ATRAGENE.

- zeylanica.** 1. **ATRAGENE** cirrhis diphyllis, petalis exterioribus quaternis. *Fl. zeyl.* 226.\*  
*Atragene. Amzn. acad.* 1. p. 405.  
*Clematis zeylanica, floribus obsoletis parvis. Herm. zeyl.* 35.  
*Habitat in Zeylona.* §
- alpina.** 2. **ATRAGENE** foliis duplicato-ternatis serratis, petalis exterioribus quaternis.  
*Clematis foliis cordatis crenatis indivisis trilobis ternatisque. Roy. lugdb.* 487.  
*Clematis alpina geraniifolia. Bauh. pin.* 300. *prodr.* 135.  
*Pluk. alm.* 109. t. 84. f. 7. *Morif. bist.* 3. f. 15. t. 2. f. ult.  
*Clematis cruciata alpina. Pon. bald.* 335.  
*Habitat in Baldi, Austriæ, Sibiria alpibus.* §. 2.  
*Folia biternata, incisa, radicalia. Scapi uniflori. Petala 4. exteriora lanceolata, magna; interiora lanceolata, obovata, parva.*
- expensis.** 3. **ATRAGENE** foliis ternatis: foliolis incisus dentatis, petalis exterioribus quinis.  
*Pulsatilla foliis trifidis dentatis, flore incarnato pleno. Burm. afric.* 148.  
*Pulsatilla apii folio rigido, flore magno. Herm. afric.* 18.  
*Habitat ad Cap. b. Spei.* §. 2.  
*Caulis fruticosus. Folia alterna, ternata, cæcisformia, trifida, acuta, dentata, nuda: intermedio longius petiolato. Scapus uniflorus, hirsutus. Involucrum triphyllum, hirsutum. Petala circiter 20, alba: horum 6. inferiora latiora, subtus villosa purpurea. Seminum capitulum ovatum, hirsutum.*

CLEMATIS.

\* *scandentes.*

1. CLEMATIS foliis compositis decompositisque ; fo- Viticella.  
liolis ovatis integerrimis. *Hort. cliff.* 225. *Roy.*  
*lugdb.* 486.  
Clematidis cærulea vel purpurea repens. *Bauh. pin.* 300.  
Clematidis altera. *Clus. hist.* 1. p. 122. *Cam. epit.*  
696.  
Clematis cærulea, flore pleno. *Bauh. pin.* 300.  
*Habitat in Italiæ, Hispaniæ, sepibus.*  
Folia infima cordata ; media triloba, superiora pinna-  
ta. Petala decidua, obtusissima, lateribus tenuiora.
2. CLEMATIS foliis compositis decompositisque : fo- Viorna.  
liolis quibusdam trifidis. *Gron. virg.* 62.  
Flammula scandens, flore violaceo clauso. *Dill. elth.*  
144. t. 118 f. 144.  
Clematis purpurea repens, petalis florum coriaceis.  
*Raj. hist.* 1928.  
Scandens caroliniana planta, viornæ folio. *Pet. succ.* 27.  
*Habitat in Virginia, Carolina.*
3. CLEMATIS foliis simplicibus ternatisque ; foliolis crispa-  
integrilobisve.  
Clematis flore crispo. *Dill. elth.* 86. t. 73. f. 84.  
*Habitat in Carolina.*
4. CLEMATIS foliis compositis : foliolis incis angu- orientalis.  
latis lobatis cuneiformibus, petalis interne villosis.  
Clematis orientalis, apii folio, flore e viridi flavescen-  
te posterius reflexo. *Tournef. cor.* 20.  
Flammula scandens, apii folio glauco. *Dill. elth.* 144.  
t. 119. f. 145.  
*Habitat in Oriente.*  
Flores cernui : petalis recurvatis, supra villosis.
5. CLEMATIS foliis ternatis integerrimis, floribus di- diolcis.  
olcis. *Amzn. acad.* 5. p. 398.  
Clematis scandens, foliis quinquenerviis ovatis nitidis  
pinnato-ternatis. *Brown. jam.* 255.  
Clematis prima s. sylvestris latifolia, foliis ternis  
*Sloan. jam.* 84. *hist.* 1. p. 199. t. 128. f. 1.  
*Habitat in America calidiore.*

6. CLE-

- virginiana.* 6. CLEMATIS foliis ternatis : foliolis cordatis sublobato-angulatis scandentibus, floribus dioicis. *Aman. acad.* 4. p. 274. \*
- Clematis *virginiana* *pannonicæ* similis. *Pluk. mant.* 51. t. 389. f. 4.
- Clematis *floridentis*, flore albo odoratissimo. *Albin. anat.* 1. p. 79. t. 7.
- Clematis *aquatica* trifoliata late scandens, floribus albis odoratis. *Gron. virg.* 62. n. 270.
- Habitat in America septentrionali.* b
- Femina gaudet staminibus, sed antheræ polline destituuntur.*
- cirrhus.* 7. CLEMATIS cirrhis scandens, foliis simplicibus. *Hort. cliff.* 226. *Roy. lugdb.* 487.
- Clematis *peregrina*, foliis pyri incis. *Bauh. pin.* 300.
- Clematis *altera bætica.* *Auf. hist.* 1. p. 123.
- Clematis *peregrina*, foliis pyri incis; nunc singularibus, nunc ternis. *Tournef. cor.* 20.
- Habitat in Bætica, arbores operiens & depressens.*
- Cirrhi oppositi, a foliis distincti.* Folia sæpius simplicia, serrata, petiolata, ovata, ex iisdem geniculis caulibus pluribus. Pedunculi axillares, uniflori; sub singulo flore calyculus remotus, hemisphæricus, minutus.
- Vitalba.* 8. CLEMATIS foliis pinnatis: foliolis cordatis scandentibus. *Hort. cliff.* 225. *Roy. lugdb.* 486. *Hall. helv.* 334. *Gron. virg.* 62. *Sauv. monsp.* 249. *Gort. gelr.* 315.
- Clematis *latifolia integra.* *Bauh. hist.* 2. p. 125.
- Clematis *tertia.* *Cam. epit.* 697.
- β. Clematis *latifolia dentata.* *Bauh. hist.* 2. p. 125.
- Vitalba* *Dod. pempt.* 404.
- γ. Clematis *sylvestris latifolia.* *Bauh. pin.* 300. *Sloan. jam.* 84.
- Habitat in sepibus Europæ australis, Virginie, Jamaicae.*
- Flammula.* 9. CLEMATIS foliis inferioribus pinnatis scandentibus laciniatis: summis simplicibus integerrimis lanceolatis. *Hort. cliff.* 225. *Roy. lugdb.* 486. *Hall. helv.* 334. *Sauv. monsp.* 249.

*Clematitis* f. *Flammula repens*. *Bauh. pin.* 300. *Raj. hist.* 621.

*Flammula*. *Dod. pempt.* 404. *Dalech. hist.* 1171.

*Habitat* in *Montspeli*, *Jenæ*, *inque Rhetiz sepibus*.

• • *erecta*.

10. *CLEMATIS* foliis pinnatis linearibus; caullibus simplicibus hexagonis. *maritima*.

*Clematis maritima repens*. *Bauh. pin.* 300. *prodr.* 135.

*Allion. nic.* 122.\*

*Habitat* ad litora maris *Adriatici*, *Venetis*, *Montspeli*.

*Planta perennis*. *Caules*, • *sarmentis*, *procumbentibus*, *phures*, *erectiusculi*, *erecti*. *Folia opposita*, *pinnata*: *foliolis sæpius 5 paribus*, *remotis*, *linearibus*, *rigidiusculis*, *subsessilibus*, *lævibus*, *indivisis* f. *rarius bi- vel trifidis*. *Flores* *C. rectæ*. *Varietatem C. Flammula* statuunt *Magnol.* & *Rajus*; *meam potius C. rectæ* judicarem *ex solo mutata*?

11. *CLEMATIS* foliis pinnatis: foliolis ovato-lanceolatis integerrimis, caule erecto, floribus pentapetalis tetrapetalisque. *recta*.

*Hort. cliff.* 225. *Hort. ups.* 155. *Roy. lugdb.* 486. *Sauv. monsp.* 249.

*Flammula recta*. *Bauh. pin.* 300.

*Flammula*. *Cam. epit.* 698. *Clus. hist.* 1. p. 124.

*Habitat* in collibus *Austriæ*, *Pannoniæ*, *Tataricæ*, *Montspeli*. 2.

*Umbellæ erectæ* & *strictæ* sunt.

12. *CLEMATIS* foliis simplicibus sessilibus ovato-lanceolatis, floribus cernuis. *integrisfolia*.

*Hort. cliff.* 225. *Hort. ups.* 156. *Roy. lugdb.* 486.

*Clematitis cærulea erecta*. *Bauh. pin.* 300.

*Clematis cærulea pannonica*. *Clus. hist.* 1. p. 123.

*Habitat* in *Hungaria*, *Tataria*. 2.

## THALICTRUM.

1. *THALICTRUM* caule simplicissimo subnudo, racemoso simplici terminali, *alpium*.

*Hort. cliff.* 227. *Fl. svec.* 455:491. *Roy. lugdb.* 485.

*Thalictrum* caule subnudo simplici, foliolis obtusis.

*Fl. lapp.* 225. \*

Tha-



- Thalictrum montanum minimum præcox*, foliis splendentibus, *Morif. hist. 3. p. 325. f. 9. t. 20. f. 14. Oed. dan. t. 11.*  
*Habitat in Alpibus Lapponiæ, Arvoniz. 2.*  
*Petala 4, Stamina 12, Pistilla 8, Racemus simplex floribus cernuis.*
- setidum. 2. **THALICTRUM** caule paniculato filiformi ramossimo folioso.  
*Thalictrum caule filiformi ramossimo in paniculam disperso subjectis foliolis. Hort. cliff. 227. Roy. lugdb. 485. Sauv. monsp. 251.*  
*Thalictrum minimum foetidissimum. Bauh. pin. 337. prodr. 147. Bauh. hist. 3. p. 488. Burf. XIX: 49. Morif. hist. 3. p. 325. f. 9. t. 20. f. 13. Pluk. alm. 367. t. 65. f. 4.*  
*Habitat Monspelii, inque Vallesia, Helvetia. 2.*
- tuberosum. 3. **THALICTRUM** floribus pentapetalis, radice tuberosa. *Hort. cliff. 226. Roy. lugdb. 486.*  
*Thalictrum minus, asphodeli radice, magno flore Tournes. inst. 271.*  
*Thalictrum minus, grumosa radice, floribus majoribus. Herm. flor. 9. Bærh. lugdb. 1. p. 45.*  
*Ranunculus thalictri folio, asphodeli radice. Morif. hist. 2. p. 438. f. 4. t. 28. f. 13.*  
*Oenanthe hederæ foliis. Bauh. pin. 163.*  
*Habitat in Hispania, Pyrenæis. 2.*
- cernui. 4. **THALICTRUM** floribus pentapetalis, radice fibrosa. *Roy. lugdb. 486.*  
*Thalictrum canadense. Corn. canad. 186.*  
*Thalictrum americanum. Park. theatr. 265.*  
*Thalictrum majus, foliis aquilegiæ, flore albo. Morif. hist. 3. p. 325.*  
*Habitat in Canada. 2.*
- dioicum. 5. **THALICTRUM** floribus dioicis.  
*Habitat in Canada. Kalm.*  
*Altitudo vix pedalis. Caulis reclus. Flores tetrapetals corollis 3 staminibus 40. albis. Antheræ fere scentes. Foliola cernua sunt, qua nota distinctissima evadit planta ex habitu. Feminae styli setacei, longiores quam in congeneribus.*

6. **THALICTRUM** foliis sexpartitis, floribus cernuis.

*Thalictrum* caule folioso, foliolis caulinis acutis, panicula divaricata, floribus nutantibus. *Fl. svec.* 454; 489. *It. gotl.* 180. \* *Dalib. parif.* 454.

*Thalictrum minus.* *Bauh. pin.* 335. *Dod. pempt.* 58.

*Ruta pratensis minor.* *Tabern. ic.* 55.

*Habitat in Europæ pratis.* 4.

*Dignoscitur apicibus foliorum purpurascens* ♂ *Caulis nebula cærulescente.* Floret præcocius cum *T. aquilegfolio*, alpino, purpurascens, angustifolio ♂ contorto.

7. **THALICTRUM** foliis tripartitis, caule foliis duplo altiore, floribus cernuis. purpura-  
scens.

*Thalictrum virginianum* elatius glaucum, staminibus purpurascens. *Morif. hist.* 3. p. 324.

*Habitat in Canada.* 4.

*Habitus Thalictri minoris, at dimidio humilius evadit, ubi illud tripedale.* Calyces purpurascens, staminibus luteis. Caulis purpurascens, abque nebula. Differt a *T. minori* foliolis minime apice purpurascens. Tempore florendi præcocius cum *T. minori*, nec cum *T. flavo*. Flos ut in præcedenti: Petalis 4, Staminibus 14, Pistillis 8.

8. **THALICTRUM** foliis tripartitis, foliolis subreflexis sibiricum, argute incis, floribus cernuis.

*Thalictrum alpinum minus saxatile*, rutæ folio, staminibus luteis. *Segu. ver.* 1. p. 476. t. 11?

*Habitat in Sibiria.* *Gmelin.*

*Habitus Th. minoris s. purpurascens, at Foliola parva, sextuplo minora, glauca colore Rutæ s. Fumariæ. Caulis viridis, maxime paniculatus. Panicula ad divaricationes fusca. Floret serius, nempe cum T. flavo* ♂ *Dioico.*

9. **THALICTRUM** foliolis lanceolato-linearibus integerrimis. angustifoli-  
um. *Hort. cliff.* 226. *Amzn. acad.* 3. p. 48. *Roy. lugdb.* 478. *Hull. helv.* 308. *Sauv. monip.* 250.

*Thalictrum pratense*, angustissimo folio. *Bauh. pin.* 337. *prodr.* 146. t. 146. *Fluk. alm.* 364. t. 65. f. 6.

*Habitat in Germania rarius.* 4.

*Petala 4, stamina 16, Pistilla 7.*

- flavum.** 10. **THALICTRUM** caule folioso sulcato, panicul; multiplici erecta. *Hort. cliff.* 226. *Fl. suec.* 453a 488. *Roy. lugdb.* 485. *Hall. helv.* 308. *Dalib. parif.* 162.  
*Thalicttrum pratense. Fl. lapp.* 224.  
*Thalicttrum majus*, siliqua angulosa f. striata. *Berk. pin.* 336.  
*Thalicttrum magnum. Ded. pempt.* 58.
- speciosum.** -β. *Thalicttrum majus flavum*, floribus luteis f. glauco folio. *Bauh. pin.* 336. *flaminibus fulvis.*  
*Habitat in Europæ septentrionalioris subhumidis*, 2. in Hispania. 2.  
*Petala* 4, *Stamina* 24, *Pistilla* 10—16.
- lucidum.** 11. **THALICTRUM** caule folioso sulcato, foliis linearibus carnosiss. *Dalib. parif.* 162. †  
*Thalicttrum minus lucidum*, libanotidis coronariæ foliis. *Fluk. alm.* 363. t. 65. f. 5.  
*Thalicttrum minus alterum parisiensium*, foliis crassioribus & lucidis. *Tournesf. inst.* 271.  
*Thalicttrum* 1, vel *pratense* 1. *Clus. hist.* 1. p. 234  
*Habitat Parisiis* & in Hispania.  
*Planta*, an satis distincta a *T. flavo*? videtur teneris filia.
- aquilegifolium.** 12. **THALICTRUM** fructibus pendulis triangularibus rectis, caule tereti.  
*Thalicttrum* seminibus triangularibus pendulis, stipulis caulinis. *Hort. cliff.* 226. *Fl. suec.* 452; 487. *Roy. lugdb.* 285. *Hall. helv.* 309.  
*Thalicttrum majus*, florum flaminibus purpurascens. *Bauh. pin.* 337.  
*Thalicttrum majus*, folliculis angulosis, caule lxx. *Bauh. hist.* 3. p. 487.  
*Habitat in Scania, Helvetia.* 2.  
*Petala* 4, *Stamina* 60, *Pistilla* 16.
- contortum.** 13. **THALICTRUM** fructibus pendulis triangularibus contortis, caule subancipiti.  
*Thalicttrum hybridum* seminibus contortis. *Amer. acc.* 3. p. 47.  
*Habitat in Sibiria. G. Demidoff.* 24.  
*Petala* 4, *Stamina* 60, *Pistilla* 8. antecedenti fructu sed humilior floribus albis.

14. **THALICTRUM** scapo subumbellato, florum filamentis oblancheolatis coloratis anthera latioribus. petalodeum.

*Habitat in Sibiria.* 24

Folia radicalia duo, supradecomposita ut in congeneribus, sed reliquis longe minora. Foliolis ovatis, obtusis, sæpe trilobis. Scapus vix pedalis, st. iatus, aphyllus, raro unico folio decompositio & pedunculo laterali. Flores fere in umbellam terminalem congesti. Calyx tetraphyllus, deciduus, rotundatus. Stamina numerosa: Filamentis membranaceis, superne magis dilatatis, coloratis, incarnatis, Antheris latioribus. Germina sessilia.

### ADONIS.

1. **ADONIS** floribus pentapetalis, fructibus ovatis. æstivalis.

*Mill. lex.*

Adonis sylvestris, flore phoeniceo, eju. que foliis longioribus. *Bauh. pin.* 178.

Adonis flore pallido. *Cam. epit.* 648.

*Habitat inter segetes Europæ australis.* ☉

*Affinis nimum A. autumnalis.*

2. **ADONIS** floribus octopetalis, fructibus subcylindricis. autumnalis.

*Hort. upi.* 156.

Adonis radice annua. *Hort. cliff.* 321. *Roy. lugdb.* 488.

*Dalh. paris.* 162. *Sauv. monf.* 253.

Adonis flore majore. *Bauh. pin.* 178.

Adonis hortensis, flore minore atro-rubente. *Bauh. pin.* 178.

Flos adonis. *Clus. hist.* 1. p. 336.

*Habitat inter segetes Europæ australis.* ☉

3. **ADONIS** flore dodecapetalo, fructu ovato. vernalis.

Adonis radice perenni. *Hort. cliff.* 231. *Fl. fvec.*

456; 492. *It. xl.* 73. *Roy. lugdb.* 488. *Sauv. monf.* 253.

Helleborus niger tenuifolius, buphthalmi flore. *Bauh. pin.* 186.

Buphthalmum. *Dod. pempt.* 261.

*Habitat in Oelandiæ, Borussia, Bohemiæ collibus apricis.* 24

Hæc Soror *A. apennina*, ut *A. æstivalis* est *A. autumnalis*.

apennina.

4. ADONIS floribus pentdecapetalis, fructu ovato. Helleborus niger ferulaceus, caule geniculato, flore magno tulipæ minoris instar. *Menz. pug. t. 3. f. 1.*

*Habitat in Sibiria, Apenninis. 2.*

*Difficillime distinguuntur plantæ 4 & 3. simillimæ. Caulis 3 adpersus pilis; 4 punctis farinosis pellucidis. Rami 3 ex caulis parte inferiore; 4 ex alis superioribus. Pedunculus 3 brevissimus; 4 longior. Folia in 3 minus pinnata; 4 magis divisa, latiora, nitidiora. Calyx in 3 griseus striatus villosus concavus; 4 luteo-virescens, glaber, avenius, planus. Petala in 3 lanceolata, sæpius duodecim; 4 ovata, imbricata, sæpius quindecim. Flos in 3 unicus; in 4 plures. Stamina in 4 valde reflexa. Planta 4 distinctius æstate durat, 3. vero vernalis est.*

capensis.

5. ADONIS floribus pentdecapetalis, fructibus depressis, foliis duplicato ternatis foliolis incisus dentatis.

*Actæa trifoliata hirsuta, foliis scabris, flore sulphureo rariore. Burm. afr. 147. t. 51.*

*Christophoriana africana ranunculoides, foliis rigidis. Boerh. lugdb. 2. p. 62.*

*Ranunculus æthiopicus, foliis rigidis, floribus ex luteo-virescentibus. Comm. hort. 1. p. 1. t. 1. Rar. suppl. 316.*

*Ranunculus perelegans Franqueville, foliis aculeatis. Lob. illustr. 135.*

*Astrantia africana, laserpitii foliis latioribus rigidis & spinosis. Moris. hist. 3. p. 279.*

*Imperatoria ranunculoides africana enneaphyllos, laserpitii lobatis foliis rigidis margine spinosis. Fluk. atm. 198. t. 95. f. 2.*

*Habitat ad Cap. b. Spei. 24*

*Caulis subdivisus, umbelligerus. Folia radicalia. 4 ternata: intermedio foliolo partiali sæpe pinnato-pentaphyllo. Calyx pentaphyllus. Petala sæpe 15, longa; horum 5 exteriora. Planta caustica variat glabra aut hirta.*

## RANUNCULUS.

*\* Foliis simplicibus.*

- Flammula. 1. RANUNCULUS foliis ovato-lanceolatis pet.

is, caule declinato. *Hort. cliff.* 228. *Fl. suec.* 458; 493. *Roy. lugdb.* 489. *Hall. helv.* 322. *Dalib. parif.* 163.

*Ranunculus foliis ovato-oblongis integerrimis, caule procumbente. Fl. lapp.* 325.

*Ranunculus longifolius palustris minor. Bauh. pin.* 180.

*Flammula Ranunculus. Dod. pempt.* 432.

β. *Ranunculus palustris, foliis ferratis. Bauh. pin.* 180.  
*Habitat in Europæ pascuis udis.* 24

2. *RANUNCULUS foliis linearibus, caule repente. reptans. Fl. lapp.* 236. t. 3. f. 5. *Fl. suec.* 459; 495.

*Ranunculus flammeus minimus, spergulæ folio. Buxb. cent.* 5. app. 41.

*Ranunculus repens, gramineis foliis, e singulis geniculis radices agens. Amm. ruth.* 80. t. 13. f. 1.

*Habitat in Svecia, Russia ad ripas lacuum, antecedenti valde affinis & forte varietas.*

2. *RANUNCULUS foliis lanceolatis, caule erecto. Lingua. Hort. cliff.* 228. *Fl. suec.* 457. *Roy. lugdb.* 489. *Dalib. parif.* 163.

*Ranunculus longifolius palustris major. Bauh. pin.* 34.

*Ranunculus lanceolatus major. Tabern. ic.* 48.

β. *Ranunculus flammeus, latiori plantaginis folio marginibus pilosis. Pluk. alm.* 312.

*Habitat in Europæ borealis fossis, aquis limosis.* 24

4. *RANUNCULUS foliis ovatis petiolatis, floribus nodifloris, sessilibus. Hort. cliff.* 228. *Dalib. parif.* 164.

*Ranunculus parisiensis pumilus, plantaginellæ folio.*

*Pet. gaz.* 40. t. 25. f. 4. *Vahl. atl.* 1719. p. 52. t. 4. f. 4.

β. *Ranunculus ficulus, folio rotundo vix serrato. Pet. gaz.* 39. t. 24. f. 9.

*Habitat Parisiis & in Siciliæ locis paludosis.*

5. *RANUNCULUS foliis lanceolato-linearibus sessilibus, caule erecto, radice bulbosa. Hort. cliff.* 228. *Roy. lugdb.* 489. *Dalib. parif.* 163. *Sauv. monsp.* 75.

*Ranunculus, gramineo folio, bulbosus. Bauh. pin.* 181.

*Ranunculus pumilus*, gramineis foliis. *Bauh. hist.* 3. p. 866.

*Hydropiper lanceolatum*. *Dalech. hist.* 1038.

β. *Ranunculus montanus*, folio gramineo. *Bauh. pin.* 180.

γ. *Ranunculus montanus*, folio gramineo, multiplex.  
*Bauh. pin.* 181.

*Habitat in Pyrenæis & Galliæ prætis aridis.* 2.

parnassifolius.

6. **RANUNCULUS** foliis subovatis nervosis integerrimis petiolatis.

*Ranunculus montanus*, graminis parnassii folio. *Tournef. inst.* 286.

*Habitat in Europa australi.*

*Gemma radicalis promit Folia duo, lanceolata, membranacea, supra terram eminentia. Folia ovata, obtusa, coriacea, nervosa, integerrima, supra pubescentia, imprimis basi involuta, longius petiolata. Caulis simplicissimus, indivisus, aphyllus. Involucrum terminale, diphyllum. Pedunculi 2—4, uniflori, pubescentes. Calyx purpureus. Corolla alba vel purpurea, nervis purpureis.*

amplexicaulis.

7. **RANUNCULUS** foliis ovatis acuminatis amplexicaulis, caule multifloro, radice fasciculata. *Hort. cliff.* 229. *Roy. lugdb.* 484. *Sauv. monsp.* 76.

*Ranunculus montanus*, folio plantaginis. *Bauh. pin.* 180. *Morif. hist.* 2. p. 444. f. 4. t. 30. f. 36.

*Ranunculus dulcis*, foliis latis rapistræ perfoliatis, floribus albis. *Menz. pug. t.* 8. f. 6.

*Habitat in Pyrenæis, Apenninis.* 2.

bullatus.

8. **RANUNCULUS** foliis ovatis serratis, scapo nudo unifloro. *Hort. cliff.* 229.

*Ranunculus latifolius bullatus*, asphodeli radice. *Bauh. pin.* 181.

*Ranunculus lusitanicus*. *Dod. pempt.* 429.

*Ranunculus grumosa* radice 1. *Clus. hist.* 1. p. 238.

β. *Ranunculus latifolius bullatus autumnalis*, flore pleno & prolifero. *Morif. hist.* 2. p. 447. f. 4. t. 31. j. 49, 50, 51.

*Ranunculus lusitanicus*, folio subrotundo, parvo flore. *Tournef. inst.* 286.

*Habitat in Lusitania, Creta.* 2.

vicaria.

9. **RANUNCULUS** foliis cordatis angulatis petiolatis.

*Fl. sicc.* 460; 496. *Hort. cliff.* 228. *Mat. m.* 1.

274. *Roy. lugdb.* 490. *Dalib. paris.* 167.  
*Chelidonia rotundifolia minor. Bauh. pin.* 309.  
*Chelidonium minus. Fuchj. hist.* 866.  
 β. *Chelidonia rotundifolia major. Bauh. pin.* 309. *prodr.* 137.  
*Habitat in Europæ raderatis, umbrosis, spongiosis.* 24
10. **RANUNCULUS** foliis reniformibus subtrilobis *Thora.*  
 crenatis: caulino sessili, floralibus lanceolatis, caule subbifloro. *Jacq. virid.* 249. \*  
*Aconitum pardalianches, f. Thora major. Bauh. pin.* 184.  
*Thora major. Cam. epit.* 825.  
*Pthora valdensium, Lœb. ic.* 604. *Clus. hist.* 1. p. 239.  
 β. *Aconitum pardalianches alterum, f. Thora minor.*  
*Bauh. pin.* 184.  
*Thora minor. Cam. epit.* 826.  
*Lineum pardalianches genus, uno tantum folio. Dalech.*  
*hist.* 1738.  
*Ranunculus foliis radicalibus cordato-subrotundis, caulinis sessilibus lanceolatis integerrimis. Hall. helv.* 327.\*  
*Habitat in Alpibus Helveticis, Pyrenaicis.* 24  
 Varietates α β conjunguntur ab Hallero, distinguuntur a Seguiero.
- \* \* *Foliis dissectis & divisis.*
11. **RANUNCULUS** foliis radicalibus reniformibus *creticus,*  
 crenatis sublobatis, caulinis tripartitis lanceolatis integerrimis, caule multifloro.  
*Ranunculus foliis radicalibus cordato-subrotundis, caulinis tripartitis sessilibus lanceolatis integerrimis. Roy.*  
*lugdb.* 490.  
*Ranunculus, asphodeli radice, creticus. Bauh. pin.* 181.  
*Ranunculus creticus latifolius. Clus. hist.* 1. p. 239.  
*Habitat in Creta.* 24
12. **RANUNCULUS** foliis radicalibus subrotundo *castubicus.*  
 cordatis crepatis; caulinis digitatis dentatis, caule multifloro.  
*Ranunculus rotundifolius vernus sylvaticus major vel castubicus, folio thoræ calthæ. Breyn. prodr.* 1. p. 45.  
*Habitat in Castubia, Sibiria.* 24
13. **RANUNCULUS** foliis radicalibus reniformibus *auricomus.*  
 natis incilis; caulinis digitatis linearibus, caule multifloro. *Hort. cuss.* 229. *Fl. suec.* 462; 498. *Roy. lugdb.*  
*490. Hall. helv.* 327. *Dalib. paris.* 167. *Leys. bal.* 487.



*Ranunculus nemorosus* f. *sylvaticus*, folio subrotundo. *Bauh. pin.* 178

*Ranunculus* 1. *sylvestris*. *Dalech. hist.* 1028.

*Habitat in Europæ pascuis humidiusculis.* 2.

Flores nobis mense frondescentiæ apetalæ, Florescentiæ di-f. tripetalæ, Grossificationis pentapetalæ.

aborivus.

14. *RANUNCULUS* foliis radicalibus cordatis crenatis, caulinis ternatis angulatis, caule subtrifloro.

*Ranunculus* foliis radicalibus crenatis, caulinis digitatis petiolatis. *Roy. lugdb.* 490.

*Ranunculus* foliis radicalibus reniformibus crenatis petiolatis; caulinis paucis digitatis sessilibus. *Gron. virg.* 166.

*Ranunculus virginianus*, flore parvo, molliori folio.

*Herm. lugdb.* 514.

*Habitat in Virginia, Canada.*

sceleratus.

15. *RANUNCULUS* foliis inferioribus palmatis; summis digitatis, fructibus oblongis. *Hort. cliff.* 230.

*Fl. fvec.* 463; 499. *Roy. lugdb.* 490. *Hall. helv.* 323.

*Gron. virg.* 63. *Dalib. parif.* 167.

*Ranunculus palustris*, apii folio, lævis. *Bauh. pin.* 180.

*Ranunculus sylvestris* 1. *Dod. pempt.* 426.

*Habitat ad Europæ fossas & paludes.* ☉

aconitifolius,

16. *RANUNCULUS* foliis omnibus quinatis lanceolatis inciso-ferratis. *Hort. cliff.* 229. *Hort. ups.* 156

*Fl. fvec.* 2. n. 497. *Roy. lugdb.* 490. *Hall. helv.* 325.

*Ranunculus montanus*, aconiti folio, albus: flore majore. *Bauh. pin.* 182.

β. *Ranunculus montanus*, aconiti folio, albus flore minore. *Bauh. pin.* 182.

*Ranunculus montanus.* 4. *Clus. hist.* 1. p. 236.

γ. *Ranunculus* folio aconiti, flore albo multiplici. *Bauh. pin.* 179.

*Ranunculus* pleno flore albo. *Clus. hist.* 1. p. 236.

*Habitat in Alpibus Helveticis, Austriacis, Pyrenæis, Norvegicis, Horto Dei.* 2.

illyricus,

17. *RANUNCULUS* foliis ternatis integerrimis lanceolatis. *It. al.* 71. *Fl. fvec.* 471; 500.

*Ranunculus* foliis tripartitis: laciniis linearibus, caule multifloro, radice tuberosa. *Hort. cliff.* 230. *Roy. lugdb.* 491. *Sauv. mansp.* 181.

Ra-

*Ranunculus lanuginosus angustifolius*, grumosa radice, major & minor. *Bauh. pin.* 181.

*Ranunculus grumosa radice* 4. *Clus. hist.* 1. p. 240.

*Habitat in* Oelandia, Hungaria, Narboina, Italia. 2.

18. *RANUNCULUS* foliis ternatis biternatisque: foliolis trifidis incis, caule inferne ramofo. *Mill. dict.* 1. 216. *asiaticus.*

*Ranunculus* foliis tripartitis laciniatis, caule inferne ramofo, radice tuberosa. *Hort. cliff.* 230. *Hort. ups.* 156. *Roy. lugdb.* 490.

*Ranunculus*, grumosa radice, ramosus. *Bauh. pin.* 181.

*Ranunculus asiaticus polyclonos* s. grumosa radice. 2. *Clus. hist.* 1. p. 241.

- β. *Ranunculus grumosa radice*, flore flavo vario. *Bauh. pin.* 181.

γ. *Ranunculus grumosa radice*, flore albo. *Bauh. pin.* 181.

δ. *Ranunculus grumosa radice*, flore albo leniter crenato. *Bauh. pin.* 181.

ε. *Ranunculus grumosa radice*, flore niveo. *Bauh. pin.* 181.

ζ. *Ranunculus grumosa radice*, flore phœniceo minimo simplici. *Bauh. pin.* 181.

η. *Ranunculus asphodeli radice*, flore sangvineo. *Bauh. pin.* 181.

θ. *Ranunculus asphodeli radice*, flore subphœniceo rubente. *Bauh. pin.* 181.

*Ranunculus*, asphodeli radice, proliſer miniatus. *Bauh. pin.* 181.

*Habitat in* Asia & Mauritania. 2.

19. *RANUNCULUS* foliis ſupradecompoſitis, caule ſimpliciſſimo unifollio unifloro, radice tuberoſa. *Hort. cliff.* 230. *Roy. lugdb.* 491. *rutefolius.*

*Ranunculus rutaceo folio*, flore ſuave rubente. *Bauh. pin.* 181. *Moriſ. hiſt.* 2. p. 448. f. 4. t. 31. f. 54.

*Ranunculus præcox* 1, rutæ folio. *Clus. hiſt.* 1. p. 232.

*Habitat in* Alpibus Auſtriæ, Delphinatus. 2.

20. *RANUNCULUS* calycibus hirsutis; caule biſloro, foliis multifidis. *Fl. ſicc.* 464; 501. *glacialis.*

*Ranunculus caule biſloro*, calyce hirsuto. *Fl. lapp.* 233. t. 3. f. 1.

*Ranunculus montanus purpureus*, calyce villoſo.

*Bauh. hiſt.* 3. p. 868. *Scheuch. alp.* 339. t. 20. f. 1.

*Oed. dan.* t. 19.

- Ranunculus alpinus*, coriandri folio, flore albo-purpureascente. *Barr. ic.* 456.  
*Habitat in Alpibus Lapponiæ, Helvetiæ, Delphinatus.* 4
- sivalis.** 21. *RANUNCULUS* calyce hirsuto, caule unifloro, foliis radicalibus palmatis: caulinis multipartitis sessilibus. *Fl. lapp.* 232. t. 3. f. 2. *Fl. suec.* 465; 502. *Hall. helv.* 326.  
*Ranunculus minimus alpinus luteus.* *Bauh. hist.* 3. p. 861. f. interior.
- β. *Ranunculus idem pygmæus.* *Fl. lapp.* 232. t. 3. f. 3.  
*Habitat in Alpibus Lapponiæ, Helvetiæ.* 4  
*Planta tota glaberrima nequit esse R. monspeliacus.*
- alpestris.** 22. *RANUNCULUS* foliis radicalibus subcordatis obtusis tripartitis: lobis trilobatis; caulino lanceolato integerrimo, caule unifloro.  
*Ranunculus caule aphylo unifloro, foliis subrotundis semitrifidis.* *Hall. helv.* 326. *Sauv. monsp.* 205.  
*Ranunculus alpinus humilis rotundifolius, flore majore & minore.* *Bauh. pin.* 181.  
*Ranunculus alpinus humilis albus, folio subrotundo.* *Segu. ver.* 1. p. 489. t. 12. f. 1.  
*Ranunculus montani* 1. species 1. & 2. *Clus. hist.* 1. p. 234.  
*Habitat in Alpibus Austriacis, Helveticis.* 4
- lapponicus.** 23. *RANUNCULUS* foliis tripartitis lobatis obtusis, caule subnudo unifloro. *Fl. suec.* 461; 503. \*  
*Ranunculus caule unifolio unifloro, foliis tripartitis.* *Fl. lapp.* 231. t. 3. f. 4. \*  
*Habitat in Alpibus Lapponicis.*
- monspeliacus.** 24. *RANUNCULUS* foliis tripartitis crenatis, caule simplici villoso subnudo unifloro. *Sauv. monsp.* 181. †  
*Ranunculus saxatilis, magno flore.* *Bauh. pin.* 182. *prodr.* 96.  
*Habitat Monspellii.*
- bulbosus.** 25. *RANUNCULUS* calycibus retroflexis, pedunculis sulcatis, caule erecto, foliis compositis. *Fl. suec.* 469; 504. *Dalib. parif.* 165.  
*Ranunculus radice simplici globosa.* *Hort. cliff.* 230.  
*Roy. lugdb.* 491. *Hall. helv.* 323.  
*Ranunculus bulbosus.* *Lob. ic.* 666. *Fl. lapp.* 229. R2.

*Ranunculus pratensis*, radice verticillimodo rotunda.

*Bauh. pin.* 179

*Habitat in Europæ pratis, pascuis.* 24

26. *RANUNCULUS* calycibus patulis, pedunculis sulcatis, sarmentis repentibus, foliis compositis. *Fl. suec.* 468; 505. *repens.*

*Ranunculus* foliis ternatis, foliolis petiolatis trifidis: medio productiore, caule multifloro. *Hort. cliff.*

230. *Gron. virg.* 166. *Roy. lugdb.* 491.

*Ranunculus pratensis repens hirsutus.* *Bauh. pin.* 179. *Fl. lapp.* 230.

*Ranunculus hortensis* 1. *Dod. pempt.* 425.

*Habitat in Europæ cultis.* 24

*Varietas potius præcedentis authore Hallero, licet Calyx non retroflexus; Radix fibrosa nec bulbosa; Caules numerosi late diffusi repentes, nec erecti.*

27. *RANUNCULUS* calycibus patulis, pedunculis polyantho-fulcatis, caule erecto, foliis multipartitis. *Fl. suec.* 467; 506. *mos.*

*Ranunculus polyanthemos simplex.* *Lob. ic.* 666.

*Ranunculus sylvestris.* *Taberu. ic.* 42. *hist.* 117.

*Habitat in Europæ borealis graminosis.* 24

28. *RANUNCULUS* calycibus patulis, pedunculis acriteretibus, foliis tripartito-multifidis: summis linearibus. *Fl. suec.* 466; 507. *Dalib. parif.* 164

*Ranunculus* foliis peltatis quinquangularibus multipartitis: laciniis linearibus, caule multifloro. *Hort. cliff.* 231. *Roy. lugdb.* 492.

*Ranunculus pratensis erectus acris.* *Bauh. pin.* 178. *Fl. lapp.* 228.

*Ranunculus hortensis.* *Dod. pempt.* 426.

29. *Ranunculus hortensis erectus, flore pleno.* *Bauh. pin.* 179.

*Habitat in Europæ pratis, pascuis.* 24

29. *RANUNCULUS* calycibus patulis, pedunculis teretibus, caule petiolisque hirtis, foliis trifidis lobatis crenatis holosericeis. *lanuginosus.*

*Ranunculus* foliis radicalibus pinnatis. *Sauv. monf.* 227.

*Ranunculus montanus lanuginosus, foliis ranunculi pratensis repentis.* *Bauh. pin.* 182. *prodr.* 96.

Ra-

*Ranunculus magnus* valde hirsutus, flore luteo. *Bauh. hist.* 3. p. 417.

β. *Ranunculus montanus* subhirsutus, geranii folio. *Bauh. pin.* 180.

*Ranunculus montanus* subhirsutus latifolius. *Bauh. prodr.* 96. t. 96.

*Ranunculus nemorosus* hirsutus, foliis caryophyllat. *Læs. pruss.* 220. t. 71.

*Habitat* Monspelii, in Helvetia, Lusatia. 24

Caulis altus, hirtus, erectus. Folia *Acomiti* s. *Ran. acris*, sed magis incisa. Semina acuta, compressa. Pedunculi longi, teretes. Macula pallida ad finis foliorum, quæ tactu crassiora sentiuntur.

chærophyl-  
los

30. *RANUNCULUS* calycibus retroflexis, pedunculis sulcatis, caule erecto unifloro, foliis compositis lineari-multifidis. *Guett. stamp.* 275. *Dalib. parif.* 166.

*Ranunculus* foliis compositis, caule simplici villosa subnudo unifloro, radice tuberosa. *Zinn. gart.* 129.

*Ranunculus grumosa* radice, foliis *Ranunculi bulbosi* *Bauh. pin.* 181. *prodr.* 96.

*Ranunculus chærophyllos*, asphodeli radice. *Bauh. pin.* 181. *Barr. ic.* 581.

*Ranunculus montanus* leptophyllos, asphodeli radice. *Col. ecphr.* 1. p. 312. t. 311.

*Habitat* in Gallia, Italia. 24

arvensis.

31. *RANUNCULUS* feminibus aculeatis, foliis superioribus decompositis linearibus. *Hort. cliff.* 229. *Fl. svec.* 470; 508. *It. al.* 104. \*. *Roy. lugdb.* 491. *Dalib. parif.* 166.

*Ranunculus arvensis* echinatus. *Bauh. pin.* 179.

*Ranunculus sylvestris* 3. *Dod. pempt.* 427.

*Habitat* in Europæ australioris agris. ☉

rugicatus.

32. *RANUNCULUS* feminibus aculeatis, foliis simplicibus lobatis obtusis glabris, caule diffuso. *H. cliff.* 229; *Hort. ups.* 157. *Roy. lugdb.* 491. *Sax. monsp.* 204.

*Ranunculus palustris* echinatus. *Bauh. pin.* 180. *prodr.* 95.

*Ranunculus creticus* echinatus latifolius. *Alp. exot.* 263. t. 262.

*Habitat* in Europæ australis fossis & humentibus. ☉

parviflorus.

33. *RANUNCULUS* feminibus rugicatis, foliis simplicibus

plcibus laciniatis acutis hirsutis, caule diffuso. *Huds. angl.* 212.

*Ranunculus hirsutus* annuus, flore minimo. *Pluk. alm.* 311. t. 55. f. 1. *Raj. angl.* 3. p. 248. t. 12. f. 1. *Morif. hist.* 2. p. 440. f. 4. t. 28. f. 21.

*Habitat in Europa australi.* ☉

*Differt a R. Muricato: flore parvo; Caule magis prostrato; Foliis profunde lacinulatis; Seminibus late-re tantum tuberculis, nec aculeis muricatis.*

24. *RANUNCULUS* seminibus spinoso-subulatis re- orientalis.  
curvis, calycibus reflexis, foliis multifidis.

*Habitat in Oriente.*

25. *RANUNCULUS* caule erecto bifolio, foliis mul- grandifo-  
tifidis; caulinis alternis sessilibus. *Roy. lugdb.* 492. rus.  
*Ranunculus orientalis*, aconiti folio, flore maximo.

*Tournef. cor.* 22.

*Habitat in Oriente.* ☿

26. *RANUNCULUS* foliis filiformi-ramosis, semini- falcatus.  
bus falcatis, scapo nudo unifloro. *Hort. upf.* 157.  
*Sauv. monsp.* 205. *Grøn. orient.* 69. *Jacq. virid.* 250.

*Myosures foliis ramosis.* *Hort. cliff.* 117. *Roy. lugdb.* 492.

*Melampyrum luteum minimum.* *Bauh. pin.* 234.

*Ranunculus ceratocephalus*, seminibus falcatis in spi-  
cam adactis. *Morif. hist.* 2. p. 440. f. 4. t. 28. f. 22.

*Ranunculus alopecuroides*, ajugæ foliis. *Boc. ficc.* 28.

*Habitat inter segetes Europæ australis* ☿ *Orientis.* ☉

27. *RANUNCULUS* foliis subrotundis trilobis inte- hederaceus.  
gerrimis, caule repente. *Hort. cliff.* 231. *Roy. lugdb.*  
492. *Dalib. parif.* 167. *Gort. gelr.* 325.

*Ranunculus aquaticus hederaceus luteus.* *Bauh. pin.*

180. *Morif. hist.* 2. p. 441. f. 4. t. 29. f. 29.

*Habitat in aquis vadosis Angliæ, Belgii.*

28. *RANUNCULUS* foliis submersis capillaceis, emer- aquatilis.  
sis subpeltatis. *Fl. suec.* 2. n. 509.

*Ranunculus foliis inferioribus capillaceis, superioribus*  
*peltatis.* *Fl. lapp.* 234. *Roy. lugdb.* 492. *Hall. helv.* 328.

*Ranunculus aquaticus*, folio rotundo & capillaceo.

*Bauh. pin.* 120.

*Ranunculus aquatilis.* *Dod. pempt.* 387.

29. *Ranunculus foliis submersis capillaceis.* *Fl. suec.* 1. n.  
472. *Dalib. parif.* 168.

- Ranunculus** foliis omnibus capillaceis circumscriptione rotundis. *Hort. cliff.* 231. *Roy. lugdb.* 492. *Hall. helv.* 328.
- Ranunculus** aquaticus albus, circinatis tenuissime divis foliis. *Fluk. alm.* 311. t. 52. f. 2.
- Millefolium** aquaticum cornutum majus. *Bauh. pin.* 141.
7. **Ranunculus** foliis capillaceis circumscriptione v. brevioribus. *Hall. helv.* 328.
- Ranunculus** aquaticus capillaceus. *Bauh. pin.* 141.
- Millefolium** aquaticum, foliis abrotani, ranunculi re & capitulo. *Bauh. pin.* 141.
- Ranunculus** trichophyllos aquaticus medio luteus. *Col. ecphr.* 1. p. 315. t. 316.
8. **Ranunculus** foliis omnibus capillaceis circumscriptione oblongis. *Roy. lugdb.* 492.
- Ranunculus** albus fluitans, peucedani foliis. *Hort. lugdb.* 517.
- Millefolium** aquaticum, foliis fœniculi, ranunculi re & capitulo. *Bauh. pin.* 141.
- Habitat in Europæ aquis undosis, fossis, rivulis.*
- Conf: Ranunculus aquaticus, cotyledonis umbilicæ folio. Col. ecphr.* 1. p. 315. t. 316, quem, cum *T. lantio* facile hujus generis speciem dicere, nisi *T. drocotylis* illum adscriberet. *Till. pif.* - - ? determinat boni itali.

## TROLLIUS.

- europæus. 1. **TROLLIUS** corollis conniventibus, nectariis longitudine staminum.
- Helleborus** flore clauso erecto petiolato, caule simplicissimo. *Fl. suec.* 474; 510.
- Helleborus** foliis angulatis multifidis, flore globoso. *Hort. cliff.* 227. *Roy. lugdb.* 484.
- Helleborus** caule simplici, flore pedunculato. *Fl. lapp.* 214.
- Ranunculus** montanus, aconiti folio, flore globoso. *Bauh. pin.* 182.
- Ranunculus** flore globoso. *Dod. pempt.* 430.
- Habitat in Alpibus & subalpinis Sueciæ, Germaniæ, Angliæ.* 2.
- asiaticus. 2. **TROLLIUS** corollis patentibus, nectariis longitudine petalorum.
- Trollius** humilis, flore patulo. *Buxb. cent.* 1. p. 15. t. 2.
- Helleborus** niger orientalis, ranunculi folio, flore nequaquam globoso. *Tournef. cor.* 20.

*Helleborus aconiti folio, flore globoſo croceo. Amm.*

*ruth. 101.*

*Habitat in Sibiria, cappadocia 24*

## ISOPYRUM.

1. ISOPYRUM ſtipulis ſubulatis, petalis acutis. *Hort. fumarol-*  
*upf. 157. des.*

*Helleborus fumaricæ foliis. Amm. ruth. 74. t. 12.*

*Habitat in nemoribus Sibiricæ. ☉*

*Confer. Aquilegiæ corniculis, Moſchatellinæ foliis,*  
*planta puſilla ex inſula cheuſan. Pluk. amalth. 3. p.*  
*360. f. 3.*

2. ISOPYRUM ſtipulis ovatis, petalis obtuſis. *Scop. thaliſtri-*  
*carn. 555. des.*

*Thaliſtrum batrachiodes, flore albo, italicum. Bocc.*

*muſ. 84. t. 79. f. 1. \**

*Ranunculus nemoroſus, thaliſtri folio. Bauh. pin. 178.*

*Raj. hiſt. 584. Morif. hiſt. 2. p. 437. f. 4. t. 28. f. 12.*

*Ranunculus præcox 11; thaliſtri folio. Cluſ. hiſt. 1.*  
*p. 233.*

*Habitat in Alpibus Italicis, Gratianopoli, inque agro*  
*Viennenſi, Carniolæ, locis umbroſis; vernalis.*

3. ISOPYRUM ſtipulis obſoletis. *aquilegiol-*

*Aquilegia montana, flore parvo, thaliſtri folio. Bauh.*

*pin. 144. prodr. 75. Bauh. hiſt. 3. p. 484. Raj. hiſt.*

*707. Morif. hiſt. 3. p. 458. f. 12. t. 11. f. 5. Hall.*

*heiv. 310.*

*Habitat in Alpibus Helveticis, Tridentinis, Apenninis.*

## HELLEBORUS.

1. HELLEBORUS flore folio inſidente. *Hort. cliſt. hycmalis.*  
*227. Hort. upf. 158. Roy. lugdb. 484.*

*Helleborus ranunculoides præcox tuberoſus, flore lu-*

*teo. Morif. hiſt. 3. p. 459. f. 12. t. 2. f. 4.*

*Aconitum uniſolium bulbolum. Bauh. pin. 182. Hill*  
*anat. t. 11.*

*Habitat in Lombardia, Italia, Apenninis. 24*

2. HELLEBORUS ſcapo ſubuniſoro ſubnudo, foliis pe- *niger.*  
*datis. Hort. upf. 157. Roy. lugdb. 484. Mat. med. 273.*

*Helleborus ſcapo florifero ſubnudo, petiolo commu-*  
*ni bipartito. Hort. cliſt. 227.*

*Helleborus niger, flore roſeo. Bauh. pin. 186. Hill.*

*anat. t. 1.*



*Elleborus niger*, legitimus. *Clus. hist.* 1. p. 275.  
*Habitat in Austris, Hetruriæ, Apennini asperis.* 2  
*Folia consistentia, hyeme peremantia.*

- viridis.** 3. **HELLEBORUS** caule multifloro folioso, foliis digitatis.  
*Helleborus caule æquali folioso, foliis radicalibus caulem tandem superantibus.* *Hort. cliff.* 227. *Hort. upf.* 158. *Roy. lugdb.* 484.  
*Helleborus niger hortensis, flore viridi.* *Bauh. pin.* 185.  
*Elleborum nigrum alterum.* *Cam. epit.* 941.  
*Habitat in montibus Viennensibus, Euganeis.* 2
- foetidus.** 4. **HELLEBORUS** caule multifloro folioso, foliis pedatis,  
*Helleborus caule inferne angustato multifolio multifloro, foliis caule brevioribus.* *Hort. cliff.* 227. *Roy. lugdb.* 484. *Dalib. parif.* 169. *Saur. monsp.* 180.  
*Helleborus niger foetidus.* *Bauh. pin.* 185.  
*Helleboraster maximus.* *Lob. ic.* 679.  
*Helleborus femina.* *Sterb. fung.* 372. t. 36. f. C.  
 β. *Helleborus niger trifolius.* *Morif. hist.* 3. p. 460. f. 12. t. 4. f. 7.  
*Habitat in Germania, Helvetia, Gallia.* ♂. 2  
*Caudis infra folia nudus.*
- trifolius.** 5. **HELLEBORUS** scapo unifloro, foliis ternatis. *Amz. acad.* 2. p. 356. t. 4. f. 18. *Kalm. it.* 3. p. 379.  
*Habitat in Canadæ, Sibiris sylvis nemorosis cum Oxalide, Circea.*

## C A L T H A.

- palustris.** 1. **CALTHA.** *Fl. lapp.* 227. *Fl. svec.* 473; 511. *Hort. cliff.* 228. *Roy. lugdb.* 484. *Clayt. virg.* 166. *Dalib. parif.* 169.  
*Caltha palustris, flore simplici.* *Bauh. pin.* 276.  
*Populago major.* *Tabern. ic.* 750.  
 α. *Populago minor.* *Tabern. ic.* 750.  
 γ. *Caltha palustris, flore pleno.* *Bauh. pin.* 276.  
*Habitat in humidiusculis Europæ.* 2

## H Y D R A S T I S.

- canadensis.** 1. **HYDRASTIS.** *Ellis.*  
*Hydrophyllum verum canadensium.* *Sp. pl.* 1. p. 146.  
*Habitat in Canadæ aquis.* 2  
*Similis Hydrophylo. Folia bina, petiolata, basi emarginata, palmata, serrata: lobis utrinque lobulato-terali.*





RUG - IBM BELGIUM N.V. - N 220 B/Recto

[illegible]

**CENTRALE BIBLIOTHEEK RUG VOORLOPIGE BOEKKAART**

65

